

# The Bcl10/paracaspase signalling complex is functionally conserved since the last common ancestor of planulozoa.

Jens Staal<sup>1,2</sup>, Yasmine Driege<sup>1,2</sup>, Paco Hulpiau<sup>1,2</sup>, Rudi Beyaert<sup>1,2</sup>

<sup>1</sup> Inflammation Research Center, VIB, Ghent, Belgium

<sup>2</sup> Department of Biomedical Molecular Biology, Ghent University, Ghent, Belgium

## Abstract

A single MALT1, cleavage and CLAP : an old story. The type 1 paracaspases are defined by their domain composition with an N-terminal Death-domain, immunoglobulin domains and a caspase-like (paracaspase) domain. Type 1 paracaspases originated in the edicarian geological period before the last common ancestor of bilaterans and cnidarians (planulozoa). Cnidarians have several paralog type 1 paracaspases, type 2 paracaspases, and a homolog of Bcl10 (CLAP). Notably in bilaterans, lineages like nematodes and insects lack Bcl10 whereas other lineages such as vertebrates, annelids, molluscs and acorn worms do contain Bcl10. There seems to be a correlation where invertebrates with Bcl10 have type 1 paracaspases which are more similar to the paracaspases found in vertebrates. A proposed evolutionary scenario includes two ancestral type 1 paracaspase paralogs in the bilateran last common ancestor, where one paralog usually is dependent on Bcl10 for its function. Functional analyses of invertebrate type 1 paracaspases and Bcl10 homologs support this scenario and indicate an ancient origin of the Bcl10/paracaspase signalling complex.

## Introduction

The paracaspase MALT1 (PCASP1) was originally identified in humans as an oncogenic fusion with IAP2 in low-grade antibiotic-resistant MALT lymphomas <sup>1</sup>. Later, it was discovered that MALT1 is a critical component in T and B cell antigen receptor signalling as part of the CARMA1-Bcl10-MALT1 (CBM) complex<sup>2 3 4</sup>. More studies made it clear that MALT1 plays a role in several different CARD\*-Bcl10-MALT1 complexes in many different types of signalling pathways, where currently known CARD\* components are CARD9, CARD11 (CARMA1), CARD14 (CARMA2) and CARD10 (CARMA3). The use of the different CARD proteins in the CBM complexes is most likely mostly dependent on cell-type specific expression <sup>5</sup>. Despite being identified as a “paracaspase” due to sequence similarity with the true caspases<sup>6</sup>, it was not until much later that a proteolytic activity of MALT1 was established <sup>7 8</sup>. In contrast to true caspases, but similar to metacaspases, the paracaspase MALT1 cleaves substrates specifically after an arginine residue <sup>9 10 11</sup>. Lately some protein substrates have been identified which are cleaved after a lysine by the API2-MALT1 oncogenic fusion <sup>12</sup>. The proteolytic activity of MALT1 was found to also be critical for another type of B-cell lymphoma, ABC-DLBCL <sup>13</sup>. MALT1 cleaves itself <sup>14</sup> and its interacting adaptor protein Bcl10 <sup>8</sup>, the anti-inflammatory deubiquitinases A20 <sup>7</sup> and CYLD <sup>15</sup>, the NF-κB member RelB <sup>16</sup>, the ubiquitin ligase HOIL-1 <sup>17 18 19</sup> and the specific RNA degrading enzymes Regnase <sup>20</sup> and Roquin <sup>21</sup>. The anti-inflammatory role of many of the known protease substrates coupled with the critical role for MALT1 in inflammatory signalling has sparked an interest in targeting MALT1 protease activity as a therapeutic strategy treatment of autoimmune diseases. Although MALT1 has been clearly associated to NF-κB activity, its protease activity plays a more subtle role being specifically required for c-Rel activation <sup>22 23 16 14</sup>. There are mounting evidence that MALT1 also regulate other pathways, such as JNK/AP-1 <sup>15</sup>, mTORC1 <sup>24</sup> and possibly WNT <sup>25</sup>. MALT1 belongs to the type 1 paracaspase family, which consist of an N-terminal Death domain, immunoglobulin domains and a paracaspase domain <sup>26</sup>. The type 1 family of paracaspases originated

sometime during the edicarian geological period, preceding the last common ancestor of bilaterans and cnidarians<sup>26 27</sup>. The cnidarians (e.g. jellyfish, anemone, hydra, coral, ...) and bilaterans (e.g. vertebrates, insects, nematodes, molluscs, roundworms, ...) form the planulozoan clade<sup>28</sup>. In our previous survey of paracaspases and MALT1-associated proteins, type 1 paracaspases and Bcl10 could not be found outside planulozoa<sup>26</sup>. Cnidarians typically contain several paralogs of both type 1 and the ancient type 2 paracaspases whereas bilaterans typically contain a single copy of a type 1 paracaspase. A notable exception is the jawed vertebrates, where the type 1 paracaspase got triplicated. Subsequently, two paralogs were lost in the mammalian lineage leaving PCASP1 (MALT1) as the single paracaspase in mammals<sup>26</sup>. Importantly, some organisms such as the nematode *Caenorhabditis elegans* contain a conserved type 1 paracaspase but lack NF- $\kappa$ B<sup>29</sup>, which indicate that other roles or mechanisms might be responsible for the conservation of the general domain organization of the type 1 paracaspases<sup>26</sup>. On the other hand, despite the remarkable conservation of type 1 paracaspases, there are examples of bilaterans that have lost its paracaspase – most notable the group of flies that include *Drosophila melanogaster*<sup>26</sup>. This indicates that alternative mechanisms can take over the unknown role filled by the type 1 paracaspases in most planulozoan organisms. Apart from functional studies of MALT1 in human and mouse models, investigating the evolutionary history of the type 1 paracaspases and its interacting proteins in alternative model systems could provide important clues to yet-unknown roles and functions of MALT1<sup>26</sup>. Finding those alternative functions of MALT1 could be important for future MALT1 inhibitor-based therapies<sup>30</sup>.

## Materials & Methods

### Sequences of type 1 paracaspases and Bcl10

Protein sequences of type 1 paracaspase and Bcl10 homologs were retrieved from NCBI (<https://www.ncbi.nlm.nih.gov>), Ensembl (<http://www.ensembl.org>), JGI (<http://genome.jgi.doe.gov/>) and ICMB (<https://imcb.a-star.edu.sg>) using BLASTP<sup>31</sup>. All sequences used in the analyses can be found in supplemental material.

### Sequence alignment and phylogenetic analysis

Sequence alignment was performed on the full sequence, using the T-coffee (<http://www.tcoffee.org/>)<sup>32</sup> algorithm. Phylogenetic analysis was performed with the MrBayes (<http://mrbayes.sourceforge.net/>)<sup>33</sup> method. Both alignments and phylogenetic analyses were performed using UGENE (<http://ugene.net/>)<sup>34</sup> on Arch (<http://www.archlinux.org>) Linux<sup>35</sup>. The PCASP2 branch was manually rotated in the scalable vector graphics (svg) file without changing any branch lengths using inkscape (<https://inkscape.org>) to avoid overlapping lines in the tree.

### Cloning of Bcl10 and paracaspase homologs

Plasmids of the cloned genes were deposited in the BCCM/LMBP plasmid collection along with detailed descriptions of cloning strategy and plasmid sequence (<http://bccm.belspo.be/about-us/bccm-lmbp>). The starlet sea anemone (*Nematostella vectensis*) type 1 paracaspase paralog “A” (LMBP: 9589) and zebrafish PCASP3 (LMBP: 9573) were cloned previously<sup>26</sup>. The *Nematostella* type 1 paracaspase paralogs “A” (LMBP: 9636) and “B” (LMBP: xxxx) and pacific oyster (*Crassostrea gigas*, LMBP: xxxx) were cloned behind the human ETV6 HLH domain for dimerization-induced activation as described previously<sup>36 14 26</sup>. Human (LMBP: 9637), zebrafish (LMBP: 9665), pacific oyster (LMBP: 9666) and *Nematostella* (LMBP: xxxx) Bcl10 were cloned in the pCAGGS vector with an N-terminal E-tag.

### Cell culture, transfection and expression analysis

MALT1 KO HEK293T cells (clone #39)<sup>26</sup> were grown under standard conditions (DMEM, 10% FCS, 5% CO<sub>2</sub>, 37 °C) and transfected with the calcium phosphate method<sup>37</sup>. For

evaluation of the conservation of cleavage activity, the HLH-fused paracaspase constructs were co-transfected with wild-type CYLD (LMBP: 6613) or uncleavable CYLD-R324A (LMBP: 6645) mutant. For evaluation of conservation of NF- $\kappa$ B induction, the three HLH paracaspase fusions were co-transfected with a NF- $\kappa$ B luciferase reporter (LMBP: 3249) and actin promoter-driven  $\beta$ -galactosidase (LMBP: 4341) as transfection control. For evaluation of the functional conservation of the Bcl10 homologs, the Bcl10 homologs were transfected with the NF- $\kappa$ B luciferase reporter and  $\beta$ -galactosidase in the MALT1 KO HEK293T cells with or without reconstitution with human MALT1 (LMBP: 5536). Detection of cleaved CYLD was done with the E10 antibody (Santa Cruz Biotechnology) recognizing the C-terminal 70kDa cleavage band or anti-E-tag (ab66152, Abcam) recognizing the 40kDa N-terminal cleavage band. Expression of the fused paracaspases was determined with anti-Flag (F-3165, Sigma). Human MALT1 was detected by the EP603Y antibody (Abcam) and the E-tagged Bcl10 clones with anti-E-tag. All western blots were developed on an Odyssey scanner (LI-COR).

## Results & Discussion

### *Correlation vertebrate-like type 1 paracaspases and presence of Bcl10.*

Alignment of several type 1 paracaspases show that type 1 paracaspases from species that contain Bcl10 (molluscs, annelids, hemichordates) cluster closer to the vertebrate paracaspases (Figure 1), indicating a conserved common Bcl10-dependent ancestor. In contrast, Bcl10 sequences appear to evolve in a manner similar to how the species have diverged throughout evolution (Figure 2).

### *Functional conservation of invertebrate type 1 paracaspases and Bcl10*

Based on BLASTP, mollusc paracaspases were identified as the non-deuterostome homologs most closely resembling vertebrate type 1 paracaspases, and the pacific sea oyster (*Crassostrea gigas*) was used as a model for the molluscs. Conversely, the most distantly related species where type 1 paracaspases and Bcl10 could be found are cnidaria<sup>26</sup>. The cnidarian starlet sea anemone (*Nematostella vectensis*) was used as a source for as divergent proteins as possible. In order to investigate the functional conservation of invertebrate type 1 paracaspases, we evaluated artificially activated paracaspases fused to the ETV6 HLH domain<sup>14</sup>. As positive control, the currently most distantly related vertebrate paracaspase with conserved activity (zebrafish PCASP3)<sup>26</sup> was used. In an NF- $\kappa$ B luciferase assay, only the activated zebrafish PCASP3 could induce the reporter, indicating that the pacific oyster (CgPCASP) and the two starlet sea anemone type 1 paracaspase paralogs (NvPCASP-t1A, NvPCASP-t1B) could not recruit critical downstream signalling components such as TRAF6 (Figure 3A). In contrast, evaluation of protease activity revealed that the pacific oyster paracaspase specifically cleaves human CYLD at R324, just like vertebrate paracaspases (Figure 3B). The "A" paralog from starlet sea anemone could not cleave CYLD at all and the "B" paralog appeared to cleave CYLD, but at a different residue, indicating that paracaspase substrate specificity is not conserved in the cnidarians.

To further investigate the functional conservation of the Bcl10/paracaspase co-evolution, we transfected human, zebrafish, pacific oyster and starlet sea anemone Bcl10 in MALT1 KO HEK293T cells with or without reconstitution with human MALT1. Strikingly, the starlet sea anemone Bcl10 could induce human MALT1-mediated NF- $\kappa$ B induction and appears to be modified and possibly cleaved in presence of MALT1, which is similar to human and zebrafish Bcl10 (Figure 3C). In contrast, the pacific oyster Bcl10 failed to induce any NF- $\kappa$ B reporter activity, which might be due to its small size. It will be interesting to see if future annotations of the mollusc genomes will establish a longer Bcl10 transcript encoding for a functional Bcl10 homolog.

### *Future challenges*

Previous studies have shown that the MALT1-like activities are conserved at least as far

back as the last common ancestor of the three vertebrate type 1 paracaspase paralogs<sup>26</sup>. Similarly, also Bcl10 has been shown to be functionally conserved as far back as zebrafish<sup>38</sup>. We have now shown that Bcl10 and MALT1-like activities from type 1 paracaspases are considerably older. The observation that invertebrate type 1 paracaspases from organisms that also contain Bcl10 are more similar to the vertebrate paracaspases provides a new interesting perspective on the functional evolution of MALT1. We still don't know how far back that MALT1-like activities such as TRAF6 interaction and NF- $\kappa$ B induction, protease activity and specificity are conserved. With the observation that mollusc paracaspases have conserved protease activity and specificity, but fail to induce NF- $\kappa$ B in a human cellular background, we are starting to unravel the sequence of evolutionary events leading to the current MALT1 activities in humans. The observation that cnidarian Bcl10 can activate human MALT1 indicates a highly conserved interaction surface between the two proteins. This type of conservation could be used to further model the interaction surfaces using evolutionary data<sup>39</sup>. A major future challenge will be to collect and functionally evaluate more invertebrate type 1 paracaspase and Bcl10 homologs to verify the proposed correlation of a Bcl10-dependent ancestral type 1 paracaspase paralog with MALT1-like activity. There are several aspects that are yet not clear, for example can no Bcl10 homolog currently be found in lancelets, which clearly have a PCASP3 ortholog. The current model proposes an ancient parallel evolution of a Bcl10-dependent and a Bcl10-independent paracaspase. An alternative scenario is that Bcl10-independence has evolved several times independently. In order to further clarify this, more invertebrate sequences from informative phyla are needed<sup>40</sup>. Several proteins associated to MALT1 in humans are conserved as far back as cnidarians, such as Bcl10, TRAF6, TRAF2 and CYLD<sup>26</sup>. Investigating early-diverging biological systems such as the cnidarian model organism *Nematostella vectensis*<sup>41</sup> for protein interactions and signal transduction mechanisms could further pin-point the original and most conserved functions.

## Acknowledgements

*Nematostella vectensis* cDNA was kindly provided by Ismail Gul (Prof Frans Van Roy, Ghent University), Zebrafish cDNA was kindly provided by Prof. Kris Vleminckx (Ghent University). A pacific oyster (*Crassostrea gigas*) cDNA library was kindly provided by Prof. Pascal Favrel (UMR BOREA Biologie des ORganismes et Ecosystèmes Aquatiques, Institut de Biologie Fondamentale et Appliquée, Université de Caen Basse-Normandie).

## Supplemental material

Supplemental text 1 : Species key : 2-letter abbreviations

Supplemental text 2 : FASTA sequences of type 1 paracaspases used in phylogeny

Supplemental text 3 : FASTA sequences of Bcl10 homologs used in phylogeny

## References

1. Dierlamm, J. *et al.* The apoptosis inhibitor gene API2 and a novel 18q gene, MLT, are recurrently rearranged in the t(11;18)(q21;q21) associated with mucosa-associated lymphoid tissue lymphomas. *Blood* **93**, 3601–3609 (1999).
2. Che, T. *et al.* MALT1/paracaspase is a signaling component downstream of CARMA1 and mediates T cell receptor-induced NF-kappaB activation. *J. Biol. Chem.* **279**, 15870–15876 (2004).
3. Ruefli-Brasse, A. A., French, D. M. & Dixit, V. M. Regulation of NF-kappaB-dependent lymphocyte activation and development by paracaspase. *Science* **302**, 1581–4 (2003).
4. Ruland, J., Duncan, G. S., Wakeham, A. & Mak, T. W. Differential requirement for Malt1 in T and B cell antigen receptor signaling. *Immunity* **19**, 749–58 (2003).
5. Scudiero, I., Vito, P. & Stilo, R. The Three CARMA Sisters: So Different, So Similar. A Portrait of the Three CARMA Proteins and their Involvement in Human Disorders. *J. Cell. Physiol.* (2013). doi:10.1002/jcp.24543
6. Uren, A. G. *et al.* Identification of paracaspases and metacaspases: two ancient families of caspase-like proteins, one of which plays a key role in MALT lymphoma. *Mol Cell* **6**, 961–7 (2000).
7. Coornaert, B. *et al.* T cell antigen receptor stimulation induces MALT1 paracaspase-mediated cleavage of the NF-kappaB inhibitor A20. *Nat Immunol* **9**, 263–71 (2008).
8. Rebeaud, F. *et al.* The proteolytic activity of the paracaspase MALT1 is key in T cell activation. *Nat Immunol* **9**, 272–81 (2008).
9. Hachmann, J. *et al.* Mechanism and specificity of the human paracaspase MALT1. *Biochem. J.* **443**, 287–295 (2012).
10. Yu, J. W., Jeffrey, P. D., Ha, J. Y., Yang, X. & Shi, Y. Crystal Structure of the Mucosa-Associated Lymphoid Tissue Lymphoma Translocation 1 (MALT1) Paracaspase Region. *Proc. Natl. Acad. Sci.* **108**, 21004–21009 (2011).
11. Wiesmann, C. *et al.* Structural Determinants of MALT1 Protease Activity. *J. Mol. Biol.* **419**, 4–21 (2012).
12. Nie, Z. *et al.* Conversion of the LIMA1 tumour suppressor into an oncogenic LMO-like protein by API2-MALT1 in MALT lymphoma. *Nat. Commun.* **6**, 5908 (2015).
13. Fontan, L. *et al.* MALT1 small molecule inhibitors specifically suppress ABC-DLBCL in vitro




and in vivo. *Cancer Cell* **22**, 812–824 (2012).

14. Baens, M. *et al.* MALT1 auto-proteolysis is essential for NF- $\kappa$ B-dependent gene transcription in activated lymphocytes. *PloS One* **9**, e103774 (2014).
15. Staal, J. *et al.* T-cell receptor-induced JNK activation requires proteolytic inactivation of CYLD by MALT1. *EMBO J.* **30**, 1742–1752 (2011).
16. Hailfinger, S. *et al.* Malt1-Dependent RelB Cleavage Promotes Canonical NF- $\kappa$ B Activation in Lymphocytes and Lymphoma Cell Lines. *Proc. Natl. Acad. Sci.* **108**, 14596–14601 (2011).
17. Elton, L. *et al.* MALT1 cleaves the E3 ubiquitin ligase HOIL-1 in activated T cells, generating a dominant negative inhibitor of LUBAC-induced NF- $\kappa$ B signaling. *FEBS J.* (2015).  
doi:10.1111/febs.13597
18. Klein, T. *et al.* The paracaspase MALT1 cleaves HOIL1 reducing linear ubiquitination by LUBAC to dampen lymphocyte NF- $\kappa$ B signalling. *Nat. Commun.* **6**, 8777 (2015).
19. Douanne, T., Gavard, J. & Bidère, N. The paracaspase MALT1 cleaves the LUBAC subunit HOIL1 during antigen receptor signaling. *J. Cell Sci.* (2016). doi:10.1242/jcs.185025
20. Uehata, T. *et al.* Malt1-Induced Cleavage of Regnase-1 in CD4<sup>+</sup> Helper T Cells Regulates Immune Activation. *Cell* **153**, 1036–1049 (2013).
21. Jeltsch, K. M. *et al.* Cleavage of roquin and regnase-1 by the paracaspase MALT1 releases their cooperatively repressed targets to promote T(H)17 differentiation. *Nat. Immunol.* **15**, 1079–1089 (2014).
22. Ferch, U. *et al.* MALT1 directs B cell receptor-induced canonical nuclear factor-kappaB signaling selectively to the c-Rel subunit. *Nat. Immunol.* **8**, 984–991 (2007).
23. Gringhuis, S. I. *et al.* Selective C-Rel activation via Malt1 controls anti-fungal T(H)-17 immunity by dectin-1 and dectin-2. *PLoS Pathog.* **7**, e1001259 (2011).
24. Hamilton, K. S. *et al.* T cell receptor-dependent activation of mTOR signaling in T cells is mediated by Carma1 and MALT1, but not Bcl10. *Sci. Signal.* **7**, ra55 (2014).
25. Bognar, M. K. *et al.* Oncogenic CARMA1 couples NF- $\kappa$ B and  $\beta$ -catenin signaling in diffuse large B-cell lymphomas. *Oncogene* (2016). doi:10.1038/onc.2015.493
26. Hulpiau, P., Driège, Y., Staal, J. & Beyaert, R. MALT1 is not alone after all: identification of novel paracaspases. *Cell. Mol. Life Sci. CMLS* (2015). doi:10.1007/s00018-015-2041-9
27. Peterson, K. J. *et al.* Estimating metazoan divergence times with a molecular clock. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 6536–6541 (2004).

28. Dunn, C. W., Giribet, G., Edgecombe, G. D. & Hejnol, A. Animal Phylogeny and Its Evolutionary Implications. *Annu. Rev. Ecol. Evol. Syst.* **45**, 371–395 (2014).
29. Sullivan, J. C. *et al.* Two Alleles of NF- $\kappa$ B in the Sea Anemone *Nematostella vectensis* Are Widely Dispersed in Nature and Encode Proteins with Distinct Activities. *PLoS ONE* **4**, (2009).
30. Demeyer, A., Staal, J. & Beyaert, R. Targeting MALT1 Proteolytic Activity in Immunity, Inflammation and Disease: Good or Bad? *Trends Mol. Med.* (2016).  
doi:10.1016/j.molmed.2015.12.004
31. Johnson, M. *et al.* NCBI BLAST: a better web interface. *Nucleic Acids Res.* **36**, W5–9 (2008).
32. Notredame, C., Higgins, D. G. & Heringa, J. T-Coffee: A novel method for fast and accurate multiple sequence alignment. *J. Mol. Biol.* **302**, 205–217 (2000).
33. Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinforma. Oxf. Engl.* **19**, 1572–1574 (2003).
34. Okonechnikov, K., Golosova, O., Fursov, M. & UGENE team. Unipro UGENE: a unified bioinformatics toolkit. *Bioinforma. Oxf. Engl.* **28**, 1166–1167 (2012).
35. Torvalds, L. The Linux Edge. *Commun ACM* **42**, 38–39 (1999).
36. Malinverni, C. *et al.* Cleavage by MALT1 induces cytosolic release of A20. *Biochem. Biophys. Res. Commun.* **400**, 543–547 (2010).
37. Calcium phosphate-mediated transfection of eukaryotic cells. *Nat. Methods* **2**, 319–320 (2005).
38. Mazzone, P. *et al.* Functional characterization of zebrafish (*Danio rerio*) Bcl10. *PloS One* **10**, e0122365 (2015).
39. Hopf, T. A. *et al.* Sequence co-evolution gives 3D contacts and structures of protein complexes. *eLife* **3**, (2014).
40. Scientists, G. C. of. The Global Invertebrate Genomics Alliance (GIGA): Developing Community Resources to Study Diverse Invertebrate Genomes. *J. Hered.* **105**, 1–18 (2014).
41. Darling, J. A. *et al.* Rising starlet: the starlet sea anemone, *Nematostella vectensis*. *BioEssays News Rev. Mol. Cell. Dev. Biol.* **27**, 211–221 (2005).

## Figure 1


Phylogenetic tree of the type 1 paracaspases.



Bcl10-containing species like molluscs and hemichordates have vertebrate-like type 1 paracaspases, and cluster with lancelets and tunicates. The paracaspase from the Bcl10-containing Annelid could not be resolved with the currently available sequence information.

## Figure 2

Phylogenetic tree of Bcl10




The hemichordate acorn worm cluster with the other invertebrates. Vertebrate Bcl10 roughly follows the evolutionary relationship at species-level. Annelid Bcl10 similar to mollusc Bcl10.




### Figure 3

#### Functional conservation of invertebrate paracaspase and Bcl10



**A)** NF- $\kappa$ B induction by activated HLH-paracaspase fusions **B)** CYLD cleavage by activated HLH-paracaspase fusions **C)** human MALT1-dependent NF- $\kappa$ B induction by different Bcl10 homologs.

**Figure 4**  
Proposed model



A model that proposes 2 ancient type 1 paracaspases, one Bcl10-dependent and one Bcl10-independent. The Bcl10-dependent type 1 paracaspase shows MALT1-like activities. We don't know when the paralogs originated but it needs to have been before or at the planulozoan last common ancestor (LCA). Deuterostomia (including vertebrates and hemichordates like the acorn worm), annelids and molluscs inherited the Bcl10-dependent type 1 paracaspase (B) whereas most other bilateran invertebrates kept the Bcl10-independent type 1 paracaspase (A). The model is based on currently available reliable sequence information and might change with additional data.