

# Widespread Historical Contingency in Influenza Viruses

Jean Claude Nshogozabahizi<sup>1,2</sup>, Jonathan Dench<sup>1,2</sup> and Stéphane Aris-Brosou<sup>1,2,\*</sup>

**1 Department of Biology, Center for Advanced Research in Environmental Genomics, University of Ottawa, Ottawa, ON, Canada**

**2 Department of Mathematics and Statistics, University of Ottawa, Ottawa, ON, Canada**

**These authors contributed equally to this work.**

**\*sarisbro@uottawa.ca**

## Abstract

In systems biology and genomics, epistasis characterizes the impact that a substitution at a particular location in a genome can have on a substitution at another location. This phenomenon is often implicated in the evolution of drug resistance or to explain why particular ‘disease-causing’ mutations do not have the same outcome in all individuals. Hence, uncovering these mutations and their location in a genome is a central question in biology. However, epistasis is notoriously difficult to uncover, especially in fast-evolving organisms. Here, we present a novel statistical approach that relies on a model developed in ecology and that we adapt to analyze genetic data in fast-evolving systems such as the influenza A virus. We validate the approach using a two-pronged strategy: extensive simulations demonstrate a low-to-moderate sensitivity with an excellent specificity, while analyses of experimentally-validated data recover known interactions, including in a eukaryotic system. We further evaluate the ability of our approach to detect correlated evolution during antigenic shifts or at the emergence of drug resistance. We show that in all cases, correlated evolution is prevalent in influenza A viruses, involving many pairs of sites linked together in chains, a hallmark of historical contingency. Strikingly, interacting sites are separated by large physical distances, which entails either long-range conformational changes or functional tradeoffs, for which we find support with the emergence of drug resistance. Our work paves a new way for the unbiased detection of epistasis in a wide range of organisms by performing whole-genome scans.

## Introduction

One of the most fundamental questions in biology is about the emergence of new structures and new functions, in particular at the molecular and genetic level (Lynch, 2007). As such, a large body of experimental work has accumulated over the past decade to unravel the mutational history at the origin of simple phenotypes. For instance, one particular bacterial drug resistance is conferred by five mutations, but out of the  $5! = 120$  possible ways in which these mutations can accumulate, only a handful of mutational trajectories are experimentally possible (Weinreich *et al.*, 2006). This line of work suggests that some mutations are required in order for subsequent mutations to occur. Further work demonstrates that such permissive mutations are not limited to bacteria, as they are also found in vertebrate (Ortlund *et al.*, 2007), yeasts (Sorrells *et al.*, 2015) and viral systems (Gong *et al.*, 2013). However, while such chains of

dependent or conditional substitutions —called historical contingency (Mohrig *et al.*, 1995; Harms and Thornton, 2014)— are expected to lead to mutational trajectories, their shape and ramifications are not completely elucidated.

As the experimental determination of these trajectories can be tedious, taking > 20 years in the case of the Long-Term Evolution Experiment (Blount *et al.*, 2008), computational solutions were sought to reconstruct historical contingencies and the mutational correlations they imply. Initial solutions relied on protein sequence alignments to compute sitewise vectors of amino acid frequencies, from which pairs of co-evolving residues could be identified (Neher, 1994; Taylor and Hatrick, 1994). While this general approach is still used in statistical physics to predict protein folds (Shindyalov *et al.*, 1994; Sutto *et al.*, 2015), numerous refinements were brought either through the use of metrics such as mutual information (Korber *et al.*, 1993; Atchley *et al.*, 2000; Gloor *et al.*, 2005) or by correcting for shared evolutionary history. One of the first methods to detect correlated evolution while accounting for phylogeny was given in the general context of the evolution of discrete morphological characters (Pagel, 1994). Further leveraging on Schöniger and von Haeseler (1994), the method was quickly extended to analyze RNA molecules by modeling dinucleotides (Muse, 1995; Rzhetsky, 1995) and to map the correlated residues hence detected onto a three-dimensional protein structure (Pollock *et al.*, 1999; Poon *et al.*, 2007b,a). More recently, a full evolutionary model was proposed to detect epistatic sites in a Bayesian framework (Nasrallah and Huelsenbeck, 2013). However, such approaches rely on complex models and Bayesian computing tools, and may scale up poorly with increasingly large data sets (Poon *et al.*, 2008; Aris-Brosou and Rodrigue, 2012).

Here we build on these developments to describe a novel, yet intuitive, statistical method for detecting correlated evolution among pairs of AAs in the typically fast-evolving influenza A virus (Worobey *et al.*, 2014). Our method takes inspiration from an approach developed in ecology and aimed at detecting correlated evolution among phenotypic traits (Pagel, 1994; Pagel and Meade, 2006). We validate our approach using a two-pronged procedure based on both extensive simulations and analyses of experimentally-validated data sets (both in viral and in eukaryotic systems). Analysis of several large data sets lead us to reconsider the nature of epistasis in influenza viruses. We find evidence that interacting AAs form networks of sites undergoing substitutions that are most likely to be permissive, as they occur in a temporal sequence. These networks cover large physical distances among interacting AAs suggesting long-range structural and/or functional effects.

## Materials and Methods

### General approach to detect epistasis

**Repurposing of BayesTraits.** In order to model correlated evolution at the molecular level, we employed the maximum likelihood model implemented in BayesTraits (Pagel, 1994; Pagel and Meade, 2006). See Talavera *et al.* (2015) for a similar model. This model was originally developed as a time-homogeneous Markov process with discrete states in continuous time in order to investigate the coevolution of discrete binary traits on phylogenetic trees. This is achieved by testing whether a dependent model of trait evolution fits the data better than an independent model. The dependent model allows two traits to coevolve as the rate of change at one trait depends on the state at the other trait, while the independent model does not place any restriction on rates of change (Pagel, 1994) (see Figure S1). In both cases, the likelihood function is optimized by summing (integrating) over all unobserved pairs of character states at internal nodes. As these two models are nested, a likelihood ratio test can be

employed for model selection. The test statistic, twice the log-likelihood difference, is assumed to follow a  $\chi^2$  distribution with four degrees of freedom.

This general framework further assumes a phylogenetic tree, with a known topology and branch lengths proportional to the amount of evolution separating each node; both assumptions can be addressed as described below, either by a bootstrap analysis or by resorting to trees sampled from their posterior distribution.

**Data recoding.** The model described above was implemented for binary traits. Here however, our goal is to analyze the coevolution of pairs of sites (DNA or AA) along a sequence alignment. In the case of proteins, each site has twenty possible AAs, which represent the states of our system. To avoid resorting to tensor kernels, data recoding is therefore necessary to reconcile the data with the approach. In this context, we evaluated two strategies. First, AAs were partitioned according to their physicochemical properties. Binary properties (side-chain group types) were naturally recoded “0”/“1”. For those with  $k > 2$  states, we compared each of the  $k_i$  states against the other ones ( $k_j$  where  $i \neq j$ ). For instance, in the case of charge, we first assigned state “0” to negative and state “1” to non-negative AAs, and circled through the two other states (Figure S2). This recoding was based on the physicochemical properties as in the R package `protr` ver. 0.2-1 (Xiao *et al.*, 2014). Second, AAs were classified as either being in the outgroup or ingroup consensus state: at each position of an alignment, the outgroup state was defined as the consensus AA present in a clade used to root the tree. Here, this rooting clade was defined either as the one containing the oldest sequences and this clade was removed for downstream analysis, or as the basal clade in a relaxed molecular clock analysis (see below). The ingroup state was then defined as any AA that differs from the consensus outgroup AA.

**Code optimization.** To discover pairs of AAs that are potentially interacting, we need to run the above model on all  $\frac{n(n-1)}{2}$  pairs of sites in an alignment of length  $n$ . This computation can be prohibitively long even with an alignment of modest size. For instance, the influenza H3N2 nucleoprotein has 498 AAs, which leads to analyzing 123,753 pairs of sites under both the dependent and the independent model (*i.e.*, 247,506 models need to be run). To decrease the computational cost, we first compressed the alignment into site patterns (Yang, 2006) (p.105). Then, as pairs of sites can be independently compared, we parallelized the code using R’s `foreach` ver. 1.4.2 and `doMC` (ver. 1.3.3) (Analytics and Weston, 2013) packages to take advantage of multicore / multiprocessor architectures (Figure S3). To account for multiple testing, we computed the False Discovery Rate (FDR) according to the Benjamini-Hochberg (BH) procedure (Benjamini and Hochberg, 1995). We used R (v3.0.2) for all analyses (Team *et al.*, 2013). In the analyses presented below, the pairs of AAs identified to be interacting were subsequently mapped on three-dimensional protein models predicted by homology modeling with SWISS-MODEL (Biasini *et al.*, 2014) and plotted using KiNG (Chen *et al.*, 2009).

## Validation based on simulations

To validate our method for detecting correlated pairs of sites, we followed two approaches: an extensive simulation study and the analysis of data sets in which epistasis was experimentally confirmed. We first present our simulation strategies that, in order to avoid biasing our results, were based on two ways to simulate correlated evolution. The Supplementary text presents additional simulations.

**Simulations with PHASE.** First, sequences were simulated with PHASE 2.0 (Gowri-Shankar and Jow, 2006), which allowed us to simulate two categories of sites: those that evolve independently and those that evolve in a correlated manner, *i.e.*, under epistasis. Sites that evolve independently (main sequence of length  $l_i$ ) were simulated using the General Time-Reversible (GTR) model of nucleotide substitution (Tavaré, 1986) (see also Aris-Brosou and Rodrigue, 2012). Both nucleotide frequencies and transition probabilities were arbitrarily set to reflect published values for *Pseudomonas aeruginosa* (Garrity *et al.*, 2004), although this particular setting has no impact on downstream analyses. With PHASE, sites that evolve in a correlated manner (epistatic sites  $l_e$ ) were simulated under the RNA7D model of dinucleotide substitution (Tillier and Collins, 1998). We modified the transition probabilities such that double substitutions occurred for 95% of changes, single substitutions for 5%, while mismatch substitutions were not permitted. Equilibrium dinucleotide frequencies were all equal (set to  $\frac{1}{16}$ ). As total sequence length was determined ahead of time (part of our factorial design), when simulating correlated evolution, we first simulated a number  $l_e$  of pairs of epistatic sites, which were then concatenated to nucleotide sites simulated under the independent model of length  $l_i$  to obtain a final sequence of length  $l_s = 2l_e + l_i$ . Where correlated evolution was not simulated  $l_s = l_i$ .

**Simulations with Coev.** Our second simulation strategy employed Coev (Dib *et al.*, 2014), which produces correlated evolution by randomly defining a dinucleotide profile (two-letter nucleotide states, *e.g.*: AA, AT, CG, *etc.*) and simulating evolution under the Jukes-Cantor (Jukes and Cantor, 1969) substitution model. The transition rates for each nucleotide of the dinucleotide may be set independently; however, as we have no *a priori* reason to assume either evolves with a different rate, we left these rates equal to their default setting. The strength of selection for the defined profile is governed by the ratio of the parameters  $d$  and  $s$  (*i.e.*,  $d/s$ ), which represent the likelihood of a dinucleotide evolving toward or away from the defined dinucleotide profile, respectively. When  $d/s = 1$ , there is no selection and sites evolve independently; we varied the strength of selection from independent ( $d/s = 1$ ) to the default setting offered by Coev-web (Dib *et al.*, 2015) ( $d/s = 100$ ), with intermediate strengths ( $d/s = \{2, 33, 66\}$ ). PHASE was still used to simulate sites that evolved independently ( $l_i$ ) to be consistent in use of a GTR substitution model.

**Sensitivity and specificity.** In both sets of simulations, we assessed the impact of branch lengths, tree shape, and number of sequences, both in the presence or absence of epistasis (Figure S4B). With PHASE, branch lengths ( $b$ ) were varied across a  $\log_2$  scale for  $b \in (-12, -1)$ . To reduce possible factorial combinations, all branches of a single simulated tree were the same length. Two tree shapes were used, simulated tree topologies ( $\tau$ ) being either symmetrically bifurcating or pectinate (Figure S4). All simulated trees were ultrametric. Each tree contained a number of sequences ( $n_s$ ) equal to 16, 32, 64 or 128. This led to a full factorial design containing 192 simulation conditions ( $12b \times 2\tau \times 4n_s \times 2$  for with or without epistasis). Each simulation condition was replicated 100 times and  $l_s$  was set to 100 bp. When epistasis was simulated, the number of epistatic pairs was set to 3. An ANOVA was used to assess the significance of each of these factors. All tests are conducted at the  $\alpha = 0.01$  (1%) significance threshold.

With Coev, branch lengths ( $b$ ) were varied across a  $\log_2$  scale for  $b \in \{-12, -10, -8, -6 - 4\}$ . Each tree contained a number of sequences ( $n_s$ ) equal to 32, or 128. The same two tree shapes were used, ( $\tau$ ) being either symmetrically bifurcating or pectinate. This led to a full factorial design containing 100 simulation conditions ( $5b \times 2\tau \times 2n_s \times 5d/s$ ). Each simulation condition was replicated 100 times

and  $l_s$  was set to 100 bp. When epistasis was simulated, the number of epistatic pairs was set to 3. All tests are conducted at the  $\alpha = 0.01$  (1%) significance threshold unless otherwise stated.

## Validations based on previous evidence

**Previous computational analyses.** As a first validation of our approach on actual data, we reanalyzed those studied in a previous computational study (Kryazhimskiy *et al.*, 2011). The four data sets obtained from the original authors consist of: 1,219 HA and 1,836 NA sequences from H1N1 viruses, as well as 2,149 HA and 2,339 NA sequences from H3N2 viruses. These data sets are here denoted KDBP11-H1 (HA in H1N1), KDBP11-N1 (NA in H1N1), KDBP11-H3 (HA in H3N2) and KDBP11-N2 (NA in H3N2), respectively.

**Experimental evidence.** As computational studies make predictions that are not always tested or validated, we reanalyzed data sets in which epistasis was experimentally confirmed. A number of recent studies reported evidence for epistasis and we present results on three of these.

First, we reanalyzed data published by Gong and coworkers (Gong *et al.*, 2013) which comprised 424 H3N2 human influenza A nucleoprotein (NP) sequences spanning 42 years between 1968-2010. This gene was originally chosen because it evolves relatively slowly and is hence amenable to experimental validation as all substitutions can be easily tested by site-directed mutagenesis (Gong *et al.*, 2013). The viral sequences that they used were downloaded from the IVR database (Bao *et al.*, 2008). This data set is here denoted Gong13NP.

We also retrieved a second data set as analyzed by Duan and coworkers (Duan *et al.*, 2014), who used an alignment that contained 1366 human influenza A H1N1 neuraminidase (NA) collected between 1999 and 2009; as in their analysis, the 2009 H1N1 pandemic sequences were excluded. This data set is denoted Duan14NA.

Finally, a recent study performed a combinatorial analysis based on mutations of the eukaryotic tRNA<sup>Arg<sup>CCT</sup></sup> gene to detect epistasis (Li *et al.*, 2016). We ran our computational analysis on an alignment of the eukaryotic tRNA gene constructed in a manner similar to the original authors (SI text: Phylogenetic data of the tRNA genes): we downloaded all available eukaryotic tRNA<sup>Arg<sup>CCT</sup></sup> genes from GtRNAdb (Chan and Lowe, 2016), aligned the sequences with the *cmalign* tool from Infernal (Nawrocki and Eddy, 2013), and kept the 75 nucleotide region that spanned the conserved portion within *Saccharomyces sp.*. As our method requires a phylogenetic tree, we used a phylogeny of eukaryotes (Hedges *et al.*, 2015), keeping only those tips corresponding to the taxa in our alignment. Lastly, as the branch lengths of this tree were in millions of years, we rescaled these to expected number of substitutions using the best available eukaryotic tRNA molecular clock (Soares *et al.*, 2009). While the original authors did not perform a fully exhaustive analysis of all pairs of sites, and our analysis requires sites to be polymorphic, we present our comparison for those pairs of sites tested in both of the analyses. For this analysis only, we used a statistical threshold of  $\alpha = 0.05$  as in the original paper (Li *et al.*, 2016). This data set is denoted Li2016.

## Nature of correlations in influenza evolution

After performing these validations, both on simulations and on experimentally-validated data, we set out to investigate the nature of these interactions by testing two hypotheses. First, we revisited the work by Koel and coworkers (Koel *et al.*, 2013) that experimentally validated the existence of AA substitutions involved in changes of antigenic clusters. For this, we used 877 H3N2 human influenza hemagglutinin (HA)

sequences as in (Koel *et al.*, 2013), collected between 1968 and 2003, to test if these substitutions responsible for antigenic changes also showed evidence for correlated evolution. This data set is denoted Koel13HA.

Second, we tested if our statistical approach could detect some evidence for correlated evolution at pairs of sites involving the S31N substitution, which is responsible for conferring resistance to the anti-influenza drug Adamantane (Abed *et al.*, 2005). For this, we retrieved 668 H3N2 human influenza A matrix protein 2 (M2) sequences that were collected between 1968 and 2003. This data set is denoted Adam03M2.

## Phylogenetic analyses

Maximum likelihood was employed to estimate phylogenetic tree for the KDBP11-H1, KDBP11-N1, KDBP11-H3, KDBP11-N2, Gong13NP, Duan14NA and Adam03M2, alignments with FastTree ver. 2.1.7 (Price *et al.*, 2010) under the WAG + $\Gamma_4$  model to account for among-site rate variation. Trees were rooted using the earliest sequences, which were AAD17229/USA/1918, CAA24269/Japan/1968, AAF77036/USA/1918, ABI92283/Australia/1968, Aichi68-NPA/Aichi/2/1968, A/Victoria/JY2/1968 and A/Albany/17/1968, respectively. The R package APE was used to visualize the trees (Paradis, 2006). For the Koel13HA data set, a rooted tree was reconstructed using BEAST ver. 1.8.0 under a relaxed molecular clock assuming an uncorrelated lognormal prior (Drummond *et al.*, 2006) and a constant-size coalescent prior under the FLU + $\Gamma_4$  substitution model. Analyses were run in duplicate to check convergence, for a total of 100 million steps with a thinning of 5,000; log files were combined with LogCombiner after conservatively removing the first 10% of each chain as a burn-in period, as checked with Tracer ver. 1.5 ([tree.bio.ed.ac.uk/software/tracer](http://tree.bio.ed.ac.uk/software/tracer)).

Phylogenetic uncertainty was taken into account by running our algorithm on bootstrapped trees (or trees sampled from the posterior distribution). Because only the Gong13NP data set showed a large proportion of SH-like aLRT (Anisimova and Gascuel, 2006) node support values in the low (0.0, 0.8) range (Figure S5-S12), the results of the bootstrap analyses are only shown for this case. All data sets and the AEGIS script (Analysis of Epistasis & Genomic Interacting Sites) used in this work are available at [github.com/sarisbro/AEGIS](https://github.com/sarisbro/AEGIS).

## Results and Discussion

### Simulation studies

**Excellent specificity but mediocre sensitivity.** As a first means to validating our approach, we conducted a fully factorial simulation study. In order to avoid biasing our results, the simulation models differ from the analysis models. Our simulation results with PHASE demonstrate that alignments containing fewer than 32 sequences have a poor ability to detect epistasis, with a sensitivity generally  $\leq 20\%$  (Figure S13). For this reason, we henceforth focus on alignments of at least 32 sequences.

The specificity ( $Sp$ ) of our approach is never below 99.09%, which occurs for the symmetric tree shape with the largest number of sequences and rather short branch length of  $2^{-10}$  substitutions per site (Figure 1A). Variation in specificity was so low that we report the value on a  $-\log_{10}(1 - Sp)$  scale in order to emphasize the performance of our approach against thresholds set by the minimum, mean and maximum number of pairwise comparisons made. These thresholds represent the number of true negatives calculable in our analyses (*i.e.*, (# pairwise comparisons - # epistatic pairs)/# pairwise comparisons) and thus performance above these thresholds demonstrates that high

specificity results from low false positive detection rate, and not simply a large dataset with unduly large numbers of true negatives. In spite of the low variation in specificity, tree shape explains most of the variance (Table S1). While we find a two-way interaction with tree shape and with the number of sequences  $n_s$ , there was no direct interaction between branch length and  $n_s$  (Table S1), hereby confirming that, for a given number of sequences, tree shape is driving specificity, which is higher for the pectinate tree ( $F = 7.905$ ,  $df = 1$ ,  $P < 2 \times 10^{-16}$ ; see Figure 1A).

While specificity is excellent, we find that sensitivity is mediocre (Figure 1B), which is consistent with previous studies (Poon *et al.*, 2007b). Our approach's sensitivity is a function of branch lengths and the number of sequences (Table S1), with the existence of an optimal branch length where sensitivity could become excellent, reaching close to 100%, before degrading quickly (on a log scale). This optimal response reflects that short branch lengths carry no information while long ones have random site patterns and thus convey no information (Yang, 1998). Pectinate trees show an optimum for shorter branch lengths than symmetric trees, and an ANOVA confirms an interaction between tree shape and branch length with respect to sensitivity (Table S1). Lastly, we note that larger alignments appear to have better sensitivity (Figure 1A). Yet we find no statistical significance for the interaction of these terms (Table S1). It is possible that the large variance in sensitivity, due to our method of simulating epistasis, has confounded this result.

Due to the low sensitivity of our method, we further investigated the impact of false discovery with two approaches. First, we plotted precision (Figure 1C), which demonstrates that false positives are never more abundant than true positives (minimum threshold). Notably, for alignments with  $> 64$  sequences and with biologically realistic branch lengths ( $2^{-6} \geq$  substitutions per site), our precision is above 95%.

To confirm that our simulation results were not biased by our method of simulating epistasis, we compared our detection method to simulated evolution generated by Coev (Dib *et al.*, 2014). These results (see Supplementary Text) show that both specificity and precision are higher than with the PHASE simulations, while sensitivity is reduced (Figure S14). These findings reflect the particularities of how epistasis was simulated and confirm the excellent specificity of our method. Importantly, these properties are found even when epistasis is simulated at very weak levels (Figure S14).

## Real data analyses

**Little overlap with previous computational results.** As a first evaluation of our algorithm on real data, we reanalyzed four large data sets ( $> 2000$  sequences) previously analyzed with another computational method designed to detect epistasis (Kryazhimskiy *et al.*, 2011). The AA data were recoded as outgroup / ingroup character states to match the above simulations: at each position of the alignment, the “outgroup” state represents the consensus AA in the outgroup sequences and the “ingroup” state represents all the AAs types that differ from the outgroup consensus AA. Overall, our approach not only detects fewer epistatic sites than the previous method, but the two approaches also show little overlap (Figure S15). This minimal overlap is even found before FDR correction (Figure S16, Tables S3-S6), so that lack of power is an unlikely explanation of the difference. Our extensive simulations suggest that this difference may be the result of the low-to-average sensitivity and of the excellent specificity of our method, so that the AAs pairs that we detect may be actually coevolving sites among the truly epistatic pairs. This result begs the question as to whether the few pairs of sites we detect would have any experimental evidence supporting epistasis.

**Extensive overlap with experimentally-validated data.** Because the previous data sets were only examined from a computational point of view, we turned to additional data that have been experimentally validated.

First, we analyzed the Gong13NP data set of 424 influenza NP protein sequences (Gong *et al.*, 2013). As this is the smallest data set in our study, we assessed two ways of recoding the data into binary character states. We first partitioned AAs according to their physicochemical properties. With this recoding strategy, four pairs of sites were detected before FDR correction (Table S7), but none of them matching the pairs detected by Gong and coworkers. Note that after FDR, no interactions were significant (Table S7).

The estimated phylogenetic tree for this data set shows a pectinate (asymmetric) shape and short branch lengths (Figure 2). Our simulation results show that, under these conditions, with  $> 100$  sequences, we can expect a sensitivity  $\geq 80\%$  (Figure 1A), so that the Gong13NP data set fulfills all the conditions for detecting true interactions. This suggests that even if the physicochemical recoding seemed *a priori* to be a good idea, capitalizing on the chemistry of life, it wastes statistical power on multiple three-way tests (Figure S2).

To better mimic our simulation conditions, we then recoded AAs as outgroup/ingroup states. With this recoding strategy, seven pairs of sites were detected: 259:334, 421:425, 246:470, 217:334, 217:343, 259:421 and 186:259 (Figure 2A, Table S8). Among these, the L259S and N334H substitutions both occurred in 1973 and, critically, were experimentally shown to coevolve (Gong *et al.*, 2013). However, we detected six additional pairs of sites that were never shown to be epistatic. These could either be false positives—but again our simulations suggest that our approach is extremely specific (Figure 1B)—or simply coevolving pairs of sites that are not epistatic (coevolution is necessary but not sufficient for epistasis to exist).

While we find multiple pairs of correlated sites, almost all these pairs are all linked to the experimentally confirmed L259S and N334H substitutions (Figure 2B), hereby forming a network of interacting sites. Can we say anything about the nature of these interactions? If they were physical, we would expect that interacting sites would be in close spatial proximity on the folded protein, as in the case of compensatory mutations in RNA molecules (Kimura, 1985; Chen *et al.*, 1999). However, the spatial distribution of these epistatic pairs of sites on the protein structure does not conform to this prediction: while two residues are considered physically linked when their distance is  $\leq 8.5$  Å (Atilgan *et al.*, 2004), we find that the average distance between interacting AAs is 25.9 Å ( $sd = 18.1$ ; Figure 2C). This distribution strongly suggests that chained epistasis is not linked by spatially-close physical interactions (Figure S17), so that thermodynamic (Thomas *et al.*, 2010) or compensatory changes of the 3D structure (Weinreich *et al.*, 2006) act at very long spatial ranges.

To further validate our approach, we also analyzed the Duan14NA data set, comprising 1366 NA sequences of pre-pandemic H1N1 viruses (Duan *et al.*, 2014). We identify the epistatic pair 275:354 (Figure 3, Table S9), which was experimentally proven to confer oseltamivir resistance and that dominated the population in 2008-2009 (Duan *et al.*, 2014). In their study, these authors showed that D354G was the main mutation responsible for maintaining the function of NA after alteration of enzyme activity by H275Y (H274Y in N2 numbering), so that this is potentially the strongest existing interaction. However, our approach fails to identify the five other mutations (V234M, R222Q, K329E, D344N and D354G) that were further identified by Duan and coworkers to be interacting with H275Y. The estimated H1N1 tree is more symmetrical in shape than the one estimated for the Gong13NP data and has shorter average branch lengths (see scale bar in Figure 3). Our simulation results suggest that in this case, the sensitivity of our approach can be very small (Figure 1A). This low sensitivity might



explain why we fail to detect the five additional sites interacting with position 275.

In spite of this negative result, we find that the interacting pair is, again, a long-range interaction (23.7 Å; Figure 3, inset). The objective of the next two paragraphs is to explore more systematically the nature of epistasis in influenza viruses, focusing more specifically on (i) the prevalence of chained long-range epistasis and (ii) the potential nature of these long-range interactions.

Lastly, we ran our approach on the Li2016 data set. While our analysis found a signal for correlated evolution in 25.7% of the pairs of sites tested by the original authors, 90.0% of our significant pairs of sites were identified as epistatic by Li *et al.* (2016) (Figure S18A). We also found a small number of site pairs (37) that were not detected to be epistatic by the original authors. We note however that (i) four of these 37 site pairs could not be statistically tested as the original study had only one biological replicate, (ii) all the site pairs we identify as correlated have epistasis measures that fall within the range of those site pairs deemed significant by the original authors (Figure S18B) and (iii) one site pair we identified forms a Watson-Crick base pair in the folded tRNA molecule. It is therefore possible that the variability in fitness measures in the original high throughput experiment be at least partially responsible for these discrepancies. In any case, this comparison supports the results of our simulation studies that show that our method has excellent specificity and mediocre sensitivity.

**Correlated networks follow a temporal pattern.** In a fifth study, the original authors investigated the mutations involved in changes of antigenic clusters, and found that double mutations could suffice to explain such cluster changes (Koel *et al.*, 2013). The high mutation rate of this virus' antigene however does not explain why these cluster changes do not occur more often than the observed 3.3 years, which led the original authors to postulate that “co-mutations” may be required to maintain viral fitness. In light of our study, the immediate interpretation of these results would be that correlated evolution is involved during cluster change. We re-analyzed these data in order to test this hypothesis.

By doing so, we find that some of the substitutions previously (and experimentally) implicated in cluster change are indeed involved in epistasis (Figure 4A-B, Table S10). In particular, as in the Gong13NP data set, we find evidence for networks of correlations, either as short chains such as position 155 interacting with both 158 and 146, which are involved in two consecutive cluster changes, but we also find a much larger network of interactions involving 20 sites, some of which are also involved in the last four cluster changes (Figure 4B). Again, as in the Gong13NP data set, a temporal sequence of substitutions along this network can be found: G124D, found at the SI87/BE89 transition, interacts with K299R, G172D and E82K; G172D interacts with sites involved in the next transition, BE89/BE92, such as G135K, which is again involved in the next transition, BE92/WU95, where G172E interacts with N262S and V196A, which is itself interacting with K156Q, involved in the WU95/SY97 transition; finally, K156Q interacts with T192I, involved in the SY97/FU02 transition. It is tentative to propose that such chained interactions reflects permissive substitutions and may provide an explanation, as an evolutionary constraint, to the paradox of high mutation rate and slow antigenic evolution. Yet, can we delve further into the nature of these constraints?

At first inspection, Figure 4C suggests that all these interactions are located in the head of the HA protein and hence might respond to steric constraints, *i.e.*, physically-mediated. However, Figure S19 shows that the strength of the association is not related to physical distance between sites of each epistatic pair. Again, the average distance between pairs of interacting sites is 23.7 Å ( $sd = 10.4$ ), which is much larger than the canonical 8.5 Å for close proximity. Can we obtain some evidence about the nature of such long-range interactions?

**Long-range interactions can be functionally-mediated.** The analysis of a sixth data set, Adam03M2, sheds some light on this question. Influenza viruses are resistant to M2 inhibitors, such as adamantane, and this resistance is associated with the S31N amino acid substitution, which is found in more than 95% of the currently circulating viruses (Wang *et al.*, 2013; Garcia and Aris-Brosou, 2014). There is evidence supporting that the spread of S31N may be unrelated to drug selection pressure and instead results from its interaction with advantageous mutations located elsewhere in the viral genome (Simonsen *et al.*, 2007) or maybe just in the M2 gene. To test this epistatic hypothesis, we used our approach to analyze a data set of M2 sequences.

The results show that only one pair of epistatic sites (S31N×V51I) is detected (Figure 5, Table S11). Again, it is a long-range interaction (35.96 Å), but the reason why this data set is illuminating is that the mutation at position 51 has been shown to play a role in virus replication by stabilizing the amphipathic helices of the M2 protein (Stewart and Pekosz, 2011). Thus, V51I may enhance the fitness of M2 protein to increase the frequency of adamantane resistance associated with S31N mutation. The reversion I51V that appeared in few sequences in 2000 (red clade in Figure 5) was apparently quickly lost, which supports the hypothesis that V51I mutation is permissive of S31N. This reversion also supports that, even in the face of high mutation rates, (i) our algorithm still maintains high specificity (Figure 1) and (ii) epistasis can be a very powerful force.

## Conclusions

With historical contingency, the accumulation of epistatic substitutions can be seen as a coevolutionary process, where what happens at one AA site depends on what happened at another site. Here, it is this very idea of coevolution that we harnessed by co-opting a method developed in ecology to test for the correlated evolution of phenotypic traits (Pagel, 1994; Pagel and Meade, 2006): instead of treating pairs of phenotypic traits as such, we repurposed the method to deal with pairs of AA sites. Because the original method was developed to handle binary traits, we explored two ways of recoding data and showed that treating AAs as outgroup/ingroup consensus states was a more sensible (albeit less intuitive) option than using physicochemical properties. Extensive simulations demonstrated mediocre sensitivity, but excellent specificity so that pairs of AAs that are detected can be assumed to be actually coevolving. We then validated our approach against other computational results, showing little overlap, and against experimentally-validated results, showing extensive overlap, hereby suggesting that detected pairs of AAs are genuinely interacting.

With this good statistical behavior, further analyses of independent influenza data sets showed a consistent pattern: (i) many pairs of correlated sites are involved in epistatic interactions, (ii) these pairs of sites form extensive networks of sites, consistent with Poon *et al.* (2007b), but that are also affected by substitutions that occur sequentially – showing evidence for historical contingency in fast-evolving organisms – and (iii), more intriguingly, that these epistatic pairs of site form long-range spatial interactions. This latter point precludes the idea of a close physical link as in the case of tRNA molecules (Kimura, 1985; Chen *et al.*, 1999), so that these long-range interactions must bring about stability (Thomas *et al.*, 2010) and/or conformational (Mitraki *et al.*, 1991; Newcomb *et al.*, 1997; Harms and Thornton, 2014) and/or functional changes, as in the case of the M2 data set or in the case of the Ebola virus (Ibeh *et al.*, 2016). While there is evidence that epistasis can be prevalent in RNA viruses (at least 31% in (Shapiro *et al.*, 2006)) and in bacteria (15% in (Weinreich *et al.*, 2006)), it is not impossible that epistasis reflects an evolutionary constraint stronger in RNA viruses than in organisms with larger and more redundant genomes: because these viruses have a small genome, mutations are expected to have large fitness effects, which can be

alleviated by compensatory mutations (Sanjuán and Elena, 2006).

Our choice of focusing on a segmented RNA virus such as influenza may be problematic, in particular in the case of H3N2 viruses, which show a pattern of punctuated evolution that can be interpreted as the result of clonal interference (Illingworth and Mustonen, 2012; Strelkowa and Lässig, 2012). In absence of recombination, each segment of the virus evolves as a clone, and each clone accumulates different beneficial mutations, only one of which becomes fixed, alongside hitchhiking deleterious mutations that, in our context, would show a correlation pattern. While (i) this process has to date only be found in H3N2 viruses, and (ii) we find evidence for chained epistasis in H1N1 viruses (KDBP11-H1, KDBP11-N1, Duan14NA) as well as in eukaryotic tRNA gene (Li2016), clonal interference remains a problem for our approach. On the other hand, the use of influenza has allowed us to limit our analysis to searching for evidence of epistasis within genes, *contra* among genes. This way of analyzing data intra-genically might make sense in the case of such viruses, as different segments code for proteins involved in relatively different functions. Yet, experimental evidence in other organisms, such as yeasts, shows that epistatic interactions can involve multiple genes and hence be *inter-genic* (Sorrells *et al.*, 2015). Although a method to detect epistasis in segmented genomes was proposed (Neverov *et al.*, 2015), the computational costs of whole-genome scans can seem prohibitive, assessing the genomic prevalence of epistasis remains an unexplored area – but one that we are currently investigating.

In this context, how can we explain the existence of large networks of interacting sites? One possibility would be that a changing environment creates new adaptive landscapes, and that natural populations (*contra* those from *in vitro* studies) do not climb peaks on the landscape but rather chase moving targets (Gavrilets, 2004, p. 36). While it is not clear whether such landscapes are robust to changing environments (Hartl, 2014), they are certainly a reality in the world of viruses, where vaccination regularly alters the adaptive landscape, hereby leading to chained networks of epistatic interactions – a mere by-product of evolution. But then, one can wonder if the metaphor of a landscape itself is appropriate when all mutational trajectories are not accessible from particular genomic backgrounds (Weinreich, 2010; Sorrells *et al.*, 2015). This may be one of the reasons why evolution is so difficult to predict (Weinreich, 2010; Sandie and Aris-Brosou, 2014).

## Acknowledgments

We thank Jeremy Dettman, Neke Ibeh, Rees Kassen, Nicolas Rodrigue and Alex Wong for fruitful discussions, Sergey Kryazhimskiy and Joshua Plotkin as well as Chuan Li and Jianzhi Zhang for sharing data with us. We are grateful to Compute Canada and to Ontario’s High Performance Computing Virtual Laboratory for giving us compute time. This work was supported by the Natural Sciences Research Council of Canada and by the Canada Foundation for Innovation (SAB) and by the University of Ottawa (JCN, JD).

## References

- Abed, Y., N. Goyette, and G. Boivin, 2005 Generation and characterization of recombinant influenza A (H1N1) viruses harboring amantadine resistance mutations. *Antimicrob Agents Chemother* **49**: 556–9.
- Analytics, R. and S. Weston, 2013 foreach: Foreach looping construct for r. R package version **1**: 2013.

- Anisimova, M. and O. Gascuel, 2006 Approximate likelihood-ratio test for branches: A fast, accurate, and powerful alternative. *Syst Biol* **55**: 539–52.
- Aris-Brosou, S. and N. Rodrigue, 2012 The essentials of computational molecular evolution. *Methods Mol Biol* **855**: 111–52.
- Atchley, W. R., K. R. Wollenberg, W. M. Fitch, W. Terhalle, and A. W. Dress, 2000 Correlations among amino acid sites in bhlh protein domains: an information theoretic analysis. *Mol Biol Evol* **17**: 164–78.
- Atilgan, A. R., P. Akan, and C. Baysal, 2004 Small-world communication of residues and significance for protein dynamics. *Biophysical Journal* **86**: 85–91.
- Bao, Y., P. Bolotov, D. Dernovoy, B. Kiryutin, L. Zaslavsky, T. Tatusova, J. Ostell, and D. Lipman, 2008 The influenza virus resource at the national center for biotechnology information. *Journal of Virology* **82**: 596–601.
- Benjamini, Y. and Y. Hochberg, 1995 Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**: 289–300.
- Biasini, M., S. Bienert, A. Waterhouse, K. Arnold, G. Studer, T. Schmidt, F. Kiefer, T. Gallo Cassarino, M. Bertoni, L. Bordoli, and T. Schwede, 2014 Swiss-model: modelling protein tertiary and quaternary structure using evolutionary information. *Nucleic Acids Res* **42**: W252–8.
- Blount, Z. D., C. Z. Borland, and R. E. Lenski, 2008 Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*. *Proc Natl Acad Sci U S A* **105**: 7899–906.
- Chan, P. P. and T. M. Lowe, 2016 GtRNADB 2.0: an expanded database of transfer RNA genes identified in complete and draft genomes. *Nucleic Acids Res* **44**: D184–9.
- Chen, V. B., I. W. Davis, and D. C. Richardson, 2009 King (kinemage, next generation): a versatile interactive molecular and scientific visualization program. *Protein Science* **18**: 2403–2409.
- Chen, Y., D. B. Carlini, J. F. Baines, J. Parsch, J. M. Braverman, S. Tanda, and W. Stephan, 1999 RNA secondary structure and compensatory evolution. *Genes & Genetic Systems* **74**: 271–286.
- Dib, L., X. Meyer, P. Artimo, V. Ioannidis, H. Stockinger, and N. Salamin, 2015 Coev-web: a web platform designed to simulate and evaluate coevolving positions along a phylogenetic tree. *BMC Bioinformatics* **16**: 394.
- Dib, L., D. Silvestro, and N. Salamin, 2014 Evolutionary footprint of coevolving positions in genes. *Bioinformatics* **30**: 1241–9.
- Drummond, A. J., S. Y. Ho, M. J. Phillips, and A. Rambaut, 2006 Relaxed phylogenetics and dating with confidence. *PLoS Biology* **4**: e88.
- Duan, S., E. A. Govorkova, J. Bahl, H. Zaraket, T. Baranovich, P. Seiler, K. Prevost, R. G. Webster, and R. J. Webby, 2014 Epistatic interactions between neuraminidase mutations facilitated the emergence of the oseltamivir-resistant H1N1 influenza viruses. *Nature Communications* **5**.
- Garcia, V. and S. Aris-Brosou, 2014 Comparative dynamics and distribution of influenza drug resistance acquisition to protein M2 and neuraminidase inhibitors. *Mol Biol Evol* **31**: 355–63.


- Garrity, G. M., J. A. Bell, and T. G. Lilburn, 2004 Taxonomic outline of the prokaryotes. Bergey's manual of systematic bacteriology. Springer, New York, Berlin, Heidelberg .
- Gavrilets, S., 2004 *Fitness landscapes and the origin of species*, volume 41. Princeton University Press, Princeton, N.J.
- Gloor, G. B., L. C. Martin, L. M. Wahl, and S. D. Dunn, 2005 Mutual information in protein multiple sequence alignments reveals two classes of coevolving positions. *Biochemistry* **44**: 7156–65.
- Gong, L. I., M. A. Suchard, and J. D. Bloom, 2013 Stability-mediated epistasis constrains the evolution of an influenza protein. *Elife* **2**.
- Gowri-Shankar, V. and H. Jow, 2006 PHASE: a software package for phylogenetics and sequence evolution. Online - URL: <http://www.bioinf.man.ac.uk/resources/phase/>.
- Harms, M. J. and J. W. Thornton, 2014 Historical contingency and its biophysical basis in glucocorticoid receptor evolution. *Nature* **512**: 203–7.
- Hartl, D. L., 2014 What can we learn from fitness landscapes? *Curr Opin Microbiol* **21**: 51–7.
- Hedges, S. B., J. Marin, M. Suleski, M. Paymer, and S. Kumar, 2015 Tree of life reveals clock-like speciation and diversification. *Mol Biol Evol* **32**: 835–45.
- Ibeh, N., J. C. Nshogozabahizi, and S. Aris-Brosou, 2016 Both epistasis and diversifying selection drive the structural evolution of the Ebola virus glycoprotein mucin-like domain. *J Virol* **90**: 5475–84.
- Illingworth, C. J. R. and V. Mustonen, 2012 Components of selection in the evolution of the influenza virus: linkage effects beat inherent selection. *PLoS Pathog* **8**: e1003091.
- Jukes, T. H. and C. R. Cantor, 1969 Evolution of protein molecules. *Mammalian protein metabolism* **3**: 132.
- Kimura, M., 1985 The role of compensatory neutral mutations in molecular evolution. *Journal of Genetics* **64**: 7–19.
- Koel, B. F., D. F. Burke, T. M. Bestebroer, S. van der Vliet, G. C. Zondag, G. Vervaet, E. Skepner, N. S. Lewis, M. I. Spronken, C. A. Russell, *et al.*, 2013 Substitutions near the receptor binding site determine major antigenic change during influenza virus evolution. *Science* **342**: 976–979.
- Korber, B. T., R. M. Farber, D. H. Wolpert, and A. S. Lapedes, 1993 Covariation of mutations in the V3 loop of human immunodeficiency virus type 1 envelope protein: an information theoretic analysis. *Proc Natl Acad Sci U S A* **90**: 7176–80.
- Kryazhimskiy, S., J. Dushoff, G. A. Bazykin, and J. B. Plotkin, 2011 Prevalence of epistasis in the evolution of influenza A surface proteins. *PLoS genetics* **7**: e1001301.
- Li, C., W. Qian, C. J. Maclean, and J. Zhang, 2016 The fitness landscape of a tRNA gene. *Science* **352**: 837–840.
- Lynch, M., 2007 *The origins of genome architecture*. Sinauer Associates, Sunderland, Mass.
- Mitraki, A., B. Fane, C. Haase-Pettingell, J. Sturtevant, and J. King, 1991 Global suppression of protein folding defects and inclusion body formation. *Science* **253**: 54–8.

- Mohrig, J. R., K. A. Moerke, D. L. Cloutier, B. D. Lane, E. C. Person, and T. B. Onasch, 1995 Importance of historical contingency in the stereochemistry of hydratase-dehydratase enzymes. *Science* **269**: 527–9.
- Muse, S. V., 1995 Evolutionary analyses of DNA sequences subject to constraints of secondary structure. *Genetics* **139**: 1429–39.
- Nasrallah, C. A. and J. P. Huelsenbeck, 2013 A phylogenetic model for the detection of epistatic interactions. *Molecular biology and evolution* **30**: 2197–2208.
- Nawrocki, E. P. and S. R. Eddy, 2013 Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics* **29**: 2933–5.
- Neher, E., 1994 How frequent are correlated changes in families of protein sequences? *Proc Natl Acad Sci U S A* **91**: 98–102.
- Neverov, A. D., S. Kryazhimskiy, J. B. Plotkin, and G. A. Bazykin, 2015 Coordinated evolution of influenza A surface proteins. *PLoS Genet* **11**: e1005404.
- Newcomb, R. D., P. M. Campbell, D. L. Ollis, E. Cheah, R. J. Russell, and J. G. Oakeshott, 1997 A single amino acid substitution converts a carboxylesterase to an organophosphorus hydrolase and confers insecticide resistance on a blowfly. *Proc Natl Acad Sci U S A* **94**: 7464–8.
- Ortlund, E. A., J. T. Bridgham, M. R. Redinbo, and J. W. Thornton, 2007 Crystal structure of an ancient protein: evolution by conformational epistasis. *Science* **317**: 1544–8.
- Pagel, M., 1994 Detecting correlated evolution on phylogenies: a general method for the comparative analysis of discrete characters. *Proceedings of the Royal Society of London B: Biological Sciences* **255**: 37–45.
- Pagel, M. and A. Meade, 2006 Bayesian analysis of correlated evolution of discrete characters by reversible-jump Markov chain Monte Carlo. *The American Naturalist* **167**: 808–825.
- Paradis, E., 2006 *Analysis of phylogenetics and evolution with R*. Springer, New York.
- Pollock, D. D., W. R. Taylor, and N. Goldman, 1999 Coevolving protein residues: maximum likelihood identification and relationship to structure. *J Mol Biol* **287**: 187–98.
- Poon, A. F., F. I. Lewis, S. D. Frost, and S. L. K. Pond, 2008 Spidermonkey: rapid detection of co-evolving sites using Bayesian graphical models. *Bioinformatics* **24**: 1949–1950.
- Poon, A. F. Y., F. I. Lewis, S. L. K. Pond, and S. D. W. Frost, 2007a Evolutionary interactions between N-linked glycosylation sites in the HIV-1 envelope. *PLoS Comput Biol* **3**: e11.
- Poon, A. F. Y., F. I. Lewis, S. L. K. Pond, and S. D. W. Frost, 2007b An evolutionary-network model reveals stratified interactions in the V3 loop of the HIV-1 envelope. *PLoS Comput Biol* **3**: e231.
- Price, M. N., P. S. Dehal, and A. P. Arkin, 2010 Fasttree 2—approximately maximum-likelihood trees for large alignments. *PloS one* **5**: e9490.
- Rzhetsky, A., 1995 Estimating substitution rates in ribosomal RNA genes. *Genetics* **141**: 771–83.


- Sandie, R. and S. Aris-Brosou, 2014 Predicting the emergence of H3N2 influenza viruses reveals contrasted modes of evolution of HA and NA antigens. *J Mol Evol* **78**: 1–12.
- Sanjuán, R. and S. F. Elena, 2006 Epistasis correlates to genomic complexity. *Proc Natl Acad Sci U S A* **103**: 14402–5.
- Schöniger, M. and A. von Haeseler, 1994 A stochastic model for the evolution of autocorrelated dna sequences. *Mol Phylogenet Evol* **3**: 240–7.
- Shapiro, B., A. Rambaut, O. G. Pybus, and E. C. Holmes, 2006 A phylogenetic method for detecting positive epistasis in gene sequences and its application to RNA virus evolution. *Mol Biol Evol* **23**: 1724–30.
- Shindyalov, I. N., N. A. Kolchanov, and C. Sander, 1994 Can three-dimensional contacts in protein structures be predicted by analysis of correlated mutations? *Protein Eng* **7**: 349–58.
- Simonsen, L., C. Viboud, B. T. Grenfell, J. Dushoff, L. Jennings, M. Smit, C. Macken, M. Hata, J. Gog, M. A. Miller, *et al.*, 2007 The genesis and spread of reassortment human influenza A/H3N2 viruses conferring adamantane resistance. *Molecular biology and evolution* **24**: 1811–1820.
- Soares, P., L. Ermini, N. Thomson, M. Mormina, T. Rito, A. Röhl, A. Salas, S. Oppenheimer, V. Macaulay, and M. B. Richards, 2009 Correcting for purifying selection: an improved human mitochondrial molecular clock. *Am J Hum Genet* **84**: 740–59.
- Sorrells, T. R., L. N. Booth, B. B. Tuch, and A. D. Johnson, 2015 Intersecting transcription networks constrain gene regulatory evolution. *Nature* **523**: 361–5.
- Stewart, S. M. and A. Pekosz, 2011 Mutations in the membrane-proximal region of the influenza A virus M2 protein cytoplasmic tail have modest effects on virus replication. *Journal of Virology* **85**: 12179–12187.
- Strelkova, N. and M. Lässig, 2012 Clonal interference in the evolution of influenza. *Genetics* **192**: 671–82.
- Sutto, L., S. Marsili, A. Valencia, and F. L. Gervasio, 2015 From residue coevolution to protein conformational ensembles and functional dynamics. *Proc Natl Acad Sci U S A* **112**: 13567–72.
- Talavera, D., S. C. Lovell, and S. Whelan, 2015 Covariation is a poor measure of molecular coevolution. *Mol Biol Evol* **32**: 2456–68.
- Tavaré, S., 1986 Some probabilistic and statistical problems in the analysis of DNA sequences. In: *Lectures on Mathematics in the Life Sciences* **17**: 57–86.
- Taylor, W. R. and K. Hatrick, 1994 Compensating changes in protein multiple sequence alignments. *Protein Eng* **7**: 341–8.
- Team, R. D. C. *et al.*, 2013 R: A language and environment for statistical computing.
- Thomas, V. L., A. C. McReynolds, and B. K. Shoichet, 2010 Structural bases for stability-function tradeoffs in antibiotic resistance. *J Mol Biol* **396**: 47–59.
- Tillier, E. R. M. and R. A. Collins, 1998 High apparent rate of simultaneous compensatory base-pair substitutions in ribosomal rna. *Genetics* **148**: 1993–2002.

- Wang, J., Y. Wu, C. Ma, G. Fiorin, J. Wang, L. H. Pinto, R. A. Lamb, M. L. Klein, and W. F. DeGrado, 2013 Structure and inhibition of the drug-resistant S31N mutant of the M2 ion channel of influenza A virus. *Proceedings of the National Academy of Sciences* **110**: 1315–1320.
- Weinreich, D. M., 2010 Predicting molecular evolutionary trajectories in principle and in practice. In *Encyclopedia of Life Sciences*, pp. 1–9, John Wiley & Sons, Ltd: Chichester.
- Weinreich, D. M., N. F. Delaney, M. A. DePristo, and D. L. Hartl, 2006 Darwinian evolution can follow only very few mutational paths to fitter proteins. *Science* **312**: 111–114.
- Worobey, M., G.-Z. Han, and A. Rambaut, 2014 A synchronized global sweep of the internal genes of modern avian influenza virus. *Nature* **508**: 254–7.
- Xiao, N., Q. Xu, and D. Cao, 2014 protr: Protein sequence descriptor calculation and similarity computation with r. R package version 0.2-1, URL <http://CRAN.R-project.org/package=protr> .
- Yang, Z., 1998 On the best evolutionary rate for phylogenetic analysis. *Syst Biol* **47**: 125–33.
- Yang, Z., 2006 *Computational molecular evolution*, volume 21. Oxford University Press Oxford.







**Figure 1. Specificity, sensitivity and precision results, from simulated data, of our novel epistasis detection method.** Results are shown for alignments with 32 (squares), 64 (circles) and 128 (triangles) sequences. Tree shapes are color-coded (symmetric in red; pectinate in blue). Branch lengths were varied on a  $\log_2$  scale. All y-axes show the mean of  $-\log_{10}(1 - \langle \text{summary statistic} \rangle)$  to highlight performance of our method as a *summary statistic* approaches 1; when this value is 1, we arbitrarily assigned it a value 10% larger than the largest finite value within the set. Panel (A) shows specificity (true negative rate) with a grey shaded polygon which illustrates thresholds for excellent specificity in parameter space. The thresholds were established by subtracting three (the number of epistatic pairs simulated) from the minimum (lower dashed line), mean (middle dashed line) and maximum (upper dashed line) number of pairwise comparisons performed across all simulations with the same branch length. These thresholds represent the number of calculable true negatives and allow us to demonstrate our method's excellent specificity. Panels (B) and (C) show sensitivity (true positive rate) and precision (positive predictive value) respectively. Each panel includes a grey shaded polygon which illustrates thresholds for 50% (lower dashed line), 95% (middle dashed line) and 99% (upper dashed line) detection. These thresholds were arbitrarily chosen to demonstrate idealized benchmarks of performance.




**Figure 2. Epistatic pairs of amino acids detected in the Gong13NP data set with the outgroup / ingroup recoding.** (A) The epistatic mutations that we detected are plotted on the NP phylogenetic tree. The substitutions in red were experimentally validated (Gong *et al.*, 2013). (B) Chained epistasis of interacting AAs. (C) The epistatic sites are mapped on three dimensional NP protein structure (based on template 3ZDP). The numbers show the AA positions experimentally validated (in red) and those detected only in this study (in black). The numbers in blue show the physical distance between epistatic sites (in Å).



**Figure 3. Epistatic pairs of amino acids detected in the Duan14NA data set.** The epistatic mutations that we detected are plotted on the NA phylogenetic tree. Inset: the epistatic sites are mapped on three dimensional NA protein structure (based on template 1HA0). The numbers show the AA positions experimentally validated (in red). The numbers in blue show the physical distance between epistatic sites (in Å).



**Figure 4. Epistatic pairs of amino acids detected in the Koel13HA data set.** (A) The epistatic mutations that we detected are plotted on the HA phylogenetic tree. The substitutions in red were experimentally validated to be responsible for cluster change (Koel *et al.*, 2013). The antigenic clusters are named after the first vaccine strain in the cluster, with letters and digits referring to location and year of isolation (HK, Hong Kong; EN, England; VI, Victoria; TX, Texas; BK, Bangkok; SI, Sichuan; BE, Beijing; WU, Wuhan; SY, Sydney; FU, Fujian). (B) The thickness of the links is proportional to  $-\log_{10} P$ -value, the strength of evidence supporting the interaction. (C) The epistatic sites are mapped on three dimensional HA protein structure (based on template 3WHE). The numbers show the AA positions experimentally validated (in red) and those detected only in this study (in black). The numbers in blue show the physical distance between epistatic sites (in Å).



**Figure 5.** Epistatic pairs of amino acids detected in the Adam03M2 data set. The epistatic mutations that we detected are plotted on the M2 phylogenetic tree. Inset: the epistatic sites are mapped on three dimensional M2 protein structure (based on template 2KIH). The numbers show the AA positions detected (in red). The numbers in blue show the physical distance between epistatic sites (in Å).