

Protein Collapse is Encoded in the Folded State Architecture

Himadri S. Samanta,¹ Pavel I. Zhuravlev,² Michael Hinczewski,³

Naoto Hori,¹ Shaon Chakrabarti,² and D. Thirumalai^{1,2}

¹*Department of Chemistry, University of Texas at Austin, TX 78712*

²*Biophysics Program, Institute for Physical Science and Technology,
University of Maryland, College Park, MD 20742*

³*Department of Physics, Case Western Reserve University, OH 44106*

Abstract

The propensity of single domain globular proteins, the workhorses in cells, to be compact is the key reason that their folded states achieve high packing density. It is known that the radius of gyration, R_g , of both the folded and unfolded (created by adding denaturants) states increase as N^ν where N is the number of amino acids in the protein. The values of the celebrated Flory exponent ν are, respectively, $\approx \frac{1}{3}$ and ≈ 0.6 in the folded and unfolded states, which coincide with those found in homopolymers in poor and good solvents. However, the extent of compaction of the unfolded state of a protein under low denaturant concentration, conditions favoring the formation of the folded state, is unknown. This problem which goes to the heart of how proteins fold and has implications for the evolution of foldable sequences is unsolved. We develop a theory based on polymer physics concepts that uses the contact map of proteins as input to quantitatively assess collapsibility of proteins. The model, which includes only two-body excluded volume interaction and interactions reflecting the strength of the contact map, has only expanded and compact states. Surprisingly, we find that although protein collapsibility is universal, the propensity to be compact depends on the protein architecture. Application of the theory to over two thousand proteins shows that the extent of collapsibility depends not only on N but also on the contact map reflecting the native fold structure. A major prediction of the theory is that β -sheet proteins are far more collapsible than structures dominated by α -helices. The theory fully resolves the apparent controversy between conclusions reached using different experimental probes assessing the extent of compaction of a couple proteins. In addition, it reveals that there are considerable similarities between the physical mechanisms of homopolymer and protein collapse. The theory provides quantitative insights into the reasons why single domain proteins are small and the physical reasons for the origin of multi-domain proteins. We also show that non-coding RNA molecules, whose collapsibility is similar to proteins with β -sheet structures, must undergo collapse prior to folding, adding support to “Compaction Selection Hypothesis” proposed in the context of RNA compaction.

INTRODUCTION

Folded states of globular proteins, which are evolved (slightly) branched heteropolymers made from twenty amino acids, are roughly spherical and are nearly maximally compact with high packing densities [1–3]. Despite achieving high packing densities in the folded states, globular proteins tolerate large volume substitutions while retaining the native fold [4]. This is explained in a couple of interesting theoretical studies [5, 6], which demonstrated that there is sufficient free volume in the folded state to accommodate mutations. Collectively these and related studies show that folded proteins are compact. When they unfold, which can be achieved upon addition of high concentrations of denaturants (or applying a mechanical force), they swell adopting expanded conformations. The radius of gyration (R_g) of a folded globular protein is well described by the Flory law with $R_g \approx 3.3N^{\frac{1}{3}} \text{ \AA}$ [7] whereas in the swollen state $R_g \approx a_D N^\nu$ where a_D is an effective monomer size and the Flory exponent $\nu \approx 0.6$ [8]. Thus, viewed from this perspective we could surmise that proteins must undergo a coil-to-globule transition [9, 10], a process that is reminiscent of the well characterized equilibrium collapse transition in homopolymers [11, 12]. The latter is driven by the interplay of conformational entropy and intra-polymer interaction energy resulting in the collapsed globular state. The swollen state is realized in good solvents (interaction between monomer and solvents is favorable) whereas in the collapsed state monomer-monomer interactions are preferred. The coil-to-globule transition in large homopolymers is akin to a phase transition. The temperature at which the interactions between the monomers roughly balance monomer-solvent energetics is the θ temperature. By analogy we may identify high (low) denaturant concentrations with good (poor) solvent for proteins.

Despite the expected similarities between the equilibrium collapse transition in homopolymers and the compaction of proteins, it is still debated whether the unfolded states of proteins are more compact under folding conditions compared to their states at high denaturant concentrations. If polypeptide chain compaction is universal, is collapse in proteins essentially the same phenomenon as in homopolymer collapse or is it driven by a different mechanism [13–17]? Surprisingly, this fundamental question in the protein folding field has not been answered satisfactorily [10, 18]. In order to explain the plausible difficulties in quantifying the extent of compaction, let us consider a protein, which undergoes an apparent two-state transition from an unfolded (swollen) to a folded (compact) state as the

denaturant concentration (C) is decreased. At the concentration, C_m , the populations of the folded and unfolded states are equal. A vexing question, which has been difficult to unambiguously answer in experiments, is: what is the size, R_g , of the unfolded state under folding conditions ($C < C_m$)? Small Angle X-ray Scattering (SAXS) experiments on some proteins show practically no change in the unfolded R_g as C is changed[19]. On the other hand, from experiments based on single molecule Fluorescence Resonance Energy Transfer (smFRET) it has been concluded that the size of the unfolded state is more compact below C_m compared to its value at high C [20, 21]. The so-called smFRET-SAXS controversy has remained unresolved. Resolving this apparent controversy is not only important in our understanding of the physics of protein folding but also has implications for the physical basis of the evolution of natural sequences.

The difficulties in describing the collapse of unfolded states as C is lowered could be attributed to the following reasons. (1) Following de Gennes[22], homopolymer collapse can be pictured as formation of a large number of blobs driven by local interactions between monomers on the scale of the blob size. Coarsening of blobs results in the equilibrium globule formation with the number of maximally compact conformations whose number scales exponentially with the number of monomers. Other scenarios resulting in the formation of fractal globules enroute to the formation of equilibrium maximally collapsed structures have also been proposed [23]. The globule formation is driven by non-specific interactions between the monomers or the blobs. Regardless of how the equilibrium globule is reached it is clear that it is largely stabilized by local interactions, because contacts between monomers that are distant along the sequence are entropically unfavorable. In contrast, even in high denaturant concentrations proteins could have residual structure, which likely becomes prominent at $C < C_m$. At low C there are specific favorable interactions between residues separated by a few or several residues along the sequence. As their strength grows, with respect to the entropic forces, the specific interactions may favor compaction in a manner different from the way non-specific local interactions induce homopolymer collapse. In other words, the dominant native-like contacts also drive compaction of unfolded states of proteins. (2) A consequence of the impact of the native-like contacts (local and non-local) on collapse of unfolded states is that specific energetic considerations dictate protein compaction resulting in the formation of minimum energy compact structures (MECS)[24]. The number of MECS, which are not fully native, is small, scaling as $\ln N$ with N being the number of amino

acid residues. Therefore, below C_m their contributions to R_g have to be carefully dissected, which is more easily done in single molecule experiments than in ensemble measurements such as SAXS. (3) Single domain proteins are finite-sized with N rarely exceeding ~ 200 . Most of those studied experimentally have $N < 100$. Thus, the extent of change in R_g of the unfolded states is predicted to be small, requiring high precision experiments to quantify the changes in R_g as C is changed. For example, in a recent study[25], we showed that in PDZ2 domain the change in R_g of the unfolded states as the denaturant concentration changes from 6 M guanidine chloride to 0 M is only about 8%. Recent experiments have also established that changes in R_g in helical proteins are not too large [20].

In homopolymers there are only two possible states, coil and globule, with a transition between the two occurring at T_θ . On the other hand, even in proteins that fold in a two-state manner one can conceive of at least three states (we ignore intermediates here): (i) the unfolded state \mathbf{U}_D at high C ; (ii) the compact but unfolded state \mathbf{U}_C , which could possibly exist below C_m ; (iii) the native state. Do the sizes of \mathbf{U}_D and \mathbf{U}_C differ? This question requires a clear answer as it impacts our understanding of how proteins fold, because the very nature of the unfolded states of proteins plays a key role in determining protein foldability [26–28].

Given the flexibility of proteins (persistence length on the order of 0.5 – 0.6 nm), we expect that the size of the extended polypeptide chain must gradually decrease as the solvent quality is altered. Experiments on a number of proteins show that this is the case[29–31]. However, in some SAXS experiments the theoretical expectation that $R_g^{U_C} < R_g^{U_D}$ for one protein was not borne out [10, 19], precipitating a more general question: are chemically denatured proteins compact at low C ? The absence of collapse is not compatible with inferences based on smFRET [21] and theory [26]. Here, we create a theory to not only resolve the smFRET-SAXS controversy but also provide a quantitative description of how the propensity to be compact is encoded in the native topology. The theory, based on polymer physics concepts, includes specific attractive interactions (mimicking interactions accounting for native contacts in the Protein Data Bank (PDB)) and a two-body excluded volume repulsion. By construction the model does not have a native state. There are only two states (analogues of \mathbf{U}_D and \mathbf{U}_C) in the model. The formation of \mathbf{U}_C is driven by the contact map of the folded state. Thus, chain compaction is driven in much the same way as in homopolymers, altered only by specific interactions that differentiate proteins from

homopolymers.

Our theory predicts quantitatively how the extent of compaction (collapsibility) is determined by the strength and the number of the native contacts and their locations along the chain. We use a large representative selection of proteins from the PDB to establish that collapsibility is an inherent characteristic of evolved protein sequences. A major outcome of this work is that β -sheet proteins are far more collapsible than structures dominated by α -helices. Our theory suggests that there is an evolutionary pressure on proteins for being compact as a pre-requisite for kinetic foldability, as we predicted over twenty years ago [26]. We come to the inevitable conclusion that the unfolded state of proteins must be compact under native conditions, and the mechanism of polypeptide chain compaction has similarities as well as differences to collapse in homopolymers. As a by-product of this work we also establish that certain non-coding RNA molecules must undergo compaction prior to folding as their folded structures are stabilized predominantly by long-range tertiary contacts.

THEORY

We start with an Edwards Hamiltonian for a polymer chain [32]:

$$\mathcal{H} = \frac{3k_B T}{2a_0^2} \int_0^N \left(\frac{\partial \mathbf{r}}{\partial s} \right)^2 ds + k_B T \mathcal{V}(\mathbf{r}(s)), \quad (1)$$

where $\mathbf{r}(s)$ is the position of the monomer s , a_0 the monomer size, and N is the number of monomers. The first term in Eq.(1) accounts for chain connectivity, and the second term represents volume interactions and favorable interactions between select monomers given by $\mathcal{V}(\mathbf{r}(s))$,

$$\mathcal{V}(\mathbf{r}(s)) = \frac{v}{(2\pi a_0^2)^{3/2}} \sum_{s=0}^N \sum_{s'=0}^N e^{-\frac{(\mathbf{r}(s)-\mathbf{r}(s'))^2}{2a_0^2}} - \frac{\kappa}{(2\pi\sigma^2)^{3/2}} \sum_{\{s_i, s_j\}} e^{-\frac{(\mathbf{r}(s_i)-\mathbf{r}(s_j))^2}{2\sigma^2}} \quad (2)$$

The first term in Eq.(2) accounts for the homopolymer (non-specific) two-body interactions. It is well established in the theory of homopolymers that in good solvents with $v > 0$ the polymer swells with $R_g \sim aN^\nu$ ($\nu \approx 0.6$). In poor solvents ($v < 0$) the polymer undergoes a coil-globule transition with $R_g \sim aN^\nu$ ($\nu \approx 1/3$). These are the celebrated Flory laws. Here, we consider only the excluded volume repulsion case ($v > 0$).

The second term in Eq.(2) requires an explanation. The generic scenario for homopolymer collapse is based on an observation by de Gennes, who pictured the collapse process as being driven by the initial formation of blobs that arrange to form a sausage-like structure. At later stages the globule forms to maximize favorable intra-molecular contacts while simultaneously minimizing surface tension. Compaction in proteins, although sharing many features in common with homopolymer collapse, could be different. A key difference is that the folded states of almost all proteins are stabilized by a mixture of local contacts (interaction between residues separated by less than say ~ 8 but greater than 3 residues) as well as non-local (> 8 residues) contacts. Note that the demarcation using 8 between local and non-local contacts is arbitrary, and is not germane to the present argument. These specific interactions also dominate the enthalpy of formation of the compact, non-native state $\mathbf{U}_{\mathbf{C}}$, playing an important role in its stability. Therefore, the topology of the folded state could dictate collapsibility (the extent to which the $\mathbf{U}_{\mathbf{D}}$ state becomes compact as the denaturant concentration is lowered) of a given protein. In combination with the finite size of single domain proteins ($N \sim 200$), the extent of protein collapse could be small. In order to assess chain compaction under native conditions we should consider the second term in Eq.(2).

We note in passing (with discussion to follow) that a number of studies have considered the effect of crosslinks on the shape of polymer chains [33–39]. Polymers with crosslinks have served as models for polymer gels and rubber elasticity[40–42]. In these studies the contacts were either random, leading to the random loop model [34], or explicit averages over the probability of realizing such contacts were made [33, 43], as may be appropriate in modeling gels. These studies inevitably predict a coil-to-globule phase transition as the number of crosslinks increases.

In contrast to models with random crosslinks, in our theory attraction is only between specific residues, described by the second term in Eq.(2), where the sum is over the set of interactions (native contacts) involving pairs $\{s_i, s_j\}$. We use the contact map of the protein (extracted from the PDB structure) in order to assign the specific interactions (their total number being N_{nc}). The contact is assigned to any two residues s_i and s_j if the distance between their C_α atoms in the PDB entry is less than $R_c = 8\text{\AA}$ and $|s_i - s_j| > 2$. We use Gaussian potentials in order to have short (but finite) range attractive interactions. For the excluded volume repulsion, this range is on the order of the size of the monomer, $a_0 = 3.8\text{\AA}$. For the specific attraction, the range is the average distance in the PDB entry between C_α

atoms forming a contact (averaged across a selection of proteins from the PBD). We obtain $\sigma = 6.3\text{\AA}$.

By changing the value of κ , and hence the strength of attraction, there is a transition between the extended and compact states. Decreasing κ is analogous to chemically denaturing proteins, although the connection is not precise. At high denaturant concentrations ($\kappa \approx 0$, good solvent) the excluded volume repulsion (first term in Eq.(2)) dominates the attraction, while at low C (high κ , poor solvent) the attractive interactions are important. The point where attraction balances repulsion is the θ -point, and the value of $\kappa = \kappa_\theta$. Although reserved for the coil-to-globule transition in the limit of $N \gg 1$ in homopolymers, we will use the same notation (θ -point) here. In our model, at the θ -point, the chain behaves like an ideal chain. To describe the globular state, a three-body repulsion needs to be added to the Hamiltonian (Eq.(2)), but we focus on the region between the extended coil and the θ -point because our interest is to access only the collapsibility of proteins. If κ_θ is very large then significant chain compaction would only occur at very low ($C \ll C_m$) denaturant concentrations, implying low propensity to collapse. Conversely, small κ_θ implies ease of collapsibility. Note that the ground state ($\kappa \gg 1$) of the Hamiltonian in Eq.(2) is a collapsed chain whose R_g is on the order of the monomer size. In other words, a stable native state does not exist for the model described in Eq.(2). Thus, we define protein collapse as the propensity of the polypeptide chain to reach the θ -point as measured by the κ_θ value, and use the changes in the radius of gyration R_g as a measure of the extent of compaction.

Assessing collapsibility: For our model, which encodes protein topology without favoring the folded state, we calculate $\langle R_g^2 \rangle$ using the Edwards-Singh (ES) method [44]. The ES method is a variational type calculation that represents the exact Hamiltonian by a Gaussian chain, whose effective monomer size is determined as follows. Consider a virtual chain without excluded volume interactions, with the radius of gyration $\langle R_g^2 \rangle = Na^2/6$ [44], described by the Hamiltonian,

$$\mathcal{H}_v = \frac{3k_B T}{2a^2} \int_0^N \left(\frac{\partial \mathbf{r}}{\partial s} \right)^2 ds, \quad (3)$$

where the monomer size in the virtual Hamiltonian is a . We split the deviation \mathcal{W} between the virtual chain Hamiltonian and the real Hamiltonian as,

$$\mathcal{H} - \mathcal{H}_v = k_B T \mathcal{W} = k_B T (\mathcal{W}_1 + \mathcal{W}_2), \quad (4)$$

where

$$\begin{aligned}\mathcal{W}_1 &= \frac{3}{2} \left(\frac{1}{a_0^2} - \frac{1}{a^2} \right) \int_0^N \left(\frac{\partial \mathbf{r}}{\partial s} \right)^2 ds, \\ \mathcal{W}_2 &= \mathcal{V}(\mathbf{r}(s)).\end{aligned}\quad (5)$$

The radius of gyration is $R_g^2 = \frac{1}{N} \int_0^N \langle \mathbf{r}^2(s) \rangle ds$, with the average being,

$$\langle \mathbf{r}^2(s) \rangle = \frac{\int r^2 e^{-\mathcal{H}/k_B T} \delta \mathbf{r}}{\int e^{-\mathcal{H}/k_B T} \delta \mathbf{r}} = \frac{\int r^2 e^{-\mathcal{H}_v/k_B T} e^{-\mathcal{W}} \delta \mathbf{r}}{\int e^{-\mathcal{H}_v/k_B T} e^{-\mathcal{W}} \delta \mathbf{r}} = \frac{\langle \mathbf{r}^2(s) e^{-\mathcal{W}} \rangle_v}{\langle e^{-\mathcal{W}} \rangle_v} \quad (6)$$

where $\langle \cdots \rangle_v$ denotes the average over \mathcal{H}_v .

Assuming that the deviation \mathcal{W} is small, we calculate the average to first order in \mathcal{W} . The result is,

$$\langle \mathbf{r}^2(s) \rangle \approx \frac{\langle \mathbf{r}^2(s)(1 - \mathcal{W}) \rangle_v}{\langle (1 - \mathcal{W}) \rangle_v} \approx \langle \mathbf{r}^2(s)(1 - \mathcal{W}) \rangle_v \langle (1 + \mathcal{W}) \rangle_v \quad (7)$$

and the radius of gyration is

$$\langle R_g^2 \rangle = \frac{1}{N} \int_0^N \langle \mathbf{r}^2(s) \rangle ds = \frac{1}{N} \int_0^N [\langle \mathbf{r}^2(s) \rangle_v + \langle \mathbf{r}^2(s) \rangle_v \langle \mathcal{W} \rangle_v - \langle \mathbf{r}^2(s) \mathcal{W} \rangle_v] ds, \quad (8)$$

If we choose the effective monomer size a in \mathcal{H}_v such that the first order correction (second and third terms on the right hand side of Eq.(8)) vanishes, then the size of the chain is, $\langle R_g^2 \rangle = Na^2/6$. This is an estimate to the exact $\langle R_g^2 \rangle$, and is an approximation as we have neglected \mathcal{W}^2 and higher powers of \mathcal{W} . Thus, in the ES theory, the optimal value of a from Eq. (8) satisfies,

$$\frac{1}{N} \int_0^N [\langle \mathbf{r}^2(s) \rangle_v \langle \mathcal{W} \rangle_v - \langle \mathbf{r}^2(s) \mathcal{W} \rangle_v] ds = 0. \quad (9)$$

Since $\mathcal{W} = \mathcal{W}_1 + \mathcal{W}_2$, the above equation can be written as

$$\frac{1}{N} \int_0^N [\langle \mathbf{r}^2(s) \rangle_v \langle \mathcal{W}_1 \rangle_v - \langle \mathbf{r}^2(s) \mathcal{W}_1 \rangle_v] ds = -\frac{1}{N} \int_0^N [\langle \mathbf{r}^2(s) \rangle_v \langle \mathcal{W}_2 \rangle_v - \langle \mathbf{r}^2(s) \mathcal{W}_2 \rangle_v] ds. \quad (10)$$

Evaluation of the $\langle \mathbf{r}^2(s) \mathcal{W}_1 \rangle_v$ term yields,

$$\begin{aligned}
 \langle \mathbf{r}^2(s) \mathcal{W}_1 \rangle_v &= \frac{\frac{3}{2} \left(\frac{1}{a_0^2} - \frac{1}{a^2} \right) \int \mathbf{r}^2 \int_0^N \dot{\mathbf{r}}^2 ds e^{-\frac{3}{2a^2} \int_0^N \dot{\mathbf{r}}^2 ds} \delta \mathbf{r}}{\int e^{-\frac{3}{2a^2} \int_0^N \dot{\mathbf{r}}^2 ds} \delta \mathbf{r}} \\
 &= \frac{3}{2} \left(\frac{1}{a_0^2} - \frac{1}{a^2} \right) \left[\frac{\partial}{\partial \alpha} \left(\frac{\int \delta \mathbf{r} \mathbf{r}^2 e^{\alpha \int \dot{\mathbf{r}}^2 ds}}{\int \delta \mathbf{r} e^{\alpha \int \dot{\mathbf{r}}^2 ds}} \right) \Big|_{\alpha = -\frac{3}{2a^2}} + \frac{\int \mathbf{r}^2 e^{-\frac{3}{2a^2} \int_0^N \dot{\mathbf{r}}^2 ds} \delta \mathbf{r}}{\left(\int e^{-\frac{3}{2a^2} \int_0^N \dot{\mathbf{r}}^2 ds} \delta \mathbf{r} \right)^2} \int \int_0^N \dot{\mathbf{r}}^2 ds e^{-\frac{3}{2a^2} \int_0^N \dot{\mathbf{r}}^2 ds} \delta \mathbf{r} \right] \\
 &= \left(\frac{1}{a_0^2} - \frac{1}{a^2} \right) a^2 \left(\frac{a^2 N}{6} \right) + \langle \mathbf{r}^2(s) \rangle_v \langle \mathcal{W}_1 \rangle_v
 \end{aligned} \tag{11}$$

With the help of Eq.(11) and Eq.(9) we obtain the following self-consistent expression for a ,

$$\frac{1}{a_0^2} - \frac{1}{a^2} = \frac{\langle \mathbf{r}^2(s) \rangle_v \langle \mathcal{V} \rangle_v - \langle \mathbf{r}^2(s) \mathcal{V} \rangle_v}{\frac{a^2}{N} \int_0^N ds \langle \mathbf{r}^2(s) \rangle_v}. \tag{12}$$

Calculating the averages in Fourier space, where $\tilde{\mathbf{r}}_n = \frac{1}{N} \int_1^N \cos\left(\frac{\pi n s}{N}\right) \mathbf{r}(s) ds$, $\mathbf{r}(s) = 2 \sum_{n=1}^N \cos\left(\frac{\pi n s}{N}\right) \tilde{\mathbf{r}}_n$, and $R_g^2 = 2 \sum_n \langle |\tilde{\mathbf{r}}_n|^2 \rangle$, we obtain

$$\begin{aligned}
 \frac{1}{a_0^2} - \frac{1}{a^2} &= v \frac{\left(\frac{3}{2}\right)^{5/2} \left(\frac{\pi}{2}\right)^{3/2}}{(a^2)^{5/2} N^{3/2} \left(\sum_{n=1}^N \frac{1}{n^2}\right)} \sum_{s=0}^N \sum_{s'=0}^N \frac{\sum_{n=1}^N \frac{1 - \cos[n\pi(s-s')/N]}{n^4}}{\left(\sum_{n=1}^N \frac{1 - \cos[n\pi(s-s')/N]}{n^2} + \frac{3\pi^2 a_0^2}{2a^2 N}\right)^{5/2}} \\
 &\quad - \kappa \frac{\left(\frac{3}{2}\right)^{5/2} \left(\frac{\pi}{2}\right)^{3/2}}{(a^2)^{5/2} N^{3/2} \left(\sum_{n=1}^N \frac{1}{n^2}\right)} \sum_{\{s_i, s_j\}} \frac{\sum_{n=1}^N \frac{1 - \cos[n\pi(s_i - s_j)/N]}{n^4}}{\left(\sum_{n=1}^N \frac{1 - \cos[n\pi(s_i - s_j)/N]}{n^2} + \frac{3\pi^2 \sigma^2}{2a^2 N}\right)^{5/2}}.
 \end{aligned} \tag{13}$$

The best estimate of the effective monomer size a can be obtained by numerically solving Eq.(13) provided the contact map is known. A bound for the actual size of the chain is $\langle R_g^2 \rangle = Na^2/6$. Because we are interested only in the collapsibility of proteins we use the definition of the θ -point to assess the condition for protein compaction instead of solving the complicated Eq.(13) numerically. The volume interactions are on the right hand side of Eq.(13). At the θ -point, the v -term should exactly balance the κ -term. Since at the θ -point the chain is ideal with $a = a_0$, we can substitute this value for a in the sums in the denominators of the v - and κ -terms. By equating the two, we obtain an expression for κ_θ .

Thus, from Eq.(13), the specific interaction strength at which two-body repulsion (v -term) equals two-body attraction (κ -term) is:

$$\kappa_{\theta} = \frac{4}{3}\pi a_0^3 \frac{\sum_{s=0}^N \sum_{s'=0}^N \frac{\sum_{n=1}^N \frac{1-\cos[n\pi(s-s')/N]}{n^4}}{\left(\sum_{n=1}^N \frac{1-\cos[n\pi(s-s')/N]}{n^2} + \frac{3\pi^2}{2N}\right)^{5/2}}}{\sum_{\{s_i, s_j\}} \frac{\sum_{n=1}^N \frac{1-\cos[n\pi(s_i-s_j)/N]}{n^4}}{\left(\sum_{n=1}^N \frac{1-\cos[n\pi(s_i-s_j)/N]}{n^2} + \frac{3\pi^2\sigma^2}{2a_0^2N}\right)^{5/2}}}. \quad (14)$$

The numerator in Eq.(14) is a consequence of chain connectivity and the denominator encodes protein topology through the contact map, determining the extent to which the sizes in $\mathbf{U}_{\mathbf{D}}$ and $\mathbf{U}_{\mathbf{C}}$ states change as C becomes less than C_m . The numerical value of κ_{θ} is a measure of collapsibility.

RESULTS

Native topology determines collapsibility: The central result in Eq.(14) can be used to quantitatively predict the extent to which a given protein has a propensity to collapse. We used a list of proteins with low mutual sequence identity selected from the Protein Data Bank PDBselect[45], and calculated κ_{θ} using Eq.(14) for these proteins. In all we considered 2306 proteins. For each contact (i, j) , the energetic contribution due to interaction between i and j is $k = (2\pi\sigma^2)^{-3/2}\kappa$ according to Eq.(2). Thus, $k_{\theta} = (2\pi\sigma^2)^{-3/2}\kappa_{\theta}$ is the average strength (in units of $k_B T$) of a contact at the θ -point. If κ_{θ} , calculated using Eq.(14), is too large then the extent of polypeptide chain collapse is expected to be small. It is worth noting that the theory cannot be used to determine the stability of the folded state, because in the Hamiltonian there are only two states, $\mathbf{U}_{\mathbf{D}}$ ($\kappa=0$ in Eq.(2)) and $\mathbf{U}_{\mathbf{C}}$ ($\kappa > \kappa_{\theta}$).

The strength of contacts in real proteins (excluding possibly salt bridges) is typically on the order of few $k_B T$ in the absence of denaturants. This is the upper bound for the contact strength any theory should predict, as adding denaturant only decreases the strength. If k_{θ} is unrealistically high (tens of $k_B T$) then the attractive interactions of the protein would be too weak to counteract the excluded volume repulsion even at zero denaturant concentration, resulting in negligible difference in R_g between the $\mathbf{U}_{\mathbf{D}}$ and $\mathbf{U}_{\mathbf{C}}$ states.

Fig.(1) shows a two-dimensional histogram of the PDBselect proteins in the (N, k_{θ}) plane. For the majority of small proteins (less than 150 residues) the value of κ_{θ} is less than 3

$k_B T$, indicating that the unfolded states of all of these proteins should become compact at $C < C_m$. That collapse must occur, as predicted by our theory and established previously in lattice[26], and off-lattice models of proteins[46], does not necessarily imply that it can be easily detected in standard scattering experiments, because the changes could be small requiring high precision experiments (see below).

Weight function of a contact: For a given N , the criterion for collapsibility in Eq.(14) depends on the architecture of the proteins explicitly represented in the denominator through the contact map. Analysis of the weight function of a contact, defined below, provides a quantitative measure of how a specific contact influences protein compaction. Some contacts may facilitate collapse to a greater extent than others, depending on the location of the pair of residues in the polypeptide chain. In this case, the same number of native contacts N_{nc} in the protein of the same length N might yield a lower (easier collapse) or higher (harder collapse) value of k_θ . In order to determine the relative importance of the contacts with respect to collapse, we consider the contribution of the contact between residues i and j in the denominator of Eq.(14),

$$W(i-j) = \frac{\sum_{n=1}^N \frac{1-\cos[n\pi(i-j)/N]}{n^4}}{\left(\sum_{n=1}^N \frac{1-\cos[n\pi(i-j)/N]}{n^2} + \frac{3\pi^2\sigma^2}{2a_0^2 N} \right)^{5/2}}. \quad (15)$$

A plot of $W(i-j)$ in Fig.(2) for different values of the chain length N shows that the weight depends on the distance between the residues along the chain. Contacts between neighboring residues have negligible weight, and there is a maximum in $W(i-j)$ at $i-j \approx 30$ (for $a_0/\sigma = 0.6$), almost independent of the protein length. The maximum is at a higher value for proteins with $N > 100$ residues. The figure further shows that longer range contacts make greater contribution to chain compaction than short range contacts. The results in Fig.(2) imply that proteins with a large fraction of non-local contacts are more easily collapsible than those dominated by short range contacts, which we elaborate further below.

Maximum and minimum collapsibility boundaries: Using $W(i-j)$ in Eq.(15), we can design protein sequences to optimize for ‘‘collapsibility’’. To design a ‘‘maximally collapsible’’ protein, for fixed N and number of native contacts N_{nc} , we assign each of the N_{nc} contacts one by one to the pair i, j with a maximal $W(i, j)$ among the available pairs with the criterion that $|i-j| > 2$. Such an assignment necessarily implies that the artificially

designed contact map will not correspond to any known protein. Similarly, we can design an artificial contact map by selecting i, j pairs with minimal $W(i, j)$ till all the N_{nc} are fully assigned. Such a map, which will be dominated by local contacts, are minimally collapsible structures.

The white lines in Fig.(1) show k_{θ} of chains of length N with $N_{\text{nc}}(N)$ contacts distributed in ways to maximize or minimize collapsibility. We estimated $N_{\text{nc}}(N) \approx 0.6N^{1.3}$ from the fit of the whole PDBSelect set. Since the lines are calculated for N_{nc} from the fit over the entire set, and not from N_{nc} for every protein, there are proteins below the minimal and above the maximal curve in Fig.(1). For a given protein, with N and N_{nc} defined by its PDB structure, k_{θ} for all possible arrangements of native contacts is largely in between the maximally and minimally collapsible lines in Fig.(1). The majority of proteins in our set are closer to the maximal collapsible curves, suggesting that the unfolded proteins have evolved to be compact under native folding conditions. This theoretical prediction is in accord with our earlier studies which suggested that foldability is determined by both collapse and folding transitions [26], and more recently supported by experiments[20].

β -sheet rather than α -helical proteins undergo larger compaction: The weight function W (Eq.(15) and Fig.(2)) suggests that contacts in α -helices ($|i - j| = 4$) only make a small contribution to collapse. Contacts corresponding to the maximum of W at $i - j \approx 30$ are typically found in loops and long antiparallel β -sheets. Fig.(3) shows a set of proteins with high α -helix ($> 90\%$) and a set with high content of β -sheets ($> 70\%$). The values of k_{θ} for the two sets are very distinct, so they barely overlap. We find that many of the α -helical proteins lie on or above the curve of minimal collapsibility while the rest are closer to the maximal collapsibility. The smaller β -rich proteins lie on the curve of maximal collapsibility slightly diverging from it as the chain length grows. These results show that the extent of collapse of proteins that are mostly α -helical is much less than those with predominantly β -sheet structures.

A note of caution is in order. The minimal collapsibility of most α -helical proteins in the set may be a consequence of some of them being transmembrane proteins, which do not fold in the same manner as globular proteins. Instead, the transmembrane α -helices are inserted into the membrane by the translocon, one by one, as they are synthesized. Such proteins would not have the evolutionary pressure to be compact.

Extent of compaction varies depending on the native structure: Our conclusions,

summarized in Figs.(1-3), are based on an approximate theory. To validate the approximations and illustrate the impact of the location of contacts on chain compaction we performed Monte Carlo simulations of chains of lengths $N = 36, 72$ and 153 using the energy function given by Eq.(2) with harmonic spring chain connectivity. The proteins are Villin headpiece ($N = 36$), protein-L ($N = 72$), and myoglobin ($N = 153$). The contact maps and cartoon representation of these proteins are in Fig.(4). Fig.(5) shows that the ratio between end-to-end distance (R_{ee}) and radius of gyration (R_g) decreases continuously with the interaction strength, indicative of chain compaction. Fig.(5) also shows that $R_{ee}/R_g \gg 1$, which is well known for Gaussian chains ($R_{ee}/R_g = \sqrt{6}$) and self-avoiding walks ($\kappa = 0$ limit) for which the ratio $> \sqrt{6}$. Based on κ_θ estimates we conclude that protein-L (α, β) is more collapsible than villin headpiece and myoglobin (helical proteins). Fig.(6a) shows the variations in the probability distribution of R_g , $P(R_g)$ for protein-L as a function of κ . The broader distribution with zero interaction strength corresponds to the extended chain. We find that $P(R_g)$ becomes narrower as the attractive strength increases. The continuous shift in to the compact state with gradual increase in the attractive strength shows the compaction of unfolded proteins as the denaturant concentration decreases. Thus, generally R_g of the \mathbf{U}_C state is less than that of the \mathbf{U}_D state. The end-to-end distribution, $P(R_{ee})$, for different values of interaction strength has been shown in Fig.(6b). The broad distribution at $\kappa = 0$ corresponds to the unfolded protein. Average R_{ee} decreases as attractive strength increases and the distribution becomes narrower. The results in Fig.(6) show that both R_{ee} , which can be inferred using FRET, and R_g (measurable using SAXS), are smaller in the \mathbf{U}_C state than the \mathbf{U}_D state. However, the extent of decrease is greater in R_{ee} than R_g .

RNAs are compact: There are major differences between how RNA and proteins fold [47]. In contrast to the apparent controversy in proteins, it is well established that RNA molecules are compact[48–50] at high ion concentrations or at low temperatures. Because our theory relies only on the knowledge of contact map we used it assess collapsibility in *Azoarcus* ribozyme and MMTV pseudoknot to merely illustrate collapsibility of RNA . The k_θ values (green stars in Fig.3) are close to the lower β -sheet line, indicating that these molecules must undergo compaction. This prediction from the theory is fully supported by both equilibrium and time-resolved SAXS experiments[51] on *Azoarcus* ribozyme. In this case ($N = 196$) the changes are so large that even using low resolution experiments collapse is readily observed[52]. We should emphasize that the size of different RNAs (for example

viral, coding, non-coding) vary greatly. For a fixed length, single-stranded viral RNAs have evolved to be maximally compact, which is rationalized in terms of the density of branching. Although the sizes of the viral RNAs considered in [53] are much longer than the *Azoarcus* ribozyme the notion that compaction is determined by the density of branching might be valid even when $N \sim 200$.

DISCUSSION

We have shown that polymer chains with specific interactions, like proteins (but ones without a unique native state), become compact, as the strength of the specific interaction changes. A clear implication is that the size of the \mathbf{U}_D state should decrease continuously as C decreases. In other words, the unfolded state under folding conditions is more compact than it is at high denaturant concentrations. Compaction is driven roughly by the same mechanism as the collapse transition in homopolymers in the sense that when the solvent quality is poor (below C_m) the size of the unfolded state decreases continuously. When the set of specific interactions is taken from protein native contacts in the PDB, our theory shows that the values of k_θ are in the range expected for interaction between amino acids in proteins. This implies that collapsibility should be a universal feature of foldable proteins but the extent of compaction varies greatly depending on the architecture in the folded state. This is manifested in our finding that proteins dominated by β -sheets are more collapsible compared to those with α -helical structures.

Magnitude of k_θ and plausible route to multi-domain formation: The scaling of k_θ with N allows us to provide arguments for the emergence of multi-domain proteins. In Eqs.(13) or (14) attractive (κ -) and repulsive (v -) terms have the same structure. The only difference in their scaling with N is due to the difference in the sums (over all the monomers in the repulsive term and over native contacts in the attractive term). Double summation over all the monomers gives a factor of N^2 to the repulsive term. The summation over native contacts in the attractive term scales as N_{nc} . Therefore, to compensate for the repulsion, N_{nc} should scale as N^2 . However, for a given protein with a certain length N and certain numbers of contacts, it is not clear how the denominator in Eq.(14) scales with N . Empirically we find $N_{nc}(N)$ dependence across a representative set of sequences scales as N^γ with γ less than 2 (likely on the order of 1.5). Thus, it follows from Eq.(14) that k_θ

increases without bound as N continues to increase. Because this is unphysical, it would imply that proteins whose lengths exceeds a threshold length N_C cannot become maximally compact even at $C = 0$. An instability must ensue when N exceeds N_C . This argument in part explains why single domain proteins are relatively small[54].

Scaling of N_{nc} as a power law in N^γ with $\gamma < 2$ means that as protein size grows, the value of k_θ will deviate more and more from those found in globular proteins, implying such proteins cannot be globally compact under physiologically relevant conditions. However, such an instability is not a problem because larger proteins typically consist of multiple domains. Thus, if the protein does not show collapse as a whole, the individual domains could fold independently, having lower values of k_θ for each domain of the multi-domain protein. It would be interesting to know if the predicted onset of instability at N_C provides a quantitative way to assess the mechanism of formation of multi-domain proteins. Extension of the theory might yield interesting patterns in the assembly of multi-domain proteins. For instance, one can quantitatively ascertain if the N-terminal domains of large proteins, which emerge from the ribosome first, have higher collapsibility (lower κ_θ) than C-terminal domains.

SAXS-smFRET controversy resolved: Our theory resolves, at least theoretically, the contradictory results using SAXS and FRET experiments on compaction of small globular proteins. It has been argued, based predominantly on protein-L ($N = 72$) SAXS experiments that R_g of \mathbf{U}_D and \mathbf{U}_C states are virtually the same at denaturant concentrations that are less than C_m [19]. This conclusion is not only at variance with SAXS experiments on other proteins but also with interpretation of smFRET data on a number of proteins. The present work, surveying over 2300 proteins, shows that the compact state has to exist, engendered by mechanisms that have much in common with homopolymer collapse. For protein-L, the $k_\theta = 1.7k_B T$, a very typical value, right on the peak of the heat map in Fig.(1). We have previously argued that because the change in R_g between the \mathbf{U}_D and \mathbf{U}_C states for small proteins is not large, high precision experiments are needed to measure the predicted changes in R_g between \mathbf{U}_C and \mathbf{U}_D . For protein-L the change is less than 10% [55], making its detection in ensemble experiments very difficult. Similar conclusions were reached in recent experiments [20]. A clear message from our theory is that, tempting as it may be, one cannot draw universal conclusions about polypeptide compaction by performing experiments on just a few proteins. One has to survey a large number of proteins with varying N and native

topology to quantitatively assess the extent of compaction. Our theory provides a framework for interpreting the results of such experiments.

Random contact maps, local and non-local contacts: In order to differentiate collapsibility between evolved and random proteins, we created twelve random contact maps keeping the total number of contacts the same as in protein-L (see Fig.(8) for examples). For each of these pseudo-proteins we calculated k_θ using Eq.(14). We find that for all the random contact maps the k_θ values are less than for protein-L, implying that the propensity of the pseudo-proteins to become compact is greater than for the wild type. This finding is in accord with studies based on homopolymer and heteropolymer collapse with random crosslinks. These studies showed that the polymer undergoes a collapse transition as the density of crosslinks is increased[34, 36, 37]. Of particular note is the demonstration by Camacho and Schanke[39], who showed using exact enumeration of random heteropolymers and scaling arrangements that the collapse can be either a first or second order transition depending on the function of hydrophobic residues [39].

Some time ago Abkevich et al.[56] showed, using Monte Carlo simulations of protein-like lattice polymers, that the folding transition in proteins with predominantly non-local contacts was first order like, which is not the case for proteins in which local contacts dominate. In light of this finding, it is interesting to examine how compaction is affected by local and non-local contacts. We created for $N=72$ (protein-L) a contact map with 185 (same number as with WT protein-L), predominantly local contacts (Fig.(8b)). The values of k_θ for these pseudo-proteins is considerably larger than for the WT, implying that proteins dominated by local contacts are minimally collapsible. We repeated the exercise by creating contact maps with predominantly non-local contacts (Fig.(8c)). Interestingly, k_θ values in this case are significantly less than for the WT. This finding explains why in proteins with varied α/β topology there is a balance between the number of local and non-local contacts. Such a balance is needed to achieve native state stability and speed of folding[56] with polypeptide compaction playing an integral part [26].

Based on these findings we conclude that R_g of the unfolded states of proteins dominated by non-local contacts must undergo greater compaction compared to those with that have mostly local contacts. The results in Fig. 3 also show that proteins rich in β -sheet are more collapsible than predominantly α -helical proteins. It follows that β -sheet proteins must have a larger fraction of non-local contacts than proteins rich in α -helices. In Fig. (8d) we plot

the distribution of the fraction of non-local contacts for the 2306 proteins. Interestingly, there is a clear separation in the distribution of non-local contacts between α -helical rich and β -sheet rich proteins. The latter have substantial fraction of non-local contacts which readily explains the findings in Fig. (8c) and the predictions in Fig. (3).

CONCLUSIONS

We have created a theory to assess collapsibility of proteins. The major implications of the theory are the following. (i) Because single domain proteins are small, the changes in the radius of gyration of the unfolded states as the denaturant concentration is lowered are often small. Thus, it has been difficult to detect the changes in SAXS experiments in a couple of proteins raising the question if unfolded polypeptide chains become compact below C_m . Here, we have solved this long-standing problem showing that the unfolded states of single-domain proteins do become compact as the denaturant concentration decreases, sharing much in common with the physical mechanisms governing homopolymer collapse. By adopting concepts from polymer physics, and using the contact maps that reflect the topology of the native states, we established that proteins are collapsible. (ii) Based on a survey of over two thousand proteins we surmise that there is evolutionary pressure for collapsibility is universal although the extent of collapse can vary greatly, because this ensures that the propensity to aggregate is minimized even if environmental fluctuations under cellular conditions transiently populate unfolded states. Two factors contribute to aggregation. First, the rate of dimer formation by diffusion controlled reaction would be enhanced if a pair of \mathbf{U}_D rather than \mathbf{U}_C molecules collided due cellular stress because the contact radius in the former would be greater than in the latter. Second, the fraction of exposed hydrophobic residues in \mathbf{U}_D is much greater than in \mathbf{U}_C , thus greatly increasing the probability of aggregation. The second factor is likely to be more important than the first. Consequently, transient population of \mathbf{U}_C due to cellular stress minimizes the probability of aggregation. (iii) We have also shown that the position of the residues forming the native contact greatly influences the collapsibility of β sheet proteins (containing a number of non-local contacts) showing greater compaction than α helical proteins, which are typically stabilized by local contacts.

Our theory also shows that most RNAs may have evolved to be compact in their natural

environments. Although the evolutionary pressure to be compact is likely to be substantial for viral RNAs [49, 50, 53, 57], it is apparent that even non-coding RNAs are also likely to be almost maximally compact in their natural environments. Our theory suggests that to a large extent collapsibility of RNA is similar to proteins with β -sheet structures. Both classes of biological macromolecules are stabilized by non-local contacts. Interestingly, it has been argued that the need to be compact (“Compaction selection hypothesis” [57]) could be a major determinant for evolved biopolymers to have minimum energy compact structures as their ground states.

-
- [1] F. M. Richards. The interpretation of protein structures: Total volume, group volume distributions and packing density. *J. Mol. Biol.*, 82:1–14, 1974.
 - [2] F. M. Richards and W. A. Lim. An analysis of packing in the protein-folding problem. *Q. Rev. Biophys.*, 26:423–498, 1994.
 - [3] J. L. Finney. Volume occupation, environment and accessibility in proteins. the problem of the protein surface. *J. Mol. Biol.*, 96:721–732, 1975.
 - [4] W. A. Lim and R. T. Sauer. Alternative packing arrangements in the hydrophobic core of λ -repressor. *Nature*, 339:31–36, 1989.
 - [5] J. Liang and K. A. Dill. Are proteins well-packed? *Biophys. J.*, 81:751–756, 2001.
 - [6] S. Bromberg and K. A. Dill. Side-chain entropy and packing in proteins. *Protein Sci.*, 3(7):997–1009, 1994.
 - [7] R. I. Dima and D. Thirumalai. Asymmetry in the shapes of folded and denatured states of proteins. *J. Phys. Chem. B*, 108:6564–6570, 2004.
 - [8] J. E. Kohn, I. S. Millett, J. Jacob, B. Azgrovic, T. M. Dillon, N. Cingel, R. S. Dothager, S. Seifert, P. Thiyagarajan, T. R. Sosnick, M. Z. Hasan an V. S. Pande, I. Ruczinski, S. Doniach, and K. W. Plaxco. Random-coil behavior and the dimensions of chemically unfolded proteins. *Proc. Natl. Acad. Sci. USA*, 101:12491–12496, 2004.
 - [9] E. Sherman and G. Haran. Coil-globule transition in the denatured state of a small protein. *Proc. Natl. Acad. Sci.*, 103:11539–11543, 2006.
 - [10] G. Haran. How, when and why proteins collapse: the relation to folding. *Curr. Opin. Struct. Biol.*, 22:14–20, 2012.

- [11] I. M. Lifshitz, A. Yu. Grosberg, and A. R. Khokhlov. Some problems of statistical physics of polymer-chains with volume interaction. *Rev. Mod. Phys.*, 50:683–713, 1978.
- [12] A. Yu. Grosberg and A. R. Khokhlov. *Statistical Physics of Macromolecules*. AIP Press, 1994.
- [13] O. B. Ptitsyn. How molten is the molten globule? *Nat. Struct. Biol.*, 3:488–490, 1996.
- [14] D. Thirumalai. From minimal models to real proteins: Time scales for protein folding kinetics. *J. Phys. I (Fr.)*, 5:1457–1467, 1995.
- [15] O. B. Ptitsyn and V. N. Uversky. The molten globule is a third thermodynamical state of protein molecules. *FEBS Lett.*, 341(1):15–18, 1994.
- [16] F. Ding, W. Guo, N. V. Dokholyan, E. I. Shakhnovich, and J. E. Shea. Reconstruction of the src-SH3 protein domain transition state ensemble using multiscale molecular dynamics simulations. *J. Mol. Biol.*, 350:1035–1050, 2005.
- [17] H. T. Tran, X. Wang, and R. V. Pappu. Reconciling observations of sequence-specific conformational propensities with the generic polymeric behavior of denatured proteins. *Biochemistry*, 44, 2005.
- [18] G. Ziv, D. Thirumalai, and G. Haran. Collapse transition in proteins. *Phys. Chem. Chem. Phys.*, 11(1):83–93, 2009.
- [19] T. Y. Yoo, S. P. Meisburger, J. Hinshaw, L. Pollack, G. Haran, T. R. Sosnick, and K. Plaxco. Small-angle X-ray scattering and single-molecule FRET spectroscopy produce highly divergent views of the low-denaturant unfolded state. *J. Mol. Biol.*, 418:226–236, 2012.
- [20] A. Borgis, W. Zheng, K. Buholzer, M. B. Borgia, A. Schüler, H. Hofmann, A. Soranno, D. Nettels, K. Gast, A. Grishaev, R. B. Best, and B. Schuler. Consistent view of polypeptide chain expansion in chemical denaturants from multiple experimental methods. *Submitted to J. Am. Chem. Soc.*, 2016.
- [21] B. Schuler and W. A. Eaton. Protein folding studied by single-molecule FRET. *Curr. Opin. Struct. Biol.*, 18(1):16–26, 2008.
- [22] P. G. De Gennes. Kinetics of collapse for a flexible coil. *J. Phys. Lett. (Fr.)*, 46(14):639–642, 1985.
- [23] A. Y. Grosberg, S. K. Nechaev, and E. I. Shakhnovich. The role of topological constraints in the kinetics of collapse of macromolecules. *J. de Physique*, 49:2095–2100, 1988.
- [24] C. J. Camacho and D. Thirumalai. Minimum energy compact structures of random sequences of heteropolymers. *Phys. Rev. Lett.*, 71:2505, 1993.

- [25] Z. Liu, G. Reddy, and D. Thirumalai. Folding PDZ2 domain using the molecular transfer model. *J. Phys. Chem. B*, 2016. in press.
- [26] C. J. Camacho and D. Thirumalai. Kinetics and thermodynamics of folding in model proteins. *Proc. Natl. Acad. Sci. USA*, 90:6369–6372, 1993.
- [27] M. S. Li, D. K. Klimov, and D. Thirumalai. Finite size effects on thermal denaturation of globular proteins. *Phys. Rev. Lett.*, 93:268107, 2004.
- [28] K. A. Dill and D. Shortle. Denatured states of proteins. *Ann. Rev. Biochem.*, 60(1):795–825, 1991.
- [29] S. Akiyama, S. Takahashi, T. Kimura, K. Ishimori, I. Morishima, Y. Nishikawa, and T. Fujisawa. Conformational landscape of cytochrome c folding studied by microsecond-resolved small-angle X-ray scattering. *Proc. Natl. Acad. Sci. USA*, 99(3):1329–1334, 2002.
- [30] T. Kimura, S. Akiyama, T. Uzawa, K. Ishimori, I. Morishima, T. Fujisawa, and S. Takahashi. Specifically collapsed intermediate in the early stage of the folding of ribonuclease A. *J. Mol. Biol.*, 350(2):349–362, 2005.
- [31] R. R. Goluguri and J. B. Udgaonkar. Microsecond rearrangements of hydrophobic clusters in an initially collapsed globule prime structure formation during the folding of a small protein. *J. Mol. Biol.*, 428(15):3102–3117, 2016.
- [32] S. F. Edwards. The statistical mechanics of polymers with excluded volume. *Proc. Phys. Soc.*, 85(4):613–624, 1965.
- [33] A. M. Gutin and E. I. Shakhnovich. Statistical mechanics of polymers with distance constraints. *J. Chem. Phys.*, 100:5920, 1994.
- [34] J. D. Bryngelson and D. Thirumalai. Internal constraints induce localization in an isolated polymer molecule. *Phys. Rev. Lett.*, 76:542, 1996.
- [35] D. Thirumalai, V. Ashwin, and J. K. Bhattacharjee. Dynamics of random hydrophobic-hydrophilic copolymers with implications for protein folding. *Phys. Rev. Lett.*, 77:5385, 1996.
- [36] Y. Kantor and M. Kardar. Collapse of randomly linked polymers. *Phys. Rev. Lett.*, 77:4275, 1996.
- [37] R. Zwanzig. Effect of close contacts on the radius of gyration of a polymer. *J. Chem. Phys.*, 106:2824, 1997.
- [38] C. J. Camacho and D. Thirumalai. A criterion that determines fast folding of proteins: A model study. *Europhys. Lett.*, 35:627–632, 1996.

- [39] C. J. Camacho and T. Schanke. From collapse to freezing in random heteropolymers. *Europhys. Lett.*, 37(9):603, 1997.
- [40] R. T. Deam and S. F. Edwards. The theory of rubber elasticity. *Phil. Trans. R. Soc. A*, 280(1296):317–353, 1976.
- [41] P. Goldbart and N. Goldenfeld. Rigidity and ergodicity of randomly cross-linked macromolecules. *Phys. Rev. Lett.*, 58(25):2676, 1987.
- [42] P. Goldbart and N. Goldenfeld. Microscopic theory for cross-linked macromolecules. I. Broken symmetry, rigidity, and topology. *Phys. Rev. A*, 39(3):1402, 1989.
- [43] H. E. Castillo, P. M. Goldbart, and A. Zippelius. Distribution of localisation lengths in randomly crosslinked macromolecular networks. *Europhys. Lett.*, 28:519, 1994.
- [44] S. F. Edwards and P. Singh. Size of a polymer molecule in solution. Part 1. Excluded volume problem. *J. Chem. Soc. Faraday Trans.*, 75:1001–1019, 1979.
- [45] S. Griep and U. Hobohm. PDBselect 1992–2009 and PDBfilter-select. *Nucl. Acids Res.*, 38:D318–D319, 2009.
- [46] J. D. Honeycutt and D. Thirumalai. The nature of folded states of globular proteins. *Biopolymers*, 32:695–709, 1992.
- [47] D. Thirumalai and C. Hyeon. RNA and protein folding: Common themes and variations. *Biochemistry*, 44(13):4957–4970, 2005.
- [48] C. Hyeon, R. I. Dima, and D. Thirumalai. Size, shape and flexibility of RNA structures. *J. Chem. Phys.*, 125:194905, 2006.
- [49] A. M. Yoffe, P. Prinsen, A. Gopal, C. M. Knobler, W. M. Gelbart, and A. Ben-Shaul. Predicting the sizes of large RNA molecules. *Proc. Natl. Acad. Sci. USA*, 105(42):16153, 2008.
- [50] L. T. Fang, W. M. Gelbart, and A. Ben-Shaul. The size of RNA as an ideal branched polymer. *J. Chem. Phys.*, 135(15):155105, 2011.
- [51] J. H. Roh, L. Guo, J. D. Kilburn, R. M. Briber, T. Irving, and S. A. Woodson. Multistage Collapse of a Bacterial Ribozyme Observed by Time-Resolved Small-Angle X-ray Scattering. *J. Am. Chem. Soc.*, 132(29):10148–10154, 2010.
- [52] R. Behrouzi, J. H. Roh, D. Kilburn, R. M. Briber, and S. A. Woodson. Cooperative tertiary interaction network guides RNA folding. *Cell*, 149:348–357, 2012.
- [53] A. Gopal, D. E. Egecioglu, A. M. Yoffe, A. Ben-Shaul, A. L. N. Rao, C. M. Knobler, and W. M. Gelbart. Viral RNAs are unusually compact. *PLoS One*, 9(9):e105875, 2014.

- [54] H. J. Bussemaker, D. Thirumalai, and J. K. Bhattacharjee. Thermodynamic stability of folded proteins against mutations. *Phys. Rev. Lett.*, 79:3530–3533, 1997.
- [55] H. Maity and G. Reddy. Folding of protein l with implications for collapse in the denatured state ensemble. *J. Am. Chem. Soc.*, 2016.
- [56] V. I. Abkevich, A. M. Gutin, and E. I. Shakhnovich. Impact of local and non-local interactions on thermodynamics and kinetics of protein folding. *J. Mol. Biol.*, 252(4):460–471, 1995.
- [57] L. Tubiana, A. L. Božič, C. Micheletti, and R. Podgornik. Synonymous mutations reduce genome compactness in icosahedral ssRNA viruses. *Biophys. J.*, 108(1):194–202, 2015.
- [58] W. Kabsch and C. Sander. Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers*, 22(12):2577–2637, 1983.
- [59] W. Humphrey, A. Dalke, and K. Schulten. VMD – Visual Molecular Dynamics. *J. Mol. Graph.*, 14:33–38, 1996.
- [60] C. Hyeon and D. Thirumalai. Mechanical unfolding of RNA: From hairpins to structures with internal multiloops. *Biophys. J.*, 92(3):731–743, 2007.




FIG. 1. Collapsibility quantified using k_θ (in units of $k_B T$) for a set of 2,306 PDB structures as a function of the length N of the proteins. White lines show the k_θ at the boundaries for maximally and minimally collapsible proteins (lower and upper lines respectively). Colors give a rough estimate of the number of proteins, which decreases from red to violet.




FIG. 2. Weight function W (Eq.(15)) of a contact, showing how much a contact between residues i and j contributes to the compaction of a protein. The colors are for different N values (shown in the inset). Interestingly, the location of the maximum is roughly independent of N .




FIG. 3. Dependence of k_θ on the secondary structure content of proteins. We display k_θ for α -rich ($> 90\%$) and β -rich ($> 70\%$) proteins. Proteins that are predominantly α -helical tend to be close to minimally collapsible (upper line), while β -rich proteins are close to maximally collapsible curve (lower curve). The green stars are for RNA with the left one at small N corresponding to the *Mouse Mammary Tumor Virus* (MMTV) pseudoknot ($N=34$) and the other is *Azoarcus* ribozyme ($N=196$).




FIG. 4. Native topologies of three proteins studied by Monte Carlo simulations; villin headpiece (a), protein-L (b), and myoglobin (c). Secondary structures were determined by DSSP [58] and shown in blue for helices and red for β -strands. We used VMD to draw the three-dimensional structures [59].




FIG. 5. Ratio between the end-end distance (R_{ee}) and radius of gyration (R_g) for three different chain lengths ($N=36, 72$ and 153). It decreases continuously with increasing interaction strength. The presence of contacts between residues decreases R_{ee}/R_g from the maximum value attainable for a self-avoiding walk.




FIG. 6. (a) The probability distribution of the radius of gyration, $P(R_g)$ for different values of interaction strength k for Protein-L. As k increases, the distribution becomes narrower. (b) Same as (a) except this panel shows end-to-end distribution $P(R_{ee})$ for different values of attractive strength k for Protein-L. The similarity between $P(R_{ee})$ and $P(R_g)$ shows that R_{ee} also is a reasonable measure of compaction.




FIG. 7. Native topologies of two RNA molecules, MMTV pseudoknot (a-c) and *Azoarcus* ribozyme (d-f). Three-dimensional structures (a and d), secondary structures (b and e), and contact maps (c and f) are shown for each RNA. Colors are used to distinguish secondary structures. Contact pairs in RNA are defined as any nucleotide pair i and j ($|i - j| > 2$) satisfying $R_{ij} < 14\text{\AA}$, where R_{ij} is the distance between centers of mass of the nucleotides [60]. MMTV has two stem basepairs (cyan and green in a-c), which contribute to non-local contacts (cyan and green in c). *Azoarcus* ribozyme has several hairpin basepairs (P2, P5, P6, P8 and P9) which can be seen in the vicinity of the diagonal in the contact map (f). There are also basepairs between nucleotides far along the sequence such as P3, P4 and P7, as well as tertiary interactions such as TL2-TR8 and TL9-TR5. These non-local contacts contribute to the collapsibility of the ribozyme.




FIG. 8. Collapsibility for synthetic contact maps. Two representative contact maps for each category are shown in the upper left and lower right of each square. Given the number of residues $N = 72$ and total number of contacts $N_{nc} = 185$ (same as protein-L), residue pairs (i, j) are randomly chosen to satisfy the following conditions: (a) uniformly distributed, $|i - j| \geq 3$; (b) local contacts only, $|i - j| \geq 3$ and $|i - j| < 8$; and (c) non-local contacts only, $|i - j| \geq 8$. The calculated values of k_θ are explicitly shown. The k_θ value for protein-L is $1.7k_B T$. (d) Distribution of the fraction of non-local contacts in the 2306 proteins. For each protein, the fraction is calculated as the number of non-local contacts (N_{nc}^{NL}) divided by the total number of contacts (N_{nc}). A contact between residues i and j is “non-local (NL)” if $|i - j| \geq 8$. There is a clear separation in this distribution for proteins rich in α helices compared to those that are rich in β -sheets implying that the latter are more collapsible than the former.