

# The Co-regulation Data Harvester for *Tetrahymena thermophila*: automated high-throughput gene annotation and functional inference in a microbial eukaryote

Lev M. Tsypin, Aaron P. Turkewitz\*

*Department of Molecular Genetics and Cell Biology, University of Chicago, Chicago IL, 60637*

---

## Abstract

Identifying co-regulated genes can provide a useful approach for defining pathway-specific machinery in an organism. To be efficient, this approach relies on thorough genome annotation, which is not available for most organisms with sequenced genomes. Studies in *Tetrahymena thermophila*, the most experimentally accessible ciliate, have generated a rich transcriptomic database covering many well-defined physiological states. Genes that are involved in the same pathway show significant co-regulation, and screens based on gene co-regulation have identified novel factors in specific pathways, for example in membrane trafficking. However, a limitation has been the relatively sparse annotation of the *Tetrahymena* genome, making it impractical to approach genome-wide analyses. We have therefore developed an efficient approach to analyze both co-regulation and gene annotation, called the Co-regulation Data Harvester (CDH). The CDH automates identification of co-regulated genes by accessing the *Tetrahymena* transcriptome database, determines their orthologs in other organisms via reciprocal BLAST searches, and collates the annotations of those orthologs' functions. Inferences drawn from the CDH reproduce and expand upon experimental findings in *Tetrahymena*. The CDH, which is freely available, represents a powerful new tool for analyzing cell biological pathways in *Tetrahymena*. Moreover, to the extent that genes and pathways are conserved between organisms, the inferences obtained via the CDH should be relevant, and can be explored, in many other systems.

---

\*Corresponding Author

*Email address:* [apturkew@uchicago.edu](mailto:apturkew@uchicago.edu) (Aaron P. Turkewitz)

*Keywords:* Co-expression, Co-regulation, Transcriptomics, Genomics, Bioinformatics, Tetrahymena

---

## 1. Motivation and significance

*Tetrahymena thermophila* is a ciliate, one of the best-studied members of this large group of protists [1]. Its use as a model system led to the Nobel Prize-winning discoveries of telomerase and self-splicing RNA, as well as to other breakthroughs, including the isolation of dyneins and making the link between histone modification and transcriptional regulation [2, 3, 4, 5]. These contributions to our understanding of important cellular pathways made use of classical forward and reverse genetics, as well as biochemical approaches. More recently, genomic and transcriptomic data became available for *T. thermophila*, which have been used to infer functional gene networks [6, 7, 8, 9, 10, 11].

The *T. thermophila* genome has been sequenced and assembled [6], and is available online on the *Tetrahymena* Genome Database (TGD) [7]. While the TGD collates the sequence data along with available gene annotations and descriptions, the genome overall remains incompletely annotated. An extensive transcriptomic database, the *Tetrahymena* Functional Genomics Database or *TetraFGD*, is also available for *T. thermophila* [10]. These data were collected over a well-established range of culture conditions in which *T. thermophila* undergoes large physiological changes [8, 9, 10, 11]. In addition to displaying individual expression profiles, the *TetraFGD* can indicate the statistical strength of co-expression between any two genes, as calculated using the Context Likelihood of Relatedness (CLR) algorithm [9, 12, 13]. Co-expression in *T. thermophila*, as judged based on mRNA levels, can reveal functionally significant co-regulation. Gene regulation in this species appears to predominantly occur at the level of transcription [14], and so steady-state mRNA levels may explain the majority of steady-state protein levels, as reported in other systems [15]. In this report, we will refer to genes that are listed as co-expressed in the *TetraFGD* as co-regulated.

A high-throughput analysis of *T. thermophila* gene expression profiles revealed that accurate gene networks can be inferred from co-regulation data [13], providing evidence that co-regulated genes tend to be functionally associated. This approach has been used in bacterial, mammalian, and apicomplexan systems [12, 16, 17]. There is also experimental evidence in *T. thermophila* to support the conclusion that co-regulation corresponds to functional association: Co-regulation data were used to successfully predict novel sorting factors and proteases involved in the biosynthesis of a class of

37 secretory vesicles, called mucocysts [18, 19]. These results suggest that the  
38 *T. thermophila* transcriptome may be used to bioinformatically infer factors  
39 involved in an array of cellular pathways.

40 The CDH was designed to facilitate genome-wide analyses of gene co-  
41 regulation. The CDH automatically mines co-regulation data for genes of  
42 interest, and annotates the co-regulated genes *via* forward and reciprocal  
43 BLAST searches that identify orthologs in other model organisms. The CDH  
44 provides a systematic tool for gathering and annotating genomic information  
45 from public databases, and it can allow a researcher to quickly develop a  
46 robust hypothesis about the cellular pathways or structures in which a gene  
47 of interest may be acting, based upon the genes with which it is co-regulated.

## 48 2. Software description

### 49 2.1. Software Architecture

50 The CDH was developed for Python 2.7, along with the following pack-  
51 ages: `sys`, `os`, `platform`, `logging`, `re`, `dill`, `difflib`, `csv`, `pdb`, `shutil`,  
52 `xml`, `win32com.shell`, `requests`, `BeautifulSoup4`, and `Biopython` [21].  
53 Executables for Windows (x64) and MacOS (10.6+) were made using the  
54 `Pyinstaller` library. The CDH gathers available data for a set of co-regulated  
55 genes from publicly available databases, and uses these data to predict pos-  
56 sible gene functions (Figure 1). The gathered information includes the co-  
57 regulation data from the *TetraFGD*, and the gene names, sequences, and  
58 annotations from the TGD. The available annotations come from a combi-  
59 nation of experimental results and inferences from homology [7]. The CDH  
60 predicts annotations for genes based on the annotation of their respective  
61 orthologs, which are themselves identified by a series of forward and reciprocal  
62 BLAST searches *via* the National Center for Biotechnology Information  
63 (NCBI). These predicted annotations are generated by using the Ratcliff-  
64 Obershelp algorithm [22], as implemented in the `python difflib` library, to  
65 identify common phrases in the orthologs' annotations.

### 66 2.2. Software Functionalities

67 The basic functions of the CDH are to gather available co-regulation and  
68 annotation data, perform forward and reciprocal BLAST searches and predict  
69 gene annotations, and report this gathered information in a human-readable  
70 format. The CDH interface first asks the user to enter the ID for the gene  
71 whose co-regulated factors are of interest (Figure 2). Next, the user defines:  
72 how many of the co-regulated genes should be interrogated *via* BLAST; how  
73 to process data files that had been previously generated and are relevant  
74 to the current query; whether to use the BLASTp or BLASTx algorithm;

75 and in which taxa to run the forward BLAST searches. The results of the  
76 CDH analysis are saved as a Comma Separated Values (.csv) file in the user's  
77 "Documents" folder: /Documents/CoregulationDataHarvester/csvFiles.

### 78 3. Illustrative Examples

#### 79 3.1. A CDH analysis of a factor required for programmed genome rearrange- 80 ment returns the vast majority of experimentally-verified genes involved 81 in the pathway

82 Programmed genome rearrangement is a tightly regulated process that oc-  
83 curs during the formation of the new somatic nucleus in conjugating *Tetrahy-*  
84 *mena* [23, 24]. This process is well-studied and known to be driven by a spe-  
85 cial adaptation of RNA interference, utilizing Dicer- and Piwi-like proteins,  
86 among other factors [25, 26]. *TWI1* encodes a Piwi-like protein that plays  
87 a central role in programmed genome rearrangement [27, 26]. When *TWI1*  
88 is entered as the query for the CDH, the CDH retrieves a large number of  
89 DNA and RNA-processing factors, as well as chromodomain proteins (Sup-  
90 plementary File 1). Importantly, these include the key factors known to be  
91 involved in programmed genome rearrangement (Table 1). The CDH report  
92 for this *TWI1* query is attached as Supplementary File 2. Within this report,  
93 we have highlighted the cases in which the CDH matched or expanded upon  
94 existing annotations.

95 It is also notable that specific homologs of Dicer that are not involved in  
96 programmed genome rearrangement, namely *DCR1* and *DCR2* [26], are not  
97 present in the list of genes co-regulated with *TWI1*. Similarly, while *TPB2*  
98 is a known genome rearrangement factor and is present in the CDH output  
99 [26], its paralog *TPB1* is neither involved in this process nor identified as  
100 co-regulated with *TWI1*. Thus, the CDH is a useful tool for focusing on  
101 pathway-specific paralogs within gene families.

#### 102 3.2. A CDH analysis of a mucocyst biogenesis factor enriches for mucocyst 103 cargo and maturation factors

104 Mucocysts in *Tetrahymena* are secretory organelles. Mucocysts undergo  
105 a maturation process that requires the catalyzed cleavage of cargo proteins,  
106 called GRLs [28]. The *T. thermophila* genome encodes approximately 480  
107 predicted proteases [6], but only five of these are co-regulated with GRLs, as  
108 revealed by a manual inspection of expression profiles on the *TetraFGD* [19].  
109 Two of these proteases, called *CTH3* (cathepsin 3) and *CTH4* (cathepsin 4),  
110 were subsequently shown to represent key enzymes for GRL cleavage [19, 29].

111 Using *CTH3* as a query for the CDH results in a list that includes a  
112 large number of genes known to be involved in mucocyst biogenesis (Table

113 2), and is enriched in membrane-trafficking factors and proteins with as-  
114 yet unknown functions in this organism (Supplementary File 3). Among  
115 the latter are a subunit of the *AP3* complex and a syntaxin in the *STX7*  
116 subfamily. Subsequent functional analysis of these genes showed that they  
117 are both essential for mucocyst formation, providing the best evidence to  
118 date that mucocysts are lysosome-related organelles (Kaur et al., submitted).  
119 The CDH report for this *CTH3* query is attached as Supplementary File 4.  
120 This report is also edited to indicate the cases when the CDH matched or  
121 expanded upon existing gene annotations.

#### 122 4. Impact

123 The CDH reproduces existing annotations with high accuracy, and pro-  
124 vides a large number of new annotations and expansions upon existing ones  
125 (Table 3; Supplementary Files 2 and 4). Effectively, the CDH increased the  
126 annotation coverage of the genes co-regulated with *TWI1* from 46% to 60%,  
127 and the annotation coverage of the genes co-regulated with *CTH3* from 41%  
128 to 57%. Specifying the BLAST parameters allows the user to discover the  
129 most informative functional predictions for their genes and pathways of in-  
130 terest. Limiting the CDH search to lineages outside of the ciliates is more  
131 likely to retrieve previously annotated orthologs, but runs the increased risk  
132 that weak homologs will generate spurious results. For some processes, such  
133 as programmed genome rearrangement in which *TWI1* is involved, the most  
134 informative BLAST searches may be those restricted to the ciliates. In our  
135 trials, the effectiveness of the CDH is maintained regardless of which taxa  
136 the BLAST searches are run against.

137 In addition to providing a means of quickly gathering available data about  
138 a set of co-regulated genes and inferring their functions, the CDH data can be  
139 extended to to investigate the potential overlap between components of dif-  
140 ferent cellular pathways. For example, *NUP50* encodes a gene that functions  
141 both in nuclear import at the nuclear pore complex and as part of a com-  
142 plex involved in transcription [30, 31]. Accordingly, the genes co-regulated  
143 with *NUP50* show extensive overlap with genes co-regulated with an import  
144 factor (Importin $\beta$ ) and with a gene involved in transcription (*RPB81*, an  
145 RNA polymerase II subunit), among other factors involved in both processes  
146 (Figure 3, A).

147 It is informative to compare the overlap of co-regulated genes in different  
148 pathways. The co-regulated gene sets for three factors involved in genome  
149 rearrangement, *TWI1*, *GIW1*, and *DCL1*, show almost complete overlap,  
150 suggesting that they may be involved in a single common process (Figure 3,  
151 B). In contrast to the case of programmed genome rearrangement, the co-

152 regulated gene sets for three factors required in mucocyst formation, *CTH3*,  
153 *SOR4*, and *APM3*, show partial overlap, hinting that one or more of these  
154 factors may also play roles unrelated to mucocysts (Figure 3, C). Consistent  
155 with this idea, *CTH3* is an essential gene, while mucocysts themselves are  
156 dispensable for cell viability in the laboratory [19]. Importantly, there is very  
157 little overlap between the co-regulated gene sets defined for the three differ-  
158 ent cellular processes (nuclear import/transcription, genome remodeling, and  
159 mucocyst formation) (Figure 3, D). The overlap is smallest between the genes  
160 co-regulated with mucocyst biogenesis factors and genes co-regulated with ei-  
161 ther nuclear import, transcriptional regulation, or programmed genome rear-  
162 rangement. The somewhat greater sharing of genes co-regulated with nuclear  
163 import, transcriptional regulation, and programmed genome rearrangement  
164 may reflect the fact that these pathways all take place in the nucleus and  
165 are intrinsically linked to the cell cycle. Given the ease of assembling sets  
166 of co-regulated genes using the CDH, this type of overlap analysis can be  
167 extended to many cellular pathways.

## 168 5. Conclusions

169 Protists constitute the majority of eukaryotic diversity, meaning that this  
170 group needs to be included in evolutionary analyses of cellular processes, but  
171 this diversity is largely overlooked in the standard collection of model eukary-  
172 otes [20]. We present the Co-regulation Data Harvester for *T. thermophila*  
173 (CDH) as a tool that expedites analyses of *T. thermophila* genome, tran-  
174 scriptome, and cellular biology in an evolutionary context. The CDH is  
175 freely available and provides a systematic framework for genome annotation.  
176 It quickly gathers information from disparate databases and, by optionally  
177 reusing BLAST results that had been stored during previous queries, can  
178 increase in speed with successive uses. In providing a new means to analyze  
179 transcriptomic data, the CDH makes clear the potential for using the rapidly  
180 growing amount of genomic and transcriptomic data in many organisms, to  
181 facilitate functional analysis in poorly annotated or emerging model systems.

182 Users of the CDH should keep in mind that its reports are necessar-  
183 ily limited by pre-existing data from the TGD, *TetraFGD*, and the NCBI.  
184 For example, the *TetraFGD* does not provide co-expression data for genes  
185 whose expression level falls below a set threshold. Because of this limit, some  
186 *T. thermophila* genes may be overlooked by the CDH. Executable files for  
187 the program can be found at <http://ciliate.org/index.php/show/CDH>.  
188 A manual with detailed instructions and usage examples is provided in Sup-  
189 plementary File 5.

## 190 Acknowledgements

191 We would like to thank Stefano Allesina for providing the initial impetus  
192 for this project, excellent advice, and review of this manuscript. Thanks to  
193 Chad Pearson and Alexander Stemm-Wolf for their feedback on the CDH  
194 and this manuscript. We thank Naomi Stover and Wei Miao for granting  
195 the Ciliate community access to the TGD and *TetraFGD*, without which  
196 none of this work would be possible. We thank Tokuko Haraguchi for helpful  
197 discussion about NUP50. We thank Daniela Sparvoli and Harsimran Kaur  
198 for helpful discussion and good company. This work was supported by NIH  
199 GM-105783 to APT.

- 200 [1] G. Witzany, M. Nowacki (Eds.), *Biocommunication of Ciliates*, Springer,  
201 2016.
- 202 [2] C. W. Greider, E. H. Blackburn, Identification of a specific telomere  
203 terminal transferase activity in tetrahymena extracts, *Cell* 43 (2) (1985)  
204 405 – 413. doi:[http://dx.doi.org/10.1016/0092-8674\(85\)90170-9](http://dx.doi.org/10.1016/0092-8674(85)90170-9).
- 205 [3] K. Kruger, P. J. Grabowski, A. J. Zaug, J. Sands, D. E. Gottschling,  
206 T. R. Cech, Self-splicing RNA: Autoexcision and autocyclization of the  
207 ribosomal RNA intervening sequence of tetrahymena, *Cell* 31 (1) (1982)  
208 147–157. doi:10.1016/0092-8674(82)90414-7.
- 209 [4] I. Gibbons, A. Rowe, Dynein: a protein with adenosine triphosphatase  
210 activity from cilia, *Science* 149 (3682) (1965) 424–426.
- 211 [5] J. E. Brownell, J. Zhou, T. Ranalli, R. Kobayashi, D. G. Edmond-  
212 son, S. Y. Roth, C. Allis, Tetrahymena histone acetyltransferase a:  
213 A homolog to yeast *gcn5p* linking histone acetylation to gene activa-  
214 tion, *Cell* 84 (6) (1996) 843 – 851. doi:[http://dx.doi.org/10.1016/S0092-8674\(00\)81063-6](http://dx.doi.org/10.1016/S0092-8674(00)81063-6).
- 215  
216 [6] J. A. Eisen, R. S. Coyne, M. Wu, D. Wu, M. Thiagarajan, J. R. Wort-  
217 man, J. H. Badger, Q. Ren, P. Amedeo, K. M. Jones, L. J. Tallon,  
218 A. L. Delcher, S. L. Salzberg, J. C. Silva, B. J. Haas, W. H. Ma-  
219 joros, M. Farzad, J. M. Carlton, R. K. Smith Jr., J. Garg, R. E. Pearl-  
220 man, K. M. Karrer, L. Sun, G. Manning, N. C. Elde, A. P. Turkewitz,  
221 D. J. Asai, D. E. Wilkes, Y. Wang, H. Cai, K. Collins, B. A. Stew-  
222 art, S. R. Lee, K. Wilamowska, Z. Weinberg, W. L. Ruzzo, D. Wloga,  
223 J. Gaertig, J. Frankel, C.-C. Tsao, M. A. Gorovsky, P. J. Keeling, R. F.  
224 Waller, N. J. Patron, J. M. Cherry, N. A. Stover, C. J. Krieger, C. del

- 225 Toro, H. F. Ryder, S. C. Williamson, R. A. Barbeau, E. P. Hamil-  
226 ton, E. Orias, Macronuclear Genome Sequence of the Ciliate Tetrahy-  
227 mena thermophila, a Model Eukaryote, PLoS Biol 4 (9) (2006) e286.  
228 doi:10.1371/journal.pbio.0040286.
- 229 [7] N. A. Stover, C. J. Krieger, G. Binkley, Q. Dong, D. G. Fisk,  
230 R. Nash, A. Sethuraman, S. Weng, J. M. Cherry, Tetrahymena Genome  
231 Database (TGD): a new genomic resource for Tetrahymena ther-  
232 mophila research, Nucleic Acids Res 34 (suppl 1) (2006) D500–D503.  
233 doi:10.1093/nar/gkj054.
- 234 [8] W. Miao, J. Xiong, J. Bowen, W. Wang, Y. Liu, O. Braguinets,  
235 J. Grigull, R. E. Pearlman, E. Orias, M. A. Gorovsky, Microarray Analy-  
236 ses of Gene Expression during the Tetrahymena thermophila Life Cycle,  
237 PLoS ONE 4 (2) (2009) e4429. doi:10.1371/journal.pone.0004429.
- 238 [9] J. Xiong, X. Y. Lu, Y. M. Lu, H. H. Zeng, D. X. Yuan, L. F. Feng,  
239 Y. Chang, J. Bowen, M. Gorovsky, C. J. Fu, W. Miao, Tetrahymena  
240 Gene Expression Database (TGED): A resource of microarray data and  
241 co-expression analyses for Tetrahymena, Science China Life Sciences  
242 54 (1) (2011) 65–67. doi:10.1007/s11427-010-4114-1.
- 243 [10] J. Xiong, Y. Lu, J. Feng, D. Yuan, M. Tian, Y. Chang, C. Fu,  
244 G. Wang, H. Zeng, W. Miao, Tetrahymena functional genomics database  
245 (TetraFGD): An integrated resource for Tetrahymena functional ge-  
246 nomics, Database 2013 (2013) 6–11. doi:10.1093/database/bat008.
- 247 [11] J. Xiong, X. Lu, Z. Zhou, Y. Chang, D. Yuan, M. Tian, Z. Zhou,  
248 L. Wang, C. Fu, E. Orias, W. Miao, Transcriptome Analysis of the Model  
249 Protozoan, Tetrahymena thermophila, Using Deep RNA Sequencing,  
250 PLoS One 7 (2) (2012) e30630. doi:10.1371/journal.pone.0030630.
- 251 [12] J. J. Faith, B. Hayete, J. T. Thaden, I. Mogno, J. Wierzbowski,  
252 G. Cottarel, S. Kasif, J. J. Collins, T. S. Gardner, Large-Scale Map-  
253 ping and Validation of Escherichia coli Transcriptional Regulation from  
254 a Compendium of Expression Profiles, PLoS Biol 5 (1) (2007) 1–13.  
255 doi:10.1371/journal.pbio.0050008.
- 256 [13] J. Xiong, D. Yuan, J. S. Fillingham, J. Garg, X. Lu, Y. Chang, Y. Liu,  
257 C. Fu, R. E. Pearlman, W. Miao, Others, Gene network landscape of  
258 the ciliate Tetrahymena thermophila, PLoS One 6 (5) (2011) e20124.

- 259 [14] L. A. Stargell, K. M. Karrer, M. A. Gorovsky, Transcriptional regula-  
260 tion of gene expression in *Tetrahymena thermophila*, *Nucleic Acids Res*  
261 18 (22) (1990) 6637–6639. doi:10.1093/nar/18.22.6637.
- 262 [15] G. Csárdi, A. Franks, D. S. Choi, E. M. Airoidi, D. A. Drummond,  
263 Accounting for Experimental Noise Reveals That mRNA Levels, Am-  
264 plified by Post-Transcriptional Processes, Largely Determine Steady-  
265 State Protein Levels in Yeast, *PLoS Genet* 11 (5) (2015) e1005206.  
266 doi:10.1371/journal.pgen.1005206.
- 267 [16] C. Gurkan, H. Lapp, C. Alory, A. I. Su, J. B. Hogenesch, W. E. Balch,  
268 Large-scale profiling of Rab GTPase trafficking networks: the mem-  
269 brome, *Mol Biol Cell* 16 (8) (2005) 3847–3864.  
270 URL <http://www.molbiolcell.org/content/16/8/3847.full>
- 271 [17] M. S. Behnke, J. C. Wootton, M. M. Lehmann, J. B. Radke, O. Lucas,  
272 J. Nawas, L. D. Sibley, M. W. White, Coordinated progression through  
273 two subtranscriptomes underlies the tachyzoite cycle of *Toxoplasma*  
274 *gondii*, *PLoS One* 5 (8) (2010) 1–20. doi:10.1371/journal.pone.0012354.
- 275 [18] J. S. Briguglio, S. Kumar, A. P. Turkewitz, Lysosomal sorting receptors  
276 are essential for secretory granule biogenesis in *Tetrahymena*, *J Cell Biol*  
277 203 (3) (2013) 537–550.
- 278 [19] S. Kumar, J. S. Briguglio, A. P. Turkewitz, An aspartyl cathepsin,  
279 CTH3, is essential for proprotein processing during secretory granule  
280 maturation in *Tetrahymena thermophila*., *Mol Biol Cell* 25 (16) (2014)  
281 2444–60. doi:10.1091/mbc.E14-03-0833.
- 282 [20] M. Lynch, M. C. Field, H. V. Goodson, H. S. Malik, J. B. Pereira-  
283 Leal, D. S. Roos, A. P. Turkewitz, S. Sazer, Evolutionary cell biology:  
284 two origins, one objective, *Proc Natl Acad Sci U S A* 111 (48) (2014)  
285 16990–16994.
- 286 [21] P. J. A. Cock, T. Antao, J. T. Chang, B. A. Chapman, C. J. Cox,  
287 A. Dalke, I. Friedberg, T. Hamelryck, F. Kauff, B. Wilczynski, M. J. L.  
288 De Hoon, Biopython: Freely available Python tools for computational  
289 molecular biology and bioinformatics, *Bioinformatics* 25 (11) (2009)  
290 1422–1423. doi:10.1093/bioinformatics/btp163.
- 291 [22] J. W. Ratcliff, D. E. Metzener, Pattern-matching-the gestalt approach,  
292 *Dr Dobbs Journal* 13 (7) (1988) 46.

- 293 [23] E. H. Blackburn, K. M. Karrer, Genomic reorganization in ciliated pro-  
294 tozoans, *Annu Rev Genet* 20 (1) (1986) 501–521.
- 295 [24] K. M. Karrer, Nuclear dualism, *Methods Cell Biol* 109 (2012) 29–52.
- 296 [25] M.-C. Yao, J.-L. Chao, RNA-Guided DNA Deletion in Tetrahy-  
297 mena: An RNAi-Based Mechanism for Programmed Genome  
298 Rearrangements, *Annu Rev Genet* 39 (1) (2005) 537–559.  
299 doi:10.1146/annurev.genet.39.073003.095906.
- 300 [26] M.-C. Yao, J.-L. Chao, C.-Y. Cheng, Programmed Genome Rearrange-  
301 ments in Tetrahymena, *Microbiology Spectrum* 2 (6).
- 302 [27] K. Mochizuki, N. A. Fine, T. Fujisawa, M. A. Gorovsky, Anal-  
303 ysis of a piwi-Related Gene Implicates Small RNAs in Genome  
304 Rearrangement in Tetrahymena, *Cell* 110 (6) (2002) 689–699.  
305 doi:http://dx.doi.org/10.1016/S0092-8674(02)00909-1.
- 306 [28] A. T. Cowan, G. R. Bowman, K. F. Edwards, J. J. Emerson, A. P. Turke-  
307 witz, Genetic, Genomic, and Functional Analysis of the Granule Lattice  
308 Proteins in Tetrahymena Secretory Granules, *Mol Biol Cell* 16 (9) (2005)  
309 4046–4060. doi:10.1091/mbc.E05-01-0028.
- 310 [29] S. Kumar, J. S. Briguglio, A. P. Turkewitz, Secretion of Polypeptide  
311 Crystals from Tetrahymena thermophila Secretory Organelles (Muco-  
312 cysts) Depends on Processing by a Cysteine Cathepsin, Cth4p, *Eukaryot*  
313 *Cell* 14 (8) (2015) 817–833. doi:10.1128/EC.00058-15.
- 314 [30] M. E. Lindsay, K. Plafker, A. E. Smith, B. E. Clurman, I. G.  
315 Macara, Npap60/Nup50 is a tri-stable switch that stimulates importin-  
316 alpha:beta-mediated nuclear protein import., *Cell* 110 (3) (2002) 349–  
317 360. doi:S009286740200836X [pii].
- 318 [31] A. L. Buchwalter, Y. Liang, M. W. Hetzer, Nup50 is required for cell  
319 differentiation and exhibits transcription-dependent dynamics., *Mol Biol*  
320 *Cell* 25 (16) (2014) 2472–84. doi:10.1091/mbc.E14-04-0865.
- 321 [32] M. A. Nikiforov, J. F. Smothers, M. A. Gorovsky, C. D. Al-  
322 lis, Excision of micronuclear-specific DNA requires parental expres-  
323 sion of Pdd2p and occurs independently from DNA replication in  
324 Tetrahymena thermophila, *Genes Dev* 13 (21) (1999) 2852–2862.  
325 doi:10.1101/gad.13.21.2852.

- 326 [33] M. A. Nikiforov, M. A. Gorovsky, C. D. Allis, A novel chromodomain  
327 protein, pdd3p, associates with internal eliminated sequences during  
328 macronuclear development in *Tetrahymena thermophila*, *Mol Cell Biol*  
329 20 (11) (2000) 4128–4134. doi:10.1128/MCB.20.11.4128-4134.2000.
- 330 [34] R. M. Schwoppe, D. L. Chalker, Mutations in Pdd1 reveal distinct re-  
331 quirements for its chromodomain and chromoshadow domain in direct-  
332 ing histone methylation and heterochromatin elimination, *Eukaryot Cell*  
333 13 (2) (2014) 190–201. doi:10.1128/EC.00219-13.
- 334 [35] C. D. Malone, A. M. Anderson, J. A. Motl, C. H. Rexer, D. L. Chalker,  
335 Germ Line Transcripts Are Processed by a Dicer-Like Protein That  
336 Is Essential for Developmentally Programmed Genome Rearrangements  
337 of *Tetrahymena thermophila*, *Mol Cell Biol* 25 (20) (2005) 9151–9164.  
338 doi:10.1128/MCB.25.20.9151-9164.2005.
- 339 [36] K. Mochizuki, M. A. Gorovsky, A Dicer-like protein in *Tetrahymena* has  
340 distinct functions in genome rearrangement, chromosome segregation,  
341 and meiotic prophase, *Genes Dev* 19 (1) (2005) 77–89.
- 342 [37] T. Noto, H. M. Kurth, K. Kataoka, L. Aronica, L. V. DeSouza,  
343 K. W. M. Siu, R. E. Pearlman, M. A. Gorovsky, K. Mochizuki,  
344 The *Tetrahymena* Argonaute-Binding Protein Giw1p Directs a Mature  
345 Argonaute-siRNA Complex to the Nucleus, *Cell* 140 (5) (2010) 692–703.  
346 arXiv:NIHMS150003, doi:10.1016/j.cell.2010.02.010.
- 347 [38] C. M. Carle, H. S. Zaher, D. L. Chalker, A Parallel G Quadruplex-  
348 Binding Protein Regulates the Boundaries of DNA Elimination  
349 Events of *Tetrahymena thermophila*, *PLoS Genet* 12 (3) (2016) 1–22.  
350 doi:10.1371/journal.pgen.1005842.
- 351 [39] C. H. Rexer, D. L. Chalker, Lia1p, a novel protein required dur-  
352 ing nuclear differentiation for genome-wide DNA rearrangements in  
353 *Tetrahymena thermophila*, *Eukaryot Cell* 6 (8) (2007) 1320–1329.  
354 doi:10.1128/EC.00157-07.
- 355 [40] A. W. Y. Shieh, D. L. Chalker, LIA5 Is Required for Nuclear Reor-  
356 ganization and Programmed DNA Rearrangements Occurring during  
357 *Tetrahymena* Macronuclear Differentiation, *PLoS One* 8 (9) (2013) 1–  
358 15. doi:10.1371/journal.pone.0075337.
- 359 [41] M.-C. Yao, C.-H. Yao, L. M. Halasz, P. Fuller, C. H. Rexer, S. H.  
360 Wang, R. Jain, R. S. Coyne, D. L. Chalker, Identification of novel

- 361 chromatin-associated proteins involved in programmed genome rear-  
362 rangements in *Tetrahymena*., *J Cell Sci* 120 (Pt 12) (2007) 1978–1989.  
363 doi:10.1242/jcs.006502.
- 364 [42] J. Bednenko, T. Noto, L. V. DeSouza, K. W. M. Siu, R. E. Pearlman,  
365 K. Mochizuki, M. a. Gorovsky, Two GW repeat proteins interact with  
366 *Tetrahymena thermophila argonaute* and promote genome rearrange-  
367 ment., *Mol Cell Biol* 29 (18) (2009) 5020–30. doi:10.1128/MCB.00076-  
368 09.
- 369 [43] L. Aronica, J. Bednenko, T. Noto, L. V. DeSouza, K. W. M. Siu,  
370 J. Loidl, R. E. Pearlman, M. A. Gorovsky, K. Mochizuki, Study of  
371 an RNA helicase implicates small RNA-noncoding RNA interactions in  
372 programmed DNA elimination in *Tetrahymena*, *Genes and Development*  
373 22 (16) (2008) 2228–2241. doi:10.1101/gad.481908.
- 374 [44] A. Vogt, K. Mochizuki, A Domesticated PiggyBac Transposase Interacts  
375 with Heterochromatin and Catalyzes Reproducible DNA Elimination in  
376 *Tetrahymena*, *PLoS Genet* 9 (12). doi:10.1371/journal.pgen.1004032.
- 377 [45] I.-T. Lin, J.-L. Chao, M.-C. Yao, An essential role for the DNA  
378 breakage-repair protein Ku80 in programmed DNA rearrangements  
379 in *Tetrahymena thermophila*., *Mol Biol Cell* 23 (11) (2012) 2213–25.  
380 doi:10.1091/mbc.E11-11-0952.
- 381 [46] Y. Liu, S. D. Taverna, T. L. Muratore, J. Shabanowitz, D. F. Hunt,  
382 C. D. Allis, RNAi-dependent H3K27 methylation is required for het-  
383 erochromatin formation and DNA elimination in *Tetrahymena*, *Genes*  
384 *Dev* 21 (12) (2007) 1530–1545. doi:10.1101/gad.1544207.




Figure 1: CDH architecture. Beginning with a single *T. thermophila* gene as a query, the CDH identifies all genes that are co-regulated with it, via the TetraFGD. Next, the CDH uses the TGD to gather the annotation and sequence data for each gene in the co-regulated set. For each gene in the co-regulated set, the CDH then runs forward and reciprocal BLAST searches, through the NCBI and TGD, to identify likely orthologs. A phrase matching algorithm, based on the Ratcliff-Obershelp algorithm [22], as implemented by the python difflib library, is then used to summarize the annotations of the putative orthologs for each *T. thermophila* gene in the co-regulated set. These summaries, which provide predictions about the function (e.g., relevant biological pathway) of the *T. thermophila* gene query, are presented along with the other data gathered, in the final report.

```
What is your quest (please enter a gene ID)? TTHERM_00313130

To determine how many of the co-regulated genes should be
subject to homology analysis, please enter the lower-bound
z-score for the strength of co-regulation: 5

How should I process your query?
(1) overwrite all associated files,
(2) overwrite just the BLASTs and analysis
    as well as fill in any missing files,
(3) overwrite only the analysis and fill in any
    missing files,
(4) sanitize database errors, or
(5) run only the FGD/TGD search
Your choice: 1

Send to Dropbox?
(1) Yes, and also write new results locally.
(2) Yes, but do not write new results locally.
    Remark: if you chose option (2) or (3) above,
    some files may still be synchronized
    between the Dropbox and local directories.
(3) No, run everything locally.
Your choice: 3

What kind of NCBI BLAST algorithm would you like to run?
(1) BLASTp,
(2) BLASTx, or
(3) both?
Your choice: 1

You may choose whether to look for homologs in all organisms
outside the Ciliates, only within the Ciliates, everywhere,
or custom entrez query:
(1) BLAST outside the Ciliates
(2) BLAST within the Ciliates
(3) BLAST everywhere
(4) Custom (please use the NCBI guidelines and
    instructions for formulating the entrez query)
Your choice: 2
```

Figure 2: Setting CDH search parameters. The CDH is run through the terminal. The CDH prompts the user to define several parameters. These are: 1) the initial gene, i.e., the query; 2) the z-score threshold to be applied as cutoff for strength of co-regulation, which determines how many of the co-regulated genes will be subject to analysis *via* homology; 3) the extent to which data gathered in prior searches should be used; 4) whether results should be stored in Dropbox; 5) whether to run BLAST searches with cDNA or protein sequences; and 6) in which taxa to run the BLAST searches. For (2), the z-score threshold determines how many co-regulated genes will be included. For example, raising the threshold increases the stringency of the requirement for strength of co-regulation, so results in fewer co-regulated genes that are subsequently analyzed via BLAST, etc. For (3), the available options are: a) to run the search from scratch, overwriting any files associated with the queried gene; b) to re-use existing data for co-regulation, annotations, and sequences, but to run all of the BLAST searches from scratch; c) to re-use any existing data that are pertinent to the given query; d) to clear NCBI database errors from a previously run search and redo the associated BLAST searches; e) to only run the search for the co-regulation, annotation, and sequence. The example query in this screenshot is set to run a CDH search for the gene TTHERM.00313130 (Sortilin 4); to consider genes that are co-regulated with it with a z-score of 5 or higher; to gather all of the data *de novo*; to save all of the data locally; and to run the BLASTp searches only within the Ciliates.




Figure 3: Using CDH outputs to assess overlap in gene function. Panels A, B, and C illustrate the overlaps in co-regulated genes for three different cellular pathways: A) nuclear import and transcriptional regulation; B) programmed genome rearrangement during cell conjugation; and C) mucocyst biogenesis. Each circle in the Venn diagrams corresponds to the full set of genes, as reported by the *TetraFGD*, that are co-regulated with the gene indicated at the periphery of the circle. (A) *NUP50* (Nucleoporin 50) plays roles both in nuclear import and in gene transcription. The dual role of *NUP50* is reflected in the overlap of genes co-regulated with *Importinβ* (an import factor) and with *RPB81* (RNA Pol II subunit), a transcription factor. *NUP50*, *RPB81*, and *Importinβ* are mutually co-regulated. The CDH identifies 214 genes co-regulated with the nucleoporin *NUP50*, 444 genes co-regulated with *RPB81*, and 200 genes co-regulated with *Importinβ*. (B) *TWI1* (*Tetrahymena* Piwi 1), *GIW1* (Gentleman in Waiting 1) and *DCL1* (Dicer-like 1) are all required for programmed genome rearrangement, and are mutually co-regulated. The CDH identifies 932 genes co-regulated with *TWI1*, 999 genes co-regulated with *GIW1*, and 814 genes co-regulated with *DCL1*. (C) *CTH3* (cathepsin 3), *APM3* ( $\mu$  subunit of the adaptin 3 complex), and *SOR4* (sortilin 4) are all required for formation of mucocysts, and are mutually co-regulated. These genes also appear to have distinct cellular functions in addition to their roles mucocyst formation. For example, mucocysts are non-essential organelles, yet *CTH3* is an essential gene. The CDH identifies 213 genes co-regulated with *CTH3*, 201 genes co-regulated with *APM3*, and 203 genes co-regulated with *SOR4*. (D) Pooling all of the genes represented in A, B, and C demonstrates that there is no overlap in co-regulated genes between A and B or C, and limited overlap between B and C.

<b>Gene Group</b>	<b>Genes</b>	<b>Experimental Reports in <i>T. thermophila</i></b>
Chromo-domain	<i>PDD1</i> , <i>PDD2</i> , <i>PDD3</i>	<i>PDD1</i> and <i>PDD2</i> are essential for programmed genome rearrangement; <i>PDD3</i> is reported to co-localize with <i>PDD1</i> , <i>PDD2</i> , and other necessary factors [32, 33, 34].
Dicer-like	<i>DCL1</i>	<i>DCL1</i> is essential for programmed genome rearrangement [35, 36].
Piwi-Associated	<i>GIW1</i>	<i>GIW1</i> is essential for programmed genome rearrangement and has no known conserved function [37].
Localized in nuclear anlagen	<i>LIA1</i> , <i>LIA2</i> , <i>LIA3</i> , <i>LIA4</i> , <i>LIA5</i> , <i>LIA6</i>	<i>LIA</i> proteins co-localize with <i>PDD1</i> at DNA rearrangement foci; <i>LIA1</i> and <i>LIA5</i> have been shown to be necessary for genome rearrangement, and <i>LIA3</i> is required for precise excision of eliminated sequences [38, 39, 40, 41].
Zinc knuckle	<i>cnjB</i>	A double-knockout of <i>cnjB</i> and <i>WAG1</i> inhibits the formation of DNA elimination structures [42].
Nucleic acid helicase	<i>EMA1</i>	<i>EMA1</i> is necessary for the association of <i>TWI1</i> with chromatin [43].
Transposase	<i>TPB2</i>	<i>TPB2</i> catalyzes DNA elimination [44].
Ku70/Ku80 beta-barrel domain	<i>TKU80</i>	<i>TKU80</i> is necessary for both <i>PDD1</i> complex assembly and DNA repair after programmed DNA elimination [45, 26]
Histone lysine methyltransferase	<i>EZL1</i>	<i>EZL1</i> is necessary for H3K27 methylation and programmed DNA elimination [46].

Table 1: A subset of the programmed genome rearrangement factors that were identified by a CDH query for *TWI1*.

Gene Group	Genes	Experimental Reports in <i>T. thermophila</i>
Sortilins	<i>SOR1</i> , <i>SOR2</i> , <i>SOR4</i>	<i>SOR2</i> and <i>SOR4</i> are essential for mucocyst biogenesis [18].
Cathepsins	<i>CTH1</i> , <i>CTH2</i> , <i>CTH4</i>	<i>CTH1</i> and <i>CTH2</i> are involved in mucocyst biogenesis [19].
Granule cargo	<i>GRL1</i> , <i>GRL2</i> , <i>GRL3</i> , <i>GRL4</i> , <i>GRL6</i> , <i>GRL7</i> , <i>GRL9</i> , <i>GRT1</i> , <i>IGR6</i> , <i>IGR7</i>	These genes, belonging to two gene families, encode mucocyst contents [28].
Adaptin complex subunits	<i>AP3</i> $\mu$ and $\beta$ subunits	<i>AP3</i> $\mu$ subunit ( <i>APM3</i> ) is necessary for mucocyst biogenesis (Kaur et al., submitted).
Syntaxins	<i>STX7L1</i>	<i>STX7L1</i> is essential for mucocyst biogenesis (Kaur et al., submitted).

Table 2: Mucocyst biogenesis and cargo factors that were identified by a CDH query for *CTH3*.

Queried Gene	<i>TWI1</i>	<i>CTH3</i>
Number co-regulated genes	932	213
Taxa BLASTed	Only Ciliates	Excluding Ciliates
Number previously annotated	430	88
Annotations matched	299	43
Annotations expanded upon	19	19
Novel annotations	126	34

Table 3: The CDH accurately reproduces existing annotations and provides new annotations at a high rate.

385 **Required Metadata**

386 **Current code version**

387 **Current executable software version**

Nr.	Code metadata description	Please fill in this column
C1	Current code version	v1.1.0
C2	Permanent link to code/repository used for this code version	<i>https</i> : <i>//bitbucket.org/ltsypin/cdhproject/</i>
C3	Legal Code License	GNU GPL v3
C4	Code versioning system used	git
C5	Software code languages, tools, and services used	python
C6	Compilation requirements, operating environments & dependencies	any system that can run python 2.7
C7	If available Link to developer documentation/manual	<i>http</i> : <i>//ciliate.org/index.php/show/CDH</i>
C8	Support email for questions	coregulationdataharvester@gmail.com

Table 4: Code metadata (mandatory)

Nr.	(Executable) software metadata description	Please fill in this column
S1	Current software version	1.1.0
S2	Permanent link to executables of this version	<i>http</i> : <i>//ciliate.org/index.php/show/CDH</i>
S3	Legal Software License	GNU GPL v3
S4	Computing platforms/Operating Systems	OS X (10.6+) and Windows (x64) Vista or later
S5	Installation requirements & dependencies	
S6	If available, link to user manual - if formally published include a reference to the publication in the reference list	<i>http</i> : <i>//ciliate.org/index.php/show/CDH</i>
S7	Support email for questions	coregulationdataharvester@gmail.com

Table 5: Software metadata (optional)