

Dynamic Current Clamp Experiments Define the Functional Roles of I_{K1} and $I_{to,f}$ in Human Induced Pluripotent Stem Cell Derived Cardiomyocytes

Short title: $I_{to,f}$ in human iPSC-derived cardiomyocytes

Scott B. Marrus, MD, PhD¹, Steven Springer, PhD^{2,3}, Eric Johnson, PhD^{2,3}, Rita J. Martinez⁴, Edward J. Dranoff^{2,3}, Rebecca Mellor^{2,3}, and Kathryn Yamada, PhD^{2,3}

¹Department of Electrical and Systems Engineering, Washington University in St Louis, St Louis MO, USA

²Department of Internal Medicine, Cardiovascular Division, ³Center for Cardiovascular Research,

⁴Department of Psychiatry, Washington University School of Medicine

Corresponding author:

Scott Marrus, MD, PhD

Washington University in St Louis

Campus Box 1042

1 Brookings Drive

St Louis MO 63130

314-935-4261

smarrus@wustl.edu

Abstract

The transient outward potassium current (I_{to}) plays a key, albeit incompletely defined, role in cardiomyocyte physiology and pathophysiology. In light of the technical challenges of studying adult human cardiomyocytes, this study examines the use of induced pluripotent stem cell-derived cardiomyocytes (iPSC-CMs) as a system which potentially preserves the native cellular milieu of human cardiomyocytes. iPSC-CMs express a robust I_{to} with slow recovery kinetics and fail to express the rapidly recovering $I_{to,f}$ which is implicated in human disease. Overexpression of the accessory subunit KChIP2 (which is not expressed in iPSC-CMs) resulted in restoration of a rapid component of recovery. To define the functional role of I_{to} , dynamic current clamp was used to introduce computationally modeled currents into iPSC-CMs while recording action potentials. However, iPSC-CMs exhibit action potentials with multiple immature physiological properties, including slow upstroke velocity, heterogeneous action potential waveforms, and the absence of a phase 1 notch, thus potentially limiting the utility of these cells as a model of adult cardiomyocytes. Importantly, the introduction of modeled inwardly rectified current (I_{K1}) ameliorated these immature properties by restoring a hyperpolarized resting membrane potential. In this context of normalized action potential morphologies, dynamic current clamp experiments introducing $I_{to,f}$ demonstrated that there is significant cell-to-cell heterogeneity and that the functional effect of $I_{to,f}$ is highly sensitive to the action potential plateau voltage in each cell.

Introduction

Normal cardiac function requires organized, unidirectional propagation of electrical activity, a process dependent on the activation and inactivation of multiple ion channels resulting in the sequential generation of depolarizing and repolarizing currents.(1) Abnormalities of a key repolarizing cardiac potassium current, the fast transient outward current ($I_{to,f}$), encoded by members of the Kv4 family, are associated with both heart failure and Brugada syndrome, a congenital arrhythmia syndrome characterized by precordial ST-segment elevation and a risk of malignant arrhythmias.(2, 3) However, the functional role of $I_{to,f}$ in normal human cardiac rhythm as well as arrhythmogenesis remains incompletely defined.

$I_{to,f}$ has several inter-related roles in cardiac myocyte function; it contributes to the action potential (AP) waveform and rate-dependent action potential properties as well as to excitation-contraction coupling by influencing calcium influx.(4-8) The role of $I_{to,f}$ in shaping cardiomyocyte action potentials varies across species, reflecting species-specific differences in other ion channel properties. In rodents with brief action potentials, $I_{to,f}$ contributes importantly to rapid repolarization.(1) In large mammals with prolonged action potentials, the role of $I_{to,f}$ is more complex. The rapid activation and inactivation of the current appears to confine its role to the phase 1 notch of the action potential. However, canine experimental and modeling studies indicate that changes in $I_{to,f}$ alter $I_{Ca,L}$, and thereby exert a complex influence on the AP waveform; this relationship results in a non-linear relationship between $I_{to,f}$ density and action potential duration.(5, 7, 9)

Due to the complex role of $I_{to,f}$ in shaping action potential waveforms, it is important to examine the functional role of the current in the context of the other native currents present in cardiomyocytes. The recent development of induced pluripotent stem cell technology and the refinement of methods to drive cardiomyocyte differentiation from pluripotent cells offer a novel and promising avenue for the study of cardiomyocyte cellular physiology. However, the utility of induced pluripotent stem cell derived cardiomyocytes (iPSC-CMs) and embryonic stem cell derived cardiomyocytes (ESC-CMs) naturally hinges on how closely they recapitulate adult human cardiomyocyte physiology.

Broadly speaking, pluripotent stem cell derived cardiomyocytes (PSC-CMs) appear to most closely resemble embryonic or immature cardiomyocytes. Over the course of *in vitro* differentiation, iPSC-CMs and ESC-CMs develop many features of mature cardiomyocytes, including expression of cardiac specific transcription factors, contractile proteins, and excitation-contraction coupling.(10) However, the morphology of the cells remains immature, sarcomeres are disorganized, and t-tubules are absent.(11) Furthermore, iPSC-CMs exhibit action potentials with a wide variety of waveforms, generally interpreted as a mixture of ventricular, atrial, and nodal action potential waveforms.(12) However, the action potential exhibit several properties which are not typical of mature cardiomyocytes, including spontaneous diastolic depolarization, slow upstroke velocity, and prolonged action potential duration.(11)

Previous studies have examined the ionic conductances which underlie the action potential waveforms.(12) To date, thorough characterization of the biophysical properties of all of the ion channels expressed in iPSC-CM remains incomplete, but the properties of the ion channels appear to be largely similar to the ion channels expressed in adult ventricular myocytes albeit with a few notable exceptions, including a smaller magnitude inwardly rectifying current (I_{K1}) and a larger magnitude funny current (I_f).(11)

In this study, we demonstrate that iPSC-CMs fail to exhibit $I_{to,f}$ due to the absence of the essential subunit KChIP2 whereas the closely related but kinetically distinct current $I_{to,s}$ is robustly expressed. Although spontaneous action potentials in iPSC-CMs exhibited highly heterogeneous and immature

features, iPSC-CM action potential morphology was rendered essentially mature with the introduction of a computationally modeled I_{K1} . This maneuver provided a reasonable model of adult human cardiomyocytes action potentials in which the functional role of $I_{to,f}$ could be examined. Using dynamic current clamp, we demonstrate that the effect of $I_{to,f}$ on action potential waveforms varies from cell to cell and is markedly sensitive to the context of other ionic conductances in the cell.

Methods

IPSC generation and cardiomyocyte differentiation

Induced pluripotent stem cells were generated as previously described.(13) Briefly, primary fibroblast lines were established from both a commercially available foreskin fibroblast line (ATCC) and dermal fibroblasts isolated from a skin biopsy of a healthy 36 yo male. Each line was transduced utilizing the non-integrating CytoTune-iPS Sendai Reprogramming Kit (A1378001, Life Technologies). Individual iPSC lines were generated by manual picking and expansion of individual clones in a feeder-free defined system (mTeSR1, StemCell Technologies) and Matrigel (354277, Corning). A minimum of two clones were frozen down for each line and characterization was performed for karyotype, pluripotency and tri-lineage germ layer analysis. All lines had a normal karyotype and expressed endodermal, mesodermal, and epidermal transcripts after spontaneous embryoid body differentiation (data not shown). One clone each from neonatal fibroblasts and adult dermal fibroblasts were used for further studies. No differences were identified between iPSC cells or iPSC-CMs derived from neonatal fibroblasts (line BJFF.6) or adult dermal fibroblasts (line F10336.6) and data from both lines were subsequently pooled. Written consent was obtained from all participants; the process for consent and acquisition of adult dermal fibroblasts was approved by the Human Research Protection Office at Washington University School of Medicine (protocol #201104172).

Cardiomyocyte differentiation was carried out as previously described using small molecule inhibitors of GSK3 and Wnt signaling; optimal concentrations for both inhibitors were determined based on percentage of beating cells.(14) IPS cells were passaged using accutase (Innovative Cell Technologies). One well of a 6-well plate was seeded with 1×10^6 iPS cells in mTeSR1 with 5 μ M ROCK inhibitor (Y27632, Tocris) and grown to confluence. On Day 0 of the differentiation, cells were fed 4 ml per well with RPMI media supplemented with B27 without insulin (Gibco/Life Technologies). GSK3 inhibitor (CHIR99021, Tocris, 12 μ M for BJFF.6 and 6 μ M for F10336.6) was added to the media on Day 0. On Day 1, precisely 24 hours after addition of CHIR99021, cells were fed with RPMI+B27 without insulin. On Day 3, 2 ml of conditioned media per well were mixed with 2 ml fresh RPMI+B27 without insulin; the Wnt signaling inhibitor IWP2 (Tocris, 5 μ M for BJFF.6 and 10 μ M for F10336.6) was added. On Day 5, cells were fed with RPMI+B27 without insulin. On Day 7, and subsequently every 2-3 days, cells were fed with RPMI+B27 with insulin.

On day 20 of differentiation, non-cardiomyocytes were eliminated by feeding with DMEM without glucose (Gibco) supplemented with 4 mM lactate for 3-5 days.(15) Between day 30 and 50 of differentiation, cells were harvested for analysis. Cardiomyocytes were digested with two 10 minute incubations with 0.02% EDTA (Sigma) followed by a brief (~3 min) incubation in 0.25% trypsin; digestion was halted with the addition of RPMI+B27 supplemented with 10% fetal bovine serum. After low-speed centrifugation, cells were replated at low density on plates prepared with 0.1% gelatin in RPMI+B27 with 10% FBS. Cells were fed with RPMI+B27 every 2-3 days and used for electrophysiological recording at least 5-7 days after digestion.

Electrophysiological recording

To confirm cells' identity as cardiomyocytes, the mitochondrial dye TMRM (5-10 nM, Molecular Probes, Life Technologies) was added to each plate 15 minutes prior to recording and only cells exhibiting

red fluorescence were studied.(16) Recording pipettes fashioned from borosilicate glass (WPI, TW150F-4) had resistances of 2-3 M Ω when filled with recording solution. Whole-cell K⁺ currents were recorded at room temperature (20-23°C) using previously described solutions and protocols.(17, 18) Action potentials were recorded at 37°C using pipettes containing (in mM): K aspartate 76; KCl 20; MgCl₂ 2.5; HEPES 10; NaCl 4; CaCl₂ 6; EGTA 10; K₂ATP 10; and NaGTP 0.1. Calculated free calcium was 73 nM and free magnesium was 0.36 mM (calculated with maxchelator.stanford.edu). Experiments were controlled by and data were collected using an Axopatch 1D, digitized with a Digidata 1322A, and stored on a PC running Clampex 9.2 (Molecular Devices).

Dynamic current clamp

Using the same recording conditions as above for action potentials, the cell membrane voltage was simultaneously digitized with a Digidata 1322A for storage by Clampex 9.2 and digitized with a NI PCI-6010 DAQ card (National Instruments) installed on a separate PC running a 64-bit real-time Linux kernel (RTAI). A BNC-2110 (National Instruments) connected the amplifier output to the input of the DAQ card. Simulated currents were computed based on the cell membrane voltage (sampled at 10 kHz) using RTXI software(19) and the output of the simulation converted to an analog signal by the PCI-6010 DAQ card. The RTXI output and the Clampex output (used to deliver a brief, depolarizing action potential trigger at 1 Hz) were combined using a custom-made voltage summing junction and the combined signal delivered to the external command of the amplifier. The liquid junctional potential was calculated using Clampex 9.2 and added to the measured transmembrane voltage signal prior to performing the modeled current calculations. Equations for the computational models of human ventricular I_{K1} and $I_{to,f}$ were published by O'Hara et al.(20)

Transcript analysis

Total RNA was isolated using the Qiagen RNeasy Mini Kit (74104) with DNase treatment. Reverse transcription and SYBR Green real time PCR were carried out as previously described.(21) Primers used for RT-PCR are summarized in Table 1. RNA expression levels were normalized first to GAPDH and second to troponin (to normalize for the variable cardiomyocyte content of each differentiation). Tissue from non-failing human hearts was obtained from donor hearts deemed unsuitable for transplantation. Immediately after explantation, small pieces of tissue were placed in RNAlater (Life Technologies) for at least 24 hours, after which RNAlater was removed and the tissue frozen at -80°C until use. Tissue was obtained from donor hearts deemed unsuitable for transplantation; written consent was obtained from the next-of-kin according to the policies of Mid-America Transplant Services. The study was approved by the Human Research Protection Office at Washington University School of Medicine (protocol #201107099).

Table 1. Sequence-specific primers used in SYBR Green RT-PCR

Gene	Protein	Forward primer	Reverse primer
<i>KCND3 long</i>	Kv 4.3 (long isoform)	5'-GATGATCCCCTGTTATCTGTACGA	5'-ACTACGACGGGAGCAGCAGG
<i>KCND3 short</i>	Kv 4.3 (short isoform)	5'-TCGAGAGCCAGCATCATCAC	5'-ACTACGACGGGAGCAGCAGG
<i>KCNA4</i>	Kv 1.4	5'-AAGATTGTCCGGTCCCTGTGT	5'-CAATCACTGGCACTGGCAAA
<i>KCNIP1</i>	KChIP1	5'-CCCCAGTGGTGTGGTCAAC	5'-CCATGAGGGAAAACTGAGCAT
<i>KCNIP2</i>	KChIP2	5'-GCAGGAGCAAACCAATTAC	5'-TTCTGAAGCCCCGGTACAG
<i>KCNIP3</i>	KChIP3	5'-GGCACAGTCCACGAGAAGCT	5'-ATGTAGCCATCCTTGTAAATGTCGTA
<i>KCNIP4</i>	KChIP4	5'-CCACAATGGAGCTGTGAGTTTC	5'-CCCCGGAGCAAATGGA

Data analysis

Analysis of electrophysiological data was performed using MatLab (Mathworks). Whole-cell membrane capacitance was calculated by integrating the capacitive transient elicited by ± 5 mV steps from a -70 mV holding potential. The decay phase of the elicited current was fit with a single exponential function; in cells with a biphasic decay, a two exponential function was used and the relative contribution of each component calculated from the weighting of each exponent, as previously described(22). Action potentials were analyzed for resting membrane potential, peak upstroke velocity, the presence and magnitude of a phase 1 notch, and action potential duration. All values are reported as mean \pm SEM.

Results

IPSC-CMs express a transient outward potassium current

All IPSC-CMs ($n=39$) exhibited a rapidly activating, rapidly inactivating transient outward potassium current (Figure 1). The transient outward current decay exhibited a fast component with a time constant (τ_{inact}) of 74.9 ± 5.4 ms (at 40 mV), similar to values reported for adult LV cardiomyocytes ($\sim 45-65$ ms).(23, 24) A subset of cells (20 out of 39) also exhibited a more slowly inactivating component (τ_{inact} 635 ± 163 ms) which has not been described in adult human LV cardiomyocytes. Subsequent analysis therefore focuses on only the fast-inactivating component of the transient outward current. The current density of the rapidly inactivating current, I_{to} , was 6 ± 0.9 pA/pF (at 40 mV), a value similar to the magnitude of the transient outward current reported in adult LV cardiomyocytes ($\sim 2-10$ pA/pF).(23, 24)

Figure 1. iPSC-CMs express a rapidly activating and inactivating transient outward current. A. Representative whole-cell Kv currents (normalized to cell capacitance) recorded from iPSC-CMs in response to voltage steps from 40 mV to -10 mV in -5 mV increments from a holding potential of -70 mV. Capacitive transients have been removed for clarity. Scale bars: 5 pA/pF, 200 ms. B. Current-voltage relationship for fast-inactivating component of the whole cell current recorded in iPSC-CMs ($n=39$, mean \pm SEM).

I_{to} in IPSC-CMs exhibits slow recovery kinetics

Two phenotypically (and molecularly) distinct transient outward currents have been described in various species, a fast I_{to} ($I_{to,f}$) characterized by rapid recovery from inactivation (on the order of 10s of ms) and a slow I_{to} ($I_{to,s}$) characterized by slow recovery from inactivation (on the order of 1000s ms).(22, 25) I_{to} in iPSC-CMs exhibited predominantly slow recovery consistent with ($I_{to,s}$) (Figure 2), with a time constant of 3735 ± 352 ms. In a minority of cells (2/14), a faster component of recovery (although slower than reported for human $I_{to,f}$) was also evident with a time constant of 298 ± 213 ms.

Figure 2. I_{to} in iPSC-CMs recovers slowly from steady state inactivation. After inactivating the current during 1.5 sec. prepulse steps to +40mV, cells were hyperpolarized to -70 mV for times ranging from 0.1 to 10 sec. before a second depolarization to +40 mV was applied to assess the extent of recovery. A. Typical whole-cell Kv current recorded during the inactivation recovery protocol. Scale bars: 5 pA/pF, 1 sec. B. The amplitude of I_{to} after each recovery interval was normalized to the amplitude of the current during the prepulse. The time course of inactivation recovery (mean \pm SEM at each timepoint) is best fit with a single exponential function with a time constant of 3735 ± 352 ms.

Expression of KCHIP2 subunit restores I_{to} rapid recovery kinetics

Transcripts for the pore-forming subunits *KCND3* (long and short splice variants) encoding Kv4.3 and the *KCNA4* transcript encoding Kv1.4, underlying $I_{to,f}$ and $I_{to,s}$, respectively, are expressed at comparable levels in both iPSC-CMs and adult left ventricular tissue (Figure 3). However, the *KCNIP2* transcript encoding the accessory subunit KCHIP2 was expressed at significantly lower levels in iPSC-CMs compared to adult left ventricular tissue (Figure 3). KCHIP2 has been previously shown to be an essential component of $I_{to,f}$ in murine hearts.(21) Thus, the low expression level of KCHIP2 transcript suggests a

reason for the lack of rapidly recovering I_{to} whereas Kv1.4 likely underlies the observed, slowly recovering I_{to} .

Figure 3. Expression of transcripts encoding subunits underlying I_{to} in human iPSC-CMs. Transcript expression levels were measured in individual samples of iPSC-CMs (n=6-10) and adult LV apical tissue (n=3), normalized to GAPDH to control for sample loading and subsequently normalized to troponin-T to control for variable numbers of cardiomyocytes in iPSC-CM samples. A. No significant differences exist between iPSC-CMs and adult human ventricular tissue in transcript expression levels of the pore-forming subunits Kv 4.3L (long splice variant), Kv 4.3S (short splice variant), or Kv1.4. B. KChIP2 is the predominant KChIP subunit expressed in adult LV tissue and is expressed at significantly (*p=0.005) lower levels in iPSC-CMs; expression in both cell types of transcripts for KChIP1,2, and 4 were negligible.

To directly test the contribution of KChIP2 to $I_{to,f}$ in human iPSC-CMs, KChIP2 was overexpressed in iPSC-CMs. Overexpression of KChIP2 resulted in an increase in I_{to} density from 5.9 ± 0.9 pA/pF to 14.1 ± 2.4 pA/pF (p=0.0003). In the presence of KChIP2, $40 \pm 5\%$ of I_{to} recovered rapidly ($\tau=98 \pm 8$ ms) while the remaining current recovered slowly ($\tau=3713 \pm 224$ ms, p=NS compared to recovery in the absence of KChIP2) (Figure 4). The time course of rapid recovery is similar to the value previously reported for $I_{to,f}$ (22-24)

Figure 4. Overexpression of KChIP2 in iPSC-CMs results in a rapid component of I_{to} recovery from inactivation. A. The amplitude of I_{to} after each recovery interval was normalized to the amplitude of the current during the prepulse. The timecourse of inactivation recovery (mean \pm SEM at each timepoint) is best fit with a bi-exponential function (compare to Figure 2B). B. After overexpression of KChIP2, the time constant of the slow component of inactivation recovery remained unchanged but a significantly faster component (*p=0.0003) was present.

IPSC-CMs exhibit a wide range of spontaneous AP waveforms

The majority (52/63) of iPSC-CMs exhibited either spontaneous or evoked action potentials. Approximately half (32/63) of the iPSC-CMs recorded exhibited spontaneous action potentials. These cells exhibited a wide range of minimum diastolic potentials (MDP) (-69 ± 11 mV), peak upstroke velocity (dV/dt) (57 ± 11 V/s), and action potential duration (APD90) (619 ± 412 ms) (Figure 5). Spontaneous APs could be grouped into nodal-like, embryonic atrial-like, and embryonic ventricular-like cells on the basis of action potential waveforms.(26) Nodal-like cells (n=4) exhibited an upstroke velocity <10 mV/ms (5 ± 2 mV/ms). Cells with an upstroke velocity >10 mV/ms were further identified as ventricular-like (n=17) or atrial-like (n=11) on the basis of the presence or absence, respectively, of a plateau phase. The presence of a plateau phase was quantified as the ratio of the slope of the AP waveform between APD70 and APD90 and between APD30 and APD50 (referred to here as the “plateau ratio”); a ratio greater than 1.5 indicated a plateau phase with a more shallow slope than the terminal repolarization.

Addition of computationally modeled I_{K1} results in mature action potential parameters

A prominent feature of iPSC-CMs is a depolarized membrane potential, attributed at least in part to a reduced inwardly rectified current I_{K1} (11) resulting from reduced expression of the underlying Kir2.1. Dynamic current clamp was used to introduce a computationally modelled I_{K1} (the mathematical formulation of human cardiac I_{K1} published by O’Hara et al. (20) was used) at the current density reported in adult human cardiomyocytes which resulted in the cessation of spontaneous action potentials and modification of action potential parameters (Figure 5). The addition of modeled I_{K1} resulted in significant membrane hyperpolarization compared to the spontaneous MDP (-90 ± 1 mV vs. -69 ± 2 mV, p<0.0001) (Figure 5). Additionally, the peak upstroke velocity (dV/dt) was increased from 57 ± 11 mV/ms to 313 ± 34 mV/ms (p<0.0001), similar to the value reported in adult human cardiomyocytes. The change in action potential morphology was evident by a decrease in the action potential duration (APD90, 619 ± 74 ms vs.

336±37 ms, $p=0.003$). Importantly, after the addition of I_{K1} all action potentials exhibited a ventricular-like waveform with a plateau ratio >1.5 (Figure 5). These results indicate that the addition of modelled I_{K1} render iPSC-CM action potentials more similar to adult ventricular cardiomyocytes.

Figure 5. Introduction of computationally modeled I_{K1} using dynamic current clamp significantly normalizes action potential waveforms. A. Representative waveforms of spontaneous action potentials and evoked action potentials after addition of I_{K1} ; each pair of traces are from the same cell. B-E. Values from individual cells are represented by grey circles; mean±SEM are indicated by the superimposed bars. Addition of I_{K1} results in a significantly hyperpolarized resting membrane potential (B), increased maximum upstroke velocity (C) and decreased APD90 (D). E. After addition of I_{K1} all cells exhibited a plateau ratio > 1.5 (indicated by the dotted line) indicative of a ventricular action potential waveform. ** $p<0.0001$, * $p<0.005$

Given the expression of a robust I_{to} , it is striking that only 3/29 cells exhibited spontaneous action potentials with rapid phase 1 repolarization and/or a phase 1 notch. One possible reason for this discrepancy is that, due to the depolarized resting membrane potential (RMP), I_{to} is mostly inactivated. Indeed, restoration of a hyperpolarized resting membrane potential by addition of modeled I_{K1} resulted in a phase 1 notch in 15/23 cells.

Addition of computationally modeled $I_{to,f}$ elucidates the functional role of $I_{to,f}$

After the addition of modeled I_{K1} , iPSC-CMs exhibit action potentials similar to adult human LV myocytes, suggesting that these cells represent an appropriate system in which to examine the functional role of $I_{to,f}$ on action potential duration and waveform. To examine the functional role of $I_{to,f}$, computationally modeled $I_{to,f}$ (based on (20)) was introduced using dynamic current clamp (while simultaneously also delivering modeled I_{K1}). The modeled current was tailored to individual cell size; the model specified a conductance density of 0.02 mS/ μ F and the cell capacitance was manually entered into the calculations. As expected, the addition of modeled $I_{to,f}$ resulted in the appearance of a phase 1 notch (4/9 cells without a baseline notch) or a deeper phase 1 notch (5/9 cells with a baseline notch). The voltage of the action potential plateau (measured 25 ms after the peak) was consistently lower after addition of 11 ± 1 pA/pF modeled $I_{to,f}$ although this failed to reach statistical significance (18 ± 3 mV vs 7 ± 6 mV, $p=0.1$). There was, however, a clear relationship between the baseline plateau voltage and the magnitude of the effect resulting from $I_{to,f}$ addition (Figure 6); cells with a more depolarized plateau voltage exhibited little effect of $I_{to,f}$ addition whereas cells with a less depolarized plateau voltage exhibited a marked decrease in plateau voltage after addition of $I_{to,f}$. The effect of $I_{to,f}$ on action potential duration was similarly heterogeneous; 3 of 9 cells exhibited no effect while the remaining cells exhibited action potential shortening (APD90 = $75\pm 5\%$ of baseline after addition of 11 ± 1 pA/pF modeled $I_{to,f}$). These results demonstrate that the functional role of $I_{to,f}$ is highly dependent on the precise cellular context.

Figure 6. Dynamic current clamp addition of $I_{to,f}$ to iPSC-CM results in cell-specific changes in plateau membrane potential. A. Representative action potential waveform (upper panel) at baseline (black) and after the addition of $I_{to,f}$ (red). The modeled $I_{to,f}$ is shown in the lower panel. B. The effect of modeled $I_{to,f}$ on the plateau membrane potential is greatest in cells with a less depolarized plateau potential at baseline.

Discussion

I_{to} in iPSC-CMs

The utility of iPSC-CMs as a model of adult ventricular cardiomyocytes naturally depends on the degree to which they recapitulate the electrophysiological properties of adult human cardiomyocytes. Previous work has demonstrated that iPSC-CMs express most of the major ionic conductances present in

adult cardiomyocytes and that the time- and voltage-dependent properties of most (12) but not all (27) ionic currents are similar to the currents recorded in adult human ventricular myocytes. Previous characterization of I_{to} in iPSC-CMs by Cordeiro et al. reported a strikingly slow recovery from inactivation ($\tau=2380$ ms at 37°C , corresponding to approximately 8000 ms at room temperature(23)).(28) In contrast, we observed a recovery timecourse which, albeit slow, was significantly faster than that reported by Cordeiro et al., with a time constant of 3735 ms at room temperature. Of note, Cordeiro et al. used embryoid body differentiation whereas this study used monolayer differentiation; this observation highlights the potential for physiological differences in cells arising from different protocols.

Both this study and Cordeiro et al. found an absence of the accessory subunit KCHIP2 in iPSC-CMs; this study further demonstrated that over-expression of KCHIP2 resulted in a rapidly recovering component of I_{to} . It has been previously shown that KCHIP2 is an essential subunit underlying $I_{to,f}$ in the mouse.(21) The present results provide the most direct evidence to date of a similar essential role of KCHIP2 in the generation of human $I_{to,f}$. Both studies confirm the absence of functional $I_{to,f}$ in iPSC-CMs, highlighting a limitation of this model for adult cardiomyocytes. In both canine and rodent hearts, KCHIP2 expression increases during post-natal development in parallel with the emergence of $I_{to,f}$; it is therefore plausible that iPSC-CMs reflect an earlier developmental state.(29)

Action potential morphology in iPSC-CMs

Multiple previous studies have reported immature action potential properties in iPSC-CMs as well as a depolarized resting membrane potential.(11, 12) Previous theoretical(28) and experimental(30) studies have suggested that restoration of normal resting membrane potential would normalize action potential waveforms. In this study, dynamic current clamp was used to introduce modelled I_{K1} which restored a normal resting membrane potential, rapid phase 0 upstroke, rapid phase 1 repolarization and reduced the action potential duration. These results complement those of Bett et al.(30) who reported that two distinct AP waveforms (atrial and ventricular) were recorded after introduction of I_{K1} whereas we observed only ventricular waveforms. One possible explanation for this difference is that Bett et al. studied iCell (Cellular Dynamics) cardiac myocytes (derived from spontaneous differentiation in embryoid bodies) whereas this study used directed cardiomyocyte differentiation. These results highlight the fact that, although introduction of I_{K1} can largely normalize action potential waveforms, different cardiomyocyte differentiation protocols potentially yield cells with different properties.

Functional role of $I_{to,f}$

The fast, transient outward potassium current, $I_{to,f}$, plays an important albeit complex role in cardiomyocyte physiology and pathophysiology. Computational simulations have predicted that increasing $I_{to,f}$ conductance approximately 1.5-fold results in moderate action potential prolongation and slightly depolarized membrane potential during the action potential plateau while further increases result in action potential collapse.(7) In contrast, a dynamic current clamp study on canine ventricular cardiomyocytes reported essentially no impact on either plateau voltage or APD; the effect of modelled I_{to} was confined to the phase 1 repolarization.(9) Here, we show that introducing modeled $I_{to,f}$ to human iPSC-CMs consistently affects the phase 1 repolarization but that the effect on plateau voltage varies significantly among cells and is sensitive to the baseline plateau voltage (which in turn reflects the baseline conductances in the particular cell). This finding, in conjunction with previously published results, highlight the fact that the functional role of $I_{to,f}$ is exquisitely sensitive to the balance of other ionic currents in the cell and is likely to be cell-type specific.

Acknowledgements

The authors thank Dr. Michael Pasque for surgical assistance with the collection of human cardiac tissue. The authors thank Translational Cardiovascular Biobank and Repository at Washington University School of Medicine for access to human cardiac tissue.

References

1. Nerbonne JM, Kass RS. Molecular physiology of cardiac repolarization. *Physiol Rev.* 2005;85(4):1205-53.
2. Antzelevitch C. Brugada syndrome. *Pacing Clin Electrophysiol.* 2006;29(10):1130-59.
3. Janse MJ. Electrophysiological changes in heart failure and their relationship to arrhythmogenesis. *Cardiovasc Res.* 2004;61(2):208-17.
4. Bassani RA. Transient outward potassium current and Ca²⁺ homeostasis in the heart: beyond the action potential. *Brazilian journal of medical and biological research = Revista brasileira de pesquisas medicas e biologicas / Sociedade Brasileira de Biofisica [et al].* 2006;39(3):393-403.
5. Decker KF, Heijman J, Silva JR, Hund TJ, Rudy Y. Properties and ionic mechanisms of action potential adaptation, restitution, and accommodation in canine epicardium. *Am J Physiol Heart Circ Physiol.* 2009;296(4):H1017-26.
6. Dong M, Sun X, Prinz AA, Wang HS. Effect of simulated I_(to) on guinea pig and canine ventricular action potential morphology. *Am J Physiol Heart Circ Physiol.* 2006;291(2):H631-7.
7. Greenstein JL, Wu R, Po S, Tomaselli GF, Winslow RL. Role of the calcium-independent transient outward current I_(to1) in shaping action potential morphology and duration. *Circ Res.* 2000;87(11):1026-33.
8. Sah R, Ramirez RJ, Oudit GY, Gidrewicz D, Trivieri MG, Zobel C, et al. Regulation of cardiac excitation-contraction coupling by action potential repolarization: role of the transient outward potassium current (I_(to)). *J Physiol.* 2003;546(Pt 1):5-18.
9. Sun X, Wang HS. Role of the transient outward current (I_{to}) in shaping canine ventricular action potential--a dynamic clamp study. *J Physiol.* 2005;564(Pt 2):411-9.
10. Sinnecker D, Goedel A, Laugwitz KL, Moretti A. Induced pluripotent stem cell-derived cardiomyocytes: a versatile tool for arrhythmia research. *Circ Res.* 2013;112(6):961-8.
11. Knollmann BC. Induced pluripotent stem cell-derived cardiomyocytes: boutique science or valuable arrhythmia model? *Circ Res.* 2013;112(6):969-76; discussion 76.
12. Ma J, Guo L, Fiene SJ, Anson BD, Thomson JA, Kamp TJ, et al. High purity human-induced pluripotent stem cell-derived cardiomyocytes: electrophysiological properties of action potentials and ionic currents. *Am J Physiol Heart Circ Physiol.* 2011;301(5):H2006-17.
13. Fusaki N, Ban H, Nishiyama A, Saeki K, Hasegawa M. Efficient induction of transgene-free human pluripotent stem cells using a vector based on Sendai virus, an RNA virus that does not integrate into the host genome. *Proceedings of the Japan Academy Series B, Physical and biological sciences.* 2009;85(8):348-62.
14. Lian X, Zhang J, Azarin SM, Zhu K, Hazeltine LB, Bao X, et al. Directed cardiomyocyte differentiation from human pluripotent stem cells by modulating Wnt/beta-catenin signaling under fully defined conditions. *Nat Protoc.* 2013;8(1):162-75.
15. Tohyama S, Hattori F, Sano M, Hishiki T, Nagahata Y, Matsuura T, et al. Distinct metabolic flow enables large-scale purification of mouse and human pluripotent stem cell-derived cardiomyocytes. *Cell Stem Cell.* 2013;12(1):127-37.
16. Hattori F, Chen H, Yamashita H, Tohyama S, Satoh YS, Yuasa S, et al. Nongenetic method for purifying stem cell-derived cardiomyocytes. *Nature methods.* 2010;7(1):61-6.
17. Foeger NC, Marionneau C, Nerbonne JM. Co-assembly of Kv4 {alpha} subunits with K⁺ channel-interacting protein 2 stabilizes protein expression and promotes surface retention of channel complexes. *J Biol Chem.* 2010;285(43):33413-22.
18. Yang KC, Foeger NC, Marionneau C, Jay PY, McMullen JR, Nerbonne JM. Homeostatic regulation of electrical excitability in physiological cardiac hypertrophy. *J Physiol.* 2010;588(Pt 24):5015-32.

19. Lin RJ, Bettencourt J, Wha lte J, Christini DJ, Butera RJ. Real-time experiment interface for biological control applications. *Conf Proc IEEE Eng Med Biol Soc.* 2010;2010:4160-3.
20. O'Hara T, Virag L, Varro A, Rudy Y. Simulation of the undiseased human cardiac ventricular action potential: model formulation and experimental validation. *PLoS computational biology.* 2011;7(5):e1002061.
21. Foeger NC, Wang W, Mellor RL, Nerbonne JM. Stabilization of Kv4 protein by the accessory K(+) channel interacting protein 2 (KChIP2) subunit is required for the generation of native myocardial fast transient outward K(+) currents. *J Physiol.* 2013;591(Pt 17):4149-66.
22. Xu H, Guo W, Nerbonne JM. Four kinetically distinct depolarization-activated K⁺ currents in adult mouse ventricular myocytes. *J Gen Physiol.* 1999;113(5):661-78.
23. Nabauer M, Beuckelmann DJ, Uberfuhr P, Steinbeck G. Regional differences in current density and rate-dependent properties of the transient outward current in subepicardial and subendocardial myocytes of human left ventricle. *Circulation.* 1996;93(1):168-77.
24. Wettwer E, Amos GJ, Posival H, Ravens U. Transient outward current in human ventricular myocytes of subepicardial and subendocardial origin. *Circ Res.* 1994;75(3):473-82.
25. Patel SP, Campbell DL. Transient outward potassium current, 'I_{to}', phenotypes in the mammalian left ventricle: underlying molecular, cellular and biophysical mechanisms. *J Physiol.* 2005;569(Pt 1):7-39.
26. Zhang J, Wilson GF, Soerens AG, Koonce CH, Yu J, Palecek SP, et al. Functional cardiomyocytes derived from human induced pluripotent stem cells. *Circ Res.* 2009;104(4):e30-41.
27. Jonsson MK, Vos MA, Mirams GR, Duker G, Sartipy P, de Boer TP, et al. Application of human stem cell-derived cardiomyocytes in safety pharmacology requires caution beyond hERG. *Journal of molecular and cellular cardiology.* 2012;52(5):998-1008.
28. Cordeiro JM, Nesterenko VV, Sicouri S, Goodrow RJ, Jr., Treat JA, Desai M, et al. Identification and characterization of a transient outward K⁺ current in human induced pluripotent stem cell-derived cardiomyocytes. *J Mol Cell Cardiol.* 2013;60:36-46.
29. Nerbonne JM. Regulation of voltage-gated K⁺ channel expression in the developing mammalian myocardium. *J Neurobiol.* 1998;37(1):37-59.
30. Bett GC, Kaplan AD, Lis A, Cimato TR, Tzanakakis ES, Zhou Q, et al. Electronic "expression" of the inward rectifier in cardiocytes derived from human-induced pluripotent stem cells. *Heart Rhythm.* 2013;10(12):1903-10.











