

1 **Isolation and Gene Flow in a Speciation Continuum in Newts**

2

3 Maciej Pabijan^{a,b,*}, Piotr Zieliński^a, Katarzyna Dudek^a, Michał Stuglik^{a,c}, Wiesław Babik^{a*}

4 ^a *Institute of Environmental Sciences, Jagiellonian University, ul. Gronostajowa 7, 30-387,*

5 *Kraków, Poland* katarzyna2.dudek@uj.edu.pl, piotr.zielinski@uj.edu.pl

6 michal.stuglik@uj.edu.pl

7 ^b Permanent address: *Department of Comparative Anatomy, Institute of Zoology and Biomedical*

8 *Research, Jagiellonian University, ul. Gronostajowa 9, 30-387, Kraków, Poland*

9 ^c *Scotland's Rural College, Integrative Animal Sciences, Easter Bush Campus, Midlothian*

10 *EH25 9RG, Scotland, UK*

11 * Corresponding authors:

12 Maciej Pabijan, Department of Comparative Anatomy, Institute of Zoology and Biomedical

13 Research, Jagiellonian University, ul. Gronostajowa 9, 30-387, maciej.pabijan@uj.edu.pl

14 Wiesław Babik, Institute of Environmental Sciences, Jagiellonian University, ul. Gronostajowa

15 7, 30-387, Kraków, Poland; wieslaw.babik@uj.edu.pl

16 **Abstract**

17 Because reproductive isolation often evolves gradually, differentiating lineages may retain the
18 potential for genetic exchange for prolonged periods, providing an opportunity to quantify
19 and understand the fundamental role of gene flow during speciation. Here we delimit taxa,
20 reconstruct the phylogeny and infer gene flow in newts of the *Lissotriton vulgaris* species
21 complex based on 74 nuclear markers sampled from 127 localities. We demonstrate that
22 distinct lineages along the speciation continuum in newts exchange nontrivial amounts of
23 genes, affecting their evolutionary trajectories. By integrating a wide array of methods, we
24 delimit nine taxa and show that two principle factors have driven their genetic differentiation:
25 time since the last common ancestor determining levels of shared ancestral polymorphism,
26 and shifts in geographic distributions determining the extent of secondary contact. Post-
27 divergence gene flow, indicative of evolutionary non-independence, has been most extensive
28 between sister and non-sister taxa in Central Europe, while four southern European lineages
29 have acquired the population genetic hallmarks of independent species (*L. graecus*, *L.*
30 *kosswigi*, *L. lantzi*, *L. schmidtleri*). We obtained strong statistical support for widespread
31 mtDNA introgression, previously suggested by discordance between mtDNA phylogeny and
32 morphology. Our study suggests that long-term evolution in structured populations that may
33 periodically exchange genes may be common: although some of these populations may
34 become extinct or fuse, others will acquire complete reproductive isolation and will carry
35 signatures of this complex history in their genomes.

36 Key Words: delimitation, reticulate evolution, taxonomy, amphibian

37 1. Introduction

38 Speciation, regardless of the mechanisms or geographic settings in which it occurs, is
39 typically a gradual process (Coyne and Orr, 2004). The consequences of this are manifold.
40 First, we see various stages of speciation in nature which makes delimitation of separately
41 evolving units challenging (Huang and Knowles, 2016; Wiens, 2007). Second, completion of
42 speciation is by no means certain, other outcomes such as fusions or extinction are also likely
43 (e.g. Rhymer and Simberloff, 1996; Rudman and Schluter, 2016). Gene flow may
44 homogenize diverging gene pools (Petit and Excoffier, 2009; Slatkin, 1985), thereby limiting
45 or even reversing differentiation. Third, it is unclear how much gene flow occurs between
46 differentiating lineages, whether contact between such lineages causes massive introgression,
47 and to what extent introgression is heterogeneous throughout the genome (Cruikshank and
48 Hahn, 2014; Wolf and Ellegren, 2016).

49 Although theory provides some guidelines, the preceding questions are mostly
50 empirical and need to be addressed by studying natural systems. Recent spectacular adaptive
51 radiations may be less informative and more exceptional in this respect than commonly
52 assumed, because extinction may not have yet had sufficient time to prune the more
53 ephemeral products of divergent selection (Cutter and Gray, 2016). Therefore, careful
54 analyses of taxa composed of metapopulation lineages at various stages of divergence (de
55 Queiroz, 2007), which can be referred to as a speciation continuum, are needed. Because the
56 necessary analyses lie at the interface of population genetics and phylogenetics, they require
57 integration of tools from both fields. This is especially important for studies that explicitly
58 incorporate gene flow into the equation, because population genetic approaches for inferring
59 post-divergence gene flow are better developed than methods co-estimating phylogeny and
60 gene flow (Sousa and Hey, 2013). Likewise, accounting for post-divergence gene flow is a
61 major challenge for current species delimitation methods (Petit and Excoffier, 2009).

62 European newts from the *Lissotriton vulgaris* group provide an empirical example of a
63 speciation continuum influenced by genetic exchange between taxa. The Carpathian newt, *L.*
64 *montandoni* (Boulenger, 1880) and the smooth newt, *Lissotriton vulgaris* (Linnaeus, 1758),
65 have parapatric distributions in central Europe (Fig. 1a; Macgregor et al., 1990; Rafiński and
66 Arntzen, 1987). The former is endemic to the Carpathian and easternmost Sudetes mountains,
67 whereas the latter has a very large range in Central and Northern Europe, with a disjunct
68 population surrounding the Greater Caucasus mountains (Fig. 1a). At the intraspecific level,
69 *L. montandoni* is morphologically uniform across its range, reflected by only shallow genetic
70 substructuring (Zieliński et al., 2014). In sharp contrast, *L. vulgaris* is morphologically
71 differentiated into at least seven subspecies (Fig. 1a; Raxworthy, 1990). The smooth newt is
72 found in a variety of habitats across the collective ranges of the subspecies (Bour et al., 2002;
73 Bousbouras and Ioannidis, 1997; Schmidtler and Franzen, 2004; Skorinov et al., 2008) from
74 deciduous woodlands, farmlands and pastures in European lowlands and mountainous
75 regions, forest-steppe and taiga in western Siberia and Kazakhstan and pockets of humid
76 habitat in otherwise dry woodland or scrub in southern Europe and Anatolia. Despite over a
77 century of study, the status of the *L. vulgaris* subspecies is still controversial (Dubois and
78 Raffaëlli, 2009; Schmidtler and Franzen, 2004; Speybroeck et al., 2010; Wielstra et al., 2015).
79 The disagreement has been exacerbated by a general lack of concordance between
80 morphologically assessed subspecific boundaries and spatial patterns of mtDNA variation
81 (Fig. 1a, Fig. S1; Babik et al., 2005; Nadachowska and Babik, 2009; Pabijan et al., 2015). The
82 traits used to distinguish the smooth newt subspecies pertain almost exclusively to male
83 secondary sexual characters, which include the presence/absence and extent of dorsal crests
84 and tail fins, toe flaps, tail filament, dorso-lateral ridges and pigmentation patterns
85 (Raxworthy, 1990) that develop in aquatic habitats during the breeding season. Despite
86 divergent courtship behavior (Pecio and Rafiński, 1985) that leads to strong but incomplete

87 prezygotic sexual isolation between *L. montandoni* and *L. vulgaris* (Michalak and Rafiński,
88 1999; Michalak et al., 1997), these two species hybridize in and around the Carpathian
89 mountains forming a bimodal hybrid zone (Babik and Rafiński, 2004; Babik et al., 2003;
90 Kotlík and Zavadil, 1999). Hybridization has led to the replacement of the original mtDNA of
91 *L. montandoni* by introgressed mtDNA of several *L. vulgaris* lineages (Babik et al., 2005;
92 Zieliński et al., 2013). Moreover, genetic exchange among the different subspecies of *L.*
93 *vulgaris* is suggested because five of the subspecific taxa do not form monophyletic mtDNA
94 lineages (Fig. S1; Babik et al., 2005; Nadachowska and Babik, 2009; Pabijan et al., 2015).
95 Areas of subspecies intergradation with morphological intermediates have been reported in
96 central and southeastern Europe (e.g. Krizmanic et al., 1997; Schmidtler and Franzen, 2004;
97 Schmidtler and Schmidtler, 1983; Wielstra et al., 2015). These findings suggest that
98 introgressive hybridization both between *L. vulgaris* and *L. montandoni*, and among the
99 various subspecies of *L. vulgaris*, has the potential to blur boundaries among the different
100 taxa.

101 Here, we provide a multi-faceted perspective on divergence and gene flow in the *L.*
102 *vulgaris* complex. First, we postulate that morphologically diagnosable taxa within the
103 complex constitute separately evolving metapopulation lineages (de Quieroz, 2007) and test
104 this assertion using a clustering approach based on allele frequencies, a Bayesian multispecies
105 coalescent delimitation method, and the genealogical sorting index. Second, we hypothesize
106 that gene flow has blurred phylogenetic relationships and misled phylogeny reconstruction
107 methods that do not take reticulation explicitly into account (Leaché et al., 2014; Solís-Lemus
108 et al., 2016). Third, we predict that the extent of historical gene flow was most extensive
109 among lineages inhabiting Central Europe, and use an Approximate Bayesian Computation
110 framework to model gene flow between pairs of taxa. Finally, we apply the inferred nuclear
111 phylogeny to statistically test the previously proposed hypothesis of extensive mtDNA

112 introgression among morphologically diagnosable taxa. We demonstrate that isolation in
113 space and time as well as post-divergence gene flow shape the genetic background of a set of
114 metapopulations that have diverged in ecology and morphology including male epigamic
115 traits.

116 **2. Material and Methods**

117 2.1 Newt sampling

118 Tissues (tail-tips) were sampled from 127 newt populations (i.e. breeding sites, Table S1 and
119 Fig. 1b). Our sampling focused on capturing the genetic variation present in all
120 morphologically defined taxa within the *L. vulgaris* group and all potential Pleistocene
121 refugia (Babik et al., 2005; Nadachowska and Babik, 2009; Pabijan et al., 2015) and was
122 therefore mostly limited to southeastern and central Europe. Because the results of the current
123 study support the elevation of four taxa to the species level (see Discussion), we use this
124 updated nomenclature throughout the text. One individual was sampled per locality.
125 Altogether we analysed sequence data for 128 individuals (plus outgroup) which included 42
126 reported in Zieliński et al. (2016) and 87 sequenced for this study (Table S1).

127 2.2 Laboratory procedures and datasets

128 DNA from ethanol preserved tissues was extracted using either a standard phenol/chloroform
129 technique or the Wizard Genomic DNA extraction kit (Promega). Amplification and
130 sequencing of a panel of 74 nuclear, mostly 3'UTR markers, followed Zieliński et al. (2014).
131 This procedure resulted in physically phased sequences of both haplotypes for most markers
132 in most individuals. We used four different datasets for analyses (Table S2) i) dataset 1 with
133 sequences of all 74 loci and all 128 individuals, ii) dataset 2 with sequences of all 74 loci but
134 with 118 individuals (excluded 9 admixed individuals and one of two newts from locality
135 Kapakli), iii) dataset 3 with sequences of all 128 individuals but 70 loci (excluded 4 loci with

136 > 10% missing sequences), iv) dataset 4 consisting of SNPs extracted from 71 loci (excluded
137 loci with >10% missing data at nucleotide positions) and 120 individuals (including
138 outgroup).

139 2.3 Species delimitation

140 Species delimitation was based on identifying distinct genetic groups from haplotype
141 frequency data and validating the genetic clusters using multispecies coalescent-based and
142 genealogical sorting methods.

143 Sequence alignments (dataset 1) were converted into input for STRUCTURE v.2.3.4
144 (Pritchard et al., 2000) by coding each haplotype as a unique integer in a custom Python
145 script. If nucleotide sequences contained over 10 Ns or gaps, they were coded as missing data,
146 if they had less than 10 Ns or gaps, then columns with missing data were removed from the
147 alignment. This procedure ensured that haplotype identity was based on nucleotide
148 differences between sequences (and not on e.g. indels or low-quality data) and provided a
149 conservative estimate of the number of haplotypes per locus. STRUCTURE was run under the
150 admixture model ($\alpha = 1.0$) with independent allele frequencies among populations ($\lambda = 1.0$)
151 for K from 1 to 12, iterated 10 times each. The Markov chain in each analysis was set to 1
152 million burnin steps; a further 2 million steps were used for parameter value estimation.
153 STRUCTURE HARVESTER (Earl and vonHoldt, 2012) and CLUMPAK (Kopelman et al.,
154 2015) were used process the STRUCTURE output and to evaluate the most probable number
155 of clusters (K) by examining $\log\text{Pr}(Data)$.

156 We validated the genetic clusters identified by STRUCTURE by applying joint
157 Bayesian species delimitation and species tree estimation using the program BPP v3.1 (Yang,
158 2015). This approach is based on the multispecies coalescent and compares different models
159 of species delimitation and species phylogeny in a Bayesian framework, accounting for

160 incomplete lineage sorting and gene tree-species tree conflict (Rannala and Yang, 2013; Yang
161 and Rannala, 2010, 2014). Input alignments were constructed from dataset 2 by taking 10
162 sequences per marker for each of the 9 STRUCTURE-delimited operational taxonomic units
163 (hereafter OTUs). This sampling strategy was based on the minimum number of sequences
164 available per taxon (imposed by 5 specimens of *L. v. meridionalis*), with random subsampling
165 of 10 sequences (without replacement) in all other taxa. Admixed individuals (as defined by
166 STRUCTURE) were excluded from this analysis. We explored a wide variety of priors on the
167 population size parameters (θ s) and divergence time at the root of the species tree (τ_0) (see
168 Extended Methods in the Supplementary Material).

169 Next, we estimated gene trees in MrBayes 3.2 (Ronquist et al., 2012) and used them as
170 input in Bayesian concordance analysis and an assessment of genealogical sorting (see
171 below). We applied dataset 3; outgroups consisted of a single, randomly chosen haplotype per
172 locus of a *Lissotriton helveticus* individual, or if unavailable due to lack of amplification, a
173 single haplotype from either *L. italicus* (for 3 markers) or *L. boscai* (for 9 markers) from
174 Zieliński et al. (2014). Nucleotide substitution models for each marker were determined in
175 jModeltest 2.1.4 (Darriba et al., 2012) under the Bayesian information criterion. Due to the
176 low phylogenetic information content of many of the markers, we did not partition the
177 alignments into codon positions for those containing coding regions. We implemented two
178 runs each with four chains for 3 million generations with trees and parameters sampled every
179 3000 generations. Convergence was assessed by comparing the $-\ln L$ values between runs and
180 by examining parameter values for a subset of markers in Tracer (Rambaut and Drummond,
181 2007). We discarded half of recorded gene trees as burnin, combined runs and constructed an
182 extended majority tree in MrBayes (allcompat command).

183 We quantified the genealogical distinctiveness of the OTUs by measuring exclusive
184 ancestry across gene trees with the genealogical sorting index (*gsi*) of Cummings et al.

185 (2008). This statistic measures the genealogical exclusivity of a grouping on a scale from 0
186 (random distribution of sequences among groups) to 1 (complete monophyly), and an
187 ensemble statistic (*egsi*) is used to integrate across gene trees. Because the *gsi* is sensitive to
188 disparity in group size, we randomly pruned leaves to a sample size of 10 (the minimum
189 number of sequences available per group). Admixed individuals were also pruned from gene
190 trees. We used the R implementation of the *gsi* (genealogicalSorting v.0.92). The null
191 hypothesis that the degree of exclusive ancestry of each group was observed by chance was
192 evaluated by 10^6 permutations.

193 2.4 Phylogeny estimation

194 Our first phylogenetic approach involved using individuals as taxonomic units and
195 concatenating the sequences of all loci. This procedure necessitates the selection of a single
196 haplotype per individual per marker and becomes non-trivial if that individual is heterozygous
197 at multiple loci, because the choice of which haplotype to concatenate into the matrix may
198 influence both the topology and branch support of resultant trees (Weisrock et al., 2012). For
199 each individual in dataset 2, we randomly merged haplotypes from each gene into two
200 concatenated sequences (i.e., each individual was represented by two entries of 36,918 bp
201 consisting of 74 concatenated markers). We then pooled haplotypes for each candidate species
202 and randomly selected half of the haplotypes from each group for analysis. We followed
203 Weisrock et al. (2012) in conducting ten replicate analyses to assess consistency. We ran
204 mixed-model phylogenetic analyses in MrBayes 3.2 on the CIPRES Science Gateway
205 platform (Miller et al., 2010). We divided the alignments into 74 partitions corresponding to
206 markers and assigned the optimal nucleotide substitution models to these partitions as
207 previously determined by jModeltest. Each analysis consisted of two runs with 8 chains for 20
208 million generations with trees and parameters sampled every 5000 generations; burnin was set
209 at 50%. Convergence between runs was assessed by examining effective samples sizes and

210 parameter values between runs in Tracer, and by comparing topologies in AWTY (Nylander
211 et al., 2008).

212 The second phylogenetic method involved using BUCKy (Ané et al., 2007; Larget et
213 al., 2010) to estimate levels of concordance in reconstructed topologies among the posterior
214 distributions of gene trees generated in MrBayes. Because the number of unique tips per gene
215 tree was prohibitive (see Extended Methods), we pruned the gene trees in the MrBayes
216 posterior distributions by randomly selecting a single tip from each OTU using an R script
217 from Weisrock et al. (2012), modified to accommodate our dataset and subsampling
218 requirements. Admixed individuals were also pruned from gene trees. We tested a range of
219 priors and parameter values for BUCKy (see Extended Methods). We pruned the original
220 gene trees to 10 OTUs (9 candidate species plus outgroup) and repeated the subsampling and
221 pruning routine 50 times, using BUCKy to calculate the primary concordance trees for each
222 replicate. We measured clade support by calculating the mean (\pm SD) sample-wide
223 concordance factors for clades across all 50 replicates. We also measured clade support by
224 computing the frequency with which each clade was recovered over all replicates.

225 We constructed a maximum likelihood tree using TreeMix 1.12 (Pickrell and
226 Pritchard, 2012) which uses a covariance matrix based on allele frequencies of SNPs to infer
227 the evolutionary relationships among pre-specified populations. The algorithm then identifies
228 populations more closely related than predicted by the phylogeny and adds a “migration
229 edge” between them, representing the direction and magnitude of gene flow (Pickrell and
230 Pritchard, 2012). We used the nine OTUs and an outgroup as populations. Note that allele
231 frequencies for our outgroup “population” were based on two gene copies because only one
232 outgroup individual was sequenced per marker (see single gene analyses above). We extracted
233 and sorted biallelic SNPs at each of the 74 markers according to minor allele frequency
234 (MAF). Five SNPs with the highest MAF, i.e. most informative at the level of the entire

235 dataset, were retrieved per marker, under the condition that they contained <10% missing data
236 at a nucleotide position; these positions were then converted to TreeMix format using a
237 custom Python script. A total of 71 out of 74 markers met our criteria, resulting in a matrix of
238 355 SNPs (dataset 4). TreeMix was invoked with $-k$ 5 to account for linkage disequilibrium
239 for SNPs from the same marker. In consecutive analyses, we added from one to three
240 migration edges to allow for gene flow in the tree. We also ran TreeMix with an alternative
241 dataset using only 1 SNP per marker with the highest MAF.

242 2.5 Gene flow among OTUs

243 We first summarized the pattern of shared variation in our dataset by counting the numbers of
244 shared haplotypes between OTUs in a custom Python script using dataset 2. We also used this
245 dataset to calculate d_{xy} values among the new OTUs in DnaSP v5 (Librado and Rozas, 2009).

246 We estimated gene flow between all pairs of OTUs (36 pairwise comparisons) in an
247 Approximate Bayesian Computation (ABC) framework. We followed Zielinski et al. (2016)
248 in removing all fully coding markers from analyses, excluding all missing data by removing
249 loci or individuals in pairwise comparisons and excluding all admixed individuals; final
250 datasets included 60-66 loci with between 5 and 24 sequences per taxon (Table S3). The
251 information present in the data was summarized for each pairwise comparison with a set of 9
252 summary statistics (Table S3 and Extended Methods).

253 We tested four simple scenarios of pairwise species divergence and the extent and
254 timing of interspecific gene flow: i) a no gene flow (NGF) model in which an ancestral
255 population splits into two at time T_{SPLIT} with no genetic exchange between descendant
256 populations, ii) an admixture (ADM) model in which a fraction of genes is moved
257 instantaneously from one population to the other; this can occur once independently in each
258 direction at any time after divergence, iii) a constant gene flow (CGF) model in which gene

259 flow is continuous and bidirectional between descendent taxa after T_{SPLIT} , and iv) a recent
260 gene flow (RGF) model in which continuous and bidirectional gene flow between descendent
261 taxa was constrained to have occurred in the last 200,000 years (approximately 50,000 newt
262 generations).

263 Coalescent simulations were done in fastsimcoal2.01 (Excoffier et al., 2013), and the
264 ABC analysis was conducted within the ABCtoolbox (Wegmann et al., 2010); detailed prior
265 and parameter values are given in the Extended Methods. A total of 10^6 datasets were
266 simulated under each demographic model from which we retained 1% (10^4) best simulations
267 and computed the marginal likelihood of the observed and retained datasets under the
268 generalized linear model (Leuenberger and Wegmann, 2010). To control the type I error for
269 multiple comparisons, we used a Bonferroni correction and removed models with the nominal
270 $P < 0.0125$. The best fitting model was then selected via Bayes factors.

271 We estimated the power to distinguish between the four models within the ABC
272 framework by randomly picking 1000 pseudo-observed datasets from all simulations
273 generated for each model and checking how often the ABC procedure correctly predicted the
274 true model (the one that produced the dataset). Each pseudo-observed dataset was treated as
275 the observed data and used to calculate marginal densities of all compared models; Bayes
276 factors were used to select the best model. Because identical prior distributions were used for
277 all pairwise comparisons, we conducted a single power analysis using populations with the
278 fewest sequences (*L. kosswigi* - *L. v. meridionalis*). This can be considered the minimum
279 estimate as all other comparisons (with larger sample sizes) will have more power in
280 distinguishing between models.

281 Next, we formally tested whether the sharing of mtDNA haplotypes among *Lissotriton*
282 groups could be accounted for by ILS (incomplete lineage sorting) or is due to hybridization.

283 We assumed that the partially resolved nuclear phylogeny based on the concordant signal
284 from concatenated/BUCKy/TreeMix analyses (further referred to as the species tree) reflects
285 true relationships among the evolutionary lineages within *L. montandoni*/*L. vulgaris* and
286 followed the statistical approach of Joyce et al. (2011). If mtDNA similarities among
287 evolutionary lineages result from ILS, then mtDNA sequences simulated within the true
288 species tree should replicate the observed mtDNA data. If mtDNA introgression has occurred,
289 sequences simulated on alternative species trees, assuming closest relationships between the
290 lineages showing highest mtDNA similarity, should better reproduce the observed data.
291 Because the relationships between the evolutionary lineages have not been resolved
292 completely, both tests were based on four lineages with relationships well-supported by
293 nuclear data: i) N *L. v. vulgaris*, *L. v. ampelensis*, *L. kosswigi* and *L. montandoni*; ii) S *L. v.*
294 *vulgaris*, *L. v. meridionalis*, *L. schmidtleri* and *L. graecus* (Fig. 2). For *L. montandoni* two
295 scenarios were evaluated, allowing introgression from either *L. v. ampelensis* or N *L. v.*
296 *vulgaris*. For each evolutionary lineage we randomly sampled one sequence from each
297 locality for which mtDNA sequences were available (Babik et al., 2005; Pabijan et al., 2015;
298 Zieliński et al., 2013). These sequences were used to calculate parameters (times of
299 divergence and mutation-scaled effective population sizes of extant and ancestral taxa) of
300 competing species trees in BPP v3.1. A thousand gene trees were simulated for each species
301 tree with MCcoal. Sequences were simulated along gene trees using SeqGen (Rambaut and
302 Grass, 1997) assuming the HKA + gamma model of sequence evolution with parameter
303 values estimated in MEGA6 (Tamura et al., 2013). For each scenario mean and minimum
304 uncorrected (p) distances between evolutionary lineages were calculated for 1000 simulated
305 datasets and compared to the observed distances to obtain *P* values for individual
306 comparisons. *P* values from individual tests were combined using Fisher's method.

307 2.6 Data archiving

308 Input data and custom scripts used for data transformation and analysis are available in Dryad
309 digital repository with doi:[to be submitted upon manuscript acceptance]. DNA sequence
310 alignments for all markers were submitted to GenBank: accession nos. [to be submitted upon
311 manuscript acceptance]. Additional analyses are provided as Supplementary Data.

312 **3. Results**

313 The largest dataset 1 contained 128 individuals and 74 loci with an average of 72.1 ± 20.29
314 haplotypes/locus (range 29-117) and 1.9 ± 3.87 % missing data per locus. Markers were of
315 similar length (498.9 ± 6.79 bp) and highly polymorphic with an average of 91.8 ± 34.56
316 segregating sites per marker, one third of these (29.1 ± 9.52) were singleton polymorphisms
317 (Table S4).

318 3.1 Species delimitation

319 Bayesian clustering solutions at $K \leq 6$ had low probabilities. $\text{Pr}(K)$ plateaued at $K = 7$ (Fig.
320 S2) and therefore we further only considered analyses at $K = 7-12$ (60 analyses in total). Most
321 of these 60 analyses contained clusters that grouped all individuals of a given subspecies or
322 species together (Fig. 1B, Table S5). The top ten analyses with highest lnP all contained $K = 9$
323 groups with identical arrangements of individuals in clusters. Analyses with $K > 9$ contained
324 at least one cluster that was assigned zero or trivial fractions of the data, indicating that under
325 high K values, the number of meaningful clusters was lower than the assumed K . We thus
326 inferred 9 clusters with high certainty (Fig. 1B, Table S5). Five of these clusters correspond to
327 morphologically designated species or subspecies: *L. montandoni*, *L. graecus*, *L. v.*
328 *meridionalis*, *L. kosswigi* and *L. lantzi*. The nominal subspecies was subdivided into two
329 clusters that we designate North *L. v. vulgaris* (N *L. v. vulgaris*) and South *L. v. vulgaris* (S *L.*
330 *v. vulgaris*), with the former living predominantly to the north and east of the Carpathian arc
331 and the latter to the west and south of the Danube. The genetic cluster encompassing *L.*

332 *schmidleri*, a taxon described from western Anatolia that is morphologically similar to the
333 nominal subspecies, was also found in the southeastern Balkans. The genetic cluster including
334 *L. v. ampelensis* contained four individuals from eastern Slovakia, extreme western Ukraine
335 and Romania classified as *L. v. vulgaris* according to morphology. Following Rosenberg et al.
336 (2001), we consider that the frequency with which each taxon formed an exclusive cluster
337 over all 60 analyses is a reliable measure of its support (Table S5). This ranged from 70% in
338 *L. kosswigi* to 88% in *L. graecus* and *L. schmidleri* and considerably exceeded the frequency
339 with which particular taxa clustered together or were split into two or more groups. Some
340 evidence of additional substructure was found in South *L. v. vulgaris* and *L. graecus* (Fig. S3).

341 Admixture was consistently detected in nine individuals (7.0%; Fig. 1B, see Table S1
342 for estimated admixture proportions), five of which originated from the western part of the
343 range of *L. vulgaris* (admixed N *L. v. vulgaris*/S *L. v. vulgaris*, morphologically classified as
344 *L. v. vulgaris*), a single individual from northern Serbia (*L. v. ampelensis*/S *L. v. vulgaris*,
345 morphological *L. v. vulgaris*), one from Kosovo (S *L. v. vulgaris*/*L. graecus*, morphologically
346 undetermined), and two morphologically undetermined newts from the Romanian Carpathians
347 composed of two (N *L. v. vulgaris*/*L. v. ampelensis*) or three (N *L. v. vulgaris*/*L. v.*
348 *ampelensis*/*L. montandoni*) genetic clusters.

349 We conducted joint inference of species delimitation and phylogeny using BPP 3 with
350 the nine OTUs defined in STRUCTURE. The posterior probabilities of each of these equaled
351 1.00 in all analyses under a wide range of priors and different starting trees (Table 1). All
352 models in the 95% credibility sets contained the nine OTUs. The multispecies coalescent
353 model in BPP recovered a number of different species trees under the applied parameter
354 settings (Table 1); their topologies and posterior probabilities varied depending on the prior
355 specification, and to a limited extent, on seed number. Although the latter may suggest a lack

356 of convergence, species trees with highest support were mostly congruent across analyses.
357 Notably, *L. montandoni* and *L. v. ampelensis* were consistently recovered as sister taxa.

358 We used the gene trees calculated in MrBayes to estimate the genealogical exclusivity
359 of each candidate species. Typically, each genealogy in the post-burnin posteriors had a
360 different topology. This was expected because there was limited phylogenetic information
361 contained within each marker and many tips (257 including outgroup) in the trees.
362 Unsurprisingly, consensus gene trees calculated for each marker were mostly unresolved, and
363 showed high levels of discordance (data not shown). Nonetheless, the majority of the
364 genealogies showed a nonrandom clustering pattern for the delimited *Lissotriton* groups as
365 indicated by the highly significant *egsi* (Table 2). We inferred high exclusivity for *L. kosswigi*
366 and *L. lantzi* and with about 40% of genes monophyletic, and moderate *egsi* values for *L.*
367 *graecus*, *L. montandoni*, *L. schmidtleri* and *L. v. meridionalis*. The values of *egsi* were
368 relatively low but still highly significant for *L. v. ampelensis*, S *L. v. vulgaris* and N *L. v.*
369 *vulgaris*.

370 3.2 Phylogenetic results

371 In concatenation analyses, we found that the individuals within OTUs always formed clades
372 with posterior probabilities (PP) of 1.0, we therefore collapsed individuals into groups and
373 show a consensus tree over all replicates (Fig. 2A) and separately for each of the 10
374 concatenation replicates (Fig. S4). The positions of some clades were consistent across all
375 replicates and were always fully supported (PP = 1.0). These included the basal placement of
376 *L. montandoni*, the subsequent split of clade 1 composed of *L. graecus* and *L. kosswigi*, and
377 the sister group relationship of *L. v. meridionalis* and S *L. v. vulgaris*. Moreover, the
378 concatenated analyses invariably identified clade 2 composed of *L. lantzi*, *L. schmidtleri*, S *L.*
379 *v. vulgaris*, N *L. v. vulgaris*, *L. v. ampelensis* and *L. v. meridionalis*. Our replicate analyses

380 differed in the choice of sequences at heterozygous loci during the concatenation process; the
381 results show that this influenced the branching patterns of the concatenated phylogenetic
382 analyses by altering the positions of 4 taxa within clade 2 (*L. lantzi*, *L. schmidtleri*, *N L. v.*
383 *vulgaris* and *L. v. ampelensis*). Nearly all alternative arrangements were strongly supported in
384 one or more concatenation replicates (Fig. S4).

385 A consensus of the 50 replicate primary concordance trees from BUCKy (Fig. 2B)
386 shows that all *L. vulgaris* taxa grouped together (to the exclusion of *L. montandoni*) across all
387 replicates. The mean concordance factor represented by the mean number of genes supporting
388 this clade was 20.5 ± 2.2 out of a total of 70 genes. Clades 1 and 2 as well as the sister group
389 relationship between *S L. v. vulgaris* and *L. v. meridionalis* were relatively well-supported,
390 although concordance factors were generally low (on average <15 genes), suggesting that
391 more than one tree may equally well describe the history of these taxa. We examined this
392 issue by extracting concordance factors from each replicate BUCKy analysis for clades with
393 relatively high support (mostly encompassing those with two terminal taxa, Fig. S5). Primary
394 histories were well-discernible in the cases of *L. montandoni*, *L. v. meridionalis*, *S L. v.*
395 *vulgaris*, *L. lantzi*, *L. graecus* and *L. kosswigi*. However, between 3 and 6 primary histories
396 could be discerned for *N L. v. vulgaris*, *L. v. ampelensis* and *L. schmidtleri*, shown by similar
397 mean concordance factors and largely overlapping quartiles (Fig. S5).

398 The maximum likelihood topologies recovered by TreeMix based on 1 or 5 SNPs per
399 marker (Fig. 2C) were very similar to the consensus topology of the concatenated analysis
400 (Fig. 2A). After adding migration edges, we recovered gene flow between *L. v. ampelensis*
401 and *L. montandoni*, followed by gene flow of lesser magnitude between *L. graecus* and *L.*
402 *schmidtleri* and the ancestor of *S L. v. vulgaris* and *L. v. meridionalis* (Fig. 2D and Fig. S6).

403 3.3 Gene flow among lineages

404 We found a gradient in divergence values (d_{xy}) among newt OTUs ranging between 1.14-
405 1.59% (Fig. 3, Table S6). Each pair of OTUs shared haplotypes at some markers (Fig. 3),
406 although in many cases this amounted to less than 2% and may be indicative of retained
407 ancestral variation. The highest proportions were shared between N *L. v. vulgaris* and *L. v.*
408 *ampelensis* (9.1%), *L. montandoni* and two other lineages (7.5 and 6.7% for N *L. v. vulgaris*-
409 *L. montandoni* and *L. v. ampelensis*-*L. montandoni*, respectively) and three pairs involving S
410 *L. v. vulgaris* and neighboring clades *L. v. ampelensis*, N *L. v. vulgaris* and *L. v. meridionalis*
411 (5.2, 4.6 and 3.3%, respectively). We note that all instances of relatively high levels of
412 haplotype sharing were inferred for Central European taxa that at present occur parapatrically.

413 In most pairwise comparisons of OTUs within the ABC framework at least one model
414 could reproduce the observed data (Tables S7 and S8). The analyses had good power to
415 discriminate between the four models (Table S8). Posterior probabilities for models varied
416 considerably among pairwise taxa comparisons (Fig. 4A) with a clear predominance of the
417 admixture (ADM) model and only 6 comparisons in which recent (in the last 200 kya) or
418 continuous gene flow prevailed, all confined to currently parapatric, Central European
419 lineages. However, in other cases episodes of gene flow in the ADM model were timed to
420 have occurred deep in the past (e.g. modal values for posteriors >1 Ma), making ADM and
421 NGF equivalent in the sense of ruling out recent and constant gene flow (e.g. in *L. kosswigi*-*L.*
422 *v. meridionalis*, *L. kosswigi*-*L. graecus*, *L. kosswigi*-*L. lantzi*, *L. montandoni*-*L. lantzi*, *L.*
423 *montandoni*-S *L. v. vulgaris*; Table S9). In other pairwise comparisons modal values for
424 admixture times in the ADM model were inferred to have occurred relatively recently, close
425 to the upper limit of the prior (e.g. *L. montandoni*-*L. v. ampelensis*, N *L. v. vulgaris*-*L.*
426 *kosswigi*), making the ADM and RGF models comparable. However, the posterior
427 distributions for the admixture time parameters were in general very broad, indicating that
428 data were not very informative in dating admixture events.

429 Phylogenies estimated from mtDNA under the assumption of ILS substantially
430 differed from trees estimated assuming mtDNA introgression (i.e. the topology was
431 incongruent with the species tree based on nuclear data, Fig. 5). Under ILS entire trees and
432 internal branches were both very short. Minimum mtDNA divergence between lineages
433 readily distinguished between ILS and hybridization. For both *L. montandoni* and *L. graecus*,
434 scenarios assuming hybridization fit the data well and were strongly preferred over ILS
435 (Table 3). One of the two hybridization scenarios involving *L. montandoni* fit the data better
436 than the other, although both hybridization scenarios had much better fit than ILS. Apparently
437 the scenario assuming introgression between N *L. v. vulgaris* and *L. montandoni* received
438 higher support because of more extensive mtDNA haplotype sharing between these two than
439 between *L. montandoni* and *L. v. ampelensis*. Mean mtDNA divergence between lineages was
440 not informative in distinguishing between ILS and hybridization as both scenarios generated
441 mean divergence that was not significantly different from the observed values (data not
442 shown).

443 **4. Discussion**

444 This study documents a series of lineages representing a continuum of genetic divergence
445 with ample evidence for historical contact and gene flow between lineages. We demonstrate
446 the importance of geographic conditions (allopatry vs. parapatry) and historical contingencies
447 such as time since divergence and timing of secondary contact in determining the continuum
448 of divergence that can be quantified in the present. Moreover, we show that a combination of
449 population genetic and phylogenetic approaches is required to describe and measure this
450 continuum. In the following sections we discuss our findings for *Lissotriton* newts, their more
451 general implications and outline research areas where methodological advances are needed.

452 **4.1 Robust delimitation despite substantial gene flow**

453 All applied delimitation methods, based on allele frequencies, multispecies coalescent and
454 genealogical concordance, robustly delimited nine evolutionary lineages. Some of these
455 lineages correspond to described morphological subspecies, while others are morphologically
456 cryptic (*N L. v. vulgaris*, *S L. v. vulgaris*) or include two distinct morphologies (*L. v.*
457 *ampelensis*). This result is not an artifact of limited sampling under isolation by distance,
458 because admixed individuals are rare and boundaries between lineages in well-sampled areas
459 are sharp. The delimited entities are distinguishable in both allele frequency-based and
460 genealogical frameworks, but are at various stages of divergence (Fig. 3). Substantial
461 admixture between *N L. v. vulgaris* and *S L. v. vulgaris* in the postglacial expansion area in
462 western Europe indicates that some lineages retain considerable potential for genetic
463 exchange. We found evidence for recent gene flow between five other lineage pairs, all of
464 which occur parapatrically in Central Europe (Figs 1 and 4b; see Zieliński et al., 2016, for a
465 detailed assessment of genetic exchange between *L. montandoni* and two *L. vulgaris*
466 lineages). Moreover, episodes of gene flow between lineages in the more distant past may
467 have been commonplace, given the overall weak support for the divergence with no gene flow
468 model in the ABC analyses (Fig. 4), two cases of confirmed old mtDNA introgression (Table
469 3), and several other instances of mtDNA paraphyly (Babik et al., 2005; Nadachowska and
470 Babik, 2009; Pabijan et al., 2015). Thus, it is by no means certain that all delimited newt
471 lineages will continue to evolve independently. The newt system is a good example of a
472 broader phenomenon: as reproductive isolation progresses, the incipient species experience a
473 prolonged phase of parapatric/allopatric structured populations which may episodically
474 exchange genes. In some instances, secondary contact may lead to the fusion of non-sister
475 lineages over parts of their collective range (e.g. *N L. v. vulgaris* and *S L. v. vulgaris* in
476 western Europe). This interpretation is in line with recent genetic and molecular studies of
477 speciation showing that post-divergence gene flow is common, e.g. it was an important

478 feature of hominin history (Pääbo, 2015), and sometimes occurs at high levels (e.g. Muir et
479 al., 2012; Nevado et al., 2011; Osborne et al., 2016; Sun et al., 2012).

480 4.2 Newt phylogeny and evolution of male nuptial morphology

481 The species tree of the *L. vulgaris* group could not be fully resolved (Fig. 2A). Relationships
482 that received high support included *L. montandoni* as the outgroup to all *L. vulgaris* taxa, *L.*
483 *graecus* and *L. kosswigi* (clade 1) as the sister group to other *L. vulgaris* lineages (clade 2),
484 and the sister group relationship between S *L. v. vulgaris* + *L. v. meridionalis*. Although the
485 monophyly of clade 2 is supported, the relationships among OTUs within this group are
486 uncertain. However, we acknowledge that only one (TreeMix) of the four applied
487 phylogenetic methods specifically modelled gene flow among OTUs. Nonetheless, well-
488 supported nodes in the concatenation and concordance analyses were compatible with the
489 topology of the maximum likelihood tree from TreeMix, suggesting that gene flow among
490 OTUs did not overwhelm the phylogenetic signal over all parts of the tree. However, species
491 tree estimation in BPP under the multispecies coalescent consistently recovered a sister group
492 relationship between *L. montandoni* and *L. v. ampelensis* (Table 1). Likewise, analyses using
493 *BEAST (Heled and Drummond, 2010), which experienced overall convergence issues,
494 produced similar results (data not shown). Given the strong evidence for recent gene flow
495 among these two taxa (Fig. 4; Zieliński et al., 2016), we consider this relationship as highly
496 unlikely. Coalescent-based species tree methods are sensitive to violations of the assumption
497 of no gene flow, particularly between non-sister taxa (Leaché et al., 2013; Solís-Lemus and
498 Ané, 2016), and are thus unsuited to study the relationships within the *L. vulgaris* complex.
499 Given the increasing number of studies reporting reticulate evolution, there is an urgent need
500 for progress in methods co-estimating phylogeny and gene flow in multispecies and
501 multilocus datasets; phylogenetic multilocus network methods (Solís-Lemus and Ané, 2016;
502 Yu et al., 2013, 2014) appear promising in this respect.

503 The species tree for the *L. vulgaris* group nonetheless sheds light on morphological
504 evolution in newts. The male nuptial morphology of *L. montandoni* consists of a smooth and
505 low crest, prominent tail filament and dorso-lateral ridges. Similar male morphologies are also
506 present in the more distantly related *L. helveticus*, *L. italicus* and *L. boscai*. This particular
507 suite of phenotypic traits may be an adaptation for the effective transmission of waterborne
508 pheromones (Pecio and Rafiński, 1985) and could signify the predominance of olfactory cues
509 for species recognition (Secondi et al., 2005). On the other hand, the derived condition of
510 prominent crests and overall larger male body size in most *L. vulgaris* lineages indicates a
511 shift towards visual cues (Secondi et al., 2012), although other explanations are possible such
512 as divergence in female preference for larger male body size (Haerty et al., 2007; Secondi et
513 al., 2010). Whatever the underlying cause(s), our species phylogeny shows that *montandoni*-
514 like male morphology was present in the ancestral *L. vulgaris* population, conserved in the *L.*
515 *graecus/L. kosswigi* clade, and replaced by the derived morphology in the sister clade giving
516 rise to other extant *L. vulgaris* lineages. The intermediate breeding male morphologies of *L. v.*
517 *ampelensis* and *L. v. meridionalis* (Raxworthy, 1990) could be interpreted as independent,
518 partial reversions to the ancestral morphology due to convergent evolution. Alternatively, and
519 possibly more likely given the evidence for substantial gene flow from neighboring taxa,
520 these intermediate morphologies could be the result of an influx of genes underlying the
521 derived morphology and dilution of *montandoni*-like ancestral traits. These two hypotheses
522 could be tested with genome-wide data.

523 4.3 Isolation and gene flow revealed by ABC modeling

524 Recognizing the limitations of existing methodologies, we explored gene flow between pairs
525 of taxa in a model-based framework using Approximate Bayesian Computation (ABC). This
526 approach, although quite simple, has been effective in unraveling the process of gene flow in
527 the newt system. The data had good power in distinguishing among the four scenarios of gene

528 flow and provided quantitative measures of support for the various models. However,
529 analyses modeling gene flow between two populations without incorporation of additional
530 populations have been criticized because ignoring such “ghost” populations may affect
531 parameter estimates or even model choice (Beerli, 2004; Eaton et al., 2015; Slatkin, 2005;
532 Strasburg and Rieseberg, 2010). We note though that pairwise comparisons among all extant
533 taxa may partially alleviate this issue. The availability of all pairwise comparisons allows the
534 identification and reinterpretation of problematic cases, such as support for recent gene flow
535 between allopatric taxa which have not likely been in recent contact, but which both may have
536 exchanged genes with other populations. In our case, inferences were largely consistent with
537 the geographic distributions of the lineages (parapatry vs. allopatry) and we accept them as
538 working hypotheses. In general, given the current state of the field, the use of simple pairwise
539 models in complexes of interbreeding species may be useful because they offer a feasible way
540 of distinguishing among different scenarios of gene flow and building more complex, but still
541 testable, hypotheses.

542 In addition to model selection, ABC estimates demographic parameters which
543 facilitate the interpretation of evolutionary history along the speciation continuum. Estimates
544 of long-term, coalescent effective population sizes (N_e) for the *Lissotriton* lineages are quite
545 large, ranging roughly between 80,000 and 800,000 (Table S9). Independent assessments
546 place the divergence time of *L. montandoni* and *L. vulgaris* between 3.7 to 6.3 mya (Stuglik
547 and Babik, 2016), while mtDNA lineages within *L. vulgaris* split in close succession during
548 the late Pliocene and early Pleistocene (Babik et al., 2005; Pabijan et al., 2015). This
549 timeframe suggests that most *Lissotriton* lineages have not attained the $4N_e$ generations when
550 roughly half of the loci in the genome achieve reciprocal monophyly (Hudson and Coyne,
551 2002; Degnan and Rosenberg, 2009), thus widespread incomplete lineage sorting (Table 2)
552 and the presence of identical haplotypes amongst all lineage pairs are expected (Fig. 3).

553 However, the lineages exchanging genes most extensively (*L. v. ampelensis*, *N L. v. vulgaris*,
554 *S L. v. vulgaris*, *L. v. meridionalis*) also have the largest N_e 's (Fig. 4B, Table S9). This
555 suggests that post-divergence gene flow contributed to elevated N_e estimates, although long-
556 term population structure within lineages may also inflate N_e . We postulate that the existence
557 of a transitional phase, in which differentiated lineages exchange genes, may explain or
558 contribute to the large ancestral population sizes inferred by various methods (IM, ABC,
559 PSMC) in many species (e.g. Duvaux et al., 2011; Nadachowska-Brzyska et al., 2015; Won et
560 al., 2005).

561 4.4 Introgression explains mtDNA paraphyly

562 With a partially resolved phylogeny, we were able to test statistically for mtDNA
563 introgression suggested previously (Babik et al., 2005; Nadachowska and Babik, 2009;
564 Pabijan et al., 2015). We confirmed introgression in two of the most striking cases (from *N L.*
565 *v. vulgaris* into *L. montandoni* and *S L. v. vulgaris* into *L. graecus*), effectively explaining the
566 discordance between mtDNA, nuclear genes and morphology, and attributing it to extensive
567 recent and ancient mtDNA introgression. While there is evidence that it has been
568 accompanied by introgression of nuclear genes in the case of *L. montandoni*, we do not have
569 strong evidence of nuclear introgression into *L. graecus*. Better power to reconstruct the
570 extent and timing of nuclear introgression would be obtained with a larger amount of data,
571 particularly from longer genomic regions to explore information contained in haplotype
572 spectra (Harris and Nielsen, 2013).

573 4.5 Taxonomic implications

574 We note that the application of a rigid taxonomy in which we appropriate scientific names to
575 all nine genetically distinct lineages delimited herein would not fully capture the intricacies of
576 the evolutionary history of *Lissotriton* newts. As stated previously (Wiens, 2007), the

577 delimitation of species poses the problem of thresholding the continuous processes of
578 speciation and lineage amalgamation. Although a number of authors have recognized some of
579 the morphological *L. vulgaris* subspecies as full species (e.g. Dubois and Raffaëlli, 2009;
580 Frost, 2016; Skorinov et al., 2014; Wielstra et al., 2015), these relegations were not based on
581 new data but on inconclusive mtDNA variation, geographic range limits, and morphology.

582 We delimited four distinct southern taxa for which our data strongly suggest species
583 status: *L. v. graecus* as *L. graecus* (Wolterstorff, 1906), *L. v. kosswigi* as *L. kosswigi*
584 (Freitag, 1955), *L. v. lantzi* as *L. lantzi* (Wolterstorff, 1914) and *L. v. schmidtleri* as *L.*
585 *schmidtleri* (Raxworthy, 1988) *L. v.* [and not *L. schmidtlerorum*, see Dubois (2007) and
586 Dubois and Raffaëlli (2009)]. We base this assertion on the concordant signals of divergence
587 and independence from allele frequency- and genealogy-based delimitation methods,
588 genealogical sorting, a nearly complete lack of hybrids in well-sampled areas and little
589 evidence for genetic exchange in the recent past. Moreover, *L. kosswigi* and *L. lantzi* are
590 allopatrically distributed (Skorinov et al., 2014; Wielstra et al., 2015), making gene exchange
591 with other lineages improbable in the near future. More extensive sampling in Bulgaria and
592 eastern Greece is needed to characterize the contact between *L. schmidtleri* and other smooth
593 new lineages. On the other hand, four central European *L. vulgaris* lineages are
594 interconnected by nontrivial amounts of gene flow. These taxa show fusion following range
595 expansion in the morphologically cryptic N *L. v. vulgaris* and S *L. v. vulgaris*, extensive
596 genetic exchange with neighboring lineages in *L. v. ampelensis*, and continuous gene flow
597 since divergence between *L. v. meridionalis* and S *L. v. vulgaris*. These results suggest that we
598 may be observing the loss of incipient divergence through hybridization in these taxa.

599 4.6 Conclusions

600 We found that *Lissotriton* newts represent a continuum of genetic divergence with varying
601 levels of shared genetic variation among taxon pairs. This pattern invokes long-term isolation
602 and independent evolution in four southern smooth newt species, genetic exchange across
603 parapatric borders for several taxa in Central Europe, and effective fusion of previously
604 separated gene pools of two lineages in the post-glacial expansion areas of Western Europe.
605 The profound discordance between taxonomic boundaries and mtDNA in *Lissotriton* is
606 explained by extensive mtDNA introgression rather than by other processes. These features
607 make the *L. vulgaris* species group a suitable system to examine the fission/fusion dynamics
608 of the speciation continuum. The newt system is relevant because many species may have
609 experienced such long-term structured populations. However, we also highlight that better
610 methods, jointly modeling incomplete lineage sorting and gene flow, are needed to analyze
611 such systems.

612 **Acknowledgements**

613 We thank Marcin Piwczyński and Ben Wielstra for insightful comments on the manuscript.
614 Piotr Grodek, Marcin Liana and Krystyna Nadachowska-Brzyska offered invaluable help in
615 the field. Pim Arntzen, Konstantinos Sotiropoulos and Ben Wielstra generously shared
616 samples and locality information. The work was funded by the Polish National Science Centre
617 (2012/04/A/NZ8/00662 to W.B) and the Jagiellonian University (DS/WBiNoZ/INoS/762/15).

618

619 **References**

- 620 Ané, C., Larget, B., Baum, D.A., Smith, S.D., Rokas, A., 2007. Bayesian estimation of
621 concordance among gene trees. *Mol. Biol. Evol.* 24, 412–426.
- 622 Babik, W., Rafiński, J., 2004. Relationship between morphometric and genetic variation in
623 pure and hybrid populations of the smooth and Montandon's newt (*Triturus vulgaris* and *T.*
624 *montandoni*). *J. Zool.* 262, 135–143.
- 625 Babik, W., Szymura, J.M., Rafiński, J., 2003. Nuclear markers, mitochondrial DNA and male
626 secondary sexual traits variation in a newt hybrid zone (*Triturus vulgaris*×*T. montandoni*).
627 *Mol. Ecol.* 12, 1913–1930.
- 628 Babik, W., Branicki, W., Crnobrnja-Isailović, J., Cogălniceanu, D., Sas, I., Olgun, K.,
629 Poyarkov, N.A., Garcia-París, M., Arntzen, J.W., 2005. Phylogeography of two European
630 newt species—discordance between mtDNA and morphology. *Mol. Ecol.* 14, 2475–2491.
- 631 Beerli, P., 2004. Effect of unsampled populations on the estimation of population sizes and
632 migration rates between sampled populations. *Mol. Ecol.* 13, 827–836.
- 633 Bour, R., Bringsøe, H., Vanderhaege, M., 2002. *Triturus vulgaris* (Linnaeus, 1758) at its
634 southern limit: distribution on the Peloponnese, Greece, with range extensions from the
635 Central and South Peloponnese. *Alytes* 20, 67–76.
- 636 Bousbouras, D., Ioannidis, Y., 1997. The distribution and habitat preferences of the
637 amphibians of Prespa National Park, in: Crivelli, A.J., Catsadorakis, G. (Eds.), *Lake*
638 *Prespa, Northwestern Greece. A Unique Balkan Wetland*. Springer, the Netherlands, pp.
639 127–133.
- 640 Coyne, J.A., Orr, H.A, 2004. *Speciation*. Sinauer Associates, Sunderland, MA.
- 641 Cruickshank, T.E., Hahn, M.W., 2014. Reanalysis suggests that genomic islands of speciation
642 are due to reduced diversity, not reduced gene flow. *Mol. Ecol.* 23, 3133–3157.
- 643 Cummings, M.P., Neel, M.C., Shaw, K.L., 2008. A genealogical approach to quantifying

- 644 lineage divergence. *Evolution* 62, 2411–2422.
- 645 Darriba, D., Taboada, G.L., Doallo, R., Posada, D. 2012. jModelTest 2: more models, new
646 heuristics and parallel computing. *Nat. Methods* 9, 772–772.
- 647 Degnan, J.H., Rosenberg, N.A., 2009. Gene tree discordance, phylogenetic inference and the
648 multispecies coalescent. *Trends Ecol. Evol.* 24, 332–340.
- 649 de Queiroz, K., 2007. Species concepts and species delimitation. *Syst. Biol.* 56, 879–886.
- 650 Dubois, A., 2007. Genitives of species and subspecies nomina derived from personal names
651 should not be emended. *Zootaxa* 1550, 49–68.
- 652 Dubois, A., Raffaëlli, J., 2009. A new ergotaxonomy of the family Salamandridae Goldfuss,
653 1820 (Amphibia, Urodela). *Alytes* 26, 1–85.
- 654 Duvaux, L., Belkhir, K., Boulesteix, M., Boursot, P., 2011. Isolation and gene flow: inferring
655 the speciation history of European house mice. *Mol. Ecol.* 20, 5248–5264.
- 656 Earl, D.A., vonHoldt, B.M., 2012. STRUCTURE HARVESTER: a website and program for
657 visualizing STRUCTURE output and implementing the Evanno method. *Conserv. Genet.*
658 *Resour.* 4, 359–361.
- 659 Eaton, D.A., Hipp, A.L., González-Rodríguez, A., Cavender-Bares, J., 2015. Historical
660 introgression among the American live oaks and the comparative nature of tests for
661 introgression. *Evolution* 69, 2587–2601.
- 662 Excoffier, L., Dupanloup, I., Huerta-Sánchez, E., Sousa, V.C., Foll, M., 2013. Robust
663 demographic inference from genomic and SNP data. *PLoS Genet.* 9, e1003905.
- 664 Frost, D.R., 2016. Amphibian Species of the World 6.0, an Online reference. Available at
665 <http://research.amnh.org/vz/herpetology/amphibia>. Accessed December 20, 2016.
- 666 Harris, K., Nielsen, R., 2013. Inferring demographic history from a spectrum of shared
667 haplotype lengths. *PLoS Genet.* 9, e1003521.
- 668 Haerty, W., Gentilhomme, E., Secondi, J., 2007. Female preference for a male sexual trait

- 669 uncorrelated with male body size in the palmate newt (*Triturus helveticus*). Behaviour 144,
670 797–814.
- 671 Heled, J., Drummond, A.J., 2010. Bayesian inference of species trees from multilocus data.
672 Mol. Biol. Evol. 27, 570–580.
- 673 Huang, J-P., Knowles, L.L., 2016. The species versus subspecies conundrum: quantitative
674 delimitation from integrating multiple data types within a single Bayesian approach in
675 Hercules beetles. Syst. Biol. 65, 685–699.
- 676 Hudson, R.R., Coyne, J.A., 2002. Mathematical consequences of the genealogical species
677 concept. Evolution 56, 1557–1565.
- 678 Joyce, D.A., Lunt, D.H., Genner, M.J., Turner, J.F., Bills, R., Seehausen, O., 2011. Repeated
679 colonization and hybridization in Lake Malawi cichlids. Curr. Biol. 21, R108–R109.
- 680 Kotlík, P., Zavadil, V., 1999. Natural hybrids between the newts *Triturus montandoni* and *T.*
681 *vulgaris*: morphological and allozyme data evidence of recombination between parental
682 genomes. Folia Zool. 48, 211–218.
- 683 Kopelman, N.M., Mayzel, J., Jakobsson, M., Rosenberg, N.A., Mayrose, I., 2015. Clumpak: a
684 program for identifying clustering modes and packaging population structure inferences
685 across K. Mol. Ecol. Resour. 15, 1179–1191.
- 686 Krizmanic, I., Mesaroš, G., Džukic, G., Kalezic, M.L., 1997. Morphology of the smooth newt
687 (*Triturus vulgaris*) in former Yugoslavia: taxonomical implications and distribution
688 patterns. Acta Zool. Acad. Sci. H. 43, 345–357.
- 689 Larget, B., Kotha, S.K., Dewey, C.N., Ané, C., 2010. BUCKy: Gene tree / species tree
690 reconciliation with the Bayesian concordance analysis. Bioinformatics 26, 2910–2911.
- 691 Leaché, A.D., Harris, R.B., Rannala, B., Yang, Z., 2014. The influence of gene flow on
692 species tree estimation: a simulation study. Syst. Biol. 63, 17–30.
- 693 Leuenberger, C., Wegmann, D., 2010. Bayesian computation and model selection without

- 694 likelihoods. *Genetics* 184, 243–252.
- 695 Librado, P., Rozas, J., 2009. DnaSP v5: A software for comprehensive analysis of DNA
696 polymorphism data. *Bioinformatics* 25, 1451–1452.
- 697 Macgregor, H.C., Sessions, S.K., Arntzen, J.W., 1990 An integrative analysis of phylogenetic
698 relationships among newts of the genus *Triturus* (family Salamandridae), using
699 comparative biochemistry, cytogenetics and reproductive interactions. *J. Evol. Biol.* 3,
700 329–373.
- 701 Michalak, P., Rafiński, J., 1999. Sexual isolation between two newt species, *Triturus vulgaris*
702 and *T. montandoni* (Amphibia, Urodela, Salamandridae). *Biol. J. Linn. Soc.* 67, 343–352.
- 703 Michalak, P., Grzesik, J., Rafiński, J., 1997. Tests for sexual incompatibility between two
704 newt species, *Triturus vulgaris* and *Triturus montandoni*: no-choice mating design.
705 *Evolution* 51, 2045–2050.
- 706 Miller, M.A., Pfeiffer, W., Schwartz, T., 2010. Creating the CIPRES Science Gateway for
707 inference of large phylogenetic trees. *Proc. Gate. Comp. Environ. Work*, 1–8.
- 708 Muir, G., Dixon, C.J., Harper, A.L., Filatov, D.A., 2012. Dynamics of drift, gene flow, and
709 selection during speciation in *Silene*. *Evolution* 66, 1447–1458.
- 710 Nadachowska, K., Babik, W., 2009 Divergence in the face of gene flow: the case of two
711 newts (Amphibia: Salamandridae). *Mol. Biol. Evol.* 26, 829–841.
- 712 Nadachowska-Brzyska, K., Li, C., Smeds, L., Zhang, G., Ellegren, H., 2015. Temporal
713 dynamics of avian populations during Pleistocene revealed by whole-genome sequences.
714 *Curr. Biol.* 25, 1375–1380.
- 715 Nevado, B., Fazalova, V., Backeljau, T., Hanssens, M., Verheyen, E., 2011. Repeated
716 unidirectional introgression of nuclear and mitochondrial DNA between four congeneric
717 Tanganyikan cichlids. *Mol. Biol. Evol.* 28, 2253–2267.
- 718 Nylander, J.A., Wilgenbusch, J.C., Warren, D.L., Swofford, D.L., 2008. AWTY (are we there

- 719 yet?): a system for graphical exploration of MCMC convergence in Bayesian
720 phylogenetics. *Bioinformatics* 24, 581–583.
- 721 Osborne, O., Chapman, M., Nevado, B., Filatov, D., 2016. Maintenance of species boundaries
722 despite ongoing gene flow in ragworts. *Genome Biol. Evol.* 8, 1038–1047.
- 723 Pabijan, M., Zieliński, P., Dudek, K., Chloupek, M., Sotiropoulos, K., Liana, M., Babik, W.,
724 2015. The dissection of a Pleistocene refugium: phylogeography of the smooth newt,
725 *Lissotriton vulgaris*, in the Balkans. *J. Biogeogr.* 42, 671–683.
- 726 Pääbo, S., 2015. The diverse origins of the human gene pool. *Nat. Rev. Genet.* 16, 313–314.
- 727 Pecio, A., Rafiński, J., 1985. Sexual behaviour of the Montandon's newt, *Triturus montandoni*
728 (Boulenger)(Caudata: Salamandridae). *Amphibia-Reptilia* 6, 11–22.
- 729 Petit, R.J., Excoffier, L., 2009. Gene flow and species delimitation. *Trends Ecol. Evol.* 24,
730 386–393.
- 731 Pickrell, J.K., Pritchard, J.K., 2012. Inference of population splits and mixtures from genome-
732 wide allele frequency data. *PLoS Genet.* 8, e1002967.
- 733 Pritchard, J.K., Stephens, M., Donnelly, P., 2000. Inference of population structure using
734 multilocus genotype data. *Genetics* 155, 945–959.
- 735 Rafiński, J., Arntzen, J.W., 1987. Biochemical systematics of the Old World newts, genus
736 *Triturus*: allozyme data. *Herpetologica* 43, 446–457.
- 737 Rambaut, A., Drummond, A.J., 2007. Tracer v1. 5. <http://tree.bio.ed.ac.uk/software/tracer>
- 738 Rambaut, A., Grass, N.C., 1997. Seq-Gen: an application for the Monte Carlo simulation of
739 DNA sequence evolution along phylogenetic trees. *Comput. Appl. Biosci.* 13, 235–238.
- 740 Rannala, B., Yang, Z., 2013. Improved reversible jump algorithms for Bayesian species
741 delimitation. *Genetics* 194, 245–253.
- 742 Raxworthy, C.J., 1990. A review of the smooth newt (*Triturus vulgaris*) subspecies, including
743 an identification key. *Herpetol. J.* 1, 481–492.

- 744 Rhymer, J.M., Simberloff, D., 1996. Extinction by hybridization and introgression. *Annu.*
745 *Rev. Ecol. Syst.* 27, 83–109.
- 746 Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B.,
747 Liu, L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: efficient Bayesian
748 phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61, 539–
749 542.
- 750 Rosenberg, N.A., Burke, T., Elo, K., Feldman, M.W., Freidlin, P.J., Groenen, M.A.M., Hillel,
751 J., Mäki-Tanila, A., Tixier-Boichard, M., Vignal, A., Wimmers, K., Weigend, S., 2001.
752 Empirical evaluation of genetic clustering methods using multilocus genotypes from 20
753 chicken breeds. *Genetics* 159, 699–713.
- 754 Rudman, S.M., Schluter, D., 2016. Ecological impacts of reverse speciation in threespine
755 stickleback. *Curr. Biol.* 26, 490–495.
- 756 Schmidler, J.F., Franzen, M., 2004. *Triturus vulgaris* (Linnaeus, 1758)–Teichmolch, in:
757 Grossenbacher, K., Thiesemeier, B. (Eds.), *Handbuch der Reptilien und Amphibien*
758 *Europas*. Aula Verlag, Wiesbaden, Germany, pp. 847–967.
- 759 Schmidler, J.J., Schmidler, J.F., 1983. Verbreitung, Ökologie und unterartliche Gliederung
760 von *Triturus vulgaris* in der adriatischen Küstengebieten (Amphibia, Salamandridae).
761 *Spixiana* 6, 229–249.
- 762 Secondi, J., Haerty, W., Lodé, T., 2005. Female attraction to conspecific chemical cues in the
763 palmate newt *Triturus helveticus*. *Ethology* 111, 726–735.
- 764 Secondi, J., Johanet, A., Pays, O., Cazimajou, F., Djalout, Z., Lemaire, C., 2010. Olfactory
765 and visual species recognition in newts and their role in hybridization. *Behaviour* 147,
766 1693–1712.
- 767 Secondi, J., Lepetz, V., Théry, M., 2012. Male attractiveness is influenced by UV
768 wavelengths in a newt species but not in its close relative. *PLoS One* 7, e30391.

- 769 Skorinov, D.V., Kuranova, V.N., Borkin, L.J., Litvinchuk, S., 2008. Distribution and
770 conservation status of the Smooth Newt (*Lissotriton vulgaris*) in Western Siberia and
771 Kazakhstan. Russian J. Herpetol. 15, 157–165.
- 772 Skorinov, D., Doronin, I., Kidov, A., Tuniev, B., Litvinchuk, S., 2014. Distribution and
773 conservation status of the Caucasian newt, *Lissotriton lantzi* (Wolterstorff, 1914). Russian
774 J. Herpetol. 21, 251–268.
- 775 Slatkin, M., 1985. Gene flow in natural populations. Annu. Rev. Ecol. Syst. 16, 393–430.
- 776 Slatkin, M., 2005. Seeing ghosts: the effect of unsampled populations on migration rates
777 estimated for sampled populations. Mol. Ecol. 14, 67–73.
- 778 Solís-Lemus, C., Ané, C., 2016. Inferring phylogenetic networks with maximum
779 pseudolikelihood under incomplete lineage sorting. PLoS Genet. 12, e1005896.
- 780 Solís-Lemus, C., Yang, M., Ané, C., 2016. Inconsistency of species-tree methods under gene
781 flow. Syst. Biol. 65, 843–851.
- 782 Sousa, V., Hey, J., 2013. Understanding the origin of species with genome-scale data:
783 modelling gene flow. Nat. Rev. Genet. 14, 404–414.
- 784 Speybroeck, J., Beukema, W., Crochet, P-A., 2010. A tentative species list of the European
785 herpetofauna (Amphibia and Reptilia)—an update. Zootaxa 2492, 1–27.
- 786 Strasburg, J. L., Rieseberg, L.H., 2010. How robust are “isolation with migration” analyses to
787 violations of the IM model? A simulation study. Mol. Biol. Evol. 27, 297–310.
- 788 Stuglik, M.T., Babik, W., 2016. Genomic heterogeneity of historical gene flow between two
789 species of newts inferred from transcriptome data. Ecol. Evol. 6, 4513–4525.
- 790 Sun, Y., Corcoran, P., Menkis, A., Whittle, C.A., Andersson, S.G., Johannesson, H., 2012.
791 Large-scale introgression shapes the evolution of the mating-type chromosomes of the
792 filamentous ascomycete *Neurospora tetrasperma*. PLoS Genetics 8, e1002820.
- 793 Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S., 2013. MEGA6: molecular

- 794 evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* 30, 2725–2729.
- 795 Wegmann, D., Leuenberger, C., Neuenschwander, S., Excoffier, L., 2010. ABCtoolbox: a
796 versatile toolkit for approximate Bayesian computations. *BMC Bioinformatics* 11, 116.
- 797 Wielstra, B., Bozkurt, E., Olgun, K., 2015. The distribution and taxonomy of *Lissotriton*
798 newts in Turkey (Amphibia, Salamandridae). *ZooKeys* 484, 11–23.
- 799 Wiens, J.J., 2007. Species delimitation: new approaches for discovering diversity. *Syst. Biol.*
800 56, 875–878.
- 801 Weisrock, D.W., Smith, S.D., Chan, L.M., Biebow, K., Kappeler, P.M., Yoder, A.D., 2012.
802 Concatenation and concordance in the reconstruction of mouse lemur phylogeny: an
803 empirical demonstration of the effect of allele sampling in phylogenetics. *Mol. Biol. Evol.*
804 29, 1615–1630.
- 805 Wolf, J.B.W., Ellegren, H., 2016. Making sense of genomic islands of differentiation in light
806 of speciation. *Nat. Rev. Genet.* 18, 87–100.
- 807 Won, Y.J., Sivasundar, A., Wang, Y., Hey, J., 2005. On the origin of Lake Malawi cichlid
808 species: a population genetic analysis of divergence. *P. Natl. Acad. Sci-Biol.* 102, 6581–
809 6586.
- 810 Yang, Z., 2015. The BPP program for species tree estimation and species delimitation. *Curr.*
811 *Zool.* 61, 854–865.
- 812 Yang, Z., Rannala, B., 2010. Bayesian species delimitation using multilocus sequence data. *P.*
813 *Natl. Acad. Sci-Biol.* 107, 9264–9269.
- 814 Yang, Z., Rannala, B., 2014. Unguided species delimitation using DNA sequence data from
815 multiple loci. *Mol. Biol. Evol.* 31, 3125–3135.
- 816 Yu, Y., Barnett, R.M., Nakhleh, L., 2013. Parsimonious inference of hybridization in the
817 presence of incomplete lineage sorting. *Syst. Biol.* 62, 738–751.
- 818 Yu, Y., Dong, J., Liu, K.J., Nakhleh, L., 2014. Maximum likelihood inference of reticulate

- 819 evolutionary histories. *Natl. Acad. Sci-Biol.* 111, 16448–16453.
- 820 Zieliński, P., Nadachowska-Brzyska, K., Wielstra, B., Szkotak, R., Covaciu-Marcov, S.D.,
821 Cogălniceanu, D., Babik, W., 2013. No evidence for nuclear introgression despite
822 complete mtDNA replacement in the Carpathian newt (*Lissotriton montandoni*). *Mol.*
823 *Ecol.* 22, 1884–1903.
- 824 Zieliński, P., Stuglik, M.T., Dudek, K., Konczal, M., Babik, W., 2014. Development,
825 validation and high-throughput analysis of sequence markers in nonmodel species. *Mol.*
826 *Ecol. Resour.* 14, 352–360.
- 827 Zieliński, P., Nadachowska-Brzyska, K., Dudek, K., Babik, W., 2016. Divergence history of
828 the Carpathian and smooth newts modelled in space and time. *Mol. Ecol.* 25, 3912–3928.

829 **Table 1.** Joint Bayesian species delimitation and species tree estimation in the *Lissotriton*
830 *vulgaris* species group using BPP. Rows show results for different analyses; letters denote
831 replicate runs differing only by seed number. Population size (θ) and divergence time (τ)
832 priors encompass a wide range of demographic and divergence scenarios. Starting guide tree
833 topologies were random unless specified otherwise (see footnotes). *N* species shows the
834 numbers of species and their posterior probabilities (PPs) in each analysis (PPs for each
835 species within analyses were always 1.00). *N* trees denotes the number of trees in the 95%
836 credibility set, and the last column shows the topology and posterior probability of the highest
837 ranked tree in the set.

838

839

Runs	Θ^a	τ^a	N species (PP)	N trees	best tree and pp of model (delimitation + tree)
1a	G(1, 35)	G(2, 500)	9 (1.00)	7	(<i>L. graecus</i> , (<i>L. lantzi</i> , ((<i>L. schmidtleri</i> , <i>L. kosswigi</i>), ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.70905
1b	G(1, 35)	G(2, 500)	9 (1.00)	4	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>L. schmidtleri</i> , (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.83335
2a^b	G(1, 35)	G(2, 500)	9 (1.00)	4	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>L. schmidtleri</i> , (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.81770
2b^b	G(1, 35)	G(2, 500)	9 (1.00)	2	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>L. schmidtleri</i> , (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.88070
3a^c	G(1, 35)	G(2, 500)	9 (1.00)	3	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>L. schmidtleri</i> , (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.82170
3b^c	G(1, 35)	G(2, 500)	9 (1.00)	2	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>L. schmidtleri</i> , (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.85825
4a^d	G(1, 35)	G(2, 500)	9 (1.00)	3	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>L. schmidtleri</i> , (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.79815
4b^d	G(1, 35)	G(2, 500)	9 (1.00)	38	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>L. schmidtleri</i> , (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.43710
5a	G(1, 10)	G(2, 2000)	9 (1.00)	9	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , (<i>L. v. meridionalis</i> , (<i>L. schmidtleri</i> , ((<i>L. montandoni</i> , <i>L. v. ampelensis</i>), (<i>N L. v. vulgaris</i> , <i>S L. v. vulgaris</i>)))))); 0.33805
5b	G(1, 10)	G(2, 2000)	9 (1.00)	5	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>L. schmidtleri</i> , (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.71930
6a	G(2,2000)	G(2,2000)	9 (1.00)	50	(<i>L. v. meridionalis</i> , (((<i>L. graecus</i> , <i>L. lantzi</i>), (<i>S L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>))), (<i>N L. v. vulgaris</i> , (<i>L. schmidtleri</i> , <i>L. kosswigi</i>))); 0.13655
6b	G(2,2000)	G(2,2000)	9 (1.00)	45	((<i>L. graecus</i> , <i>L. lantzi</i>), ((<i>L. schmidtleri</i> , <i>L. kosswigi</i>), ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>))))); 0.12370
7a	G(1, 10)	G(1, 10)	9 (1.00)	3	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>L. schmidtleri</i> , (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.66405
7b	G(1, 10)	G(1, 10)	9 (1.00)	5	(<i>L. graecus</i> , (<i>L. lantzi</i> , ((<i>L. schmidtleri</i> , <i>L. kosswigi</i>), ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.77125
8a	G(1, 10)	G(2,1000)	9 (1.00)	3	(<i>L. graecus</i> , (<i>L. lantzi</i> , ((<i>L. schmidtleri</i> , <i>L. kosswigi</i>), ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.80270
8b	G(1, 10)	G(2,1000)	9 (1.00)	9	(<i>L. graecus</i> , (<i>L. lantzi</i> , (<i>L. kosswigi</i> , ((<i>S L. v. vulgaris</i> , <i>L. v. meridionalis</i>), (<i>L. schmidtleri</i> , (<i>N L. v. vulgaris</i> , (<i>L. montandoni</i> , <i>L. v. ampelensis</i>)))))); 0.48025

840 ^a - G(1, 35), G(2, 500): empirically derived priors; G(1, 10), G(2, 2000): large populations

841 with shallow divergence; G(2,2000), G(2,2000): small populations with shallow divergence;

842 G(1, 10), G(1, 10): large populations with deep divergence; G(1, 10), G(2,1000): large

843 populations with intermediate divergence

844 ^b - starting tree according to consensus phylogeny

845 ^c - starting tree reflects gene flow among *L. montandoni*, *L. v. ampelensis* and *N L. v. vulgaris*

846 ^d - starting tree reflects gene flow among *N L. v. vulgaris* and *S L. v. vulgaris*, basal position

847 of *L. lantzi*

848 **Table 2.** Ensemble genealogical sorting index (*egsi*) for *Lissotriton* operational taxonomic
849 units (10 alleles per group) based on 70 gene trees. All of the P values for *egsi* are <0.000001.
850 The last column provides the number of gene trees in which a group reached complete
851 monophyly.

group	egsi	monophyly
<i>L. lantzi</i>	0.692	27
<i>L. kosswigi</i>	0.679	30
<i>L. graecus</i>	0.564	17
<i>L. schmidleri</i>	0.559	14
<i>L. montandoni</i>	0.548	16
<i>L. v. meridionalis</i>	0.497	7
<i>L. v. ampelensis</i>	0.349	0
South <i>L. v. vulgaris</i>	0.343	0
North <i>L. v. vulgaris</i>	0.281	0

852

853 **Table 3.** Evaluation of fit of mtDNA sequence data to scenarios of incomplete lineage sorting
 854 (ILS) and hybridization (Hyb). *Lva* – *L. v. ampelensis*; *Lg* – *L. graecus*; *Lk* – *L. kosswigi*; *Lvm*
 855 – *L. v. meridionalis*; *Lm* – *L. montandoni*; *Ls* – *L. schmidtleri*; *NLvv* – North *L. v. vulgaris*;
 856 *SLvv* – South *L. v. vulgaris*

Scenario	Pair of lineages	Minimum divergence - Observed	Minimum divergence - Expected (median of 1000 simulations)	P	Interpretation
mtDNA introgression into <i>L. montandoni</i>					
ILS	<i>Lm-Lva</i>	0.0020	0.0010	0.258	O = E
	<i>Lm -NLvv</i>	0.0000	0.0020	0.294	O = E
	<i>Lm -Lk</i>	0.0384	0.0020	0.002	O > E
	<i>Lva-NLvv</i>	0.0020	0.0010	0.220	O = E
	<i>Lva-Lk</i>	0.0207	0.0020	0.006	O > E
	<i>NLvv-Lk</i>	0.0374	0.0030	0.002	O > E
	Combined (Fisher's method)			0.00002	data do not fit scenario
Hyb1	<i>Lm -Lva</i>	0.0020	0.0010	0.034	
	<i>Lm -NLvv</i>	0.0000	0.0010	0.500	O = E
	<i>Lm -Lk</i>	0.0384	0.0098	0.046	O > E
	<i>Lva-NLvv</i>	0.0020	0.0010	0.242	O = E
	<i>Lva-Lk</i>	0.0207	0.0098	0.150	O = E
	<i>NLvv-Lk</i>	0.0374	0.0108	0.066	O = E
	Combined (Fisher's method)			0.0095	data do not fit scenario
Hyb2	<i>Lm -Lva</i>	0.0020	0.0020	0.514	O = E
	<i>Lm -NLvv</i>	0.0000	0.0000	0.318	O = E
	<i>Lm -Lk</i>	0.0384	0.0089	0.040	O > E
	<i>Lva-NLvv</i>	0.0020	0.0020	0.824	O = E
	<i>Lva-Lk</i>	0.0207	0.0089	0.110	O = E
	<i>NLvv-Lk</i>	0.0374	0.0098	0.056	O = E
	Combined (Fisher's method)			0.0561	data fit scenario
mtDNA introgression into <i>L. graecus</i>					
ILS	<i>Lg -SLvv</i>	0.0000	0.0049	0.006	O < E
	<i>Lg -Lvm</i>	0.0139	0.0059	0.120	O = E
	<i>Lg -Ls</i>	0.0208	0.0059	0.036	O > E
	<i>SLvv-Lvm</i>	0.0069	0.0039	0.164	O = E
	<i>SLvv-Ls</i>	0.0139	0.0049	0.042	O > E
	<i>Lvm-Ls</i>	0.0396	0.0069	0.022	O > E
	Combined (Fisher's method)			0.0001	data do not fit scenario
Hyb	<i>Lg -SLvv</i>	0.0000	0.0020	0.180	O = E
	<i>Lg -Lvm</i>	0.0139	0.0069	0.086	O = E
	<i>Lg -Ls</i>	0.0208	0.0118	0.212	O = E
	<i>SLvv-Lvm</i>	0.0069	0.0069	0.832	O = E
	<i>SLvv-Ls</i>	0.0139	0.0118	0.560	O = E
	<i>Lvm-Ls</i>	0.0396	0.0138	0.080	O = E
	Combined (Fisher's method)			0.1152	data fit scenario

858 **Figure 1.** Map with ranges of morphological subspecies (A). Delimitation of evolutionary
859 lineages in the *Lissotriton vulgaris* species group in STRUCTURE (B). Pie charts show mean
860 individual cluster membership coefficients. *Lg* – *L. graecus*; *Lk* – *L. kosswigi*; *Ll* – *L. lanzti*;
861 *Lm* – *L. montandoni*; *Ls* – *L. schmidtleri*; *Lva* – *L. v. ampelensis*; *Lvm* – *L. v. meridionalis*;
862 *NLv* – North *L. v. vulgaris*; *SLv* – South *L. v. vulgaris*


863 **Figure 2.** Phylogenetic relationships among STRUCTURE-delimited groups of *Lissotriton*
864 newts; most genetic groups correspond to morphologically defined subspecies (except for *L.*
865 *v. vulgaris* that is split into northeastern and southwestern lineages and *L. v. ampelensis* which
866 contains two morphologies; see text for details). (A) Majority-rule consensus tree based on 10
867 replicate analyses (Fig. S4) of 74 concatenated nuclear loci for a total of 36,918 bp; values
868 designate the frequencies of clades found across replicates. Filled circles at nodes indicate
869 congruence across concatenation, concordance and TreeMix analyses. (B) Majority-rule
870 consensus of 50 primary concordance trees. Frequency of clades over all replicates (top
871 number) and concordance factors shown as the mean number of loci (out of 70) supporting
872 the clade (\pm SD). (C) Maximum likelihood tree as inferred by Treemix based on 355 SNPs;
873 horizontal branch lengths are proportional to the amount of genetic drift that has occurred in
874 each lineage. (D) Inferred evolutionary relationships among newt groups by TreeMix with
875 one (left), two (middle) and three (right) migration events. The migration arrows are coloured
876 according to their weights; the weight is correlated with the ancestry fraction.

877 **Figure 3.** Top: pairwise d_{xy} distances among *Lissotriton* taxa. Bottom: proportion (in percent,
878 left y-axis and solid circles) of shared haplotypes out of total number of haplotypes summed
879 for each taxon pair across 74 markers; values above solid circles indicate the number of
880 alleles shared by each taxon pair. Note that taxon pairs are in the same order on both charts.
881 *Lg* – *L. graecus*; *Lk* – *L. kosswigi*; *Ll* – *L. lanzti*; *Lm* – *L. montandoni*; *Ls* – *L. schmidtleri*;

882 *Lva* – *L. v. ampelensis*; *Lvm* – *L. v. meridionalis*; *NLvv* – North *L. v. vulgaris*; *SLvv* – South *L.*
883 *v. vulgaris*


884 **Figure 4.** Pie charts representing posterior distributions of Approximate Bayesian
885 Computation analyses in pairwise taxon comparisons (A), color coded according to
886 demographic models (inset in A). *Lissotriton vulgaris* group phylogeny showing well-
887 supported instances of post-divergence gene flow (green arrows) and their corresponding
888 posterior distributions from the ABC analyses (B).

889 **Figure 5.** Parameters of species trees under competing scenarios of incomplete lineage
890 sorting (ILS) and hybridization (HYB). Both θ and τ are measured as the expected number of
891 mutations per site.




892

893 **Fig. 1.**





896



897


898 Fig. 3.

899



900

901 **Fig. 4**



902

903 **Fig. 5**

904 **SUPPORTING INFORMATION**

905 **Extended Methods:** Validation of OTUs in BPP, phylogenetic analysis using BUCKy, and
906 ABC analyses.

907 **Table S1.** A list of newt specimens with individual ID numbers, morphological species
908 designation, geographical provenance and estimated proportion of genes in STRUCTURE-
909 defined genetic clusters (c1-c9) representative of the species delimitation proposed in this
910 study.

911 **Table S2.** Datasets used for species delimitation and phylogenetic reconstruction.

912 **Table S3.** Summary statistics for pairwise comparisons of *Lissotriton* taxa used in
913 Approximate Bayesian Computation analyses.

914 **Table S4.** Summary of the nucleotide variation in 74 nuclear markers (mostly 3' UTR
915 regions) in *Lissotriton vulgaris* (including all lineages delimited herein) and *L. montandoni*.

916 **Table S5.** Summary of STRUCTURE results for $K = 7-12$, iterated 10 times each for a total
917 of 60 separate analyses.

918 **Table S6.** Pairwise matrix of d_{xy} values calculated for newt OTUs.

919 **Table S7.** Results of Approximate Bayesian Computation analyses under four demographic
920 models: no gene flow (NGF), constant gene flow after divergence (CGF), recent (<200 kya)
921 gene flow (RGF), and instantaneous admixture after divergence (ADM).

922 **Table S8.** Power analysis for Approximate Bayesian Analyses.

923 **Table S9.** Prior and posterior distributions for best model in pairwise taxon comparisons in
924 *Lissotriton* newts from ABC analyses.

925

926 **Figure S1.** Relationships among major mtDNA lineages (denoted by capital letters) in
927 *Lissotriton*, modified from Pabijan et al. (2015).

928 **Figure S2.** Estimated L_n probability of the data for a given K , from STRUCTURE
929 HARVESTER; ten replicate analyses were run for each K .

930 **Figure S3.** Additional substructure found within South *L. v. vulgaris* and *L. graecus*.

931 **Figure S4.** Consensus trees from 10 replicates of concatenated analyses in MrBayes.

932 **Figure S5.** Variation in concordance factors (expressed as the number of genes supporting a
933 particular clade) for main clades found in 50 replicate BUCKy analyses.

934 **Figure S6.** Scaled residual fit from the TreeMix analysis based on the maximum likelihood
935 tree in Fig. 2C (no migration edges).