Approximation to the distribution of fitness effects across functional categories in human segregating polymorphisms

Fernando Racimo[1,*], Joshua G. Schraiber[1],

1 Department of Integrative Biology, University of California, Berkeley, CA, USA

∗ E-mail: fernandoracimo@gmail.com

Abstract

Quantifying the proportion of polymorphic mutations that are deleterious or neutral is of fundamental importance to our understanding of evolution, disease genetics and the maintenance of variation genome-wide. Here, we develop an approximation to the distribution of fitness effects (DFE) of segregating single-nucleotide mutations in humans. Unlike previous methods, we do not assume that synonymous mutations are neutral or not strongly selected, and we do not rely on fitting the DFE of all new nonsynonymous mutations to a single probability distribution, which is poorly motivated on a biological level. We rely on a previously developed method that utilizes a variety of published annotations (including conservation scores, protein deleteriousness estimates and regulatory data) to score all mutations in the human genome based on how likely they are to be affected by negative selection, controlling for mutation rate. We map this score to a scale of fitness coefficients via maximum likelihood using diffusion theory and a Poisson random field model on SNP data. Our method serves to approximate the deleterious DFE of mutations that are segregating, regardless of their genomic consequence. We can then compare the proportion of mutations that are negatively selected or neutral across various categories, including different types of regulatory sites. We observe that the distribution of intergenic polymorphisms is highly peaked at neutrality, while the distribution of nonsynonymous polymorphisms is bimodal, with a neutral peak and a second peak at $s \approx -10^{-4}$. Other types of polymorphisms have shapes that fall roughly in between these two. We find that transcriptional start sites, strong CTCF-enriched elements and enhancers are the regulatory categories with the largest proportion of deleterious polymorphisms.

Author Summary

The relative frequencies of polymorphic mutations that are deleterious, nearly neutral and neutral is traditionally called the distribution of fitness effects (DFE). Obtaining an accurate approximation to this distribution in humans can help us understand the nature of disease and the mechanisms by which variation is maintained in the genome. Previous methods to approximate this distribution have relied on fitting the DFE of new mutations to a single probability distribution, like a normal or an exponential distribution. Generally, these methods also assume that a particular category of mutations, like synonymous changes, can be assumed to be neutral or nearly neutral. Here, we provide a novel method designed to reflect the strength of negative selection operating on any segregating site in the human genome. We use a maximum likelihood mapping approach to fit these scores to a scale of neutral and negative fitness coefficients. Finally, we compare the shape of the DFEs we obtain from this mapping for different types of functional categories. We observe the distribution of polymorphisms has a strong peak at neutrality, as well as a second peak of deleterious effects when restricting to nonsynonymous polymorphisms.

Introduction

Genetic variation within species is shaped by a variety of evolutionary processes, including mutation, demography, and natural selection. With the advent of whole-genome sequencing, we can make unprecedented inferences about these and other processes by analyzing population genomic data. An important goal is to understand the extent to which segregating genetic variants are impacted by natural selection, and to quantify the intensity of natural selection acting genome-wide. Understanding the prevalence of different modes of selection on a genomic scale has wide-ranging implications across evolutionary and medical genetics. For instance, genome-wide association studies (GWAS) are searching for mutations associated with disease in large samples of humans [1]. Because mutations associated with disease are *a priori* likely to be deleterious, quantifying the portion of mutations that are deleterious along with their average effects could have significant implications for the design and interpretation of GWAS. Moreover, the ENCODE project has recently claimed that much of the genome is involved in some form of functional activity [2,3]. However, the extent to which molecular signals identified by this project are actually produced by biological processes affecting fitness has been disputed [4,5]. Indeed, comparative genomics studies suggest that only a small proportion of the human genome $(5-10\%)$ is under purifying selection,

based on signals detectable on phylogenetic timescales [6–8]. Quantifying the DFE in noncoding regions is a first step toward understanding the fitness implications of functional activity at the genomic level.

Traditionally, studies have sought to estimate the distribution of fitness effects (DFE) for nonsynonymous mutations by using summary statistics based on the number of polymorphisms and substitutions [9–11] and/or the full frequency spectrum [12–14]. These studies typically assumed that synonymous variation is neutral or under weak selection. Many of these analyses suggest that while a large proportion of nonsynonymous mutations are nearly neutral, there is also a significant probability that an amino acid changing mutation will be strongly deleterious. While these studies were limited to analysis of protein-coding genes, recent work has focused on quantifying the DFE in regulatory regions, including short interspersed genomic elements such as enhancers [15,16] and cis-regulatory regions [17]. Reviews of these and other approaches can be found in ref. [18,19].

There are several obstacles to quantifying the DFE of new or segregating mutations genome-wide. First, inferences about the DFE are confounded by demography [20]. For example, a high proportion of low frequency derived alleles is a signature of negative selection, but can also be caused by recent population growth [21]. Hence, a well-supported demographic model must be used to appropriately control for population history when inferring the DFE. Second, most current methods rely on dividing up polymorphisms into either putatively selected sites or putatively neutral (or less selected) sites (for example, nonsynonymous and synonymous sites, respectively). These studies have relied on fitting a demographic moel to the neutral class of sites and then fitting the DFE of new mutations to a probability distribution, typically an exponential or gamma distribution [9,13] to the class of sites that are putatively under selection (e.g. nonsynonymous sites). While flexible, these distributions may miss some important features of the DFE [22]. For example, mutation accumulation experiments suggest that the DFE may be bimodal for at least some species, with most mutations either having nearly neutral or strongly deleterious effects, and very few mutations falling in between [23,24]. Thus, assuming two classes of sites may not serve to capture all the relevant information about the DFE (but see [25] for an example of fitting a multimodal DFE to population genetic data and [22,26] for nonparametric approaches to estimating the DFE of new amino-acid changing mutations). Finally, previous studies have been restricted to analyzing specific subclasses of mutations (e.g. nonsynonymous, enhancers, etc.) because until recently, no single metric existed that could serve to compare the disruptive potential of any type of variant, regardless of its genomic consequence.

Recently, Kircher et al. [27] developed a method to synthesize a large number of annotations into a single score to predict the pathogenicity or disruptive potential of any mutation in the genome. It is based on an analysis comparing real and simulated changes that occurred in the human lineage since the human-chimpanzee ancestor, and that are now fixed in present-day humans. The method relies on the realistic assumption that the set of real changes is depleted of deleterious variation due to the action of negative selection, which has pruned away disruptive variants, while the simulated set is not depleted of such variation. A support vector machine (SVM) was trained to distinguish the real from the simulated changes using a kernel of 63 annotations (including conservation scores, regulatory data and protein deleteriousness scores), and then used to assign a score (C-score) to all possible single-nucleotide changes in the human genome, controlling for local variation in mutation rates. These C-scores are meant to be predictors of how disruptive a given change may be, and are comparable across all types of sites (nonsynonymous, synonymous, regulatory, intronic or intergenic). Thus, they allow for a strict ranking of predicted functional disruption for mutations that may not be otherwise comparable. C-scores are PHRED scaled, with larger values corresponding to more disruptive effects.

Importantly, human-specific genetic variation patterns are not used as input to train the C-score SVM. In this work, we make use of the C-scores to provide a fine-grained stratification of the deleteriousness of variants segregating in modern human populations. We take advantage of the Poisson random field model [28, 29] with a realistic model of human demographic history to fit a maximum likelihood selection coefficient for each C-score, creating a mapping from C-scores to selection coefficients.

Results

A mapping from C-scores to selection coefficients

To map C-scores to selective coefficients, we obtained allele frequency information from 9 Yoruba (YRI) individuals (18 haploid sequences) sequenced to high-coverage using whole-genome shotgun sequencing as part of a dataset produced by Complete Genomics (CG) [30]. We removed sites that had a Duke Uniqueness 20bp-mapability score < 1 (downloaded from the UCSC Genome Browser, [31]), to avoid potential errors due to mismapping or miscalling in regions of the genome that are not uniquely mapable.

When inferring the DFE, we focused only on models of neutral evolution and negative selection, because C-scores are uninformative about adaptive vs. deleterious disruption (i.e. a high C-score could

either reflect a highly deleterious change or a highly adaptive change). Additionally, because we are using

polymorphism data only, positive selection should contribute little to the site-frequency spectrum [32].

We first binned polymorphisms into C-scores rounded to the nearest integer and computed the site

frequency spectrum for each bin (Figure S1). We then fit the lowest possible C-score (C= 0), presumed

to be neutral, to different models of demographic history. We computed the likelihood of the SFS in

this bin for a constant population size model, a range of exponential growth models, the model inferred

by Tennessen et al. [33] and the model inferred by Harris and Nielsen [34] from the distribution of

tracts of identity by state (IBS) (Figure S2), and used an EM algorithm to correct for ancestral state

misidentification (Figure S3, see Materials and Methods). We find that a model of exponential growth

at population-scaled rate $= 1$ for 13,000 generations is the best fit to the corrected SFS, although the

Tennessen model is almost as good a fit (Figure S2).

Using the best-fitting demography, we next fit a range of models with different selection coefficients

to the EM-corrected site frequency spectrum for each C-score bin, using maximum likelihood (Figure

1.A) (see Methods). We restricted to $C \leq 40$, because very few sites have larger C-scores, and hence

estimates of the selection coefficients for those C-scores are unreliable. We tested the robustness of the

mappings to different levels of background selection, by partitioning the data into deciles of B-scores [35]

and re-computing the C-to-s mapping for each decile. We observe that the mapping is generally robust

to background selection, with substantial differences only observed at the lowest two B-score quantiles,

which correspond to high background selection (Figure S4). For this reason, and so as to obtain reliable

DFEs at exonic sites (where background selection is generally higher than in the rest of the genome), we

also performed a neutral demographic fitting and a C-to-s mapping while restricting only to sites in the

exome (Figure 1.C). This mapping has a steeper decline than the genomic mapping, reflecting patterns

of background selection which are not fully controlled by C-scores but that affects the SFS. We therefore

show estimated DFEs using both the genome-wide and the exome-wide fittings below. After removing

the C-score bins that best fit the neutral model, we fit a smoothing spline with 20 degrees of freedom

to the remaining C-scores, so as to produce a continuous mapping of C-scores to selection coefficients

(Figure 1.A).

We were concerned that our binning-based mapping would miss important features about the dis-

tribution of coefficients within each bin. To address this, we also fitted individual gamma distributions

of selection coefficients to each of the bins. We show the mean, standard deviation (SD) and ancestral

misidentification rate of each gamma fitting in Figure S3. The shape of the fitted gammas vary from

an L-shape (Mean/SD $<$ 1) at low C bins, to a shape resembling a skewed normal distribution at in-

termediate C bins (Mean/SD $>$ 1) to a shape resembling an exponential distribution at high C bins

(Mean/SD $\approx$ 1) (Figure 1.D). We performed a likelihood ratio test comparing the gamma model to the

single-coefficient model, and find that only 4 out of the 40 bins (containing only 0.5% of all polymor-

phisms and 4.7% of nonsynonymous polymorphisms) are significantly supportive of the gamma model

(Figure 1.E). A chi-squared test of the fit to the data shows both models perform similarly well, though

both result in significant chi-squared scores at low C-score bins when using the genome-wide data (Figure

1.F). We attribute this to the large amount of data present in those bins as well as complex details

of demographic history that affect neutral sites but that we did not model in our neutral fitting. In

contrast, when mapping using only the exome, almost all bins have non-significant statistics, suggesting

that selection dominates over demography in these regions. Based on these results, we decided to use

the smoothed single-coefficient fitting for estimating the DFE in most downstream analyses, although we

may be missing a small proportion of within-bin variability. Additionally, we show the inferred DFE of

all, nonsynonymous and synonymous polymorphisms obtained from the gamma-fitted mapping in Figure

S5.

We aimed to test the robustness of the selection coefficient estimates within each bin. We were specif-

ically concerned about highly deleterious bins, which are composed of a smaller number of segregating

sites than neutral or nearly neutral bins, and could produce unstable or biased estimates. We obtained

bootstrapped confidence intervals for each bin and observe that the mappings are relatively stable up to

C= 35 (Figure 1.A). As expected, the standard deviation of the bootstrap estimates is strongly negatively

correlated with the sample-size per bin (Figure S6, Pearson correlation coefficient = -0.89). Thus, most

of the increase in the width of the confidence intervals observed at higher C-score bins can be explained

by the small number of polymorphisms available in those bins, and is likely not the result of other un-

accounted processes, such as positive selection, operating exclusively on highly scored polymorphisms.

To verify that our mapping was not an artifact of the different number of C-scores within each bin, we

also performed 100 randomizations of the C-score assignments to each SNP in the genome. For each

randomization, we re-computed the C-to-s mapping. When doing so, the bootstrap confidence intervals

increase in size with increasing C scores, but the mapping remains flat, as expected (Figure 1.B).

Further, we verified that the mapping did not change considerably when filtering for sites in regions

with low CpG density (< 0.05), defined as the proportion of CpG dinucleotides in a window of +/-
75bp around the site [27] (Figure S7.A). This is expected, as the C-score model accounts for differential
mutation rates at CpG sites and the resulting scores are generally robust to them [27]. As before, the
gamma model is a significantly better fit than the single-coefficient model at only 4 out of the 40 bins
(Figure S7.B).

Additionally, we re-mapped the scores using a constant-size model, and verified that the mapping
does not change considerably if we assume a different demographic history than the best fit (Figure S8).
The mappings are highly similar in shape, with the exception that, because the constant-size model is
depleted of singletons relative to the best-fit model, it takes more bins to reach an SFS that is deleterious
enough to map to $s \neq 0$, and so more C-scores map to s= 0.

To test the dependence of our mapping on the choice of score used to estimate selection coefficients,
we performed the same fitting procedure on a variety of other conservation and deleteriousness scores (see
Methods). We note, however, that all of these scores are included as input in the C-score SVM. Figure
S9 shows that the shape of the mapping is fairly consistent across different choices of scores, except for
highly deleterious bins, which contain very few sites. When comparing different categories of sites in the
Results, we show their distribution of selection coefficients obtained from the C-score mapping, as this
score has been shown to be a better correlate to functional disruption than all the other scores mentioned
above, and also controls for mutation rate variation across the genome, while other scores do not [27].
Additionally, we plot the mapped density of selection coefficients (with smoothing bandwidth = 0.000005)
for all polymorphisms as well as synonymous and nonsynonymous polymorphisms using each of the other
scores in Figure S10. We observe that, while all scores easily distinguish genic sites, PhastCons scores
have difficulty distinguishing between synonymous and nonsynonymous sites, which we believe is due to
PhastCons scores being regional, rather than position-specific scores. Additionally, while bimodality at
genic sites is most prominent when using C-scores, it also becomes apparent in other position-specific
scores when plotting the density with a finer smoothing bandwidth (= 0.000001, Figure S11).

The distribution of fitness effects of segregating mutations

197 Using the C-score-to-selection coefficient mapping, we obtained the DFE of segregating polymorphisms
198 in Yoruba individuals. This distribution is highly peaked when all polymorphisms are considered (Figure
199 2, black dashed line), with a large proportion of neutral changes and a long tail of deleterious mutations,

as has been observed before when estimating the DFE of coding sequences [9, 11–13, 20]. Interestingly,

we observe a pronounced drop in frequency for values of $s < -10^{-4}$. We note that this is not due to our

capping our mapping at $C = 40$ as the selection coefficients we are able to map are of a greater magnitude

than this drop (Figure 1, S12).

We partition the data by the genomic consequence of the polymorphisms, using the Ensembl Variant

Effect Predictor (v.2.5) [36]. Some classes exhibit a peak of highly deleterious changes around $s =$

-10^{-4}. This peak results in a bimodal distribution that is especially pronounced for nonsynonymous

sites (Figure 2, red line), and is almost non-existent for intergenic sites (Figure 2, pink line). Other types

of polymorphisms—like splice site, synonymous, 3' UTR, 5' UTR and regulatory mutations—have less

deleterious peaks than the one observed at nonsynonymous polymorphisms (Figure 2). We can compare

the selection coefficient distributions to the distributions of unmapped C-scores (Figure S12) which are

much less tightly peaked at intermediate C-score values and do not show a sharp decrease in density for

high values, as does the s distribution in Figure 2. We show various statistics calculated on each of the

selection coefficient distributions in Table 1 with the genome-wide mapping and in Table S1 with the

exome-wide mapping.

Next, we partitioned the data by whether the polymorphisms were found in the GWAS database [37]

or not (Figure S13, Tables 1, S1). While we observe a second deleterious peak among the GWAS SNPs

as well, these SNPs seem to be highly enriched for neutral polymorphisms.

Finally, we classified polymorphisms by different regulatory categories. We used two regulatory tracks

downloaded from the UCSC Genome Browser [31, 38]. First, we partitioned the genome by regulatory

regions identified by Regulome DB [39], which predicts regulatory activity in noncoding regions based

on different types of experimental evidence (Figure S14, Tables 1, S1). Second, we used the Segway

regulatory segment tracks [40], which are the product of an unsupervised pattern discovery algorithm

that serves to identify and label regulatory regions along the genome, including genic regions (Figure 3,

Tables 1, S1).

Discussion

The distribution of fitness effects (DFE) describes the proportion of mutations with given selection

coefficients. Knowledge of the DFE has profound implications for our understanding of evolution and

health. We infer a highly peaked distribution for all polymorphisms, with a drop in density at $s \approx -10^{-4}$,

which may be the cutoff between weakly deleterious mutations that segregate in human populations and

highly deleterious mutations that are easily pruned away by negative selection.

Our inferred non-synonymous distribution is bimodal and looks very similar to the one obtained for

nonsynonymous mutations in Drosophila in ref. [11], with a peak at neutrality and another peak at $s \approx$

-10^{-4}. Several experimental studies have also shown that non-synonymous non-lethal mutations tend to

have a multimodal DFE in model organisms [41,42] (see ref. [18] for a comprehensive review). We note that

it is impossible to obtain such kinds of distributions using a gamma or lognormal probability distribution

unless one approximates bimodality by assuming a second, separate class of nonsynonymous mutations

that are completely neutral and do not follow the best-fitting probability distribution [11, 13, 20, 25].

We also tested the precision of the C-scores by fitting gamma distributed DFEs to each C-score bin.

While only very few bins were fit by a highly peaked gamma distribution (Figure 1D), the increased

variation offered by the gamma distribution rarely improved the likelihood significantly (Figure 1E).

This suggests that the C-scores are precise quantifications of negative selection, and that mutations with

similar C-scores are likely to have similar selection coefficients.

Interestingly, we found that for many low C-score bins, a chi-squared test would reject the fit of the

demographic model to the data. This is likely because these low C-score bins have a significant number

of sites, and hence subtle features of the demography and quality control are relevant. Nonetheless, we

note that for moderate-to-high C-score bins and for exonic data, we would not be able to reject the fit

of the the predicted site frequency spectrum from the data. While these bins have fewer sites, it is also

likely that stronger selection is obscuring some of the signal of subtle demographic events.

Our novel expectation-maximization approach to quantifying ancestral state misidentification allows

us to assess differential misidentification rates across C-score categories. Ancestral state misidentification

occurs because a site is hit by more than one mutation, hence obscuring the identity of the ancestral

allele. Here, we found that low C-score bins are enriched with ancestral state misidentification, with rates

in excess of 5% for some bins (Figure S3). This may reflect a bias induced by the C-scores themselves,

because C-scores are trained to distinguish classes of sites that have relatively few substitutions between

humans and chimpanzees. Because the signal of ancestral state misidentification and positive selection

are very similar [43], it is possible that low C-score bins are enriched for positive selection, although we

did not pursue that direction any further. For larger C-score bins, we infer misidentification rates along

the lines of those obtained in simulation studies by ref. [43].

Importantly, unlike previous studies, we also obtain DFEs for other types of mutations, including
synonymous, splice site, 3' UTR, 5' UTR and regulatory polymorphisms, which exhibit bimodality to
a lesser degree than the nonsynonymous DFE. In particular, 5' UTR changes constitute the category
with the smallest proportion of neutral or nearly neutral ($|s| < 10^{-5}$) polymorphisms after nonsynony-
mous changes, likely reflecting selection on gene regulation upstream of coding sequences. Futhermore,
distributions corresponding to mutations in UTR and regulatory regions have a less pronounced trough
between the two peaks than the ones observed among coding mutations, suggesting that the magnitude
of deleterious effects is more uniformly distributed in non-coding regions. In contrast, missense mutations
appear to have more of an "all-or-nothing" effect, as would perhaps be expected when replacing an amino
acid inside a protein.

Our method does not use synonymous sites as a neutral benchmark, as do other studies [9, 11, 20]. In
fact, our inferred DFE suggests that a considerable number of synonymous polymorphisms may not be
neutral, as we observe a second deleterious peak in them too, albeit less deleterious than the one observed
at nonsynonymous polymorphisms. We caution, however, that this second peak is less promient when
using an exome-specific mapping (Figure 2) or when using strictly position-specific scores (Figures S10,
S11), which suggests that at least part of this peak may be caused by regional patterns of conservation or
background selection in the exomes. Instead, intergenic polymorphisms are the class of sites most likely to
evolve neutrally. Because this class is so abundant, most of the signal observed when all polymorphisms
are pooled together closely reflects the distribution observed for intergenic polymorphisms (Figure 2).

Our results have implications for GWAS, as we find a high proportion of GWAS SNPs to be neutral
or nearly neutral, which could suggest a high rate of false positives in this type of association studies.
On the other hand, GWAS studies only aim to find polymorphisms linked to causative variants, and so
GWAS SNPs need not have strongly deleterious effects. Alternatively, if the effect size of many GWAS
SNPs are sufficiently small, it is possible that many of them are not subject to strong selection.

Additionally, by stratifying our results based on different ENCODE categories, we can elucidate the
fitness consequences of molecular activity signals detected by ENCODE [2, 3, 39]. We find the category
with the lowest proportion of neutral polymorphisms to be the one corresponding to sites that have eQTL
evidence as well as evidence for transcription factor (TF) binding, a matched TF motif, a matched DNase
footprint and that are located in a DNase peak. In general, categories that combine many regulatory

signals tend to show lower proportions of neutral mutations than those that do not, suggesting that data integration across distinct approaches to detecting selection and functionality is likely to do better than any individual approach [44]. Moreover, this suggests that much of the molecular activity detected by ENCODE may not have significant fitness consequences.

Stratification by Segway regions allows us to look at a different aspect of regulatory activity in the genome. Interestingly, we observe that polymorphisms in Transcription Start Sites (TSS) are the ones containing the largest proportion of deleteriousness. This echoes results from analyses of variation at transcription factor binding sites, which have been found to be under stronger constraint when found near TSS than when found far from them [45]. Other regions with high proportions of deleterious polymorphisms include Gene Body (Start), strong CTCF and Enhancer / Gene Middle. This suggests that regions with strong repressor, insulator or enhancer activity, as well as near the start of genes, are particularly important for biological function, perhaps unsurprisingly given our knowledge of the molecular role that these regions play in the regulation of transcription.

There are several limitations to our method. First, we have restricted ourselves to estimating the DFE of segregating mutations that have reached appreciable frequencies in the population. An extension of this approach would be to infer the DFE of new mutations from the DFE of segregating mutations genome-wide. Second, we assumed no dominance, epistasis or positive selection, which future studies could attempt to incorporate into our approach. In addition, we have assumed sites are independent and have therefore ignored the covariance between linked sites, which likely leads to an underestimatation of confidence intervals obtained from the bootstrapping. The free-recombination assumption may also affect inference due to Hill-Robertson interference between mutations subject to selection [46] as well as linked background selection affecting the SFS of neutral sites in the human genome [35]. This may be a more important issue in our case than other genic-only approaches because we are also including intergenic mutations in our analysis, so the space between analyzed polymorphisms is on average smaller than if we were only looking at coding polymorphisms [20]. We also assume no positive selection. This, however, should not be a major problem, because we are only basing our inferences on polymorphic sites and advantageous mutations contribute little to polymorphism, assuming $N_e s > 25$ [32]. One final limitation is that the type of inference performed here is only possible in species in which C-scores have been estimated (for now, humans only) and that it therefore relies on C-scores being able to correctly rank sites by deleteriousness. Nevertheless, it should not be hard to obtain accurate C-scores for other

318 organisms in the future, although limitations on available annotations for non-human organisms may

319 make the approximation to the fitness distribution less accurate than in humans.

Materials and Methods

320

Computing the theoretical site frequency spectrum

321

322 We used the theory developed by Evans *et al.* [47] to obtain the expected population site frequency

323 spectrum with non-equilibrium demography. We assume a Wright-Fisher population in the limit of large

324 population size and use diffusion theory to model this process. Writing $f(x, t)$ for the frequency spectrum

325 at frequency x and time $t = 2N(0)\tau$ where τ is in units of generations and $g(x, t) := x(1 - x)f(x, t)$,

326 we can approximate the dynamics of $g(x, t)$ with genic selection and mutation by solving the following

327 partial differential equation:

$$\frac{\partial}{\partial t}g(x, t) = -Sx(1 - x)\frac{\partial}{\partial x}[g(x, t)] + \frac{x(1 - x)}{2\rho(t)}\frac{\partial^2}{\partial x^2}[g(x, t)] \tag{1}$$

328 subject to the boundary condition:

$$\lim_{x \to 0} g(x, t) = \theta\rho(t) \tag{2}$$

329 where S is the population-scaled selection coefficient ($S = 2N(0)s$), θ is the population-scaled mutation

330 rate ($\theta = 4N(0)\mu$) and $\rho(t) = N(t)/N(0)$ is the effective population size at time t relative to the population

331 size at time 0.

332 We assume N(0) to be 10,000 for all exponential and constant models. For the constant population size

333 model, $\rho(t) = 1$. For the exponential growth model $\rho(t) = exp(rt)$ where $r = 2N(0)R$ is the population-

334 scaled growth rate and R is the per-generation growth rate. For models taken from the literature, we use

335 the N(0) assumed by the corresponding paper. For the model of Harris and Nielsen, $\rho(t)$ is piece-wise

336 defined according to their Figure 7. The Tennessen model is similarly defined in a piece-wise fashion

337 according to their Figure 2, although it also includes periods of exponential growth, as opposed to simply

338 being piece-wise constant as in the Harris and Nielsen model.

339 We solve for $g(x, t)$ numerically in Mathematica, and can then compute the expected number of

340 segregating sites with i copies of the derived allele out of a sample of n genes. Denoting by $g_s(x, t)$ the

theoretical SFS with selection coefficient s, this quantity is

$$f_{n,i}(t) = \int_0^\infty \left(\int_0^1 \binom{n}{i} x^{i-1}(1-x)^{n-i-1} g_{-s}(x,t) dx \right) p(s) ds. \tag{3}$$

where $p(s)$ is the parameterized distribution of selection coefficients. For gamma distributed fits,

$$p(s) = \frac{x^{\alpha-1} e^{-x\beta}}{\beta^{-\alpha} \Gamma(\alpha)}$$

where α and β are the shape and rate parameters of the gamma distribution and $\Gamma(\cdot)$ is the gamma function. For a point mass at $\hat{s}$,

$$p(s) = \delta(s - \hat{s})$$

where $\delta(\cdot)$ is the usual Dirac delta function.

We focused on fitting the shape of the SFS, and hence worked with the probability that a given site in a sample of n has i copies of the derived allele,

$$p_{n,i}(t) = \frac{f_{n,i}(t)}{\sum_{j=1}^{n-1} f_{n,j}(t)}. \tag{4}$$

The Mathematica code used for obtaining the theoretical SFS can be found online at:

http://malecot.popgen.dk/schraiber/

Expectation maximization algorithm to fit ancestral state misidentification rates

We observed that the SFS showed signs of ancestral state misidentification, in particular an excess of high frequency derived alleles (Figure S2). To account for the ancestral state misidentification errors, we developed an expectation maximization (EM) algorithm. In the E step, we estimate the posterior probability that a site at frequency i out of n is misidentified given the current estimated site frequencies, $\{p_{n,i}^{(k)}, 1 \leq i < n\}$, and the current estimate of the misidentification rate, $p_{mis}^{(k)}$, as

$$m_i \propto p_{n,i}^{(k)} p_{mis}^{(k)}. \tag{5}$$

357 Then, during the M step, we re-estimate the misidentification rate as

$$p_{mis}^{(k+1)} \propto \sum_{i=1}^{n-1} x_i m_i \tag{6}$$

358 where x_i is the number of sites at frequency i. We next re-estimate either the demographic parameters **359** or the parameters of the selected model using the log-likelihood

$$\log L = \sum_{i=1}^{n-1} x_i \left(m_i \log p_{n,i}^{(k)} + (1 - m_i) \log p_{n,n-i}^{(k)} \right) \tag{7}$$

360 to obtain the theoretical SFS for the next iteration, $\{p_{n,i}^{(k+1)}, 1 \leq i < n\}$

361 Maximum likelihood fitting of demographic models

362 The exponential growth model has two free parameters, r, the population-scaled growth rate and t, the **363** total time of exponential growth. We first obtained the site frequency spectrum for all sites with $C = 0$. **364** Next we solved $g(x, t)$ for the exponential growth model across a grid of t and r, as well as the constant, **365** Harris and Tennessen models, and applied our EM algorithm to estimate the best fitting demographic **366** model, using a grid search over demographic models during the M step.

367 Maximum likelihood fitting of selection coefficients

368 To find the maximum likelihood estimate of s for each C-score bin, we first obtained the site frequency **369** spectrum corresponding to each C-score bin. Next, we solved $g(x, t)$ under the fitted demography for **370** $\log_{10}(-s) \in [-7, -1.3]$ in steps of 0.005, along with $s = 0$. To obtain an estimated SFS under the **371** assumption of gamma distributed selection coefficients, we used the trapezoid rule to numerically integrate **372** against a gamma distribution as in formula 3.

373   We used our EM algorithm to estimate the best fitting selection coefficient for each bin. When fitting **374** a single coefficient, we used a grid search during the M-step, and when fitting gamma distributed selection **375** coefficients, we used the L-BFGS-B algorithm. To plot the DFE, we used kernel density estimation with **376** smoothing bandwith $= 0.000005$, unless otherwise stated.

Mapping using different scores

To test how robust the mapping of C-scores to selection coefficients is to different types of conservation scores, we obtained PhyloP [48] and PhastCons [49] scores derived from vertebrate, mammal and primate alignments (excluding humans), as well as GERP++ rejected substitution (Gerp S) scores [50], for all YRI SNPs [27]. We attempted to equalize the range of all scores by PHRED-scaling them, i.e. converting each score to -$\log_{10}(p)$ where p is the probability of observing a change as or more disruptive / conserved (based on that particular score scale) among all polymorphic YRI sites. We note that this is different from the natural PHRED scale of C-scores (where p is the the probability of observing a score as or more disruptive among all possible, but not necessarily realized, mutations in the human genome), and so we also re-scaled the C-scores to produce a fair comparison. Then, we repeated the maximum likelihood mapping for each PHRED-scaled score in bins of 0.25 units (e.g. 0-0.125, 0.125-0.375, 0.375-0.625, etc). It is important to note that PhastCons are regional scores, while PhyloP and Gerp S are position-specific scores. Another difference is that PhastCons scores only measure the probability of negative selection, while PhyloP and Gerp S scores also measure positive selection (i.e. low/negative scores represent faster evolution than expected purely under drift), which may be why we observe an uptick at the lower end of the mapping for those scores in Figure S9.

Acknowledgments

We thank Montgomery Slatkin and Rasmus Nielsen for helpful advice and discussions, and Martin Kircher for providing the C-score data and other annotations. This work was supported by the National Institutes of Health (R01-GM40282 grant to Montgomery Slatkin).

References

1. McCarthy MI, Abecasis GR, Cardon LR, Goldstein DB, Little J, et al. (2008) Genome-wide association studies for complex traits: consensus, uncertainty and challenges. Nature Reviews Genetics 9: 356–369.

2. Dunham I, Kundaje A, Aldred S, Collins P, Davis C, et al. (2012) An integrated encyclopedia of DNA elements in the human genome. Nature 489: 57–74.

3. Eddy SR (2013) The encode project: missteps overshadowing a success. Current Biology 23: R259–R261.

4. Graur D, Zheng Y, Price N, Azevedo RB, Zufall RA, et al. (2013) On the immortality of television sets: "function" in the human genome according to the evolution-free gospel of ENCODE. Genome biology and evolution 5: 578–590.

5. Lawrie DS, Petrov DA (2014) Comparative population genomics: power and principles for the inference of functionality. Trends in Genetics 30: 133–139.

6. Siepel A, Pollard KS, Haussler D (2006) New methods for detecting lineage-specific selection. In: Research in Computational Molecular Biology. Springer, pp. 190–205.

7. Lindblad-Toh K, Garber M, Zuk O, Lin MF, Parker BJ, et al. (2011) A high-resolution map of human evolutionary constraint using 29 mammals. Nature 478: 476–482.

8. Ward LD, Kellis M (2012) Evidence of abundant purifying selection in humans for recently acquired regulatory functions. Science 337: 1675–1678.

9. Piganeau G, Eyre-Walker A (2003) Estimating the distribution of fitness effects from DNA sequence data: Implications for the molecular clock. Proceedings of the National Academy of Sciences 100: 10335–10340.

10. Sawyer SA, Kulathinal RJ, Bustamante CD, Hartl DL (2003) Bayesian analysis suggests that most amino acid replacements in Drosophila are driven by positive selection. Journal of molecular evolution 57: S154–S164.

11. Loewe L, Charlesworth B, Bartolomé C, Nöel V (2006) Estimating selection on nonsynonymous mutations. Genetics 172: 1079–1092.

12. Keightley PD, Eyre-Walker A (2007) Joint inference of the distribution of fitness effects of deleterious mutations and population demography based on nucleotide polymorphism frequencies. Genetics 177: 2251–2261.

13. Boyko AR, Williamson SH, Indap AR, Degenhardt JD, Hernandez RD, et al. (2008) Assessing the evolutionary impact of amino acid mutations in the human genome. PLoS genetics 4: e1000083.

14. Wilson DJ, Hernandez RD, Andolfatto P, Przeworski M (2011) A population genetics-phylogenetics approach to inferring natural selection in coding sequences. PLoS genetics 7: e1002395.

15. Arbiza L, Gronau I, Aksoy BA, Hubisz MJ, Gulko B, et al. (2013) Genome-wide inference of natural selection on human transcription factor binding sites. Nature genetics .

16. Gronau I, Arbiza L, Mohammed J, Siepel A (2013) Inference of natural selection from interspersed genomic elements based on polymorphism and divergence. Molecular biology and evolution 30: 1159–1171.

17. Torgerson DG, Boyko AR, Hernandez RD, Indap A, Hu X, et al. (2009) Evolutionary processes acting on candidate cis-regulatory regions in humans inferred from patterns of polymorphism and divergence. PLoS genetics 5: e1000592.

18. Eyre-Walker A, Keightley PD (2007) The distribution of fitness effects of new mutations. Nature Reviews Genetics 8: 610–618.

19. Siepel A, Arbiza L (2014) Cis-regulatory elements and human evolution. bioRxiv .

20. Eyre-Walker A, Woolfit M, Phelps T (2006) The distribution of fitness effects of new deleterious amino acid mutations in humans. Genetics 173: 891–900.

21. Williamson SH, Hernandez R, Fledel-Alon A, Zhu L, Nielsen R, et al. (2005) Simultaneous inference of selection and population growth from patterns of variation in the human genome. Proceedings of the National Academy of Sciences 102: 7882–7887.

22. Kousathanas A, Keightley PD (2013) A comparison of models to infer the distribution of fitness effects of new mutations. Genetics 193: 1197–1208.

23. Wloch DM, Szafraniec K, Borts RH, Korona R (2001) Direct estimate of the mutation rate and the distribution of fitness effects in the yeast Saccharomyces cerevisiae. Genetics 159: 441–452.

24. Sanjuán R, Moya A, Elena SF (2004) The distribution of fitness effects caused by single-nucleotide substitutions in an RNA virus. Proceedings of the National Academy of Sciences of the United States of America 101: 8396–8401.

25. Loewe L, Charlesworth B (2006) Inferring the distribution of mutational effects on fitness in Drosophila. Biology Letters 2: 426–430.

26. Keightley PD, Eyre-Walker A (2010) What can we learn about the distribution of fitness effects of new mutations from dna sequence data? Philosophical Transactions of the Royal Society B: Biological Sciences 365: 1187–1193.

27. Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, et al. (In press) A general framework for estimating the relative pathogenicity of human genetic variants .

28. Sawyer SA, Hartl DL (1992) Population genetics of polymorphism and divergence. Genetics 132: 1161–1176.

29. Bustamante C, Nielsen R, Sawyer S, Olsen K, Purugganan M, et al. (2002) The cost of inbreeding in Arabidopsis. Nature 416: 531.

30. Drmanac R, Sparks AB, Callow MJ, Halpern AL, Burns NL, et al. (2010) Human genome sequencing using unchained base reads on self-assembling dna nanoarrays. Science 327: 78–81.

31. Karolchik D, Barber GP, Casper J, Clawson H, Cline MS, et al. (2014) The ucsc genome browser database: 2014 update. Nucleic acids research 42: D764–D770.

32. Smith NG, Eyre-Walker A (2002) Adaptive protein evolution in Drosophila. Nature 415: 1022–1024.

33. Tennessen JA, Bigham AW, O'Connor TD, Fu W, Kenny EE, et al. (2012) Evolution and functional impact of rare coding variation from deep sequencing of human exomes. Science 337: 64–69.

34. Harris K, Nielsen R (2013) Inferring demographic history from a spectrum of shared haplotype lengths. PLoS genetics 9: e1003521.

35. McVicker G, Gordon D, Davis C, Green P (2009) Widespread genomic signatures of natural selection in hominid evolution. PLoS genetics 5: e1000471.

36. McLaren W, Pritchard B, Rios D, Chen Y, Flicek P, et al. (2010) Deriving the consequences of genomic variants with the ensembl api and snp effect predictor. Bioinformatics 26: 2069–2070.

37. Hindorff LA, Sethupathy P, Junkins HA, Ramos EM, Mehta JP, et al. (2009) Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. Proceedings of the National Academy of Sciences 106: 9362–9367.

38. Rosenbloom KR, Dreszer TR, Pheasant M, Barber GP, Meyer LR, et al. (2010) Encode whole-genome data in the ucsc genome browser. Nucleic acids research 38: D620–D625.

39. Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, et al. (2012) Annotation of functional variation in personal genomes using RegulomeDB. Genome research 22: 1790–1797.

40. Hoffman MM, Buske OJ, Wang J, Weng Z, Bilmes JA, et al. (2012) Unsupervised pattern discovery in human chromatin structure through genomic segmentation. Nature methods 9: 473–476.

41. Davies EK, Peters AD, Keightley PD (1999) High frequency of cryptic deleterious mutations in Caenorhabditis elegans. Science 285: 1748–1751.

42. Keightley PD (1996) Nature of deleterious mutation load in Drosophila. Genetics 144: 1993–1999.

43. Hernandez RD, Williamson SH, Bustamante CD (2007) Context dependence, ancestral misidentification, and spurious signatures of natural selection. Molecular biology and evolution 24: 1792–1800.

44. Scheinfeldt LB, Tishkoff SA (2013) Recent human adaptation: genomic approaches, interpretation and insights. Nature Reviews Genetics 14: 692–702.

45. Mu XJ, Lu ZJ, Kong Y, Lam HY, Gerstein MB (2011) Analysis of genomic variation in non-coding elements using population-scale sequencing data from the 1000 genomes project. Nucleic acids research 39: 7058–7076.

46. McVean GA, Charlesworth B (2000) The effects of Hill-Robertson interference between weakly selected mutations on patterns of molecular evolution and variation. Genetics 155: 929–944.

47. Evans SN, Shvets Y, Slatkin M (2007) Non-equilibrium theory of the allele frequency spectrum. Theoretical population biology 71: 109–119.

48. Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A (2010) Detection of nonneutral substitution rates on mammalian phylogenies. Genome research 20: 110–121.

505   49. Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, et al. (2005) Evolutionarily conserved
506   elements in vertebrate, insect, worm, and yeast genomes. Genome research 15: 1034–1050.

507   50. Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, et al. (2010) A method and
508   server for predicting damaging missense mutations. Nat Methods 7: 248–249.

 # Figures

Figure 1. Mapping of C-scores to selection coefficients. A) We first fit a single coefficient to each bin using data from all autosomes in the genome. Red dots represent the maximum likelihood selection coefficient corresponding to each C-score bin. The blue line is a smooth spline fitted to the discrete mappings of C-scores to log-scaled selection coefficients after excluding the neutral bins (degrees of freedom = 20). The grey shade is a 95% confidence interval obtained from bootstrapping the data 100 times in each bin. B) To verify the shape of the mapping was not a result of the number of sites in each bin, we randomized the C-score labels across polymorphisms, and recomputed the mapping. C) To account for exonic patterns of conservation and background selection that may not have been captured by C-scores, we computed a mapping based solely on exonic sites. D) We fitted gamma distributions of selection coefficients to each bin, and computed the mean divided by the standard deviation of each distribution, which is indicative of its shape (see Results). E) To test whether the gamma distributions were a significantly better fit than the single-coefficient mapping, we computed log-likelihood ratio statistics of the two models at each bin. The black line denotes the Bonferroni-corrected significance cutoff. F) To test how well we were fitting the data at each bin, we computed chi-squared statistics of the fitted SFS to the observed SFS at each bin. The black line denotes the Bonferroni-corrected significance cutoff.

Figure 2. Distribution of fitness effects among YRI polymorphisms in the Complete Genomics dataset, partitioned by the genomic consequence of the mutated site. The right panels show a zoomed-in version of the same distributions after removing neutral polymorphisms and log-scaling the x-axis. Top row: DFE obtained from genome-wide mapping. Bottom row: DFE obtained from exome-wide mapping. Consequences were determined using the Ensembl Variant Effect Predictor (v.2.5). If more than one consequence existed for a given SNP, that SNP was assigned to the most severe of the predicted categories, following the VEP's hierarchy of consequences. NonSyn = nonsynonymous. Syn = synonymous. Syn to unpref. codon = synonymous change from a preferred to an unpreferred codon. Syn to pref. codon = synonymous change from an unpreferred to a preferred codon. Syn no pref. = synonymous change from an unpreferred codon to a codon that is also unpreferred. Splice = splice site. CG = Complete Genomics data.

Figure 3. Distribution of fitness effects among YRI polymorphisms, partitioned by Segway regulatory segments. The right panel shows a zoomed-in version of the same distributions after removing neutral polymorphisms and log-scaling the x-axis.

Tables

Table 1. Characteristics of fitness effect estimated for YRI SNPs classified by different genomic consequence, RegulomeDB and Segway categories, using the genome-wide C-to-s mapping. We show quantiles of selection coefficients, the log base 10 of the mean selection coefficient and the log base 10 of the standard deviation of coefficients in each category.

Category	n	$|s| < 10^{-5}$	$|s| \geq 10^{-5}$	$|s| \geq 5*10^{-5}$	$|s| \geq 10^{-4}$	$\log_{10}(\overline{-s})$	$\log_{10}(SD(-s))$
All	9065398	73.77%	26.23%	0.09%	0%	-5.16	-4.95
Nonsynonymous	32522	28.88%	71.12%	2.07%	0.14%	-4.6	-4.74
Synonymous	30630	42.08%	57.92%	0.01%	0%	-4.81	-4.87
Synonymous 4-fold degenerate	16895	41.93%	58.07%	0%	0%	-4.81	-4.87
Synonymous to unpreferred codon	12498	40.01%	59.99%	0.01%	0.01%	-4.79	-4.86
Synonymous to preferred codon	4574	38.43%	61.57%	0%	0%	-4.78	-4.87
Synonymous no preference change	11844	41.80%	58.20%	0.02%	0%	-4.81	-4.87
Splice site	10122	52.11%	47.89%	0.05%	0%	-4.87	-4.84
5' UTR	23125	37.53%	62.47%	0.11%	0.02%	-4.76	-4.84
3' UTR	81605	49.05%	50.95%	0.19%	0.02%	-4.88	-4.87
Regulatory	864769	58.72%	41.28%	0.02%	0%	-4.99	-4.92
Intergenic	3544204	77.69%	22.31%	0.16%	0%	-5.22	-4.96
GWAS	8693	65.71%	34.29%	0.18%	0%	-5.04	-4.91
eQTL+TF binding+matched TF motif+matched DNase Footprint+DNase peak	240	47.08%	52.92%	0%	0%	-4.86	-4.88
eQTL+TF binding+any motif+DNase Footprint+DNase peak	1965	51.30%	48.70%	0.10%	0%	-4.91	-4.9
eQTL+TF binding+matched TF motif+DNase peak	62	66.13%	33.87%	0%	0%	-5.01	-4.89
eQTL+TF binding+any motif+DNase peak	1263	58.04%	41.96%	0.08%	0%	-4.97	-4.9
eQTL+TF binding+matched TF motif	44	65.91%	34.09%	0%	0%	-4.95	-4.83
eQTL+TF binding / DNase peak	26610	63.87%	36.13%	0.06%	0%	-5.03	-4.92
TF binding+matched TF motif+matched DNase Footprint+DNase peak	12411	50.05%	49.95%	0.10%	0%	-4.87	-4.87
TF binding+any motif+DNase Footprint+DNase peak	120642	57.03%	42.97%	0.12%	0%	-4.95	-4.89
TF binding+matched TF motif+DNase peak	5355	65.45%	34.55%	0.06%	0%	-5.05	-4.92
TF binding+any motif+DNase peak	96905	63.14%	36.86%	0.16%	0%	-5.02	-4.9
TF binding+matched TF motif	4359	73.34%	26.66%	0.07%	0%	-5.15	-4.95
TF binding+DNase peak	418548	59.95%	40.05%	0.10%	0%	-4.99	-4.91
TF binding or DNase peak	1625195	69.18%	30.82%	0.14%	0%	-5.09	-4.93
C0 - CTCF (strong)	20813	57.22%	42.78%	0.02%	0.01%	-4.98	-4.94
C1 - CTCF (weak)	61946	65.88%	34.12%	0.05%	0%	-5.08	-4.96
D - dead zone	873933	81.44%	18.56%	0.06%	0%	-5.3	-5
E/GM - enhancer/gene middle	70593	59.27%	40.73%	0.06%	0%	-4.99	-4.91
F0 - FAIRE only	1177948	71.94%	28.06%	0.11%	0%	-5.13	-4.94
F1- FAIRE only	1481766	74.45%	25.55%	0.10%	0%	-5.17	-4.95
GE0 - gene body (end)	413390	67.56%	32.44%	0.09%	0%	-5.06	-4.91
GE1 - gene body (end)	163365	63.99%	36.01%	0.10%	0%	-5.03	-4.91
GE2 - gene body (end)	54261	69.52%	30.48%	0.13%	0.01%	-5.1	-4.93
GM0 - gene body (middle)	130000	64.24%	35.76%	0.07%	0%	-5.04	-4.93
GM1 - gene body (middle)	101426	63.21%	36.79%	0.07%	0%	-5.04	-4.94
GS - gene body (start)	54940	41.21%	58.79%	0.06%	0.01%	-4.83	-4.89
H3K9me1	457492	87.22%	12.78%	0.02%	0%	-5.46	-5.07
L0 - low zone	673274	81.71%	18.29%	0.07%	0%	-5.31	-5.01
L1 - low zone	725876	70.63%	29.37%	0.15%	0%	-5.11	-4.94
N/A	53354	95.36%	4.64%	0%	0%	-5.77	-5.29
R0 - repression	716710	74.73%	25.27%	0.12%	0%	-5.17	-4.95
R1 - repression	335824	69.73%	30.27%	0.09%	0%	-5.1	-4.94
R2 - repression	447655	68.85%	31.15%	0.13%	0%	-5.08	-4.91
R3 - repression	433156	75.62%	24.38%	0.09%	0%	-5.19	-4.97
R4 - repression	205971	73.26%	26.74%	0.06%	0%	-5.15	-4.94
R5 - repression	297152	70.84%	29.16%	0.09%	0.01%	-5.13	-4.96
TF0 - transcription factor activity	346083	73.50%	26.50%	0.09%	0%	-5.17	-4.97
TF1 - transcription factor activity	327695	77.18%	22.82%	0.06%	0%	-5.22	-4.98
TF2 - transcription factor activity	127366	71.09%	28.91%	0.12%	0.01%	-5.12	-4.95
TSS - transcription start site	25144	23.16%	76.84%	0.12%	0.02%	-4.68	-4.88

 # Supplementary Figures

Figure S1. Observed SFS for sites under different C-score bins using the Complete Genomics YRI data, for all autosomes in the genome (left) and the exome (right). Note that the spectrum gets more skewed towards singletons with increasing C-scores, likely reflecting the action of negative selection on deleterious mutations.

Figure S2. Observed SFS of YRI Complete Genomics data for sites with C=0. The full SFS was corrected for ancestral state misidentification using an EM algorithm and fit to different models of neutral evolution. We show results for both the genome and the exome.

Figure S3. Features of fitted single-coefficient and gamma distributions. A) Fitted single coefficients and means of fitted gamma distributions for each C-score bin, using genome-wide or exome-wide polymorphisms. B) Standard deviation of fitted gamma distributions for each bin. C) Ancestral misidentification rate obtained from an EM algorithm used to jointly fit the data and infer this rate at each bin. SD = standard deviation.

Figure S4. C-to-s mapping stratified by B-score deciles. We partitioned the genome by deciles of B-scores [35], which reflect levels of background selection. Then, we recomputed the demographic fitting and C-to-s mapping for each decile.

Figure S5. Inferred DFEs for all, nonsynonymous and synonymous polymorphisms obtained from gamma distribution fitting of each C-score bin. The plot shows, for each category, a weighted sum of gamma distributions, where each C-score bin contributes its corresponding genome-wide best-fitting gamma distribution in proportion to the number of polymorphisms present at that bin.

Figure S6. Comparison between the size of each C-score bin and the standard deviation of single-coefficient fits obtained from 100 bootstraps of the data within each bin. Top panel: Standard deviation per C-score bin plotted as a function of sample size per bin (log-scale). Bottom panel: Same plot but with the y-axis on a log-scale.

Figure S7. Mapping of sites with low CpG density. A) We filtered for sites with low CpG density, such that the proportion of CpG sites in a +/- 75 bp window around each site was < 0.05, and then recomputed the C-to-s mapping. B) We also repeated the gamma fitting and calculated a liklelihood ratio test of the gamma model against the single-coefficient model at each C-score bin.

Figure S8. Comparison between a C-to-s mapping using the best-fit demographic model (exponential growth with t=13,000 and r=1) and a constant-size model.

Figure S9. Maximum likelihood mapping of different types of scores to a selection coefficient scale, excluding bins mapped to neutrality, using the Complete Genomics data. Before mapping, scores were re-scaled on a common PHRED scale, by converting each score to $-\log_{10}(p)$ where p is the probability of observing a change as or more disruptive / conserved (based on that particular score scale) among all polymorphic YRI sites. Some scores extend over larger PHRED scores than others because they have a finer stratification (smaller number of sites with tied scores). The wide fluctuations to the right of the figures are due to the small number of sites per bin at highly deleterious bins.

Figure S10. Distribution of fitness effects at nonsynonymous (red), synonymous (blue) and all (black) polymorphisms in Yoruba, using different types of conservation scores for mapping (smoothing bandwidth = 0.000005). We only mapped sites with PHRED-scaled scores ≤ 5, because the mappings become erratic for higher values, due to the small number of sites per bin (Figure S9).

Figure S11. Distribution of fitness effects at nonsynonymous (red), synonymous (blue) and all (black) polymorphisms in Yoruba, using different types of conservation scores for mapping (smoothing bandwidth = 0.000001). We only mapped sites with PHRED-scaled scores ≤ 5, because the mappings become erratic for higher values, due to the small number of sites per bin (Figure S9).

Figure S12. Distribution of unmapped C-scores among YRI polymorphisms, partitioned by the genomic consequence of the mutated site. Consequences were determined using the Ensembl Variant Effect Predictor (v.2.5). NonSyn = nonsynonymous. Syn = synonymous. Syn to unpref. codon = synonymous change from a preferred to an unpreferred codon. Syn to pref. codon = synonymous change from an unpreferred to a preferred codon. Syn no pref. = synonymous change from an unpreferred codon to a codon that is also unpreferred. Splice = splice site.

Figure S13. Distribution of fitness effects among YRI polymorphisms, partitioned by whether the SNPs are found in the GWAS database or not. The right panel shows a zoomed-in version of the same distributions after removing neutral polymorphisms and log-scaling the x-axis.

Figure S14. Distribution of fitness effects among different types of RegulomeDB regulatory YRI polymorphisms, obtained from various ENCODE assays. The black dashed line corresponds to the distribution of all YRI SNPs.

512 Supplementary Tables

Table S1. Characteristics of fitness effect distributions estimated for YRI SNPs classified by different genomic consequence categories, RegulomeDB categories and Segway categories, using the exome-wide C-to-s mapping. We show quantiles of selection coefficients, the log base 10 of the mean selection coefficient and the log base 10 of the standard deviation of coefficients in each category.

Category	n	$|s| < 10^{-5}$	$|s| \geq 10^{-5}$	$|s| \geq 5*10^{-5}$	$|s| \geq 10^{-4}$	$\log_{10}(\overline{-s})$	$\log_{10}(SD(-s))$
All	9065398	40.51%	59.49%	5.87%	0.92%	-4.79	-4.7
Nonsynonymous	32522	17.26%	82.74%	45.87%	11.52%	-4.28	-4.35
Synonymous	30630	22%	78%	9.77%	0.06%	-4.61	-4.74
Synonymous 4-fold degenerate	16895	22.79%	77.21%	10.70%	0.03%	-4.61	-4.73
Synonymous to unpreferred codon	12498	21.54%	78.46%	10.52%	0.10%	-4.59	-4.73
Synonymous to preferred codon	4574	19.52%	80.48%	10.08%	0%	-4.59	-4.75
Synonymous no preference change	11844	22.55%	77.45%	9.58%	0.03%	-4.61	-4.74
Splice site	10122	25.43%	74.57%	16.08%	0.42%	-4.6	-4.63
5' UTR	23125	13.95%	86.05%	18.65%	0.88%	-4.52	-4.55
3' UTR	81605	20.98%	79.02%	11.94%	1.60%	-4.6	-4.58
Regulatory	864769	23.20%	76.80%	7.42%	0.14%	-4.7	-4.75
Intergenic	3544204	44.21%	55.79%	5.35%	1.12%	-4.82	-4.69
GWAS	8693	30.16%	69.84%	8.24%	1.69%	-4.7	-4.64
eQTL+TF binding+matched TF motif+matched DNase Footprint+DNase peak	240	14.58%	85.42%	11.67%	0.83%	-4.6	-4.7
eQTL+TF binding+any motif+DNase Footprint+DNase peak	1965	18.78%	81.22%	9.97%	0.51%	-4.64	-4.69
eQTL+TF binding+matched TF motif+DNase peak	62	25.81%	74.19%	8.06%	1.61%	-4.69	-4.65
eQTL+TF binding+any motif+DNase peak	1263	22.33%	77.67%	9.11%	0.55%	-4.67	-4.7
eQTL+TF binding+matched TF motif	44	31.82%	68.18%	13.64%	2.27%	-4.62	-4.56
eQTL+TF binding / DNase peak	26610	27.75%	72.25%	7.16%	0.86%	-4.71	-4.7
TF binding+matched TF motif+matched DNase Footprint+DNase peak	12411	17.94%	82.06%	13.42%	0.54%	-4.6	-4.65
TF binding+any motif+DNase Footprint+DNase peak	120642	22.76%	77.24%	9.62%	0.82%	-4.66	-4.66
TF binding+matched TF motif+DNase peak	5355	30.51%	69.49%	7.79%	1.05%	-4.72	-4.68
TF binding+any motif+DNase peak	96905	28.49%	71.51%	8.61%	1.29%	-4.69	-4.64
TF binding+matched TF motif	4359	38.63%	61.37%	6.33%	0.99%	-4.78	-4.7
TF binding+DNase peak	418548	24.64%	75.36%	8.15%	0.87%	-4.68	-4.68
TF binding or DNase peak	1625195	33.43%	66.57%	7.02%	1.32%	-4.74	-4.66
C0 - CTCF (strong)	20813	18.32%	81.68%	6.75%	0.15%	-4.69	-4.77
C1 - CTCF (weak)	61946	27.65%	72.35%	5.21%	0.46%	-4.75	-4.77
D - dead zone	873933	52.91%	47.09%	4.24%	0.65%	-4.89	-4.75
E/GM - enhancer/gene middle	70593	23.64%	76.36%	7.94%	0.67%	-4.68	-4.7
F0 - FAIRE only	1177948	36.88%	63.12%	6.41%	1.04%	-4.76	-4.69
F1- FAIRE only	1481766	41.02%	58.98%	5.94%	1.03%	-4.79	-4.69
GE0 - gene body (end)	413390	33.96%	66.04%	8.03%	1.09%	-4.72	-4.66
GE1 - gene body (end)	163365	28.37%	71.63%	8.06%	1.03%	-4.7	-4.67
GE2 - gene body (end)	54261	32.46%	67.54%	6.38%	0.94%	-4.74	-4.66
GM0 - gene body (middle)	130000	27.23%	72.77%	7.27%	0.82%	-4.71	-4.7
GM1 - gene body (middle)	101426	24.46%	75.54%	6.44%	0.62%	-4.71	-4.73
GS - gene body (start)	54940	10.49%	89.51%	11.35%	0.45%	-4.58	-4.7
H3K9me1	457492	62.34%	37.66%	2.70%	0.27%	-4.98	-4.85
L0 - low zone	673274	52.14%	47.86%	4.02%	0.67%	-4.89	-4.76
L1 - low zone	725876	33.64%	66.36%	6.32%	1.24%	-4.75	-4.68
N/A	53354	35.18%	64.82%	0.87%	0.07%	-4.96	-5.09
R0 - repression	716710	42.48%	57.52%	6.18%	1.11%	-4.79	-4.68
R1 - repression	335824	32.41%	67.59%	6.33%	1.01%	-4.75	-4.69
R2 - repression	447655	34.45%	65.55%	7.83%	1.37%	-4.73	-4.65
R3 - repression	433156	41.40%	58.60%	5.10%	0.85%	-4.81	-4.72
R4 - repression	205971	41.10%	58.90%	6.35%	0.80%	-4.78	-4.7
R5 - repression	297152	31.70%	68.30%	5.15%	0.66%	-4.77	-4.74
TF0 - transcription factor activity	346083	35.10%	64.90%	4.81%	0.69%	-4.79	-4.75
TF1 - transcription factor activity	327695	45.07%	54.93%	4.85%	0.65%	-4.83	-4.74
TF2 - transcription factor activity	127366	33.95%	66.05%	5.69%	0.82%	-4.76	-4.68
TSS - transcription start site	25144	5.01%	94.99%	21.84%	0.89%	-4.46	-4.6