Insect immune specificity in a host-parasite model

C.E. Riddell[1], J.D. Lobaton Garces[1], S. Adams[2], D. Twell[1] & E.B. Mallon[1]

[1]Department of Biology, University of Leicester, U.K.

[2] School of Life Sciences, University of Warwick, U.K.

Abstract

Ecological studies routinely show host-parasite genotype-genotype interactions in insect systems.

The mechanisms behind these interactions are not clearly understood. Using the bumblebee *Bombus*

terrestris / trypanosome *Crithidia bombi model system,* we have carried out a transcriptome-wide

analysis of gene expression in bees during *C. bombi* infection. We have performed three analyses,

1) comparing expression in infected and non-infected bees 24 hours after infection by *Crithidia*

bombi, 2) comparing expression at 24 and 48 hours after *C.bombi* infection and 3) most importantly

searching for differential gene expression associated with the host-parasite genotype-genotype

interaction at 24 hours after infection. We found a large number of genes differentially regulated

related to numerous canonical immune pathways. These genes include receptors, signaling

pathways and effectors. We discovered a possible interaction between the peritrophic membrane

and the insect immune system in defense against *Crithidia.* Most interestingly we found differential

expression of *Dscam* related transcripts depends on the genotype-genotype interactions of the given

bumblebee colony and *Crithidia* strain.

Invertebrate ecological studies have found infection outcomes within a given host-parasite system are variable. Part of this variance is determined by the interaction of the genotype of the host and the genotype of the parasite (Schmid-Hempel 2001; Carius *et al.* 2001; Lambrechts 2010). That is this interaction between host and parasite is specific (Schmid-Hempel 2005). How is this level of specificity generated? An obvious answer would be an interaction between the parasite and the host's immune response. We cannot take this for granted. Various ecological measures of disease outcome have been used to quantify genotype-genotype interactions. These measures include host mortality, fecundity and infection rate. Such measures cannot test directly if it is the immune response that produces this level of specificity (Hauton & Smith 2007). It may be other non-immune processes could explain such outcomes.

The bumblebee, *Bombus terrestris*/trypanosome *Crithidia bombi* system displays host x parasite genotype-genotype interactions (Schmid-Hempel *et al.* 1999; Schmid-Hempel & Reber-Funk 2004). There is evidence that the immune system has a role in generating this host-parasite specific response. A number of studies have found differential immune gene expression in response to *Crithidia* (Schlüns *et al.* 2010; Riddell *et al.* 2011; Brunner *et al.* 2013; Barribeau & Schmid-Hempel 2013). We found increased *Crithidia* loads in bees whose expression of antimicrobial peptides was knocked down by RNAi (Deshwal & Mallon 2014). We have even shown that bees from different host genotypes induce differential expression of antimicrobial peptides (AMPs), according to the strain of *C. bombi* they had been infected with (Riddell *et al.* 2009), that is we found specificity in the immune response itself. A recent paper using RNA-Seq found numerous genes are differentially expressed in a genotype-genotype fashion (Barribeau *et al.* 2014).

Here, we carry out a transcriptome-wide analysis of gene expression in bees during *C.bombi* infection. We have carried out three analyses, comparing 1) expression in infected and non-infected bees 24 hours after infection by *Crithidia bombi* (Infected versus uninfected) 2) expression at 24

and 48 hours after *C.bombi* infection (24 versus 48 hour) and 3) searching for differential gene

expression associated with the host-parasite genotype-genotype interaction at 24 hours post

infection (Specificity). Enrichment analysis was also carried out on expression data to see which

categories of molecules are differentially regulated during infection. The results confirm our

previous findings of up-regulation in antimicrobial peptide expression and provide a comprehensive

overview of changes in and the specificity of gene expression after exposure to 2 strains of

C.bombi.

Methods

The samples used during this experiment are as previously described (Riddell *et al.* 2009). We have chosen samples that displayed a reciprocal pattern of expression for the three antimicrobial peptides (AMPs) tested in that paper. These were colony K5 (called K from now on) and Q1 (Q) and strains 6 and 8. K-8 showed a high AMP expression, Q-8 a low expression level, Q-6 a high level and K-6 a low level of AMP expression.

Sample Collection

Experiments were carried out on one commercially reared bumblebee colony from Koppert Biological Systems U.K. (Colony K) and one colony from wild caught queens (Colony Q). Faecal samples from these colonies were checked under a light microscope to ensure there was no *Crithidia bombi* present (Mallon *et al.* 2003). All parasite isolates used originated from wild queens collected in Spring 2008 in the botanical gardens, University of Leicester. Experiments began when the colonies had a minimum of thirty workers, approximately four weeks old. Between observations, colonies were fed *ad libitum* with pollen (Percie du sert, France) and 50% diluted glucose/fructose mix (Meliose – Roquette, France). Before and during the experiments colonies were kept at 26°C and 60% humidity in constant red light.

Infections

To prepare *C. bombi* isolates, faeces was collected from workers of naturally infected colonies, and mixed with 50% diluted Meliose to create a standardized dose of 500 *Crithidia* cells per μl of inoculum. Previous studies had shown that such inocula, prepared from different colonies, are genotypically different (Schmid-Hempel & Reber-Funk 2004) and generate specific responses in novel hosts (Schmid-Hempel *et al.* 1999). However we can not rule out the possibility that we may be confounding our results with the addition of hidden infections or gut microbiota from the donor colonies. We infected a sample of workers from each of K and Q bumblebee colonies (representing different host lines) with an inoculum of faeces from each of the two wild infected colonies (6 and 8

Crithidia strain). We also collected uninfected controls. Bees were four days old at the time of

infection. Bees were collected over several days and distributed across treatment groups (Moret &

Schmid-Hempel 2009). After infection bees were kept in colony x strain groups (1–3 individuals

depending on day collected) and fed *ad libitum*. Twenty four hours or 48 hours post infection the

bees were sacrificed by freezing in liquid nitrogen and stored at −80°C.

RNA sample preparation and sequencing

Total RNA was extracted from 23 individual homogenised abdomens using Tri-reagent (Sigma-

Aldrich, UK). Samples (Colony-Strain-Timepoint (number of replicates)) were K-6-24 (3), K-6-48

(3), K-8-24 (3), K-8-48 (3), K-Uninfected (2), Q-6-24 (3), Q-6-48 (3), Q-8-24 (2), Q-uninfected (1).

Any residual contaminants were removed from the RNA using the RNeasy mini kit (Qiagen, UK)

and manufacturer's RNA clean-up protocol. To remove residual genomic DNA, RNA samples were

treated with DNase (Sigma-Aldrich, UK). TruSeq RNA-seq libraries were made from the 23

samples at NBAF Edinburgh. Sequencing was performed on an Illumina HiSeq®2000 instrument

(Illumina, Inc.) by the manufacturer's protocol. Multiplexed 50 base single-read runs were carried

out yielding an average of 12M reads per sample.

Statistical analysis

The reference transcriptome was downloaded from

http://www.nematodes.org/downloads/databases/Bombus_terrestris/   (Colgan   *et   al.*   2011).

Functional annotation related to the transcriptome was obtained using the BLAST2GO package

(Götz *et al.* 2008). Alignment was done using GSNAP (version 2012-07-20) (Wu & Nacu 2010).

Only reads that mapped uniquely were selected for further analysis. Counts were generated per

transcript for each sample.

Differential expression analysis was performed using the edgeR (3.4.0) package (McCarthy *et al.*

2012) in R (3.0.1) (R Core Team 2013). Normalization factors were computed using the TMM

technique, after which tagwise dispersions were calculated and subjected to a generalized linear

model (GLM). Resulting *p* values were subjected to Benjamini–Hochberg multiple testing

correction to derive FDRs; only transcripts with a FDR < 0.05 were considered for further analysis.

Three separate GLMs were carried out, one looked for transcripts that are differentially expressed

upon infection with *Crithidia* at 24 hours post-infection (Infected versus uninfected)

(~0+colony+infect(yes/no)) infect here are bees either infected with strain 6 or 8, one looking at the

gene expression difference between 24 hours and 48 hours post strain 6 infection (24 versus 48

hours)(~0+colony + time) and a further GLM that looked for transcripts that were expressed in a

specific pattern at 24 hours post-infection (specifcity)(~0+colony*strain).

Using Blast2Go, we then carried out an enrichment analysis (Fisher exact test) on each of these lists

of differentially expressed genes to see which GO terms are overrepresented relative to the entire

genome. We then used REVIGO to summarize and visualise these terms (Supek *et al.* 2011).

For each of the lists of differentially expressed transcripts we also carried out a blastx analysis

against the insect innate immunity database (IIID) (Brucker *et al.* 2012). We used the BLOSUM62

matrix with a word size of 3. The results were filtered to only contain hits with an *E*-value <1e-10

and a bit score ≥30.

Results

Genes differentially expressed at 24 hours post-infection (Infected versus uninfected)

31,843 unique transcripts were mapped to the transcriptome. 489 transcripts were found to be differentially expressed 24 hours post-infection (FDR < 0.05), including 324 downregulated and 165 upregulated transcripts (See supplementary data for complete list). Reannotating the transcripts using Blast2GO (blastx against the nr database with e < 0.001), 109 had no BLAST hits. A further 68 had uninformative BLAST hits (anonymous predicted protein). The remaining 312 were used in the enrichment analysis. Figure 1 shows a summary of the enriched GO terms found (Fisher's test p < 0.05). Defense response (GO:0006952, FDR = 0.047) and chitin metabolism (GO:0006030, FDR = 0.032) were the only processes significantly enriched at a more stringent level (FDR < 0.05).

Peritrophic membrane:

The peritrophic matrix (PM) forms a layer composed of chitin and glycoproteins that lines the insect midgut lumen (Kuraishi *et al.* 2011). The PM facilitates digestion and forms a protective barrier to prevent the invasion of ingested pathogens (Lehane 1997; Kuraishi *et al.* 2011). Fibrillin 1 (BTT14121_ 1), a venom protein precursor (BTT32193_1), Neurotrypsin (BTT07956_1), Peritrophin-1-like (BTT01709_1, BTT22959_1, BTT37215_1, BTT42262_1) and four chitinase transcripts (Chitinase 3: BTT23997_1 BTT38724_1, Chitinase 4 BTT20684_1, BTT23469_1) are downregulated upon infection. Fibrillins are extracellular matrix macromolecules, ubiquitous in the connective tissues (Isogai *et al.* 2003). BTT32193_1 was classed as a venom protein, but was also very similar to Chitinase 3 (blastx e = 1e-[16]). Chitinases modulate the structure and porosity of the PM (Dinglasan *et al.* 2009). Neurotrypsin is a serine protease expressed in the nervous system (Gschwend *et al.* 1997). However in the protease domain it shares similarities with Sp22D, a chitin binding serine protease (Danielli *et al.* 2000). The chitin fibrils of the PM are assembled into a wide cross-hatched pattern connected by peritrophins (Dinglasan *et al.* 2009). A second group made up of Peritrophin-1 (BTT05886_1, BTT20661_1) and 3 further chitinase transcripts (Chitinase 2

:BTT23246_1, Chitinase 3: BTT39163_1, Chitinase 4: BTT05313_1) is upregulated. Figure 2

shows the correlation of expression patterns between these sixteen transcripts related to chitin

metabolism. There is some clustering, but not of any clear functional groups. Taken together this

differential expression suggests an important role for the repair or restructuring of the peritrophic

matrix in the bumblebees' response to *Crithidia*.

When the BLAST searches against the IIID and nr databases were combined, we found that 89

transcripts relate to canonical insect immune genes. We describe them in the order receptors, serine

proteases, signalling pathways and effectors (Schmid-Hempel 2005).

Receptors:

The Down syndrome cell adhesion molecule (Dscam), a pattern recognition receptor has come to

the forefront of research into insect immune specificity as thousands of different splice forms have

been identified and it is associated with insect immunity (Smith *et al.* 2011). We found five

downregulated transcripts annotated as immunoglobulin superfamily (*Dscam* included in hit list)

(BTT03519_1, BTT08682_1, BTT15814_1, BTT26724_1, BTT27678_1) and one upregulated

transcript (BTT03519_1).

Serine proteases:

Serine proteases are important proteolytic enzymes in many molecular pathways. When these serine

proteases are no longer needed, they are inactivated by serine protease inhibitors (Zhao *et al.* 2012).

CLIP domain serine proteases mediate insect innate immunity (Zou *et al.* 2006). Twenty one

transcripts related to serine proteases, serine protease homologues or serine protease inhibitors were

differentially expressed upon infection (see Table 1). Lipophorin receptor 2 (downregulated

BTT34617_1) binds with serpins to aid in their encytocytosis (Soukup *et al.* 2009).

Signalling pathways:

We found a transcript for *Spatzle* (BTT19738_1) downregulated at this time point. Activation of the

Toll immune pathway requires the activation of *Spatzle* (Lemaitre & Hoffmann 2007). MyD88

(upregulated BTT15687_1) is a death domain-containing adaptor activated by Toll leading to the

activation of *Pelle*. *Dorsal* (BTT25273_1) was also downregulated. The nuclear translocation of

Dorsal, a member of the NF-kB family, in the Toll pathway induces the expression of many

immune genes. We found an upregulated transcript (BTT09662_1) for *Helicase89B* part of the Toll

and Imd Pathway. It is required downstream of NF-kB for the activation of AMP genes in

Drosophila melanogaster (Yagi & Ip 2005). *ird5* codes for a catalytic subunit of an IkappaB kinase

that cleaves Relish. Relish (Imd pathway) is an essential regulator of antimicrobial peptide gene

induction. We found *ird5* (BTT03904_1) to be downregulated 24 hours post-infection.

In mammals semaphorins are crucially involved in various aspects of the immune response

(Takamatsu & Kumanogoh 2012). A semaphorin-5A-like transcript (BTT01850_1) was

downregulated 24 hours post-infection. Semaphorin regulates the activity of Ras-family small

GTPases (Takamatsu & Kumanogoh 2012). A Ras-like protein11B transcript (BTT05368_1) was

also downregulated. The Ras/MAPK pathway was found to be essential for the suppression of the

Imd immune pathway in *Drosophila* (Ragab *et al.* 2011).

The downregulated Drumstick (BTT13062_1) interacts with the JAK/STAT pathway during its'

development role (Johansen *et al.* 2003), but we could not find any information about its immune

role. Two transcripts (BTT11590_1, BTT14205_1) of *Puckered* were downregulated. *Puckered*,

which codes for a dual specificity phosphatase, is a key regulator of the c-Jun-N-terminal kinase

(JNK) immune pathway (Karkali & Panayotou 2012). Mpk2/p38a (downregulated BTT05769_1) is

involved in the JNK Pathway and JAK/STAT Pathway. Heat-shock factor activation by p38 is a

recently discovered part of antimicrobial reactions in flies (Chen *et al.* 2010). We found two heat

shock protein transcripts (BTT23758_2, BTT37030_1) and one other (BTT17701_1) that were

downregulated and upregulated respectively. These are all involved in the JAK/STAT pathway.

Effectors:

In our previous paper (Riddell *et al.* 2011) we found that antimicrobial peptides were upregulated at

24 hours post-infection. We would expect the same trend here. Indeed, we found that five

transcripts for defensin (BTT06274_2, BTT8490_1, BTT10405_1, BTT14019_1, and

BTT42034_1) and three hymenoptaecin transcripts (BTT18071_1, BTT24170_1, BTT24170_2)

were all upregulated. An apidaecin precursor (BTT33652_1) was downregulated. Apidaecin has

recently been shown to be expressed in bumblebees (Colgan *et al.* 2011). The downregulated beta-

amyloid-like protein (BTT20240_1) has been shown to be an antimicrobial peptide in mammals

(Soscia *et al.* 2010). Hemolectin (BTT15326_1, upregulated) is a clotting protein known to have a

role against gram-negative bacteria (Lesch *et al.* 2007).

Reactive oxygen species (ROS) are generated by respiration in the mitochondria or as part of the

immune response (Molina-Cruz *et al.* 2008). P450 cytochromes are oxidases acting terminally in

monooxygenase systems (Felix & Silveira 2012). Some are regulated in response to infection

possibly either as direct immune responders (Vlachou *et al.* 2005), producing nitric oxide (NO) or

other reactive oxygen radicals or as part of the host detoxification process decreasing oxidative

stress after an infection (Molina-Cruz *et al.* 2008). A number of *cytochromes P450* were

differentially expressed 24 hours post infection. Ten cytochrome p450 transcripts (*Cyp4p3*:

BTT05294_1, BTT20848_1, BTT22253_1, BTT23317_1, BTT32674_1, *cytochrome P450 4g15*:

BTT23811_1, BTT32459_1, *cytochrome P450 6k1*: BTT35547_1, BTT40653_1, *cytochrome P450*

6a14: BTT38445_1) were found to be downregulated. Three other *cytochrome P450* transcripts

(*Cyp4p3*: BTT21216_1, BTT35543_1, *cytochrome P450 315a1*: BTT26726_1) were upregulated.

Several other cytochromes (cytochrome b: BTT20524_1, BTT39776_1, BTT41896_1, and

cytochrome c: BTT05255_2) were downregulated.

Numerous other actors in the production of ROS were found to be differentially expressed. *TPX4*

(BTT13285_1), coding for a Thioredoxin-dependent peroxidase, which detoxifies H_2O_2 was

downregulated. This gene was found be differentially expressed during *Plasmodium* infection in

Anopheles gambiae (Baton *et al.* 2009). Calcineurin (BTT08150_1, BTT26273_1) was found to be

downregulated 24 hours post-infection, which agrees with our previous findings (Riddell *et al.*

2011). In infected *D. melanogaster* larvae, NO signals are enhanced by Calcineurin to promote

induction of robust immune responses via the Imd signalling pathway (Dijkers & O'Farrell 2007).

We found downregulation of sortilin-related receptor-like (BTT31654_1). In mammals, sortilin aids

in phagocytosis (Wähe *et al.* 2010). Two downregulated transcripts (BTT35021_1, BTT08756_1)

were matched to *croquemort,* which codes for a key scavenger receptor in the Imd pathway (Franc

et al. 1999). Annexin IX (downregulated BTT02025_1) has been shown to be induced by septic

injury in *Drosophila* and is thought to encode for an anticoagulant (Gregorio *et al.* 2001).

Miscellaneous:

Major royal jelly protein (BTT05317_2, BTT36365_1 upregulated) has been shown to have

antimicrobial properties and to be expressed in response to bacterial infection in honeybees

(Scharlaken *et al.* 2007; Buttstedt *et al.* 2013). Vitellogenin (downregulated BTT36006_1) is a

potent regulator of the immune response in honeybees (Amdam *et al.* 2004). Several orthologs of

putative *Drosophila* immune loci were differentially expressed 24 hours post-infection (CG12505:

BTT00934_1, CG18348: BTT04397_1, CG7296: BTT15035_1, BTT18395_1, CG8791:

BTT18908_1, CG5527: BTT35653_1, Fst: BTT11511_1). The downregulated CG4393

(BTT05817_1) is weakly analogous to TNF receptor associated factor 3 (TRAF3) that mediates

signal transduction involved in mammalian immune responses. Downregulated BTT37289_1 codes

for a putative fatty acyl-CoA reductase.

Genes differentially expressed between 24 hours post-infection and 48 hours post-infection (24

versus 48 hours)

43 transcripts were differentially expressed between 24 hours post-infection and 48 hours post-

infection. Of these 17 had no BLAST hits. A further six had uninformative BLAST hits

(anonymous predicted protein). The remaining 20 were used in the analysis. Defence response was

the only GO term significantly enriched (FDR= 0.00015), with seven transcripts. Three transcripts

correspond to Hymenoptaecin (BTT18071_1, BTT24170_1, BTT24170_2). They were all

upregulated. This suggests a continuing strong AMP production 48 hours after infection. This

agrees with other immune assays in bumblebees (Korner & Schmid-Hempel 2004). Argonaute-2, a

RNA-silencing endonuclease, is involved in antiviral defence in insects (downregulated

BTT02484_1) (van Rij *et al.* 2006). GstD8, a glutathione S-transferase, is involved in the

detoxification process (upregulated BTT04810_1) (Gerardo *et al.* 2010). Dopa decarboxylase

(upregulated BTT28048_1) converts L-dopa to dopamine during the melanisation process (Ferdig *et*

al. 2000). SCR-B9 (upregulated BTT40924_1) codes for a scavenger receptor protein. Scavenger

receptor proteins have been found to be microbial pattern recognition receptors in flies (Rämet *et al.*

2001).

Genes differentially expressed depending on host genotype – parasite genotype interactions

(Specificity)

There were 591 differentially expressed transcripts (FDR < 0.05). Reannotating the transcripts

using Blast2GO (blastx against the nr database with e < 0.001), 150 had no BLAST hits. A further

64 had uninformative BLAST hits (anonymous predicted protein). There were 109 transcripts that

had previously been found to be differentially expressed at 24 hours post infection. A

multidimensional scaling (MDS) plot of the samples based on the expression of these 591 genes.

reveals that the 11 samples group into their colony-strain interaction (Figure 3).

Of the 591 transcripts, 132 were upregulated and 459 were downregulated. Up or downregulation

does not have the same meaning here as in the infected versus uninfected model were there was a

clear baseline (uninfected). Depending on how the GLM is ordered, the reciprocal result could be

observed. Our model used colony K strain 8 as the final contrast. From our previously published

qPCR data (Riddell *et al.* 2009), we know the colony K strain 8 interaction displayed the highest

levels of AMPs (effectors). Therefore when we say a transcript is upregulated, we mean it is

upregulated in this high immune response interaction.

As with the infection data, we combined the BLAST searches against the IIID and nr databases.

Ninety transcripts correspond to canonical insect immune genes. We again describe them in the

order receptors, serine proteases, signalling pathways and effectors (Schmid-Hempel 2005).

Receptors:

Two transcripts were associated with gram-negative binding proteins (upregulated GNBP,

BTT03533_1 and downregulated *GNBP1-2* BTT35513_1). Although GNBPs are most associated

with defense against gram-negative bacteria, they have been show to have a role in response to

Plasmodium infections (Tahar *et al.* 2002). C-type lectins (CTLs) bind carbohydrates and mediate

processes including pathogen recognition (Cirimotich *et al.* 2010). CTL4 is agonist to *Plasmodium*

infections in mosquitoes (Cirimotich *et al.* 2010). A CTL4 transcript (BTT29328_1) was found to

be downregulated.

One downregulated transcript was related to *Dscam* (BTT12755_1). A further fourteen

downregulated transcripts were part of the Ig superfamily (IGFn3-1: BTT05561_1, BTT05581_1,

BTT08682_1, BTT12655_1, BTT13442_1, BTT14516_1, BTT18750_1, BTT21156_1,

BTT22598_1, BTT22819_1, BTT23339_1, BTT24070_1, IGFn3-7: BTT08109_1, BTT09498_1)

and one was upregulated (IGFn3-8: BTT03519_1). *Dscam* and most of the other Ig superfamily

transcripts cluster together in the top right of figure 4, suggesting they are similarly expressed.

Serine proteases:

28 transcripts related to serine proteases, serine protease homologues or serine protease inhibitors

were differentially expressed (see Table 2).

Signalling pathways:

The Toll-like receptor *18Wheeler* (BTT35732_1) and *Toll 10* (BTT09386_1) were both

upregulated. 18Wheeler has been shown to be important in the anti gram-negative immune response

in *Drosophila* larvae (Ligoxygakis *et al.* 2002). *Dorsal 1A* (BTT04010_1), a transcription factor

that is an important part of the Toll pathway, was downregulated. A transcript for *Spatzle 1-2* was

also downregulated (BTT10679_1).

The tyrosine kinase *Pvr* (BTT04822_1), which inhibits JNK activation (Bond & Foley 2009) was

downregulated. *Jun*, a transcription factor of the JNK pathway was downregulated (BTT13636_1).

Mpk2/p38a (downregulated BTT16580_1) and *MAPKKK9* (downregulated BTT04404_1) are

mitogen-activated protein kinases involved in the JNK Pathway and JAK/STAT pathways. We

found two heat shock protein transcripts (BTT17371_1, BTT22195_1) and one other

(BTT17701_1) that were downregulated and upregulated respectively. These are all involved in the

JAK/STAT pathway. *Akt 1* (downregulated BTT14188_1) is part of the insulin/insulin-like growth

factor 1 signaling (IIS) cascade. IIS plays a critical role in the regulation of innate immunity.

Activation of *Akt* signaling leads to a decrease in malaria infection intensity in mosquitoes (Corby-

Harris *et al.* 2010).

Effectors:

Five transcripts relate to the AMPs *defensin* (BTT06274_2, BTT42034_1) and hymenoptaecin (BTT18071_1, BTT24170_1, BTT24170_2). They were all upregulated. An apidaecin precursor (BTT20828_1) was upregulated. Hemolectin had three downregulated transcripts (BTT14194_1, BTT17013_1, BTT26614_1) and one upregulated (BTT15326_1). Argonaute-2, a RNA-silencing endonuclease, is involved in antiviral defense in insects (downregulated BTT02374_1) (van Rij *et al.* 2006).

Eater encodes a transmembrane receptor involved in phagocytosis in *Drosophila* (Kocks *et al.* 2005). A transcript (BTT11132_1) relating to *Eater* was upregulated. The melanisation process component Dopa decarboxylase (BTT19093_1) was upregulated. Another enzyme involved in melanisation, laccase was found to be downregulated (BTT20241_1, BTT33633_1) (Arakane *et al.* 2005).

Cyp4p3 transcript BTT40653_1 was upregulated. Two previously unseen *Cyp4p3* transcripts (BTT05254_1, BTT20622_2) were upregulated and one (BTT36257_1) downregulated. *TPX4* (BTT13285_1) that codes for a Thioredoxin-dependent peroxidase was downregulated.

Miscellaneous:

A small number of transcripts were related to chitin metabolism. SCRASP1 has a chitin-binding domain that has been hypothesized to sense chitin in response to injury and to transduce signals via the serine protease domain (Blumberg *et al.* 2013). We found an upregulated transcript related to *SCRASP 1* (BTT41923_1). A peritrophin precursor was also upregulated (BTT10727_1). As was a chitinase 3 transcript (BTT23246_1).

Retinoid and fatty-acid-binding protein (RfaBp) (BTT07678_1) was downregulated. RfaBp was

found to be upregulated upon injection of LPS in *Drosophila* during a proteomic study (Vierstraete

et al. 2004). *Notch* (upregulated BTT09545_1) is involved in the specification of crystal cells in

Drosophila melanogaster (Mukherjee *et al.* 2011). Finally, several orthologs of putative *Drosophila*

immune loci were found to be differentially expressed (CG5527: BTT08512_1, CG12505:

BTT00934_1, CG13323: BTT38025_1, BTT38087_1, CG17560: BTT02877_1 downregulated,

BTT05845_1 upregulated, CG18348: BTT20843_1).

Discussion

We present a comprehensive transcriptomic analysis of gene expression in this important model host-parasite system. We have identified a large number of bumblebee genes whose expression are changed upon infection with *Crithidia*. We also found a large number of genes whose expression depends on the interaction between host and parasite genotypes and therefore show specificity.

We confirmed the importance of antimicrobial peptides in the specific defence against *Crithidia* (Riddell *et al.* 2009, 2011; Deshwal & Mallon 2014). It is also clear that several other effectors including ROS and phagocytosis may be important. Several immune pathways seem to be important in the anti-*Crithidia* response. These include the Toll, Imd and JAK/STAT pathways. Toll especially seems to be important in a specific immune response.

There are a larger proportion of receptor transcripts found in the specificity analysis (3.2% 19/591) compared to the infection analysis (1.2% 6/489). This is not surprising, as it may be expected that a specific immune response to a given strain would be based mainly on how it is recognised. Although several receptors, including GNBPs and lectins, are differentially expressed, the most exciting discovery is the large number of transcripts related to *Dscam*. The Down syndrome cell adhesion molecule (Dscam), a pattern recognition receptor has come to the forefront of research into insect immune specificity as thousands of different splice forms are generated and it is associated with insect immunity (Smith *et al.* 2011). In the fruit fly *Drosophila*, silencing of *Dscam* retards the insect's capacity to engulf bacteria by phagocytosis (Watson *et al.* 2005). In *Anopheles*, the Dscam splice forms produced in response to parasite exposure differs between bacteria and *Plasmodium* and between *Plasmodium berghei* and *Plasmodium falciparum* (Dong *et al.* 2006). This has been tempered by the finding that Dscam diversity does not increase with exposure to increasing heterogeneity of *Plasmodium* falciparum genotypes (Smith *et al.* 2011). Recently it has been shown that *Dscam* specificity is mediated by the transcriptional regulation of specific splicing

factors downstream of the activation of the Toll and IMD pathways (Dong *et al.* 2012). Our results

suggest that *Dscam* related genes may be important in differentiating strains of the trypanosome

Crithidia bombi.

We found a number of genes associated with chitin metabolism. Through several pathways chitin

metabolism is fundamental to invertebrate immunity (Beckerman *et al.* 2013). As an aside, an

intriguing hypothesis is that chitin metabolism is the nexus through which defense against predators

and against parasites are traded-off (Beckerman *et al.* 2013). Our data suggests that the peritrophic

matrix may be fundamental in the bee's defence against *Crithidia.* The peritrophic matrix acts as an

immunological barrier against trypanosomes. Tsetse flies with an underdeveloped PM have lower

levels of refractoriness to trypanosome infections (Weiss *et al.* 2013). This is due to a premature

immune response; the trypanosomes get through the PM quicker and stimulate the immune

response at an earlier stage compared to refractory flies.

A similar study has recently been published (Barribeau *et al.* 2014). Although they find genotype x

genotype interactions in gene expression, it is in a much smaller number of genes. We hypothesise

that our much larger catalogue of genes, including *Dscam*, is due to our experimental design. We

used a 2x2 analysis of preselected reciporcal host-parasite interactions. This increased our

likelihood of detecting differential expression using RNA-seq compared to a less directed sample

collection.

In this paper we have shown that the expression of immune genes is associated with specific

interactions between different host and parsite genotypes in this bumblebee / trypanosome model. In

future RNAi work we will knockdown candidate genes thereby altering these specific interactions

to directly examine their biological significance.

Acknowledgements

Thanks to S. Barribeau & P. Schmid-Hempel for discussions. CR was funded by a BBSRC

studentship. This work was partially funded through a NERC NBAF pilot grant (NBAF606) to

EBM.

References

Amdam GV, Simões ZLP, Hagen A *et al.* (2004) Hormonal control of the yolk precursor

vitellogenin regulates immune function and longevity in honeybees. *Experimental Gerontology*, **39**,

767–773.

Arakane Y, Muthukrishnan S, Beeman RW, Kanost MR, Kramer KJ (2005) Laccase 2 is the

phenoloxidase gene required for beetle cuticle tanning. *Proceedings of the National Academy of*

Sciences of the United States of America, **102**, 11337–11342.

Barribeau SM, Sadd BM, Plessis L du, Schmid-Hempel P (2014) Gene expression differences

underlying genotype-by-genotype specificity in a host–parasite system. *Proceedings of the National*

Academy of Sciences, 201318628.

Barribeau SM, Schmid-Hempel P (2013) Qualitatively different immune response of the bumblebee

host, Bombus terrestris, to infection by different genotypes of the trypanosome gut parasite,

Crithidia bombi. *Infection, Genetics and Evolution*, **20**, 249–256.

Baton LA, Robertson A, Warr E, Strand MR, Dimopoulos G (2009) Genome-wide transcriptomic

profiling of Anopheles gambiae hemocytes reveals pathogen-specific signatures upon bacterial

challenge and Plasmodium berghei infection. *BMC Genomics*, **10**, 257.

Beckerman AP, de Roij J, Dennis SR, Little TJ (2013) A shared mechanism of defense against

predators and parasites: chitin regulation and its implications for life-history theory. *Ecology and*

Evolution, **3**, 5119–5126.

Blumberg BJ, Trop S, Das S, Dimopoulos G (2013) Bacteria- and IMD Pathway-Independent

Immune Defenses against Plasmodium falciparum in Anopheles gambiae. *PLoS ONE*, **8**.

Bond D, Foley E (2009) A quantitative RNAi screen for JNK modifiers identifies Pvr as a novel

regulator of Drosophila immune signaling. *PLoS pathogens*, **5**, e1000655.

Brucker RM, Funkhouser LJ, Setia S, Pauly R, Bordenstein SR (2012) Insect Innate Immunity

Database (IIID): An Annotation Tool for Identifying Immune Genes in Insect Genomes. *PLoS*

ONE, **7**, e45125.

Brunner FS, Schmid-Hempel P, Barribeau SM (2013) Immune Gene Expression in Bombus

terrestris: Signatures of Infection Despite Strong Variation among Populations, Colonies, and Sister

Workers. *PLoS ONE*, **8**, e68181.

Buttstedt A, Moritz RFA, Erler S (2013) Origin and function of the major royal jelly proteins of the

honeybee (Apis mellifera) as members of the yellow gene family. *Biological Reviews*, 255–269.

Carius HJ, Little TJ, Ebert D (2001) Genetic variation in a host-parasite association: Potential for

coevolution and frequency-dependent selection. *Evolution*, **55**, 1136–1145.

Chen J, Xie C, Tian L *et al.* (2010) Participation of the p38 pathway in Drosophila host defense

against pathogenic bacteria and fungi. *Proceedings of the National Academy of Sciences*, **107**,

20774–20779.

Cirimotich CM, Dong Y, Garver LS, Sim S, Dimopoulos G (2010) Mosquito immune defenses

against Plasmodium infection. *Developmental & Comparative Immunology*, **34**, 387–395.

Colgan TJ, Carolan JC, Bridgett SJ *et al.* (2011) Polyphenism in social insects: insights from a

transcriptome-wide analysis of gene expression in the life stages of the key pollinator, Bombus

terrestris. *BMC Genomics*, **12**, 623.

Corby-Harris V, Drexler A, Watkins de Jong L *et al.* (2010) Activation of Akt Signaling Reduces

the Prevalence and Intensity of Malaria Parasite Infection and Lifespan in Anopheles stephensi

Mosquitoes. *PLoS Pathog*, **6**, e1001003.

Danielli A, Loukeris TG, Lagueux M *et al.* (2000) A modular chitin-binding protease associated

with hemocytes and hemolymph in the mosquito Anopheles gambiae. *Proceedings of the National*

Academy of Sciences, **97**, 7136–7141.

Deshwal S, Mallon EB (2014) Antimicrobial peptides play a functional role in bumblebee anti-

trypanosome defense. *Developmental & Comparative Immunology*, **42**, 240–243.

Dijkers PF, O'Farrell PH (2007) Drosophila Calcineurin Promotes Induction of Innate Immune

Responses. *Current Biology*, **17**, 2087–2093.

Dinglasan RR, Devenport M, Florens L *et al.* (2009) The Anopheles gambiae adult midgut

peritrophic matrix proteome. *Insect Biochemistry and Molecular Biology*, **39**, 125–134.

Dong Y, Cirimotich CM, Pike A, Chandra R, Dimopoulos G (2012) Anopheles NF-κB-regulated

splicing factors direct pathogen-specific repertoires of the hypervariable pattern recognition

receptor AgDscam. *Cell Host and Microbe*, **12**, 521–530.

Dong YM, Taylor HE, Dimopoulos G (2006) AgDscam, a hypervariable immunoglobulin domain-

containing receptor of the Anopheles gambiae innate immune system. *Plos Biology*, **4**, 1137–1146.

Felix R, Silveira H (2012) The Role of Anopheles gambiae P450 Cytochrome in Insecticide

Resistance and Infection. In: *Insecticides - Pest Engineering* . InTech.

Ferdig MT, Taft AS, Smartt CT *et al.* (2000) Aedes aegypti dopa decarboxylase: gene structure and

regulation. *Insect molecular biology*, **9**, 231–239.

Franc NC, Heitzler P, B RA, Ezekowitz, White K (1999) Requirement for Croquemort in

Phagocytosis of Apoptotic Cells in Drosophila. *Science*, **284**, 1991–1994.

Gerardo NM, Altincicek B, Anselme C *et al.* (2010) Immunity and other defenses in pea aphids,

Acyrthosiphon pisum. *Genome Biology*, **11**, R21.

Götz S, García-Gómez JM, Terol J *et al.* (2008) High-throughput functional annotation and data

mining with the Blast2GO suite. *Nucleic Acids Research*, **36**, 3420–3435.

Gregorio ED, Spellman PT, Rubin GM, Lemaitre B (2001) Genome-wide analysis of the

Drosophila immune response by using oligonucleotide microarrays. *Proceedings of the National*

Academy of Sciences, **98**, 12590–12595.

Gschwend TP, Krueger SR, Kozlov SV, Wolfer DP, Sonderegger P (1997) Neurotrypsin, a Novel

Multidomain Serine Protease Expressed in the Nervous System. *Molecular and Cellular*

Neuroscience, **9**, 207–219.

Hauton C, Smith VJ (2007) Adaptive immunity in invertebrates: a straw house without a

mechanistic foundation. *Bioessays*, **29**, 1138–1146.

Isogai Z, Ono RN, Ushiro S *et al.* (2003) Latent transforming growth factor β-binding protein 1

interacts with fibrillin and is a microfibril-associated protein. *Journal of Biological Chemistry*, **278**,

2750–2757.

Johansen KA, Iwaki DD, Lengyel JA (2003) Localized JAK/STAT signaling is required for

oriented cell rearrangement in a tubular epithelium. *Development*, **130**, 135–145.

Karkali K, Panayotou G (2012) The Drosophila DUSP Puckered is phosphorylated by JNK and p38

in response to arsenite-induced oxidative stress. *Biochemical and Biophysical Research*

Communications, **418**, 301–306.

Kocks C, Cho JH, Nehme N *et al.* (2005) Eater, a Transmembrane Protein Mediating Phagocytosis

of Bacterial Pathogens in Drosophila. *Cell*, **123**, 335–346.

Korner P, Schmid-Hempel P (2004) In vivo dynamics of an immune response in the bumble bee

Bombus terrestris. *Journal of Invertebrate Pathology*, **87**, 59–66.

Kuraishi T, Binggeli O, Opota O, Buchon N, Lemaitre B (2011) Genetic evidence for a protective

role of the peritrophic matrix against intestinal bacterial infection in Drosophila melanogaster.

Proceedings of the National Academy of Sciences of the United States of America, **108**, 15966–

15971.

Lambrechts L (2010) Dissecting the Genetic Architecture of Host-Pathogen Specificity. *PLOS*

Pathogens, **6**.

Lehane MJ (1997) Peritrophic matrix structure and function. *Annual review of entomology*, **42**,

525–550.

Lemaitre B, Hoffmann J (2007) The host defense of Drosophila melanogaster. *Annual Review of*

Immunology, **25**, 697–743.

Lesch C, Goto A, Lindgren M *et al.* (2007) A role for Hemolectin in coagulation and immunity in

Drosophila melanogaster. *Developmental and comparative immunology*, **31**, 1255–1263.

Ligoxygakis P, Bulet P, Reichhart J-M (2002) Critical evaluation of the role of the Toll-like

receptor 18-Wheeler in the host defense of Drosophila. *EMBO reports*, **3**, 666–673.

Mallon EB, Loosli R, Schmid-Hempel P (2003) Specific versus nonspecific immune defense in the

bumblebee, Bombus terrestris L. *Evolution*, **57**, 1444–1447.

McCarthy DJ, Chen, Y., Smyth GK (2012) Differential expression analysis of multifactor RNA-Seq

experiments with respect to biological variation. *Nucleic Acids Research*, **40**, 4288–4297.

Molina-Cruz A, DeJong RJ, Charles B *et al.* (2008) Reactive oxygen species modulate Anopheles

gambiae immunity against bacteria and Plasmodium. *The Journal of biological chemistry*, **283**,

3217–3223.

Moret Y, Schmid-Hempel P (2009) Immune responses of bumblebee workers as a function of

individual and colony age: senescence versus plastic adjustment of the immune function. *Oikos*,

118, 371–378.

Mukherjee T, Kim WS, Mandal L, Banerjee U (2011) Interaction between Notch and Hif-alpha in

development and survival of Drosophila blood cells. *Science (New York, N.Y.)*, **332**, 1210–1213.

R Core Team (2013) *R: A language and environment for statistical computing.* R foundation for

statistical computing, Vienna, Austria.

Ragab A, Buechling T, Gesellchen V *et al.* (2011) Drosophila Ras/MAPK signalling regulates

innate immune responses in immune and intestinal stem cells. *The EMBO Journal*, **30**, 1123–1136.

Rämet M, Pearson A, Manfruelli P *et al.* (2001) Drosophila scavenger receptor CI is a pattern

recognition receptor for bacteria. *Immunity*, **15**, 1027–1038.

Riddell C, Adams S, Schmid-Hempel P, Mallon EB (2009) Differential Expression of Immune

Defences Is Associated with Specific Host-Parasite Interactions in Insects. *PLoS ONE*, **4**, e7621.

Riddell CE, Sumner S, Adams S, Mallon EB (2011) Pathways to immunity: Temporal dynamics of

the bumblebee (Bombus terrestris) immune response against a trypanosomal gut parasite. *Insect*

Molecular Biology, **20**, 529–540.

Van Rij RP, Saleh M-C, Berry B *et al.* (2006) The RNA silencing endonuclease Argonaute 2

mediates specific antiviral immunity in Drosophila melanogaster. *Genes & development*, **20**, 2985–

2995.

Scharlaken B, De Graaf DC, Memmi S *et al.* (2007) Differential protein expression in the honey

bee head after a bacterial challenge. *Archives of Insect Biochemistry and Physiology*, **65**, 223–237.

Schlüns H, Sadd BM, Schmid-Hempel P, Crozier RH (2010) Infection with the trypanosome

Crithidia bombi and expression of immune-related genes in the bumblebee Bombus terrestris.

Developmental and comparative immunology, **34**, 705–709.

Schmid-Hempel P (2001) On the evolutionary ecology of host-parasite interactions: addressing the

question with regard to bumblebees and their parasites. *Naturwissenschaften*, **88**, 147–158.

Schmid-Hempel P (2005) Natural insect host-parasite systems show immune priming and

specificity: puzzles to be solved. *Bioessays*, **27**, 1026–1034.

Schmid-Hempel P, Puhr K, Kruger N, Reber C, Schmid-Hempel R (1999) Dynamic and genetic

consequences of variation in horizontal transmission for a microparasitic infection. *Evolution*, **53**,

426–434.

Schmid-Hempel P, Reber-Funk C (2004) The distribution of genotypes of the trypanosome parasite,

Crithidia bombi, in populations of its host Bombus terrestris. *Parasitology*, **129**, 147–158.

Smith PH, Mwangi JM, Afrane YA *et al.* (2011) Alternative splicing of the Anopheles gambiae

Dscam gene in diverse Plasmodium falciparum infections. *Malaria Journal*, **10**, 156.

Soscia SJ, Kirby JE, Washicosky KJ *et al.* (2010) The Alzheimer's Disease-Associated Amyloid β-

Protein Is an Antimicrobial Peptide. *PLoS ONE*, **5**, e9505.

Soukup SF, Culi J, Gubb D (2009) Uptake of the Necrotic Serpin in Drosophila melanogaster via

the Lipophorin Receptor-1. *PLoS Genet*, **5**, e1000532.

Supek F, Bošnjak M, Škunca N, Šmuc T (2011) REVIGO Summarizes and Visualizes Long Lists

of Gene Ontology Terms. *PLoS ONE*, **6**, e21800.

Tahar R, Boudin C, Thiery I, Bourgouin C (2002) Immune response of Anopheles gambiae to the

early sporogonic stages of the human malaria parasite Plasmodium falciparum. *The EMBO Journal*,

21, 6673–6680.

Takamatsu H, Kumanogoh A (2012) Diverse roles for semaphorin−plexin signaling in the immune

system. *Trends in Immunology*, **33**, 127–135.

Vierstraete E, Verleyen P, Baggerman G *et al.* (2004) A proteomic approach for the analysis of

instantly released wound and immune proteins in Drosophila melanogaster hemolymph.

Proceedings of the National Academy of Sciences of the United States of America, **101**, 470–475.

Vlachou D, Schlegelmilch T, Christophides GK, Kafatos FC (2005) Functional genomic analysis of

midgut epithelial responses in Anopheles during Plasmodium invasion. *Current biology: CB*, **15**,

1185–1195.

Wähe A, Kasmapour B, Schmaderer C *et al.* (2010) Golgi-to-phagosome transport of acid

sphingomyelinase and prosaposin is mediated by sortilin. *Journal of Cell Science*, **123**, 2502–2511.

Watson FL, Püttmann-Holgado R, Thomas F *et al.* (2005) Extensive Diversity of Ig-Superfamily

Proteins in the Immune System of Insects. *Science*, **309**, 1874–1878.

Weiss BL, Wang J, Maltz MA, Wu Y, Aksoy S (2013) Trypanosome Infection Establishment in the

Tsetse Fly Gut Is Influenced by Microbiome-Regulated Host Immune Barriers. *PLoS Pathog*, **9**,

e1003318.

Wu TD, Nacu S (2010) Fast and SNP-tolerant detection of complex variants and splicing in short

reads. *Bioinformatics (Oxford, England)*, **26**, 873–881.

Yagi Y, Ip YT (2005) Helicase89B is a Mot1p/BTAF1 homologue that mediates an antimicrobial

response in Drosophila. *EMBO reports*, **6**, 1088–1094.

Zhao P, Dong Z, Duan J *et al.* (2012) Genome-Wide Identification and Immune Response Analysis

of Serine Protease Inhibitor Genes in the Silkworm, Bombyx mori. *PLoS ONE*, **7**, e31168.

Zou Z, Lopez DL, Kanost MR, Evans JD, Jiang H (2006) Comparative analysis of serine protease-

related genes in the honey bee genome: possible involvement in embryonic development and innate

immunity. *Insect Molecular Biology*, **15**, 603–614.

Figure 1 A summary of the enriched GO terms (based on Blast2Go annotation) found for differentially expressed genes at 24 hours (infected versus uninfected). This figure was produced using Revigo *(Supek* et al. *2011).*

Figure 2. A heatmap showing the correlations of the expression patterns of the transcripts labelled as chitin metabolism genes that where differentially expressed twenty four hours post-infection compared to uninfected samples (infected versus uninfected).

Figure 3. A multidimensional scaling (MDS) plot of the 11 samples used in the specificity analysis based on the expression of 591 differentially expressed transcripts. There are two colonies (K (red) and Q (blue)) and two *Crithidia* strains (6 (light) and 8 (dark)). A/B/C refers to replicates. Dimension 1 is the direction that best separates the samples. Dimension 2 is the next best direction, uncorrelated with the first, that separates the samples. The leading log-fold-change is the average (root-mean-square) of the largest absolute log-fold- changes between each pair of samples.The samples are grouped into their colony-strain interaction.

Figure 4. A heatmap showing the correlations of the expression patterns of the 90 transcripts labelled as immune genes in the analysis identifying genes differentially expressed depending on host genotype – parasite genotype interactions (Specifcity).

Serine Proteases	Upregulated	Downregulated
CLIPA6		BTT20125_1
CLIPA7		BTT07313_1, BTT31897_1
CLIPD5		BTT10579_1, BTT10912_1, BTT18247_1 BTT25711_1, BTT06803_1
SP24	BTT03436_1	
SP27		BTT08108_1, BTT38696_1
SP35	BTT05300_1	
Serine protease homologues	Upregulated	Downregulated
SPH54	BTT01977_1	BTT06125_1
Serine protease inhibitors	Upregulated	Downregulated
NEC-like		BTT31997_1
spn4		BTT04130_1, BTT40693_1, BTT41025_1, BTT41461_1
SRPN10		BTT04508_1, BTT20259_1

Table 1. List of transcriptions associated with serine proteases and serine protease inhibitors found to be differentially expressed twenty four hours post infection with Crithidia bombi (Infected versus uninfected).

Serine Proteases	Upregulated	Downregulated
cSp3	BTT35293_1	BTT10579_1, BTT10912_1, BTT18247_1, BTT25711_1
Sp18	BTT20808_1	
Sp27	BTT40251_1	
Sp28		BTT20637_1
Sp35	BTT05300_1	BTT10155_1
Sp40	BTT15256_1	
Tequilla/GRAL/Sp23	BTT01709_1, BTT05886_1, BTT09081_1, BTT20661_1, BTT20725_1, BTT24359_1, BTT25071_1	
Serine protease homologues	Upregulated	Downregulated
cSPH39		BTT21868_1
Sph54		BTT27769_1
Sph56	BTT17814_1	
Serine protease inhibitors	Upregulated	Downregulated
Kunitz ser-protease inhibitor		BTT14993_1
necrotic (nec)	BTT35742_1	
Spn 4		BTT04130_1
SRPN10		BTT02607_1, BTT4508_1, BTT20259_1, BTT40693_1

Table 2. List of transcriptions associated with serine proteases and serine protease inhibitorsfound to be differentially expressed in the specificity analysis.