T-lex2: genotyping, frequency estimation and re-annotation of transposable

elements using single or pooled next-generation sequencing data

Anna-Sophie Fiston-Lavier[1,2], Maite G. Barrón[3,4], Dmitri A. Petrov[1] and Josefa González[4]

[1]Department of Biology, Stanford University, Stanford, CA 94305-5020, USA

[2]Institut des Sciences de l'Evolution (ISEM), UMR5554 CNRS-Université Montpellier 2, France

[3]Genomics, Bioinformatics and Evolution Group, Institut de Biotecnologia i de Biomedicina -

IBB/Department of Genetics and Microbiology, Campus Universitat Autònoma de Barcelona, Bellaterra,

08193, Spain.

[4]Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Barcelona, 08003, Spain.

Corresponding author:
Anna-Sophie Fiston-Lavier, PhD
Assistant professor, University Montpellier2,
Adaptation Genomic team, Institute of Sciences of Evolution (ISEM),
Bâtiment 22, 1er étage,
34095 Montpellier, FRANCE
+33 0467143289
asfiston@univ-montp2.fr

22
23
24
Abbreviations:
TE, transposable element
NGS, next-generation sequencing
LTR, long-terminal repeat
TSD, target site duplication
PTS, putative target site
PE, paired-end

ABSTRACT

Transposable elements (TEs) are the most active, diverse and ancient component

in a broad range of genomes. As such, a complete understanding of genome function

and evolution cannot be achieved without a thorough understanding of TE impact and

biology. However, in-depth analyses of TEs still represent a challenge due to the

repetitive nature of these genomic entities. In this work, we present a broadly applicable

and flexible tool: T-lex2. T-lex2 is the only available software that allows routine,

automatic, and accurate genotyping of individual TE insertions and estimation of their

population frequencies both using individual strain and pooled next-generation

sequencing (NGS) data. Furthermore, T-lex2 also assesses the quality of the calls

allowing the identification of miss-annotated TEs and providing the necessary

information to re-annotate them. Although we tested the fidelity of T-lex2 using the high

quality *Drosophila melanogaster* genome, the flexible and customizable design of T-

lex2 allows running it in any genome and for any type of TE insertion. Overall, T-lex2

represents a significant improvement in our ability to analyze the contribution of TEs to

genome function and evolution as well as learning about the biology of TEs. T-lex2 is

freely available online at http://petrov.stanford.edu/cgi-bin/Tlex.html.

INTRODUCTION

The key question in genomics is how genomes vary and evolve at both large and fine scales. In order to answer such question, we need to be able to study genome variation, *i.e.* identifying and analyzing functionally relevant SNPs (Single Nucleotide Polymorphisms) and structural variants, both within and between populations. Next-generation sequencing (NGS) technology has revolutionized this field by allowing one to study variation of a large number of individuals and even cells within individuals (1-3). Unfortunately, NGS technology generally provides fairly short, even if extremely abundant, sequencing reads and thus is not perfectly designed for the study of structural variants (4).

The study of one class of structural variants, transposable elements (TEs), is particularly difficult to carry out using NGS data. TEs are repetitive, ubiquitous and dynamic components of genomes that often vary in genomic location among members of the same species. TEs are classified in three orders: DNA transposons, long-terminal repeat (LTR) elements, and non-LTR elements, and each TE order is represented by several TE families (5,6). This classification highlights the biological diversity in terms of sequence, dynamics and evolution of these repetitive elements that makes their analysis challenging.

TEs represent a large part of most of the eukaryotes genomes (7). A recent study suggests that more than two-thirds of the human genome is composed of TEs (8), and in plants, TEs may represent up to 90% (9). TEs are responsible for a large number of mutations both in populations and somatically within individuals (10,11). Although most of the TE-induced mutations are deleterious, evidence for an adaptive role of TE-induced mutations is starting to accumulate (12,13). TEs can provide active promoter,

splice site, or terminator features that can affect the expression, structure and function of

nearby genes (14). TEs are also involved in the creation of transcriptional regulatory

networks, and in the generation of chromosomal rearrangements (12,15). Thus, knowing

the role of TEs on genome dynamics and evolution, it is crucial to identify and quantify

their impact (16,17). Given the increased number of sequenced genomes, a tool that

allows routine and automatic genotyping of individual TE insertions using a large

number of NGS samples is needed.

Recently several accurate computational approaches have been developed to

analyze TE insertions using NGS data (10,18-22). Designed for specific projects, all

these tools were built with a limited number of features, *e.g.* some tools were designed

to specifically call particular families of TEs (10), while others were designed

exclusively for pooled NGS data (22). Unfortunately, none of the currently available

tools allows a complete and in-depth analysis of individual annotated TE insertions.

With the idea of providing a tool to automatically call TE insertions in multiple

genomes and to estimate their population frequency, we recently designed an integrated,

flexible pipeline called T-lex (23). While most of the available approaches only detect

the presence of TE insertions, T-lex launches two complementary and distinct TE

detection modules: one to detect the "presence" and another to detect the "absence" of a

TE insertion. Both detection approaches are based on the analysis of the junction

sequences of a TE insertion and its flanking sequences as defined by genome annotation.

Thus, the accuracy of the TE calls mainly depends on the quality of the annotation and

more specifically on the delimitation of the individual TE insertions. Unfortunately, the

detection of the precise TE insertion sites is still one of the main challenges of the TE

annotation process (24). The veracity of the TE calls also depends on the genomic

environment of the TE insertion. For instance, a TE insertion located inside a

duplication with one of the copies lacking the TE insertion can be miscalled as

heterozygote (*i.e.*, both present and absent). In addition, a TE insertion flanked by other

repeats such as low-complexity regions or other TE sequences can also be miscalled.

Here, we present T-lex2, the version 2 of the original T-lex pipeline (23).

Besides improving the accuracy of the TE frequency estimate, T-lex2 now assesses the

quality of the TE calls. To achieve this, T-lex2 uses information from (i) a new module

specifically designed to identify target site duplications (TSDs), and (ii) the genomic

context of each insertion, to identify putatively miss-annotated TE insertions highly

likely to produce wrong calls. TSDs are short duplications, from two base pairs to 20 bp,

flanking the TE sequences as a result of the transposition mechanism of most TE

families. While current TSD detection approaches require knowing the biology of the

TEs and are limited to LTR or non-LTR elements, T-lex2 TSD detection module works

without *a priori* knowledge, and for all type of TEs. This new module allows to

precisely delimit the location of each TE insertion. In this new version, frequency of TE

insertions can be estimated using individual and pooled NGS data.

We provide evidence that the new features of T-lex2 help improving the

genotyping and frequency estimate of TE insertions, and allow re-annotating TE

locations using both individual and pooled *Drosophila melanogaster* genomic data.

MATERIAL AND METHODS

T-lex2 pipeline overview

    T-lex2 is composed of five modules that can be run with individual strain or pooled NGS data (Figure 1). (i) The TE-analysis module investigates the flanking regions of the known TE insertions (*i.e.* annotated in a reference genome) to identify those likely to return wrong calls. (ii) The TE-presence detection and (iii) the TE-absence detection modules combine mapping and read depth coverage information to identify reads providing evidence for the presence and for the absence, respectively, of the known TE insertions. (iv) Then the TE-combine module combines the results of the presence and absence detection modules to genotype TEs and/or to estimate their frequencies in populations. Finally, (v) the TE-TSD detection module annotates TSDs in an unbiased and accurate way. Each of these modules can be launched independently and provides a large number of options allowing the user to carry out flexible and customizable analyses. A detailed manual describing step-by-step how to run T-lex2 and listing all Tlex-2 options is available on line at http://petrov.stanford.edu/cgi-bin/Tlex2_manual.html.

TE-analysis module. This module analyses the flanking sequences of each known TE insertion to identify features that might interfere with the presence and absence detection modules. The TE-analysis module (i) provides information about the presence of repetitive elements using RepeatMasker (25), (ii) identifies miss-annotated poly-A tails, and (iii) flags the TE insertions that are part of segmental duplications.
(i) To identify the presence of repetitive elements, the TE-analysis module extracts the flanking sequences of each TE insertion (125 bp by default; see option "-f") and

launches RepeatMasker to annotate the TEs, simple repeats and low-complexity regions

(25). The option "-s" allows specification of the name of the model organism and thus

specifies the repeat library that needs to be used by RepeatMasker (25). If at least one

flanking region shows a repeat density greater than the pre-specified value (50% by

default; option "-d") the TE insertion is flagged as flanked by repetitive elements. By

default, these TE insertions are filtered out. This filtering step can be bypassed using the

option "-noFilterTE".

(ii) A new feature of the TE-analysis module identifies putatively miss-annotated poly-A

tails by searching for stretches of A's or T's (by default more than five base pairs) at the

TE junctions, without *a priori* knowledge of the TE type. Non-LTR elements harbor

poly-A tails at their 3'end that are known to be highly variable in length. Such

sequences are therefore very difficult to annotate automatically. As RepeatMasker only

annotates repeats longer than 20 bp, it cannot be used to identify short stretches of A's

or T's that may correspond to miss-annotated poly-A tail sequences (25).

(iii) When a TE insertion is part of duplication, other copies of the same duplication may

not contain the insertion. The analysis of the flanking sequences of such TE insertions

may lead to call this TE insertion as present and absent while it should only be called as

present. We added one new feature in the analysis of the TE flanking regions to identify

the TE insertions fully or partially part of segmental duplications. For each TE insertion,

the two extracted flanking sequences are blated against a reference genome (26) (BLAT

v. 34 default parameters; http://www.soe.ucsc.edu/~kent). When more than 50% of a

flanking sequence matches at more than one location on the reference genome with

more than 90% of sequence identity, the TE insertion is flagged as part of a duplication.

TE insertions partially part of duplication (notified as "sd_left" or "sd_right") are

distinguished from the TE fully part of a duplication (notified as "sd"). When a TE is

fully part of a duplication while another copies do not contain the TE insertion, this TE

insertion is notified as "sd_noTE".

The analysis of the TE flanking sequences including the RepeatMasker output, whether

the TE is flanked by a longer than annotated poly-A tail, and the segmental duplication

analysis results, are stored in a sub-directory called "Tanalyses".

TE-presence detection module. The presence detection module detects sequencing

reads overlapping the flanking regions of the TE insertion (Figure 2A). The junction

sequences of each TE insertion are extracted and used as reference sequences. By

default, each TE junction sequence encompasses the terminal TE sequence (60 bp from

each end) and the flanking sequence (1 kilo base on each side). The lengths of the TE

sequence and the flanking sequence can be changed using the options "-b" and "-j"

respectively. While the original T-lex used a 20 bp region of the TE sequence, T-lex2

uses a 60 bp region at this step. The increased length improves sensitivity and can now

be used given the increased read lengths in the NGS data compared to even a few years

ago. The reads are mapped on these new extracted reference sequences (Figure 2A) and

are used to build a contig using Phrap (27) (version 1.090518; http://phrap.org). While

the original T-lex uses all the mapped reads at this step, T-lex2 increases the stringency

by using only the reads mapping with a minimum Phrep quality score (30 by default;

this can be changed using the option "-minQ"). The pipeline also reports now the

number of reads mapped on the TE junction at the minimum Phrep quality score (Figure

2A). Besides increasing the minimum length of the match inside and outside the TE

sequence that was required by the original version of the pipeline in order to return a

"presence" call, the new version of T-lex also requires a minimum read sequence

identity (Figure 2A). These two parameters are set at 15 bp and 95% by default and can

be modified using the options "-limp" and "-id", respectively. By default, multiple

alignments of the contigs for each TE side are also returned. The analysis and TE calls

from this module are stored in a sub-directory called "Tpresence".

TE-absence detection module. The absence module detects reads overlapping both

sides of the TE insertion, *i.e.* reads spanning across the TE insertion site (Figure 2B). It

starts by extracting the two flanking sequences for each TE insertion (125 bp each by

default; see option "–f"). The two sequences are then concatenated and the new

constructed sequence is used as the reference sequence for the absence detection (Figure

2B). This new constructed sequence approximates the ancestral sequence prior to the TE

insertion. However, note that traces of the TE insertion mechanism, called TSDs, may

also be encompassed in the new constructed sequence and located at the TE insertion

breakpoint. The length of the extracted sequences should be longer than the reads

themselves and total length of the new constructed sequence should be similar to the

insert size of the paired-end (PE) data in order to get reads and/or pairs spanning the TE

insertion breakpoints. T-lex2 uses now by default a length of 125 bp matches (instead of

100 bp in the original T-lex) for libraries of 100 bp read length and an insert size of 250

bp. The reads are then mapped on the new constructed sequence using SHRiMP2 (28)

(version 2.2.1, October 2011) as opposed to the original version of SHRiMP (29) used in

the original T-lex version. One major improvement of SHRiMP2 is that it can now

handle PE read sequencing data.

SHRiMP2, as SHRiMP, was specifically designed to handle long gaps and

polymorphisms (28,29). Such features allow the mapping of reads despite the presence

of the TSDs or despite miss-annotation. To handle long gaps, SHRiMP2 is launched

with a Smith-Waterman gap open score of -40 and a gap extension score of -1 for both

query and reference sequences. We also allow higher divergence with a Smith-

Waterman mismatch score of -20. SHRiMP2 maps the reads in every pair together (28)

(see option "-pairends"). If a pair does not map, SHRiMP2 attempts to map each read

individually (28). If the two reads of a pair map to the new constructed reference

sequence, only the pairs with reads mapping on opposite strands are selected (*i.e.,*

concordant pairs). When the two reads from the pair overlap in the mapping, the absence

module merges them and considers the pair as a single read. The absence module

specifically looks for "reads" spanning the TE junction with at least 15 bp (parameters

by default; see option "–v") of overlap on each side. T-lex2 then runs RepeatMasker on

the selected reads (see option "-noint") in order to test whether the mapping is due

primarily to simple repeats or low-complexity regions (25). After this step, T-lex2

selects the reads that have at least five non-repetitive (low-complexity or satellites) base

pairs at each end that both map to the flanking regions. Because of the stringency of our

approach, even a single read mapping to both flanking sides and thereby spanning the

insertion site of the TE is sufficient for the absence detection module to classify the TE

insertion as "absent" (Figure 2B). If no reads overlap the TE junction, two possibilities

are considered: (i) the TE is present in the strain (or fixed in the pool sample) or (ii) the

coverage of the data used is insufficient to detect reads providing evidence for the

absence. In order to distinguish between these two possibilities, the absence detection

module checks whether reads map on both flanking sides but fail to map over the

junction. If this is the case, given that the flanking regions are similar in length to the

reads themselves, the module concludes that the TE is present.

If reads do not map to both flanking sides and do not map over the junction site, the

module concludes that the coverage is insufficient and returns a "no call" as a result. By

default, the multiple alignments of the contigs for each TE breakpoint are also returned.

The analysis and TE calls from this module are stored in a sub-directory called

"Tabsence".

TE-combine module. The results from the presence and absence detection approaches

can then be combined to return definitive TE detection calls. We defined five TE call

categories based on the evidence for presence and absence. Briefly, a TE is called

"present" or "absent" when the calls from both detection modules are congruent. A TE

is called as "polymorphic" when this TE is clearly detected as present and absent (*i.e.*,

the presence detection module detects it as present and the absence detection module

detects it as absent). A TE is called as "present/polymorphic" when the presence

detection module detects it as present while the absence detection module fails to return

a call. A TE is called as "absent/polymorphic" if the TE is detected as absent by the

absence detection approach, while the presence detection fails (see Table 1 in (23)).

T-lex2 can also combine the TE calls from several NGS data from the same

population and can estimate the population frequency of each annotated TE insertion

(see options "–combRes" and "-combine"). In T-lex2, the probability for a TE to be

present is associated with each of the above categories as follows: "present" 100%,

"present/polymorphic" 75%, "polymorphic" 50%, "absent/polymorphic" 25%, "absent"

zero percent, respectively.

Finally, T-lex2 pipeline can also estimate the TE frequency using pooled NGS

data by taking into account the number of reads supporting the presence and the number

of reads supporting the absence. In this case, the user just needs to add in the command

line the option "-pooled". This approach is based on two assumptions: (i) individuals in

the pooled sample are equally represented, and (ii) the read-depth coverage is correlated

with the occurrence of a TE sequence in a population. Based on such assumptions, we

expect to observe a positive correlation between the frequency and the number of reads

providing evidence for the presence. We also expect to observe a negative correlation

between the frequency and the number of reads providing evidence for the absence. The

number of copies of TEs from the same family may also have an impact on the number

of reads mapping at the TE breakpoints suggesting that the number of reads supporting

the presence can be more biased than the number of reads supporting the absence.

Taking all these expectations into account, we designed and verified a metric based on

the local read depth coverage in the vicinity of the TE insertion to estimate TE

frequencies from pooled data: total number of reads supporting the presence (NP)

divided by the total number of reads supporting the presence (NP) and supporting the

absence (NA), *i.e*. TE frequency = NP/(NP+NA).

The file combining the TE calls from both TE detection modules is called "Tresults".

The TE frequency estimates are stored in the "Tfreq" or Tfreq_pooled" if the option "-

pooled" is specified.

TE-TSD detection module. To detect the TSD of each annotated TE insertion, this

module analyzes the read alignments generated by the absence detection module (Figure

3). As all T-lex2 modules, this new module can be launched independently although it

does require the read alignments from the absence detection module (see option "-tsd").

The TSD detection module looks for tandem duplications located on the absence

reference sequence spanning the TE breakpoint. It starts by assembling all the selected

reads supporting the absence for each TE using the Phrap program (27) (Figure 3).

Because Phrap requires a minimum of three sequences to build a contig, if fewer than

three reads are selected to support the absence call, the reads are considered

independently (Figure 3). Each contig (or read) is then re-aligned on the absence

reference sequence using BLAT v.34 program (26) (default parameters; Figure 3). Only

the absence calls with a gap larger than two base pairs and located close to the TE

breakpoint are selected and analyzed. If no gaps are observed, the TE-TSD detection

module returns "no Gap". If a gap is observed, the motif present in front of the gap is

called the putative target site or PTS (Figure 3). Using the FastaGrep program (available

from bioinfo.ut.ee/download/dl.php?file=6 and executed with the default parameters), a

tool that looks for short and conserved motifs, the TSD detection module looks for the

copy of the PTS on the other side of the breakpoint. Based on the sensitivity of this

program and the short length of the TSDs, we only trust the matches with more than

80% sequence identity (Figure 3). T-lex2 then sorts the FastaGrep hits to identify the

sequence showing the closer and highly conserved motif. If FastaGrep fails to detect a

match, the TSD detection returns "noTSD" as a result, otherwise, the TSD detection

module returns "detected". In the latter case, the PTS and its closest copy sequences are

also returned. Because several contigs can be generated for the same TE insertion,

several TSDs can also be returned. The results of the TSD detection process are stored

in a file called "TSDannot".

TE-TSD detection module flags the TE insertions for which the TSDs cannot be clearly

defined. TSD detection may fail for old TE insertions for which the TSD is too

divergent, for truncated TEs, and when the boundaries of the TE insertions are not well

annotated (Figure 4). The latter TEs can be re-annotated by the user after manual

inspection.

Re-annotation of TE insertions

Because both TE detection approaches are based on the mapping of NGS reads to the

flanking TE sequences, the accuracy of the TE calls depends on having a correct TE

annotation. TE-analysis and TE-TSD detection modules help identifying putatively

miss-annotated TE insertions and thus putatively wrong T-lex2 calls.

A particular TE insertion can be miss-annotated because it is longer than annotated

(Figure 4A.II) or shorter than annotated (Figure 4A.III) in both cases the TE-TSD

detection will fail. If the TE insertion is present in a given genome, both the presence

detection and the absence detection modules will provide the correct call even though

the TE is longer or shorter than initially annotated (Figure 4B). However, if the TE

insertion is absent, and the TE is miss-annotated the presence and the absence modules

could return a wrong call (Figure 4B). For example, if the TE sequence is longer than

the official annotation and the TE is absent, the gap in the reads providing the absence

could be longer than the TSD size and the module could return present as a call (Figure

4B.k). On the other hand, when the TE is shorter than annotated and the TE is absent,

the gap will be present in the reference sequence and not in the reads. In such situation,

the module could return present as a call (Figure 4B.l).

The Tfreq_combine file of T-lex2 provides information from both the TE-

analyses and TE-TSD detection modules, *i.e.* polyA tail longer than the official

annotation and TSD detection failures, that allows classifying these TE insertions as

putatively miss-annotated TEs. This information can be use to perform manual curation

of these insertions and re-annotate them.

Validation dataset

Veracity of T-lex2 calls was assessed by calling a total of 755 well-studied TE

insertions in *Drosophila melanogaster* strains from the Drosophila Genome Reference

Panel (DGRP) project (30,31). Estimates of TE frequency based on pooled-PCR are

available for these 755 TEs (31,32). TE frequency estimates based on pooled-PCR

approach were categorized in four classes: "very rare" (*i.e*. TE frequency < 1.5%),

"rare" (*i.e*., TE frequency ~2-15% but not absent), "common" (*i.e*., TE frequency ~10-

98%) and "fixed" (*i.e*., TE frequency > 98%) (31). The *D. melanogaster* reference

genome (release 5) was downloaded from Flybase website (http://flybase.org). The

annotations of the TE insertions were extracted from the release 5.43

(http://flybase.org). The resequencing data from 86 *D. melanogaster* DGRP single

strains from the freeze 2 were downloaded from the DGRP website (~20X;

http://dgrp.gnets.ncsu.edu/, Supplementary Table S1). We also called the same TE

insertions in the DGRP population using a high-depth sequencing data from a pool of 92

D. melanogaster DGRP strains (~60X) (33). 84 strains are common in the two datasets

(Supplementary Table S1). To investigate the effect of different read-depth coverage

and different number of flies in pooled samples, we also used T-lex2 to analyzed TE

frequencies in four other pools of lower depth coverage described in Zhu *et al*. 2012

(33). All the NGS data used in this work was generated using the *Illumina* technology.

Experimental validation of T-lex2 results was also done for 11 TE insertions known to

be polymorphic (32) in 24 single *D. melanogaster* North American DGRP strains using

a single PCR approach (Supplementary Table S1). The PCR conditions and primers

used were as described in Gonzalez *et al*. 2008 (32).

Statistical analysis

All the statistical analyses were performed using R (http://cran.r-project.org/).

Sensitivity and specificity were estimated using individual PCR frequencies as we have

the T-lex2 frequencies estimates using the same exact strains. Sensitivity and specificity

were computed such as: (i) Sensitivity = number of correct presence calls / total number

of presence calls and (ii) Specificity = number of correct absence calls / total number of

absence calls.

RESULTS

T-lex2 is an improved and expanded version of T-lex

T-lex was originally designed to genotype and estimate frequencies of the known

TE insertions in individual strain NGS data (23). The new version of T-lex incorporates

numerous new features in order to improve genotyping and frequency estimation of

known TE insertions (Table 1). T-lex2 now assesses the quality of the calls by analyzing

the genomic context of each TE insertion and detecting their TSDs that allows

identifying miss-annotated TE insertions. The new version of T-lex takes into account

the sequencing technology advances and can now handle PE sequencing and longer

reads. Unlike the previous version, T-lex2 can be run for individual strains and for

pooled NGS for the TE frequency estimates (Table 1).

Accurate TE frequency estimates using NGS data from individual strains

To test the quality of the T-lex2 calls, we compared T-lex2 TE frequency

estimates with (i) frequency estimates previously obtained using a pooled-PCR approach

(31,32,34) and (ii) frequency estimates obtained in this work by individual strain PCRs.

We first launched the pipeline to estimate the frequency of 755 TE insertions for

which we have an estimate of their population frequencies based on a pooled-PCR

approach (31,32,34). We run T-lex2 on 86 single DGRP (Supplementary Table S1; see

Material and Methods). The TE call distribution, *i.e*. proportion of TEs called as present,

absent, polymorphic, present/polymorphic, absent/polymorphic and no calls, is

consistent among the strains, excepted for 14 strains that differ drastically

(Supplementary Figure S1). For 11 out of the 14 strains, we observed more than 40% of

no calls, and for the other three strains (RAL-379, RAL-362 and RAL-313), more than

70% of TE insertions were detected as present. As we suspect these 14 strains to be

problematic (most likely a combination of poor sequence quality and extensive residual

heterozygosity), we decided to remove them from our analysis and analyze the calls in

the remaining 72 strains (Supplementary Figure S1).

Out of the 755 TE insertions, TE-analysis module detected six that were flanked

by other TE insertions, and three that were part of segmental duplications

(Supplementary Table S2). We removed these nine insertions because they are likely to

produce wrong calls (see Material and Methods). We then filtered out seven TEs for

which T-lex2 failed to return any calls, and 30 TE insertions for which T-lex2 fails to

return a call for more than 75% of the strains (Supplementary Table S2). All of these

removed TE insertions, 46 in total, are likely to be miss-annotated as we failed to call

them in all or most of the strains. We thus estimated the population frequencies of 709

TE insertions using 72 strains. Note that 65 out of the 709 TEs were identified in a first

T-lex2 run as miss-annotated based on results of the TE-TSD detection module (see

below). We manually re-annotated these 65 TE insertions, and we have re-run T-lex2 to

obtain accurate frequency estimates of these TEs.

T-lex2 frequency estimates of the 709 TEs in 72 strains are significantly and

positively correlated with previous estimates based on a pooled-PCR approach (Figure

5A: Spearman's $\varrho = 0.87$, *P-val* $\ll 0.001$). The vast majority of the very rare and fixed

TE insertions are correctly detected by T-lex2 with 92.5% (357/386) of the very rare TE

insertions absent and 85% (91/107) of the fixed TE insertions fixed in the population

(Figure 5A). As the pooled-PCR and T-lex2 frequencies categories were not build the

same way, we expected to observe intermediate profiles for the rare and common TE

insertions. As expected, most of the rare TE insertions (114/146) are detected with a

zero or low frequency, while most of the common TE insertions (67/70) appear frequent

or fixed (Figure 5A).

Overall, we identified only 1.5% (11/709) of clear discrepant estimates between pooled-

PCR and T-lex2 frequency estimates. However, after manual curation, we suspected all

these TE insertions to be miss-classified by the experimental approach or miss-

annotated. For example, insertion *FBti0020042* is classified as present at very low

frequency by T-lex2 but is detected at high frequency using the pooled-PCR approach.

However, a single strain PCR approach confirms that this TE is present at a low

frequency as estimated using T-lex2 (32). The reclassification of this TE insertion,

allows to conclude that T-lex2 returns accurate TE frequencies with only ~1% of wrong

estimates.

To further test the accuracy of T-lex2 calls, we experimentally determined the

frequency of 11 well-studied TE insertions known to be polymorphic and thus providing

all types of calls (32). We then randomly selected 24 strains part of the bigger set of 86

DGRP strains (Supplementary Table S1) and performed single strain PCR for each of

the 11 analyzed TEs. We experimentally called the 11 TE insertions in the same 24

DGRP strains (Supplementary Table S3). We first compared the frequency estimates for

the 11 TE insertions obtained by T-lex2 using all the DGRP strains (86) and the subset

of 24 strains. This analysis confirms the robustness of the frequency estimates using the

subset of 24 strains (Figure 5B: Pearson's $\varrho = 0.96$, *P-val* $<< 0.001$). Out of the 264

calls (11 detections in 24 strains), we removed 53 for which we failed to get a call

experimentally or using T-lex2 and ended up analyzing 211 TE calls. We observed only

11 calls that were different between the experimental data and the T-lex2 calls

(Supplementary Table S3). This represents 5.2% of putatively wrong calls that is close

to the error rate previously estimated for the TE detection using PCR or the previous

version of T-lex, around five percent in both cases (23,31). We finally conservatively

estimated a high TE call accuracy with 99.14% of sensitivity and 89.58% of specificity

(see Material and Methods). Thus, we overall observed accurate TE frequency estimates

comparing the T-lex2 estimates with the pooled-PCR and the individual PCR

experimental estimates.

Accurate TE population frequency estimates using pooled sequencing data

We called the presence/absence of the 709 selected TE insertions using data

from a pool of 92 DGRP strains, containing the 86 single DGRP individual strains

previously used (33) (Supplementary Table S1). For 29 out of the 709 selected TE

insertions, T-lex2 failed to return a call using the pooled sample, most probably due to

low read coverage in these particular regions of the genome. We thus ended up

estimating the population frequency of 680 TE insertions (Supplementary Table S2). As

expected, we found that the number of reads providing evidence for the absence is

negatively correlated with the experimental TE frequency estimates (Figure 6A:

Spearman's ϱ = -0.60, *P-val* << 0.001), while the number of reads providing evidence

for the presence is positively correlated (Figure 6B: Spearman's ϱ = 0.87, *P-val* <<

0.001). Interestingly, the number of reads supporting the presence does not seem to be

strongly biased by the number of copies of TEs from the same family indicating the

accuracy of our approach (see Material and Methods).

We observed significant and strong positive correlations between T-lex2 TE

frequency estimates using the pooled sequencing data and the pooled-PCR determined

frequencies (Figure 6C: Spearman's ϱ = 0.87, *P-val* << 0.001) and between the T-lex2

TE frequency estimates using the pooled sequencing data and the T-lex2 estimates using

single strains (Figure 6D: Pearson's $\varrho = 0.98$, *P-val* << 0.001). This finding supports

that the TE population frequency can be estimated with accuracy using pooled NGS

sample only based on the read depth coverage.

For 25% of the TEs, all of them polymorphic, we noted an overestimation of the TE

frequency using the pooled sample compared to the estimates using single strains

(Figure 6D). However, for the majority of TEs the difference between the two estimates

is <10%, which can be explained by the sampling effect when the pool and the

individual strain libraries are constructed. To get accurate frequency estimates, each

selected individual should be equally represented in the pooled library. Unfortunately,

those conditions are hard to set up or control during library construction. For seven

percent of the TEs, the discrepancy between the two estimates was >10% and the pool

estimate was consistently higher than the individual strain estimate. However, we run T-

lex2 in a different set of individual and pooled strains and although five percent of the

TEs showed discrepant frequencies estimates between pooled and individual samples,

these discrepancies were both over- and under-estimations (Supplementary Figure S2).

These results suggest that T-lex2 frequency estimation is not biased and that the

observed overestimation for the DGRP strains is likely due to some particular feature of

this dataset such as sequencing coverage or number of strains used in pooled *vs*

individual strains.

To assess the effect of the sampling and coverage on the TE frequency estimates using

pooled data, we compared the TE frequency estimates using five different pooled

datasets (Table 2). The read depth coverage varies from 20X to ~60X (33), the number

of strains from 42 to 92, and the number of flies sampled varies from 42 to 134. With

only 42 flies pooled and a minimum coverage of 20X, the correlation is strongly positive

(Table 2). The increase in the number of flies only does not seem to significantly

improve the TE frequency estimates (Table 2). However, the increase of the coverage

but not the number of strains improves significantly the TE frequency estimates (Table

2). Thus, the coverage matters more than the number of flies or strains pooled in the TE

population frequency estimation using pooled sequencing samples. Read-depth coverage

of 20X appears sufficient to obtain good estimates although the increase of the coverage

allows reducing the number of errors in the calling.

T-lex2 detection of TSDs is unbiased and accurate

The new TE-TSD detection module of T-lex2 allows the automatic annotation of TSDs

without *a priori* knowledge of the biology of the TEs. While only one copy of the TSD

is present in the genome without the TE insertion (*i.e.*, the real TE absence), two TSD

copies should be observed in the new constructed reference sequence used by T-lex2 for

the absence detection (*i.e.*, the TE absence reference sequence constructed by

computational removal of the TE sequence from the genome). Based on this

expectation, the new T-lex2 module post-processes the results from the absence module

to analyze the alignments of the reads spanning the TE insertion site to identify the

TSDs. This new module looks for the expected missing sequence in the alignment

flanking the TE insertion site (Figure 3). If the missing sequence corresponds strictly to

the TSD, another copy of this TSD should be detected in the vicinity of the missing

sequence. Contrary to most of the TSD detection approaches (35-37), no expected TSD

length is required for T-lex2 TSD detection.

Out of the 587 TEs with at least one read providing evidence for the absence, T-lex2

detects TSDs for 390 of them (Table 3). TSD length ranges from two base pairs to 19 bp

with an average and a median of five base pairs (Supplementary Figure S3). TSDs of the

LTR retroelements are consistently short: four base pairs long on average. TSD lengths

of DNA elements are also consistent but longer: seven base pairs on average. On the

contrary, TSDs of the non-LTR retroelements (LINE) are not conserved within families

and vary in length from four base pairs to 19 bp (Table 3). Overall, we identified for the

first time TSDs for 10 TE families: six DNA transposon families and four non-LTR

families (Table 3).

To assess the quality of the TSD annotation using T-lex2, we compared 148 TSD results

obtained by T-lex2 with the annotations obtained with *LTRharvest* software (35)

(Supplementary Table S4). For 92 out of the 148 TE insertions, the TSD annotations are

identical in length and motif. For seven TEs, TSDs have the same length but differ in

motif (Supplementary Table S4) and for the other 49 TEs TSD motifs are identical but

differ in length: seven TEs have shorter motifs and 42 TEs have longer motifs. All the

longer TSDs correspond to shifts of only one base pair except for one TSD that is three

base pairs longer than previously annotated. For 13 of these TEs, corresponding to

Copia elements, T-lex2 results are supported by experimental studies indicating that

Copia TSD are five base pairs long (38,39) and not four base pairs as detected with

LTRHarvest (Supplementary Table S4) (35).

TE-TSD detection module did not identify TSDs for 197 TEs out of the 587 with at least

one read providing evidence for the absence. TSD detection may fail for old TE

insertions for which the TSD is too divergent, for truncated TEs, and when the

boundaries of the TE insertions are not well annotated.

T-lex2 allows reannotation of TEs

Combining outputs from the TE-analysis module and the TE-TSD annotation, T-

lex2 allows manual curation and reannotation of TE insertions. To facilitate the

identification of the putatively miss-annotated TEs, information on TEs for which the

TSD detection failed and TEs with poly-A tail longer than annotated, are also given in

"Tfreq_combine" file. TSD detection fails for TEs that are longer (Figure 4A.II) and

shorter (Figure 4A.III) than annotated. We thus manually curated the 197 TEs for which

TSD could not be detected. For 65 out of the 197 TE insertions, manual curation

confirms the miss-annotation. All these 65 TEs are detected experimentally and by T-

lex2 at very low frequency (Supplementary Table S5). 64 of the 65 miss-annotated TE

insertions are longer than annotated and they all correspond to non-LTR TE insertions:

33 have a longer polyA tail than annotated, and the other 32 TEs have one or the two

junctions miss-annotated. The increase in length of these TEs ranges from five base

pairs to 40 bp (12 bp on average; Supplementary Table S5). The only TE insertion

detected as shorter than annotated is an LTR (*FBti0020118*; Supplementary Table S5).

Thus, T-lex2 allows identifying miss-annotated TEs and manual curation allows to

reannotate ~10% (65/709) of the analyzed TEs.

DISCUSSION

We present here the new version of the T-lex pipeline: T-lex2 (Figure 1). This integrated

Perl computational pipeline designed to analyze TE insertions in NGS data is freely

available and user-friendly since only one command line is necessary to run it

(http://petrov.stanford.edu/cgi-bin/Tlex.html). T-lex2 is also flexible. The pipeline is

composed of five distinct modules that can be launched independently and each one of

them includes a large number of options that allows performing flexible and

customizable analyses (see T-lex2 manual at http://petrov.stanford.edu/cgi-

bin/Tlex2_manual.html). T-lex2 runs with single-end and/or PE reads and can be

customized for NGS datasets with different read lengths, read quality and read depth.

Thus, T-lex is not only able to work with the different NGS datasets already available,

but also with upcoming datasets resulting from improvements in the NGS technology.

Briefly, T-lex2 has improved and/or expanded the four modules present in T-lex1 and

has incorporated a new module that allows detecting TSDs in an unbiased and accurate

way (Table 1). Besides individual strain NGS data, T-lex2 can now estimate TE

frequencies from pooled NGS data. This is an important new feature of T-lex2 since

sequencing pooled samples is an inexpensive and efficient means of obtaining

population genome data (22,33). Another important new feature of T-lex2 is the ability

to identify TEs likely to return wrong calls *i.e*. miss-annotated TEs, TEs part of

segmental duplications and TEs flanked by other TEs. This has been accomplished by

the expansion of the TE-filter module of T-lex1 and by the incorporation of the TSD

detection module (Figure 1). Being able to identify TEs likely to give wrong calls is

especially important for the analyses of TE population frequencies in genomes that do

not have a high-quality TE annotation. Note that even in an extremely high quality

genome as the *D*. *melanogaster* one, we re-annotated ~10% of the analyzed TEs. This

result also supports the opportunities offered by the NGS data to improve TE

annotations (40).

Although we tested the reliability of T-lex2 calls and frequency estimates using the fly

genome, the pipeline can be used for any species for which NGS data and TE annotation

is available. T-lex2 can be run for all type of TEs in any species because (i) it is a

flexible pipeline that allows to customize runs for NGS datasets of different qualities,

(ii) it is able to identify putatively wrong calls and re-annotate miss-annotate TEs, and

(iii) no information about the biology of the TE is needed to get accurate frequency

estimates.

There are other available tools that can be used to estimate TE frequencies (10,18-22).

However T-lex2 is to date the only tool that (i) combines presence and absence detection

analysis that allows the identification of heterozygotes and/or polymorphic TE

insertions; (ii) can be used for individual and pooled samples; (iii) provides the user with

an output file with the frequency of the analyzed TEs; (iv) can be easily customize to be

run with different NGS datasets in any organism; and (v) assess the quality of the calls

allowing to re-annotate miss-annotated TEs. Thus overall, T-lex2 is the most broadly

applicable and flexible tool available to date.

The flexibility of T-lex2, exemplified by the incorporation of the TE-TSD detection

module in this new version, will allow us to easily add new modules in a near future. We

are currently working on a new module designed to identify TEs not annotated in

reference genomes. However, T-lex2 can be used to genotype and estimate frequencies

of *de novo* TEs identified with already available software such as RetroSeq (19). The

ability of T-lex2 to re-annotate TEs would be specially useful to study this *de novo* TEs

as their junctions might not be as well annotated as the ones in the reference genomes.

Overall, we are providing a versatile tool that allows exploring the impact of TEs on

genome evolution as well as learning about TE biology. By analyzing the frequency of

TEs in different populations, we will be able to determine which proportion of the TE-

induced mutations are subject to strong purifying selection or are likely to be evolving

under positive selection. Additionally, accurate annotation of TE insertions will allow us

to study for example the target site preferences of different TE families. The availability

of a tool such as T-lex2 improves our ability to explore TE dynamics and TE biology in

an increasing number of species. This is important, as we cannot hope to understand

genome structure and function without a thorough understanding of its most active,

diverse, and ancient component.

ACKNOWLEDGMENTS

We would like to thanks the ScaleGenomics (http://scalegen.com/) and Proclus

(Stanford University; http://www.stanford.edu/) computational platforms that we used to

investigate this study.

FUNDING

This work was supported by grants from the National Institutes of Health

[R01GM089926 to D.A.P.], from the European Commission [PCIG-GA-2011-293860

to J.G.], and from the Spanish Government [BFU-2011-24397 to J. G.]. MGB was

supported by a PIF fellowship from the Universitat Autònoma de Barcelona.

Table 1. Comparison of the original T-lex version and T-lex2 modules and features

T-lex2 Modules	Features	T-lex (23)	T-lex2
TE-analysis	Detection of low-complexity regions	✓	✓
	Detection of repeat-rich regions	✓	✓
	Detection of terminal poly-A stretches	✗	✓
	Detection of TEs part of segmental duplications	✗	✓
TE-presence detection	Analyzes inside the TE	✓ (20bp)	✓ (60bp)
	Read quality filtered	✗	✓ (phrep score ≥ 30)
	Read depth reported	✗	✓
	Min. match length	✓ (5bp)	✓ (15bp)
	Min. sequence identity	✗	✓ (95%)
TE-absence detection	Analyzes on each side of the TE	✓ (100bp)	✓ (125bp)
	PE data analysis	✗	✓
	Read quality filtered	✗	✓ (30)
	Read depth reported	✗	✓
TE-combine	Take into account the no calls	✗	✓
	Estimates using pooled sample	✗	✓
TE-TSD	TSD detection	✗	✓
	Flag of putative miss-annotated TEs	✗	✓

653 **Table 2**. Sampling effect on the population frequency estimates of 680 TE insertions.
654

Sample identifier	Read depth coverage	Nb. of strains pooled	Nb. of flies pooled	% (Nb.) of no calls	Pearson's ϱ [IC95]
B1	20X	42	42	3.82(26)	0.968 [0.963-0.973]
B4	20X	50	50	4.70(32)	0.977 [0.973-0.980]
B3	40X	42	84	0.88(6)	0.980 [0.977-0.983]
B5	40X	92	92	0.88(6)	0.981 [0.978-0.983]
B3-B4	60X	92	134	0(0)	0.985 [0.982-0.987]

Table 3. TSD detection for the 53 TE families analyzed in this study. ND: Not determined.

| TE order | TE superfamily | TE family | T-lex2 | | | | Known TSD length (bp) |
			Nb. of TE insertions with TSD detected	TSD consensus motif[1]	Min. length (bp)	Max. length (bp)	
DNA	hAT	H	11	ACCCACAG	8	8	8[2]
DNA	Mariner/Tc1	FB	7	AAAACCGTC	9	9	ND
DNA	Mariner/Tc1	S2	1	AT	2	2	ND
DNA	P-element	1360	36	GTYYYG	6	7	7[2]
DNA	Transib	hopper	7	CAATG	5	5	5[2]
DNA	Transib	transib2	5	CAVTG	5	5	5[2]
LINE	Jockey	BS	12	✖	4	12	ND
LINE	Jockey	Doc	14	✖	4	12	ND
LINE	Jockey	F-element	6	✖	9	13	~10[3]
LINE	Jockey	G2	4	✖	5	12	~10[3]
LINE	I-element	I-element	2	✖	15	19	ND
LINE	Jockey	Ivk	1	CACATCATCTTAT	13	13	~10[4]
LINE	Jockey	jockey	7	✖	7	15	ND
LINE	Jockey	Juan	1	ACGAAACAACATTA	14	14	ND
LINE	R1	Rt1a	2	✖	6	14	ND
LINE	R1	Rt1b	1	GCTATC	6	6	ND
LINE	Jockey	X-element	1	TTTTGAAA	8	8	ND
LTR	copia	1731	1	TAAAT	5	5	4-6[5]
LTR	copia	copia	21	AAAAT	5	5	4-6[5]
LTR	Gypsy	17.6	3	ATAT	4	4	4-6[5]
LTR	Gypsy	297	20	ATAT	4	4	4[2]
LTR	Gypsy	412	16	AAAC	4	4	4[2]
LTR	Gypsy	blood	19	AACC	4	4	4-6[5]
LTR	Gypsy	Burdock	9	TRYA	4	4	4[2]
LTR	Gypsy	diver	6	AAGGG	5	5	4-6[5]
LTR	Gypsy	flea	9	TXHA	4	4	4-6[5]
LTR	Gypsy	gtwin	1	TACA	4	4	4[2]
LTR	Gypsy	gypsy	1	TACA	4	4	4[2]
LTR	Gypsy	gypsy2	1	ATA	3	3	4[2]
LTR	Gypsy	gypsy5	2	CGCG	4	4	4[2]
LTR	Gypsy	HMS-Beagle	3	TGYA	4	4	4[2]
LTR	Gypsy	HMS-Beagle2	2	✖	4	7	4-6[5]
LTR	Gypsy	invader1	2	ATG	3	4	4-6[5]
LTR	Gypsy	invader3	1	AAAT	4	4	4[2]
LTR	Gypsy	invader4	1	TATA	4	4	4-6[5]

LTR	Gypsy	invader6	1	TATA	4	4	4-6[5]
LTR	Gypsy	McClintock	2	YATAT	5	5	4-6[5]
LTR	Gypsy	mdg1	10	AAGG	4	4	4[2]
LTR	Gypsy	mdg3	9	AGTA	4	4	4[2]
LTR	Gypsy	opus	3	TAA	3	4	4[2]
LTR	Gypsy	Quasimodo	5	ATAT	4	4	4[2]
LTR	Gypsy	rover	1	ATAT	4	4	4-6[5]
LTR	Gypsy	springer	1	GTA	3	3	4-6[5]
LTR	Gypsy	Stalker	4	CAAT	4	4	4-6[5]
LTR	Gypsy	Stalker2	4	AAAT	4	4	4[2]
LTR	Gypsy	Stalker4	2	ATAC	4	4	4-6[5]
LTR	Gypsy	Tabor	2	CCTG	4	4	4[2]
LTR	Gypsy	Tirant	15	ACG	2	4	2[2]
LTR	Gypsy	Transpac	4	ATAT	4	4	4[2]
LTR	Pao	3S18	4	GYRDT	5	5	4-6[5]
LTR	Pao	Max	2	CCTCA	5	5	5[2]
LTR	Pao	roo	84	AACAG	4	7	5[2]
LTR	Pao	rooA	1	ACAAT	5	5	4-6[5]

662 [1]TSD motifs represented in the standard IUB/IUPAC nucleic acid codes. The sign "✗" indicates that no
663 consensus can be reported due to a lack of motifs or too variable motifs.
664 [2](41), [3](42), [4](43), [5](5).
665

666   **FIGURE LEGENDS**
667
Figure 1. T-lex2 pipeline.

Schematic representation of the five different T-lex modules (dark gray boxes) and of

the input files (white boxes) required to run T-lex2. SD: Segmental Duplication

Figure 2. T-lex2 TE-presence detection and TE-absence detection modules.

(A) TE-presence detection module is based on the mapping of the NGS reads on the TE

insertion junctions. (B) TE-absence detection module is based on the mapping of NGS

reads on the putative ancestral genomic sequence prior the TE insertion. Input files

required to run each module (white boxes), the different steps of the pipeline (light grey

boxes), and examples of reads providing evidence for the presence, absence, and no call

reads are depicted.

Figure 3. T-lex2 TE-TSD detection module.

Schematic representation of the procedure to identify TSDs. Input files required to run

this module and the different steps followed by the pipeline are also depicted.

Figure 4. Detection of putatively mis-annotated TE insertions.

(A) Schematic representation of a correctly annotated TE (I), a TE longer than annotated

(II) and a TE shorter than annotated (III). (B) Effect of miss-annotation on the TE calls

when the TE insertion is present and absent. When the TE insertion is absent, T-lex2

will give a correct call if the miss-annotation is short and an erroneous call in the miss-

annotation is long (B.h, B.i, B.k, and B.l). (1) If the miss-annotated region is longer than

the maximum read length, selected reads will not overlap the real TE junction(s). These

reads will be mapped inside the TE sequence. Consequently, the number of selected

reads will depend on the number of TE copies from the same family. (2) If the miss-

annotated region is longer than the maximum read length, selected reads will fully map

on the TE flanking region. The number of selected reads will be a proxy of the local

read depth coverage.

Figure 5. TE frequency estimates using single strains.

(A) Comparison of the frequency estimates of 709 TE insertions using T-lex2 and

pooled PCR approach. (B) Frequency estimates of 11 TE insertions using T-lex2 and

single PCR approach.

Figure 6. TE frequency estimates using pooled samples.

(A) Number of reads supporting the absence and (B) number of reads supporting the

presence for each of the TE frequency classes. (C) T-lex2 frequency estimates using

pooled sample *versus* TE frequency classes experimentally determined using a pooled-

PCR approach. (D) Comparison of T-lex2 frequency estimates using single and pooled

NGS data.

REFERENCES

1. Gilad, Y., Pritchard, J.K. and Thornton, K. (2009) Characterizing natural variation using next-generation sequencing technologies. *Trends in genetics : TIG*, **25**, 463-471.
2. Koboldt, D.C., Steinberg, K.M., Larson, D.E., Wilson, R.K. and Mardis, E.R. (2013) The next-generation sequencing revolution and its impact on genomics. *Cell*, **155**, 27-38.
3. Shapiro, E., Biezuner, T. and Linnarsson, S. (2013) Single-cell sequencing-based technologies will revolutionize whole-organism science. *Nature reviews. Genetics*, **14**, 618-630.
4. Treangen, T.J. and Salzberg, S.L. (2012) Repetitive DNA and next-generation sequencing: computational challenges and solutions. *Nature reviews. Genetics*, **13**, 36-46.
5. Wicker, T., Sabot, F., Hua-Van, A., Bennetzen, J.L., Capy, P., Chalhoub, B., Flavell, A., Leroy, P., Morgante, M., Panaud, O. *et al*. (2007) A unified classification system for eukaryotic transposable elements. *Nature reviews. Genetics*, **8**, 973-982.
6. Kapitonov, V.V. and Jurka, J. (2008) A universal classification of eukaryotic transposable elements implemented in Repbase. *Nature reviews. Genetics*, **9**, 411-412; author reply 414.
7. Biemont, C. (2010) A brief history of the status of transposable elements: from junk DNA to major players in evolution. *Genetics*, **186**, 1085-1093.
8. de Koning, A.P., Gu, W., Castoe, T.A., Batzer, M.A. and Pollock, D.D. (2011) Repetitive elements may comprise over two-thirds of the human genome. *PLoS genetics*, **7**, e1002384.
9. Devos, K.M., Beales, J., Ogihara, Y. and Doust, A.N. (2005) Comparative sequence analysis of the phytochrome C gene and its upstream region in allohexaploid wheat reveals new data on the evolution of its three constituent genomes. *Plant molecular biology*, **58**, 625-641.
10. Ewing, A.D. and Kazazian, H.H., Jr. (2011) Whole-genome resequencing allows detection of many rare LINE-1 insertion alleles in humans. *Genome research*, **21**, 985-990.
11. Iskow, R.C., McCabe, M.T., Mills, R.E., Torene, S., Pittard, W.S., Neuwald, A.F., Van Meir, E.G., Vertino, P.M. and Devine, S.E. (2010) Natural mutagenesis of human genomes by endogenous retrotransposons. *Cell*, **141**, 1253-1261.
12. Cowley, M. and Oakey, R.J. (2013) Transposable elements re-wire and fine-tune the transcriptome. *PLoS genetics*, **9**, e1003234.
13. Casacuberta, E. and Gonzalez, J. (2013) The impact of transposable elements in environmental adaptation. *Molecular ecology*, **22**, 1503-1517.
14. Akagi, K., Li, J. and Symer, D.E. (2013) How do mammalian transposons induce genetic variation? A conceptual framework: the age, structure, allele frequency, and genome context of transposable elements may define their wide-ranging biological impacts. *BioEssays : news and reviews in molecular, cellular and developmental biology*, **35**, 397-407.
15. Chenais, B., Caruso, A., Hiard, S. and Casse, N. (2012) The impact of transposable elements on eukaryotic genomes: from genome size increase to

genetic adaptation to stressful environments. *Gene*, **509**, 7-15.

16.   Hua-Van, A., Le Rouzic, A., Boutin, T.S., Filee, J. and Capy, P. (2011) The
struggle for life of the genome's selfish architects. *Biology direct*, **6**, 19.

17.   Makalowski, W., Pande, A., Gotea, V. and Makalowska, I. (2012) Transposable
elements and their identification. *Methods in molecular biology*, **855**, 337-359.

18.   Hormozdiari, F., Hajirasouliha, I., Dao, P., Hach, F., Yorukoglu, D., Alkan, C.,
Eichler, E.E. and Sahinalp, S.C. (2010) Next-generation VariationHunter:
combinatorial algorithms for transposon insertion discovery. *Bioinformatics*, **26**,
i350-357.

19.   Keane, T.M., Wong, K. and Adams, D.J. (2013) RetroSeq: transposable element
discovery from next-generation sequencing data. *Bioinformatics*, **29**, 389-390.

20.   Lee, E., Iskow, R., Yang, L., Gokcumen, O., Haseley, P., Luquette, L.J., 3rd,
Lohr, J.G., Harris, C.C., Ding, L., Wilson, R.K. *et al.* (2012) Landscape of
somatic retrotransposition in human cancers. *Science*, **337**, 967-971.

21.   Robb, S.M., Lu, L., Valencia, E., Burnette, J.M., 3rd, Okumoto, Y., Wessler,
S.R. and Stajich, J.E. (2013) The Use of RelocaTE and Unassembled Short
Reads to Produce High-Resolution Snapshots of Transposable Element
Generated Diversity in Rice. *G3 (Bethesda)*, **3**, 949-957.

22.   Kofler, R., Betancourt, A.J. and Schlotterer, C. (2012) Sequencing of pooled
DNA samples (Pool-Seq) uncovers complex dynamics of transposable element
insertions in Drosophila melanogaster. *PLoS genetics*, **8**, e1002487.

23.   Fiston-Lavier, A.S., Carrigan, M., Petrov, D.A. and Gonzalez, J. (2011) T-lex: a
program for fast and accurate assessment of transposable element presence using
next-generation sequencing data. *Nucleic acids research*, **39**, e36.

24.   Lerat, E. (2010) Identifying repeats and transposable elements in sequenced
genomes: how to find your way through the dense forest of programs. *Heredity*,
104, 520-533.

25.   Smit, A., Hubley, R & Green, P. . (1996-2010) RepeatMasker Open-3.0.
<Erreur ! Référence de lien hypertexte non valide.. .

26.   Kent, W.J. (2002) BLAT--the BLAST-like alignment tool. *Genome research*,
12, 656-664.

27.   Green, P. (2009) 2009. Phrap, version 1.090518. http://phrap.org. .

28.   David, M., Dzamba, M., Lister, D., Ilie, L. and Brudno, M. (2011) SHRiMP2:
sensitive yet practical SHort Read Mapping. *Bioinformatics*, **27**, 1011-1012.

29.   Rumble, S.M., Lacroute, P., Dalca, A.V., Fiume, M., Sidow, A. and Brudno, M.
(2009) SHRiMP: accurate mapping of short color-space reads. *PLoS Comput*
Biol, **5**, e1000386.

30.   Mackay, T.F., Richards, S., Stone, E.A., Barbadilla, A., Ayroles, J.F., Zhu, D.,
Casillas, S., Han, Y., Magwire, M.M., Cridland, J.M. *et al.* (2012) The
Drosophila melanogaster Genetic Reference Panel. *Nature*, **482**, 173-178.

31.   Petrov, D.A., Fiston-Lavier, A.S., Lipatov, M., Lenkov, K. and Gonzalez, J.
(2011) Population genomics of transposable elements in Drosophila
melanogaster. *Mol Biol Evol*, **28**, 1633-1644.

32.   Gonzalez, J., Lenkov, K., Lipatov, M., Macpherson, J.M. and Petrov, D.A.
(2008) High rate of recent transposable element-induced adaptation in
Drosophila melanogaster. *PLoS Biol*, **6**, e251.

33.   Zhu, Y., Bergland, A.O., Gonzalez, J. and Petrov, D.A. (2012) Empirical
validation of pooled whole genome population re-sequencing in Drosophila
melanogaster. *PloS one*, **7**, e41901.

34.   Gonzalez, J., Karasov, T.L., Messer, P.W. and Petrov, D.A. (2010) Genome-

wide patterns of adaptation to temperate environments associated with
transposable elements in Drosophila. *PLoS genetics*, **6**, e1000905.

35.   Ellinghaus, D., Kurtz, S. and Willhoeft, U. (2008) LTRharvest, an efficient and
flexible software for de novo detection of LTR retrotransposons. *BMC*
Bioinformatics, **9**, 18.

36.   Szak, S.T., Pickeral, O.K., Makalowski, W., Boguski, M.S., Landsman, D. and
Boeke, J.D. (2002) Molecular archeology of L1 insertions in the human genome.
Genome Biol, **3**, research0052.

37.   Rawal, K. and Ramaswamy, R. (2011) Genome-wide analysis of mobile genetic
element insertion sites. *Nucleic acids research*, **39**, 6864-6878.

38.   Dunsmuir, P., Brorein, W.J., Jr., Simon, M.A. and Rubin, G.M. (1980) Insertion
of the Drosophila transposable element copia generates a 5 base pair duplication.
Cell, **21**, 575-579.

39.   Rubin, G.M., Brorein, W.J., Jr., Dunsmuir, P., Flavell, A.J., Levis, R., Strobel,
E., Toole, J.J. and Young, E. (1981) Copia-like transposable elements in the
Drosophila genome. *Cold Spring Harb Symp Quant Biol*, **45 Pt 2**, 619-628.

40.   Chaparro, C. and Sabot, F. (2012) Methods and software in NGS for TE
analysis. *Methods in molecular biology*, **859**, 105-114.

41.   Linheiro, R.S. and Bergman, C.M. (2012) Whole genome resequencing reveals
natural target site preferences of transposable elements in Drosophila
melanogaster. *PloS one*, **7**, e30008.

42.   Kapitonov, V.V. and Jurka, J. (2003) Molecular paleontology of transposable
elements in the Drosophila melanogaster genome. *Proc Natl Acad Sci U S A*,
100, 6569-6574.

43.   Berezikov, E., Bucheton, A. and Busseau, I. (2000) A search for reverse
transcriptase-coding sequences reveals new non-LTR retrotransposons in the
genome of Drosophila melanogaster. *Genome Biol*, **1**, RESEARCH0012.

Figure 1.

Figure 2.

A. TE-presence detection module

B. TE-absence detection module

Figure 3.

Figure 4.

A. TE annotation and misannotation

I. TE correctly annotated

II. TE longer than annotated

misannotated
TE sequence

III. TE shorter than annotated

mis-annotated
flanking sequence

B. Effect of misannotation on TE calls

		TE insertion is PRESENT	**TE insertion is ABSENT**
TE-presence detection	**I. TE correctly annotated**	a) ref. sequences / present	g) ref. sequences / absent
	II. TE longer than annotated	b) present[1]	h) absent or no call
	III. TE shorter than annotated	c) present[2]	i) absent or present
TE-absence detection	**I. TE correctly annotated**	d) ref. sequence / present	j) ref. sequence / absent
	II. TE longer than annotated	e) present	k) gaps — absent or present
	III. TE shorter than annotated	f) present	l) absent or present

(1) If the mis-annotated region is longer than the maximum read length, selected reads will not overlap the real TE junction(s). These reads will be mapped inside the TE sequence. Consequently, the number of selected reads will depend on the number of TE copies from the same family.
(2) If the mis-annotated region is longer than the maximum read length, selected reads will fully map on the TE flanking region. The number of selected reads will be a proxy of the local read depth coverage.

Figure 5.

A.

B.

Figure 6.