The overdue promise of short tandem repeat variation for heritability.

Maximilian O. Press, Keisha D. Carlson, Christine Queitsch

Department of Genome Sciences, University of Washington, Foege Building S-250, Box 355065, 3720 15th Ave NE, Seattle WA 98195-5065.

Corresponding author: Queitsch, C. (queitsch@uw.edu).

Keywords: short tandem repeats, microsatellites, heritability, epistasis, sequencing technologies.

This article is under review (June 12, 2014) at the journal *Trends in Genetics*.

Abstract

Short tandem repeat (STR) variation has been proposed as a major explanatory factor

in the heritability of complex traits in humans and model organisms. However, we still

struggle to incorporate STR variation into genotype-phenotype maps. Here, we review

the promise of STRs in contributing to complex trait heritability, and highlight the

challenges that STRs pose due to their repetitive nature. We argue that STR variants

are more likely than single nucleotide variants to have epistatic interactions, reiterate the

need for targeted assays to accurately genotype STRs, and call for more appropriate

statistical methods in detecting STR-phenotype associations. Lastly, somatic STR

variation within individuals may serve as a read-out of disease susceptibility, and is thus

potentially a valuable covariate for future association studies.

The 'missing heritability' of complex diseases and STR variation.

Complex diseases such as diabetes, various cancers, cardiovascular disease,

and neurological disorders cluster in families, and are thus considered to have a genetic

component [1–3] (Glossary). The identification of these genetic factors has proven

challenging; although genome-wide association (GWA) studies have identified many

genetic variants that are associated with complex diseases, these generally confer less

disease risk than expected from empirical estimates of heritability. This discrepancy,

termed the 'missing heritability', has been attributed to many factors [1–6]. A trivial

explanation is that shared environments among relatives may artificially inflate

estimates of heritability. However, missing heritability may also be due to variants in the

human genome that are currently inaccessible at a population scale [1,2]. One such

class of variation is short tandem repeat (STR) copy number variation. Some have

previously suggested that adding STR variation to existing genetic models would

considerably increase the proportion of heritability explained by genetic factors in

human disease [7,8]. Three percent of the human genome consists of STRs [9] and 6%

of human coding regions are estimated to contain STR variation [10,11]. Recently, the

first catalog of genome-wide population-scale human STR variation has appeared [12],

opening up new possibilities for understanding the contribution of STRs to human

genetic diseases.

STRs consist of short (2-10 bp) DNA sequences (units) that are repeated head-

to-tail multiple times. This structure causes frequent errors in recombination and

replication that add or subtract units, leading to STR mutation rates that are 10-fold to

10^4-fold higher than those of non-repetitive loci [13,14]. Due to technical barriers, STR

variation has until very recently remained inaccessible to genome-wide assessment.

STRs are often conserved (even if their copy number or even sequence

changes), especially in coding sequences [15–18]. In both humans and the yeast

Saccharomyces cerevisiae, promoter regions are known to be dramatically enriched for

STRs [19,20]. In coding regions, STRs tend to occur in genes with roles in

transcriptional regulation, DNA binding, protein-protein binding, and developmental

processes [16,21, 22]. These consistent functional enrichments across vastly diverged

lineages suggest important functional roles for STRs.

Indeed, analysis of STR variation in the Drosophila Genetic Reference Panel

identified dozens of associations between STR variants and quantitative phenotypes in

recombinant inbred fly lines [22]. Moreover, accumulating evidence from exhaustive

genetic studies shows that STR variation has dramatic, often background-dependent

phenotypic effects in model organisms [23–27]. Together, these findings suggest that

STR variation has the potential to dramatically revise the heritability estimates

attributable to genetic factors.

The high STR mutation rate also leads to substantial somatic variation of STR

loci within individuals. In fact, this somatic variation, also called microsatellite instability

(MSI), has been used for decades as a biomarker for different classes of cancer [28].

Recent studies demonstrate that organisms exposed to various environmental stresses

and perturbations show increased genome instability, including MSI [29–32]. MSI may

be useful as a biomarker for cellular stress states that may predispose to disease.

The broad interest in STR variation has led to the development of techniques for

high-throughput genotyping of STRs [33,34] and an explosion of analysis tools for

extracting STR variation from existing sequence data [35–37]. However, the precision of

these methods remains limited, due to a combination of low effective coverage of STRs

and the lack of robust models for distinguishing technical error from somatic variation.

Attempts to use STR variation for GWA in a fashion equivalent to SNV variation may be

underpowered and confounded by the unique characteristics of this class of variants. In

this review, we discuss the latest advances in these fields, and lay out a set of priorities

for the future study of STRs.

STR variation is associated with human genetic diseases

Within coding regions, STR mutations are generally in-frame additions and subtractions of repeat units, resulting in proteins with variable, low-complexity amino acid runs [18]. These mutations can result in phenotypic effects and lead to genetic disorders; several neurological diseases (spinocerebellar ataxias, Huntington's disease, spinobulbar muscular atrophy, dentatorubral-pallidoluysian atrophy, intellectual disability, etc.) are a consequence of dramatically expanded STR alleles [7,38,39]. Many of these disease-associated STR expansions behave as dominant gain-of-function mutations [7]. However, even comparatively modest coding STR variation may confer disease risk or behavioral phenotypes, according to a variety of single-marker association studies [40–43]; for instance, variants in two separate coding STRs in RUNX2 are associated with defects in bone mineralization and higher incidence of fractures [44,45]. Noncoding STR variation in regulatory sequences can affect transcription, RNA stability, and chromatin organization. For instance, certain STR variants alter CFTR expression and thus cystic fibrosis status [13]. We take these studies as evidence that STR variation, even in the absence of large expansions, may contribute significantly to the heritability of human traits and genetic diseases.

The severity of the STR expansion-associated diseases may suggest that natural selection should eliminate STRs in functional regions, but several recent studies across many organisms indicate that variable STRs are globally maintained [16,17,21,46,47]. For example, the pre-expansion polyQ-encoding STR in the human gene SCA2 is under

positive selection, suggesting that this variable STR is actively maintained in spite of the

pathogenic expansions that do occasionally occur and cause spinocerebellar ataxia

[48]. Considering both the evidence of positive selection on STRs and the functional

enrichments of STR-containing genes, several authors have proposed that functional

STRs are maintained because they confer 'evolvability', or the capacity for fast

adaptation [18,19,49–51]. This suggestion is intriguing, in part because many STR

mutations are dominant, and, when beneficial, can quickly sweep to fixation. While we

do not further discuss these evolutionary considerations here, they underscore the

phenotypic potential of STR variation.

STR variation has dramatic background-dependent effects on phenotype

To date, the functional consequences of copy number variation in selected STRs

have been studied in plants, fungi, flies, voles, dogs, and fish [23,25,26,52–54], among

other organisms. In *Saccharomyces cerevisiae*, STR copy number in the *FLO1* gene

accurately predicts the phenotype of cell-cell and cell-substrate adhesion (flocculation);

flocculation provides protection against various stresses [54,55]. STR variation in yeast

promoters has been shown to alter gene expression [19]. In *Drosophila melanogaster*,

Neurospora crassa, and *Arabidopsis thaliana,* natural coding STR variation in circadian

clock genes alters diurnal rhythmicity and developmental timing [23–25,56]. Some have

proposed that the large phenotypic responses to selection observed in the Canidae are

a consequence of elevated STR mutation rates relative to other mammalian clades

[50,57]. We can state unambiguously that naturally variable STRs underlie dramatic

phenotypic variation in model organisms.

Beyond the observable fact that variable STRs affect phenotype, we can make

specific predictions about the components of phenotypic variation that they affect. Both

theoretical expectations and empirical data indicate that STR variants are likely to

participate in epistatic interactions, and probably more so than most SNVs. One

plausible hypothesis is that STRs act as mutational modifiers of other loci, as may be

expected intuitively from their elevated mutation rate (Box 1, Figure I).

This expectation is borne out in the handful of studies reporting exhaustive

genetic analysis of STRs. For instance, in the *Xiphophorus* genus of fish, a genetic

incompatibility has recently been attributed to the interaction between the *xmrk*

oncogene and an STR in the promoter of the tumor suppressor *cdkn2a/b* [27,58]. If the

xmrk gene product is not properly regulated by *cdkn2a/b*, fish develop fatal melanomas,

a two-locus Bateson-Dobzhansky-Muller incompatibility described in classic genetic

experiments (Figure 1A) [59–61]. Expansions in the *cdkn2a/b* promoter STR are

associated with the presence of a functional copy of the *xmrk* oncogene across species,

and are thought to functionally repress the activity of the *xmrk* gene product through

increased dosage of the tumor suppressor [27].

Similarly, we have shown that natural variation in the polyQ-encoding *ELF3* STR

significantly affects all ELF3-dependent phenotypes in the plant *A. thaliana,* with *ELF3*

STR length and phenotype showing a strikingly nonlinear relationship (Figure 1B)[23].

Some naturally occurring *ELF3* STR variants phenocopy *elf3*-loss-function mutants in a

common reference background (Figure 1B), suggesting background-specific modifiers.

Indeed, when we compare the phenotypic effects of each *ELF3* STR variant between

two divergent backgrounds Columbia (Col-0) and Wassilewskija (Ws), we find dramatic

differences. The endogenous STR alleles from these two strains (Col-0 7 units, Ws 16

units) show mutual incompatibility when exchanged between backgrounds. The ELF3

protein is thought to function as an "adaptor protein" or physical bridge in diverse protein

complexes [62,63]. We speculated that background-specific polymorphisms in these

interacting proteins underlie the *ELF3* STR-dependent background effect.

Also in *A. thaliana*, a variable STR in the promoter of the *CONSTANS* gene has

been linked to phenotypic variation in the onset of flowering [26]. *CONSTANS* encodes

a major regulatory protein that promotes flowering. Transgenic experiments

demonstrate that this regulatory STR variation affects *CONSTANS* expression and

thence onset of flowering. However, the effects of this STR variation depend on the

presence of a functional allele of *FRIGIDA*, a negative regulator of flowering that is

highly polymorphic across *A. thaliana* populations.

A dramatic example of incompatibility can be found in an intronic repeat in the

IIL1 gene in *A. thaliana*, which was found to be dramatically expanded in one strain [52].

The expansion delayed flowering under high temperatures, but when crossed into the

reference genetic background, a strongly interacting locus modifies this phenotype.

In the *Drosophila* genus, coding STR variation in the *per* gene co-evolves with

other variants [56,64]. Transgenic flies expressing chimeric *per* genes with a *D.*

melanogaster STR domain fused to a *D. pseudoobscura* flanking region (and vice

versa) have arrhythmic circadian clocks, indicating the modifying effect of flanking variation in generating an STR-based genetic incompatibility. Among STRs subjected to exhaustive genetic study, to our knowledge, only the yeast *FLO1* coding STR has no known modifiers due to variation in genetic background [54].

In addition to these exhaustive genetic studies, there are several other observations that support the role of the genetic background in controlling the phenotypic effects of STRs. For instance, experiments in *Caenorhabditis elegans* and human cells indicate that the phenotypic effects of proteins with expanded polyQ tracts are modulated by genetic background [65], or by variants in interacting proteins [66]. In humans, genetic association studies indicate the existence of genetic modifiers of polyQ expansion disorders for both Huntington's disease [67] and spinocerebellar ataxias [68]. Taken together, these experimental and observational data support our argument that functional STRs are likely to be enriched for variants in epistasis with other loci.

STRs with background-dependent phenotypic effects tend to either encode polyQ tracts or reside in promoter regions. There are good reasons to expect that these STR classes might be enriched in DNA/protein-protein interactions that could underlie epistasis. PolyQ tracts, specifically, often bind DNA surfaces [69], and an analysis of human protein interactome data found that polyQ-containing proteins engage in more physical interactions with other proteins than those without polyQs [70]. Similarly, noncoding STRs in regulatory regions may compensate for mutations in trans-acting factors, as observed for the STRs in the *cdkn2a/b* promoter in *Xiphophorus* [27] and in the *CONSTANS* promoter in *A. thaliana* [26]. We suggest that polymorphisms in protein

interaction partners or in transcriptional regulators are plausible explanations for the

observed background effects. In summary, we expect that STR variation is likely to

contribute a substantial epistatic component to heritability, which has important

implications for their use in explaining phenotypic variation.

Analytical tools and genotyping methods continue to struggle with STR-specific

challenges.

To fulfill the promise of STR variation for explaining heritability, we need

accurate, genome-wide assessment of STR variation in populations of humans and

other organisms. The scientific community has tackled this problem in a flurry of recent

studies describing methods for genotyping STRs genome-wide (Table 1). Specifically, in

the last two years, several analytical tools have been developed to call STR genotypes

from whole-genome-sequencing data [35–37]. These tools attempt to address the two

major challenges for genotyping STRs: poor mappability due to low sequence

complexity and high technical error rate due to amplification stutter.

To accurately map an STR sequence read and retrieve its copy number

genotype, the sequence read must span the STR of interest and include some unique

flanking sequence. This requirement severely limits the length of STRs that can be

accurately genotyped and dramatically decreases effective STR coverage compared to

average whole-genome-sequencing coverage (Figure 2). For this reason, much of the

existing sequencing data, which consists largely of short reads (36 bp, 50 bp, or 76 bp)

with only modest genome coverage (5-20X) is not suitable for accurate, genome-wide

calls of STR genotypes; only a fraction of STRs, mostly short ones, can be assessed

with some confidence (Figure 2).

Moreover, these analytical tools estimate technical error based on STR

genotypes from sequenced homozygous or haploid genomes, ignoring the expected

diversity of somatic alleles within individuals. In the face of presumably high somatic

STR variation and high technical error, the probability-based error models of these tools

should require substantial read coverage to call STR genotypes with confidence.

However, because of the low effective coverage of STR loci (Figure 2), STR genotype

calls are based on as few as one or two STR-spanning reads [35,36] (Table 1). This

limitation calls into question how accurate these calls are even for homozygous

germline alleles. Clearly, calling heterozygous STR genotypes remains particularly

challenging with the modest coverage of most available whole-genome-sequencing

data, such as found in the 1000 Genomes Project (Figure 2) [12]. To illustrate this

challenge, consider a heterozygous ~30 bp-STR locus and whole-genome sequencing

with 101 bp-reads at 5x coverage – this scenario is likely to yield just three STR-

spanning reads (Figure 2). These three reads may represent one, two, or three different

alleles, either true alleles, somatic alleles, or technical error, making an accurate call all

but impossible.

Others have attempted to genotype STRs using whole-genome-sequencing data

from paired-end reads (50bp) of size-selected genomic fragments [37], similar to

strategies used to detect large insertions or deletions [71–74]. This approach is limited

by the resolution of gel electrophoresis in the size selection of DNA fragments.

Consequently, this method cannot determine STR copy number genotypes, but rather

reports whether an STR is variable across samples. The authors argue that this

approach is the most accurate for population-level detection of STR variability [75], but it

is not informative for discerning the relationship between STR copy number genotype

and phenotype.

Although these analysis tools represent important and useful advances, their

limitations highlight that 'dustbin-diving' of whole-genome-sequencing data may not

suffice for accurate population-scale genotyping of STRs genome-wide. Alternative

approaches that enrich for STR-spanning sequencing reads are needed. Indeed, two

such approaches have been recently published. Both use targeted capture of STRs to

enrich for STR-spanning reads combined with high-throughput sequencing compatible

with midsize-reads (101 bp, 500 bp) [33,34]. Targeted STR capture requires the design

of STR-specific probes (or rather probes specific to their unique flanking sequences)

and involves additional sequencing, but these approaches can dramatically increase the

number of informative reads, therefore providing substantial STR coverage for accurate

genotyping calls (Table 1). For example, the SureSelect-RNA-probe capture method

reports 27% informative STR-spanning reads compared to the 0.2 % informative reads

found in whole-genome-sequencing data (Table 1). This increase in informative reads is

a major advantage over whole-genome resequencing because STRs represent only a

small fraction of the genome overall [33,34]. Although targeted capture combined with

high-throughput sequencing appears to be a cost-effective alternative for accurate STR

genotyping compared to whole-genome sequencing, distinguishing heterozygous

alleles, somatic variants, and technical error remains a challenge. We suggest that

recent innovations in single-molecule targeted capture [76] should be useful in

distinguishing these categories and in further increasing enrichment of informative,

STR-spanning reads.

Lack of statistical models for detecting STR-phenotype associations in GWA.

Assuming that we obtain accurate, population-scale genotype data for STRs, we

may not yet have statistical tools appropriate for detecting STR associations with

phenotype [8]. In diploid organisms, a biallelic SNV is typically analyzed by modeling

phenotype as a function of the number of non-reference alleles at that locus (0, 1, or 2)

in each individual. A null hypothesis of no monotonic relationship between phenotype

and the allele count is then formulated and tested [77]. This framework cannot

accommodate more than two alleles, which we would expect for many STRs. Simply

using tag SNVs linked to STRs to perform GWA is unfeasible, because linkage

disequilibrium decays very quickly between SNVs and STRs across human populations

[12].

To address these complications, a previous study attempted GWA between STR

genotypes and human disease phenotypes by comparing relative frequencies of various

alleles in pooled DNA from cases and controls [78]. By pooling samples, this approach

eases the analysis of multiallelic loci, but it loses information by ignoring specific

individuals.

In a more recent study, the authors used logistic regression and the analysis of

variance to detect associations between STR alleles and quantitative phenotypes in an

inbred *Drosophila* mapping population [22]. Given that significant associations were

detected, such approaches may be sufficiently powerful in recombinant inbred lines.

However, their strategy relied on homozygosity, and considered multiallelic STRs in a

pairwise fashion, so these straightforward methods will lose power with outbred

populations and multiallelic STRs.

The central confounder of these studies is that most STRs of appreciable

variability (and thus, interest) are multiallelic, as a simple consequence of the STR

mutational mechanism [14]. We could accommodate this multiallelic feature by treating

STR alleles categorically, but this choice entails a corresponding reduction in power,

because many alleles are rare.

Some studies have reported linear associations between STR copy number and

quantitative phenotypes [25,54], suggesting that using simple tests of linear correlation

between these variables may be a powerful option. However, this linearity (or even

monotonicity) of the relationship between STR copy number genotype and phenotype is

a poorly-supported assumption [23]. Nonetheless, STR copy number is a numerical

variable, and it would be preferable to gain power from treating it as such. For instance,

more similar STR copy number genotypes might be associated with more similar

phenotypes, but this intuition may be difficult to generalize.

Lastly, both intuition (Box 1) and the studies discussed above lead us to expect

that relatively many phenotypically relevant variable STRs will show epistasis with other

loci. This epistasis will reduce power in tests of association between STRs and

phenotype [79], given the inadequacy of the current paradigm of quantitative genetics in

detecting and modeling the effects of epistasis [79,80]. At present, targeted and

exhaustive genetic studies (as described above) are the only effective method for

understanding the effects of epistasis.

In total, these obstacles present a daunting challenge for the integration of STR

genotypes into the current genotype-phenotype maps. Overall, we call for a reappraisal

of statistical methodologies for use in GWA with STR variation, to account for these

various STR-specific confounders.

Somatic STR variation may be a sensitive marker for increased disease

susceptibility.

It has been appreciated for some time that the high STR mutation rate leads to

somatic variation within individuals in addition to germ-line variation between individuals

[71]. This somatic STR variation is particularly noticeable in tumor tissues, but is also

measurable in primary tissues [81,82]. While these findings immediately led to systems

of classification for tumor types and clones [83–85], the investigation of somatic STR

variation (or MSI) may also inform us about general phenotypic states and disease

susceptibility.

Patients with various complex diseases tend to carry a greater load of rare germ-

line variants than unaffected control groups [6]. It is widely assumed that these rare

variants contribute in some fashion to these disorders [86]; however, an alternative

interpretation holds that they are signs of stochastic genome instability, which when

increased leads to higher susceptibility to complex diseases. [6]. Increased genome

instability will increase somatic variation, which may then serve as a read-out of disease

susceptibility [6].This alternative interpretation has some support from empirical data.

For instance, perturbation of the molecular chaperone Hsp90, which stabilizes diverse

DNA repair proteins, leads to increased somatic STR mutation rates in human cells; in

various model organisms Hsp90 perturbation increases transposon mobility and

intrachromosomal homologous recombination [29–32]. Hsp90 perturbation also

increases the penetrance of many genetic variants in flies, plants, fish, worms and

yeast, suggesting that increased genome instability and increased phenotypic

heritability are associated [32]. If this association also applies to disease phenotypes,

increased genome instability may predict higher disease susceptibility.

Consequently, while somatic MSI may not be the cause of disease phenotypes, it

may serve as an biomarker for individuals who are more vulnerable to environmental

and genetic perturbations leading to disease. Again, this strategy hinges on the

development of cost-effective technologies for screening panels of STRs for somatic

mutations across many humans.

Another possibility is that somatic variation is itself phenotypically relevant, or

even plays a role in developmental processes. It is known that STRs are enriched in

genes with neuronal function [87]; some have even proposed that such somatic

mutation is a component of normal neuronal development in humans [88]. If this is the

case, then a greater appreciation of somatic variation will be necessary to understand

canonical developmental processes. Collectively, STR variation within (in addition to

between) individuals has great potential as a read-out for disease susceptibility, and

perhaps also as a cause of phenotypic variation itself.

Conclusions

The study of STRs and other underascertained genomic elements has the potential to

reshape our model of the heritability of complex diseases and traits, both in terms of the

overall proportion of heritability explained, and in terms of the components of heritability

themselves (Outstanding Questions). Experimental studies in model organisms have

taught us that the phenotypic effects of genome-wide STR variation are both dramatic

and impossible to understand without taking epistasis into account. In the future, our

understanding will be improved by 1) accurate STR population-scale and somatic

genotyping, 2) more appropriate statistical methods for analyzing STR-phenotype

associations, and 3) a broader description of epistasis between STR variation and other

loci in determining phenotype.

Aknowledgements

We thank members of the Queitsch lab for helpful discussions. We thank Wen Huang

and David Mittelman for responding to email inquiries. This work was supported by

grants from the National Human Genome Research Institute Interdisciplinary Training in

Genomic Sciences (2T32HG35-16 to MOP, T32 HG00035 to KDC) and the National

Institute of Health New Innovator Award (DP2OD008371 to CQ).

References

Manolio, T.A. *et al.* (2009) Finding the missing heritability of complex diseases. *Nature* 461, 747–53

Eichler, E.E. *et al.* (2010) Missing heritability and strategies for finding the underlying causes of complex disease. *Nat. Rev. Genet.* 11, 446–50

Gibson, G. (2011) Rare and common variants: twenty arguments. *Nat. Rev. Genet.* 13, 135–45

Zuk, O. *et al.* (2012) The mystery of missing heritability: Genetic interactions create phantom heritability. *Proc. Natl. Acad. Sci. U. S. A.* 109, 1193–8

Bloom, J.S. *et al.* (2013) Finding the sources of missing heritability in a yeast cross. *Nature* 494, 234–7

Heng, H.H.Q. (2010) Missing heritability and stochastic genome alterations. *Nat. Rev. Genet.* 11, 813

Fondon, J.W. *et al.* (2008) Simple sequence repeats: genetic modulators of brain function and behavior. *Trends Neurosci.* 31, 328–34

Hannan, A.J. (2010) Tandem repeat polymorphisms: modulators of disease susceptibility and candidates for "missing heritability". *Trends Genet.* 26, 59–65

Subramanian, S. *et al.* (2003) Genome-wide analysis of microsatellite repeats in humans: their abundance and density in specific genomic regions. *Genome Biol.* 4, R13

O'Dushlaine, C.T. *et al.* (2005) Tandem repeat copy-number variation in protein-coding regions of human genes. *Genome Biol.* 6, R69

Mularoni, L. *et al.* (2006) Mutation patterns of amino acid tandem repeats in the human proteome. *Genome Biol.* 7, R33

Willems, T.F. *et al.* (2014) The Landscape of Human STR Variation. *bioRxiv* DOI: 10.1101/004671

Eckert, K.A. and Hile, S.E. (2009) Every microsatellite is different: Intrinsic DNA features dictate mutagenesis of common microsatellites present in the human genome. *Mol. Carcinog.* 48, 379–88

Legendre, M. *et al.* (2007) Sequence-based estimation of minisatellite and microsatellite repeat variability. *Genome Res.* 17, 1787–96

Zhao, Z. *et al.* (2014) Genome-wide analysis of tandem repeats in plants and green algae. *G3* 4, 67–78

Schaper, E. *et al.* (2014) Deep Conservation of Human Protein Tandem Repeats within the Eukaryotes. *Mol. Biol. Evol.* 31, 1132–1148

Li, H. *et al.* (2012) Insight into role of selection in the evolution of polyglutamine tracts in humans. *PLoS One* 7, e41167

Gemayel, R. *et al.* (2010) Variable tandem repeats accelerate evolution of coding and regulatory sequences. *Annu. Rev. Genet.* 44, 445–77

Vinces, M.D. *et al.* (2009) Unstable tandem repeats in promoters confer transcriptional evolvability. *Science (80-.).* 324, 1213–6

Sawaya, S. *et al.* (2013) Microsatellite tandem repeats are abundant in human promoters and are associated with regulatory elements. *PLoS One* 8, e54710

Persi, E. and Horn, D. (2013) Systematic analysis of compositional order of proteins reveals new characteristics of biological functions and a universal correlate of macroevolution. *PLoS Comput. Biol.* 9, e1003346

Mackay, T.F.C. *et al.* (2012) The Drosophila melanogaster Genetic Reference Panel. *Nature* 482, 173–8

Undurraga, S.F. *et al.* (2012) Background-dependent effects of polyglutamine variation in the Arabidopsis thaliana gene ELF3. *Proc. Natl. Acad. Sci.* DOI: 10.1073/pnas.1211021109

Michael, T.P. *et al.* (2007) Simple sequence repeats provide a substrate for phenotypic variation in the Neurospora crassa circadian clock. *PLoS One* 2, e795

Sawyer, L.A. (1997) Natural Variation in a Drosophila Clock Gene and Temperature Compensation. *Science (80-.).* 278, 2117–2120

Rosas, U. *et al.* (2014) Variation in Arabidopsis flowering time associated with cis-regulatory variation in CONSTANS. *Nat. Commun.* 5, 3651

Scarpino, S. V *et al.* (2013) Evolution of a genetic incompatibility in the genus Xiphophorus. *Mol. Biol. Evol.* 30, 2302–10

Boland, C.R. *et al.* (1998) A National Cancer Institute Workshop on Microsatellite Instability for Cancer Detection and Familial Predisposition: Development of International Criteria for the Determination of Microsatellite Instability in Colorectal Cancer. *Cancer Res.* 58, 5248–5257

Kovalchuk, I. *et al.* (2003) Pathogen-induced systemic plant signal triggers DNA rearrangements. *Nature* 423, 760–2

Specchia, V. *et al.* (2010) Hsp90 prevents phenotypic variation by suppressing the mutagenic activity of transposons. *Nature* 463, 662–5

Mittelman, D. *et al.* (2010) Hsp90 modulates CAG repeat instability in human cells. *Cell Stress Chaperones* 15, 753–9

Queitsch, C. *et al.* (2012) Lessons from model organisms: phenotypic robustness and missing heritability in complex disease. *PLoS Genet.* 8, e1003041

Guilmatre, A. *et al.* (2013) Rapid multiplexed genotyping of simple tandem repeats using capture and high-throughput sequencing. *Hum. Mutat.* 34, 1304–11

Duitama, J. *et al.* (2014) Large-scale analysis of tandem repeat variability in the human genome. *Nucleic Acids Res.* DOI: 10.1093/nar/gku212

Highnam, G. *et al.* (2013) Accurate human microsatellite genotypes from high-throughput resequencing data using informed error profiles. *Nucleic Acids Res.* 41, e32

Gymrek, M. *et al.* (2012) lobSTR: A short tandem repeat profiler for personal genomes. *Genome Res.* 22, 1154–62

Cao, M.D. *et al.* (2014) Inferring short tandem repeat variation from paired-end short reads. *Nucleic Acids Res.* 42, e16

Gatchel, J.R. and Zoghbi, H.Y. (2005) Diseases of unstable repeat expansion: mechanisms and common principles. *Nat. Rev. Genet.* 6, 743–55

Poeta, L. *et al.* (2013) A regulatory path associated with X-linked intellectual disability and epilepsy links KDM5C to the polyalanine expansions in ARX. *Am. J. Hum. Genet.* 92, 114–25

Caspi, A. *et al.* (2003) Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science* 301, 386–9

Zhang, Y. *et al.* (2012) IL1 receptor antagonist gene IL1-RN variable number of tandem repeats polymorphism and cancer risk: a literature review and meta-analysis. *PLoS One* 7, e46017

Eisenegger, C. *et al.* (2013) DAT1 polymorphism determines L-DOPA effects on learning about others' prosociality. *PLoS One* 8, e67820

Inanir, A. *et al.* (2013) Association of IL-4 gene VNTR variant with deep venous thrombosis in Behçet's disease and its effect on ocular involvement. *Mol. Vis.* 19, 675–683

Morrison, N.A. *et al.* (2012) Glutamine repeat variants in human RUNX2 associated with decreased femoral neck BMD, broadband ultrasound attenuation and target gene transactivation. *PLoS One* 7, e42617

Morrison, N.A. *et al.* (2013) Polyalanine repeat polymorphism in RUNX2 is associated with site-specific fracture in post-menopausal females. *PLoS One* 8, e72740

Mularoni, L. *et al.* (2007) Highly constrained proteins contain an unexpectedly large number of amino acid tandem repeats. *Genomics* 89, 316–25

Mularoni, L. *et al.* (2010) Natural selection drives the accumulation of amino acid tandem repeats in human proteins. *Genome Res.* 20, 745–54

Yu, F. *et al.* (2005) Positive selection of a pre-expansion CAG repeat of the human SCA2 gene. *PLoS Genet.* 1, e41

Gemayel, R. *et al.* (2012) Beyond junk-variable tandem repeats as facilitators of rapid evolution of regulatory and coding sequences. *Genes (Basel).* 3, 461–80

Laidlaw, J. *et al.* (2007) Elevated Basal Slippage Mutation Rates among the Canidae. *J. Hered.* 98, 452–460

King, D.G. (2012) Indirect selection of implicit mutation protocols. *Ann. N. Y. Acad. Sci.* 1267, 45–52

Sureshkumar, S. *et al.* (2009) A genetic defect caused by a triplet repeat expansion in Arabidopsis thaliana. *Science* 323, 1060–3

Hammock, E.A.D. and Young, L.J. (2005) Microsatellite instability generates diversity in brain and sociobehavioral traits. *Science* 308, 1630–4

Smukalla, S. *et al.* (2008) FLO1 is a variable green beard gene that drives biofilm-like cooperation in budding yeast. *Cell* 135, 726–37

Verstrepen, K.J. *et al.* (2005) Intragenic tandem repeats generate functional variability. *Nat. Genet.* 37, 986–90

Peixoto, a a *et al.* (1998) Molecular coevolution within a Drosophila clock gene. *Proc. Natl. Acad. Sci. U. S. A.* 95, 4475–80

Fondon, J.W. and Garner, H.R. (2004) Molecular origins of rapid and continuous morphological evolution. *Proc. Natl. Acad. Sci. U. S. A.* 101, 18058–63

Butler, A.P. *et al.* (2007) Regulation of CDKN2A/B and Retinoblastoma genes in Xiphophorus melanoma. *Comp. Biochem. Physiol. C. Toxicol. Pharmacol.* 145, 145–55

Gordon, M. (1927) THE GENETICS OF A VIVIPAROUS TOP-MINNOW PLATYPOECILUS; THE INHERITANCE OF TWO KINDS OF MELANOPHORES. *Genetics* 12, 253–283

Kosswig, C. (1928) Über Kreuzungen zwischen den Teleostiern Xiphophorus helleri und Platypoecilus maculatus. *Z. Indukt. Abstammungs-Vererbungsl.* 47, 150–158

Meierjohann, S. and Schartl, M. (2006) From Mendelian to molecular genetics: the Xiphophorus melanoma model. *Trends Genet.* 22, 654–61

Yu, J.-W. *et al.* (2008) COP1 and ELF3 control circadian function and photoperiodic flowering by regulating GI stability. *Mol. Cell* 32, 617–30

Nusinow, D. a. *et al.* (2011) The ELF4–ELF3–LUX complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* DOI: 10.1038/nature10182

Peixoto, A. *et al.* (1993) Molecular evolution of a repetitive region within the per gene of Drosophila. *Mol. Biol. Evol.* 10, 127–139

Gidalevitz, T. *et al.* (2013) Natural genetic variation determines susceptibility to aggregation or toxicity in a C. elegans model for polyglutamine disease. *BMC Biol.* 11, 100

Metzger, S. *et al.* (2008) Huntingtin-associated protein-1 is a modifier of the age-at-onset of Huntington's disease. *Hum. Mol. Genet.* 17, 1137–46

Rubinsztein, D.C. *et al.* (1997) Genotypes at the GluR6 kainate receptor locus are associated with variation in the age of onset of Huntington disease. *Proc. Natl. Acad. Sci.* 94, 3872–3876

Zühlke, C. *et al.* (2005) Spinocerebellar ataxia type 17: report of a family with reduced penetrance of an unstable Gln49 TBP allele, haplotype analysis supporting a founder effect for unstable alleles and comparative analysis of SCA17 genotypes. *BMC Med. Genet.* 6, 27

Escher, D. *et al.* (2000) Conservation of Glutamine-Rich Transactivation Function between Yeast and Humans. *Mol. Cell. Biol.* 20, 2774–2782

Schaefer, M.H. *et al.* (2012) Evolution and function of CAG/polyglutamine repeats in protein-protein interaction networks. *Nucleic Acids Res.* 40, 4273–87

Chen, K. *et al.* (2009) BreakDancer: an algorithm for high-resolution mapping of genomic structural variation. *Nat. Methods* 6, 677–81

Hajirasouliha, I. *et al.* (2010) Detection and characterization of novel sequence insertions using paired-end next-generation sequencing. *Bioinformatics* 26, 1277–83

Qi, J. and Zhao, F. (2011) inGAP-sv: a novel scheme to identify and visualize structural variation from paired end mapping data. *Nucleic Acids Res.* 39, W567–75

Grimm, D. *et al.* (2013) Accurate indel prediction using paired-end short reads. *BMC Genomics* 14, 132

Cao, M.D. *et al.* (2014) Sequencing technologies and tools for short tandem repeat variation detection. *Brief. Bioinform.* DOI: 10.1093/bib/bbu001

Hiatt, J.B. *et al.* (2013) Single molecule molecular inversion probes for targeted, high-accuracy detection of low-frequency variation. *Genome Res.* 23, 843–54

Hayes, B. (2013) Overview of Statistical Methods for Genome-Wide Association Studies (GWAS). *Methods Mol. Biol.* 1019, 149–69

Oka, A. *et al.* (2003) Localization of a non-melanoma skin cancer susceptibility region within the major histocompatibility complex by association analysis using microsatellite markers. *Tissue Antigens* 61, 203–210

Mackay, T.F.C. (2014) Epistasis and quantitative traits: using model organisms to study gene-gene interactions. *Nat. Rev. Genet.* 15, 22–33

Nelson, R.M. *et al.* (2013) A century after Fisher: time for a new paradigm in quantitative genetics. *Trends Genet.* 29, 669–76

Armour, J.A.L. *et al.* (1989) Analysis of somatic mutations at human minisatellite loci in tumors and cell lines. *Genomics* 4, 328–334

Slagboom, P.E. *et al.* (1991) Somatic mutations and cellular aging: two-dimensional DNA typing of rat fibroblast clones. *Mutat. Res.* 256, 311–321

Lee, H.S. *et al.* (2001) Microsatellite instability in synchronous gastric carcinomas. *Int. J. Cancer* 91, 619–24

Kim, T.-M. *et al.* (2013) The landscape of microsatellite instability in colorectal and endometrial cancer genomes. *Cell* 155, 858–68

Chapal-Ilani, N. *et al.* (2013) Comparing algorithms that reconstruct cell lineage trees utilizing information on microsatellite mutations. *PLoS Comput. Biol.* 9, e1003297

Girirajan, S. *et al.* (2011) Human copy number variation and complex genetic disease. *Annu. Rev. Genet.* 45, 203–26

Bolton, K.A. *et al.* (2013) STaRRRT: a table of short tandem repeats in regulatory regions of the human genome. *BMC Genomics* 14, 795

Nithianantharajah, J. and Hannan, A.J. (2007) Dynamic mutations as digital genetic modulators of brain development, function and dysfunction. *Bioessays* 29, 525–35

Abecasis, G.R. *et al.* (2010) A map of human genome variation from population-scale sequencing. *Nature* 467, 1061–73

Cao, J. *et al.* (2011) Whole-genome sequencing of multiple Arabidopsis thaliana populations. *Nat. Genet.* 43, 956–63

Gan, X. *et al.* (2011) Multiple reference genomes and transcriptomes for Arabidopsis thaliana. *Nature* 477, 419–23

Figure I. A locus with higher mutation rates allows genetic modification of unfavorable genotypes at interacting loci. Top, a model of evolution under epistasis with only one slow mutation rate. Middle, a model of evolution under epistasis with a slow and a fast mutation rate. Boxes represent loci, stars represent SNV-type mutations, black and white checkering indicates an STR locus (*a/b*, *a*/b*, and *a*/b** signify different genotypes). Arrows with numbers represent possible mutations and their respective rates. Bottom, fitness of each genotype under both models. We expect that the model with two mutation rates will occupy the fully derived state (*a*/b**) more quickly.

Figure 1. Genetic and transgenic analysis reveals STR-mediated incompatibilities. A, the Gordon-Kosswig-Anders cross shows a genetic incompatibility

between two fish species in the *Xiphophorus* genus. Modified from Meierjohann and

Schartl [61]. F_1 hybrids back-crossed to their *X. helleri* parent yield a 3:1 ratio of viability,

where the inviables result from co-segregation of the functional *xmrk* gene and a short

STR allele in the *cdkn2a/b* promoter. Shading indicates melanism conferred by *xmrk*. **B,**

genetic background is epistatic to effects of *ELF3* STR variation in *A. thaliana.*

Expression-matched transgenic plants with various alleles of the *ELF3* STR in the

Columbia (Col-0) and Wassilewskija (Ws) backgrounds, showing endogenous,

exogenous, and synthetic ("0") alleles in each background [23]. White boxes indicate

transgenic plants carrying the *ELF3* STR endogenous to their respective background;

white arrowheads indicate early-flowering ELF3 STR genotypes (*elf3* mutants and

poorly-functioning *ELF3* STR alleles confer early flowering).

Figure 2. Effective reduction in STR coverage in whole-genome sequencing.

Expected coverage of STRs for various sequencing depths and read lengths. We

assumed 8 bp of flanking sequence on either side (per requirement for LobSTR

software [36]). Black bars indicate nominal sequencing coverage for each scenario. 4-

5X coverage (left panel) is typical for genomes in the human 1000 Genomes Project

[89]; 15-20X coverage is typical for genomes in the *A. thaliana* 1001 Genomes Project

[90,91].

2 **Figure 1.**

4 **Figure I.**

2   **Figure 2.**

4   **Table 1. Technologies for assessing STR variation by high-throughput**
5   **sequencing.**

Name	Data Source	Analysis strategy	Accepted coverage[a]	Reported accuracy	Reported efficiency	Limitations	Ref.
lobSTR	Human, whole-genome[b,c]	Align to modified reference	1 read	88%-95%	0.2% of reads are informative	Depends on depth of sequencing and length of reads	[36]
RepeatSeq	Human, whole-genome[b,d]	Align to reference, locally realigned	2 reads	92%	Not reported	Depends on depth of sequencing and length of reads	[35]
STRViper	*A. thaliana* whole-genome[b,d]	Compare insert size to reference	10 reads	74%	Not reported	Cannot call STR copy number genotypes	[37]
Array Capture	Human, array capture[b]	RepeatSeq	2 reads	88%-92%	2.2% informative reads	Low enrichment for STR-spanning reads	[33]
SureSelect RNA probe capture	Human, target enrichment, Roche 454	Locally align flanking regions	4 reads	88%-95%	27% informative reads	Expensive probe design, captured only 60% of targeted STRs	[34]

6   a: Minimum coverage of a single STR that is considered sufficient to call a genotype.

b: Sequence data from Illumina HiSeq technology.
c: data references: [82,83]
d: data references: [73,84]
e: data references: [74,75]

BOX 1: Modifier mutations leading to epistasis are expected in STRs. We have previously proposed that STRs might be more susceptible to genetic interactions [23], as we will briefly explicate here. Consider a simple two-locus haploid model under panmixis, in which loci *A* and *B* each start with a single allele (*ab*) and have the same probability *p* per generation of mutating to a second allele (*a** or *b**), with *p* also as the probability per generation of reverting mutations (Figure I). Let us further assume that *A* and *B* are in sign epistasis [58] (that is, *a*b* and/or *ab** have fitness less than *ab* and *a*b**). To escape the unfavorable *a*b* genotype, the organism may either revert to *ab* or mutate forward to *a*b**. When the *A* and *B* loci have equal mutation rates, we expect that the reversion of a single mutant is just as likely as a second mutation, and consequently that *a*b** individuals will appear only relatively rarely and slowly. However, consider a similar model, in which locus B has an elevated mutation rate $p_b > p_a$. In this case, the *a*b* genotype has a higher probability of a second, modifying mutation to *a*b** than of a reversion to *ab*. Moreover, flux along the other mutational path (*ab* → *ab** → *a*b**) will be increased. In sum, *a*b** genotypes will arise at higher rates, and will attain their equilibrium frequency much more rapidly, if either A or B has an elevated mutation rate [59] (p.131). This scenario can lead quickly to an equilibrium population in which incompatible epistatic dalleles are frequent, even though recombinants have lower fitness. Relaxing the assumption of no population structure will further speed this process. Consequently, we would expect STRs and other loci with high mutation rates to be more likely to modify other alleles than loci with lower mutation rates, as long as we assume that all loci are equally capable of genetic interactions. This process may be referred to as 'coadaptation'. For a rigorous model of the evolution of hybrid incompatibility, see Orr [57].