A framework of immunities for pathogens and hypersensitivities

Wan-Chung Hu*

MD National Taiwan University

PhD Johns Hopkins University

Department of Clinical Pathology, Far Eastern Memorial Hospital, New Taipei City, Taiwan

***Correspondence:** Wanchung.Hu09@gmail.com (W. Hu)

Keywords: Tr1, Th1. Th2, Th17, Th19, Th22, Treg, Tfh

BioRxiv preprint doi: https://doi.org/10.1101/006965

Abstract

THfh is to initiate four eradicable immunities. Treg is for four tolerable immunities. TH1/TH1like is immunity for intracellular bacteria/protozoa and type4 hypersensitivity. TH1 includes macrophage1, Tc1, IFNg T helper, and IgG3. TH1like includes macrophage2, suppressive CTL, IFNg/TGFβ T helper, and IgA1. TH2/TH9 is immunity for helminths and type1 hypersensitivity. TH2 includes iEOS, basophil/MCct, IL-4 T helper, and IgE/IgG4. TH9 includes rEOS, basophils/MCt, IL-9 T helper, and IgA2. TH22/TH17 is immunity for extracellular bacteria/fungi and type3 hypersensitivity. TH22 includes neutrophils1, IL-22 T helper, and IgG2. TH17 include neutrophils2, IL-17 T helper, and IgA2. THαβ/TH3 is immunity for viruses and type2 hypersensitivity. THαβ includes NK1, Tc2, IL10 T helper, and IgG1. TH3 includes NK2, suppressive CTL, IL-10/TGFβ T helper, and IgA1.

Background of host immunities

There are many discovered host immunological pathways including traditional TH1, TH2, TH3, TH17, TH22, THfh, Treg, TH9, and Tr1(THαβ). These identified pathways are not logically organized. Here, I will propose a detailed picture about the whole context of host immunological pathways. (Figure 1)

The traditional TH1/TH2 paradigm was proposed by Dr. Mosmann in 1986.(1) TH1 was thought the host immunity against viruses and intracellular bacteria. TH2 is the host immunity against multicellular parasites (helminthes). In my PhD thesis, I proposed a new THαβ immunological pathway against viruses that is divided from traditional TH1 immunity.(2) The TH1 immunity is then focusing on intracellular bacteria and protozoa. Then, TH3 immunity and Tr1 immunological pathways were identified later.(3, 4) Recently, additional immune responses are discovered including TH17, TH22, THfh, Treg, and TH1-like immunological pathways.(5-9)

Initiatory immune response

Follicular helper T cells (THfh) is thought to be the key helper cells for the B cell germinal centers in lymph nodes.(10) THfh cells are characterized by IL-21 producing T cells(11, 12). BCL6 is a key transcription factor for THfh. TGF beta with STAT5 signal can constrain the differentiation of the IL-21 producing helper T cells(13, 14). IL-21 production is also related to STAT1and STAT3 activation as well as STAT5 activation. Since immunosuppressive prolactin can induce STAT5a to suppress BCL6 expression.(15, 16)On the contrary, STAT5b can up-regulate BCL6.(17)STAT5a and STAT5b have distinct target genes in immune responses.(18)The transcription factor

to induce THfh should be STAT5b. BCL6 is key in THfh development.(19-

21)Follicular helper T cell can induce B cells to start to produce IgM antibody.(22)

Thus, it is the earliest T lymphocytes to begin the adaptive host immunity.(23-25)

Different STAT proteins regulate different immunological pathways.(26, 27) If the

infection is tend to be eradicable, the following host immunological pathways are

generated with other cytokines.

Eradicable immune responses

TH1 immunity is driven by IL-12. It is the host immunity against intracellular bacteria

or protozoa. The main effector cells of TH1 immunity are stimulatory macrophages

(M1), IFNg secreting cytotoxic CD8 T cells (CD28+ Tc1), IFNg secreting CD4 T

cells, and IgG3 producing B cells.(28-31)The key transcription factors for TH1

immunity is STAT4. T-bet also plays a vital role in TH1 immunological pathway. TH1

immunity against self antigen is Type 4 Delayed-type hypersensitivity such as type1

diabetes mellitus.(32, 33)

TH2 immunity is driven by IL-4. TH2 immunity is against extracellular parasites

(helminthes). The main effector cells of TH2 immunity are eosinophils (iEOS),

basophils/connective tissue mast cells(MCtc, mast cell tryptase & chymase), IL-4/IL-

5 secreting CD4 T cells, and IgG4/IgE producing B cells.(34)IgG4 activates

eosinophils and IgE activates mast cells such as in acute anaphylaxis,

respectively.(35) The function of IgG4-eosinophil is to activate eosinophil mediated

cellular immunity against parasites or insects. The function of IgE-mast cell is to

expel helminths or insects by physiological mechanism. Activated mast cells by IgE

can release histamine which causes bronchoconstriction, vomiting/nausea, rhinorrhea,

skin itchiness, stomach acidification, increased local vascular permeability, or

increased bowel movement. These actions can all help to expel helminths or insects

physiologically. The key transcription factor for TH2 immunity is STAT6. GATA3

also plays a vital role in TH2 immunological pathway. TH2 immunity against self

antigen is Type1 immediate allergy such as food/drug allergy or urticaria.(36)

THαβ is distinguished from the traditional TH1 immunity(2). THαβ immunity is

against viruses. It was called Tr1 cell by some previous researchers.(4)THαβimmunity

is driven by IFNa/b or IL-10. The main effector cells of THαβ immunity are IL-10

producing stimulatory NK cells(CD56-CD16+ NK1 cells), IL-10/IL-27 secreting CD4

T cells, IL-10 secreting cytotoxic CD8 T cells (CD28+ Tc2), and IgG1 producing B

cells.(28, 30, 33, 37, 38)CD27 molecule is important for virus immunity.(39, 40)The

key transcription factor for THαβ immunity is STAT1and STAT2.(41)THαβ immunity

against self antigen is Type 2 Antibody dependent cytotoxic hypersensitivity such as

acute stage of Myasthenia Gravis. It is worth noting that IL-10 is not merely a

immunosuppressive cytokine; it can have potent stimulatory effects on NK cells,

CTLs, and B cells.(42)

TH22 is the host innate immunity against extracellular bacteria and fungi(43, 44).

TH22 is driven by IL-6 or TNFa(45, 46). The main effector cells for TH22 immunity

are PMNs(N1), IL-22 secreting CD4 T cells, complements, pentraxins, and IgG2

producing B cells.(6, 47)The key transcription factor for TH22 is STAT3(48). AP1

and CEBP are also important. TGF beta can suppress IL-22 to skew to TH17

immunity.(49)TH22 against self antigen is Type 3 immune-complex and complement

mediated hypersensitivity such as Arthus reaction.(50) It is worth noting that the

extracellular or intracellular locations of protozoa or fungi mainly decide the host

immunological pathways.

It is interesting to know that four IgG subtypes fit the four types of acute

immunological pathways. Murine IgG antibodies also have four subclasses. There is a

correlation between murine and human IgG subtypes: Human IgG1<->Murine IgG2a;

Human IgG2<->Murine IgG3; Human IgG3<->Murine IgG2b; Human IgG4<-

>Murine IgG1.(51) hIgG1/mIgG2a is against viral antigens; hIgG2/mIgG3 is against

bacterial antigen, especially polysaccharides; hIgG3/mIgG2b is against intracellular

bacteria; and hIgG4/mIgG1 is related to parasite antigens.(52-55)

Regulatory immune responses

Treg is the host immune inhibitory mechanism(9). It is driven by IL-2 and TGF beta.

The main effector cells for Treg are TGFb producing CD4 T cell and IgA producing B

cell. The key transcription factor for Treg pathway is STAT5, especially STAT5a. But,

both STAT5a and STAT5b play non-redundant roles in Treg generation.(56) They may

act sequentially with STAT5b activation first in THfh signaling. Combined STAT5b

and STAT5a signaling induces the generation of Treg. The combination of Treg and

the above four immunological pathways is important to shift adaptive immunity to

tolerable immunity. During the initial infection, acute stage fierce cytokines can

rapidly kill pathogens as well as infected cells or tissues. However, if the pathogen

infects a lot of cells in a tissue such as liver, to kill the infected cells will total

destroyed the organ.(57) Thus, regulatory T cells STAT5 signal combining

TH1/TH2/TH22/THαβ will make CD4 T cells with less fierce cytokines.(56) Then,

TH1like/TH9/TH17/TH3 immunological pathways will be generated in more chronic

stage. It is worth noting that there are two subtypes of IgA antibodies: IgA1 and IgA2.

IgA1 is the dominant IgA antibody in serum, and IgA2 is the dominant IgA in

mucosa. TGF beta can induce either IgA1 or IgA2 which seems to be dependent on

lymphoid follicle location.(58) In GULTs or Peyer's Patch, IgA2 is the dominant IgA

antibody produced in GI mucosa there. In lymph nodes of other body locations, IgA1

is the dominant IgA antibody produced there.(59) However, IgA1 is especially related

to viral protein antigens and IgA2 is especially related to bacterial antigens such as

LPS.(60) The heavy chain locus sequence of B cell antibodies is IgM, IgD, IgG3,

IgG1, IgA1, IgG2, IgG4, IgE, and IgA2. First, B cells double express IgM and IgD.

We can view IgG3, IgG1, and IgA1 as the first group for cellular immunity. Then,

IgG2, IgG4, IgE, and IgA2 can be viewed as the second group for humoral immunity.

The gene sequence order is important, and it affects the time sequence of isotype

switch. It is also worth noting that IL-13 is also a Treg related cytokine which is pro-

fibrogenic and related to TGF beta signaling.(61, 62)

Tolerable immune responses

TH1-like cells (non-classic TH1) are initiated by TGF beta(STAT5 signaling) and

IFNg(STAT4 signaling). TH1-like cells with Foxp3+ regulatory character are

identified.(8, 63)There is a close relation to TH1 helper cells and TH1-like cells.(64,

65)TH1-like cells are the chronic host immunity of TH1 immune response. Thus, it

could be related to chronic inflammation such as long-term tuberculosis infection. The

effector cells of TH1-like immunity include suppressive macrophages (M2),

suppressive CD8 T cells (CD8 Treg CD28-), IgA1 producing B cells, and IFNg/TGFb

producing CD4 T cells.(29, 30, 66, 67)TH1-like immunity induces type4 delayed-type

hypersensitivity such as Crohn's disease.(68)

TH9 cell is driven by IL-4 (STAT6 signaling) combining TGF beta(STAT5

signaling).(69-71)Thus, TH9 cell is closely related to TH2 immunological pathway. It

is characterized by IL-9 secreting CD4 T cell. TH9 cells are found to be important in

chronic allergic condition such as asthma. Thus, TH9 helper cell is the chronic T

helper cells related to TH2 immunity. The effector cells of TH9 immunity include

regulatory eosinophils(rEOS), basophils/mucosal mast cells (MCt, mast cell tryptase),

IL-9 producing CD4 T cells, and IgA2 producing B cells.(72) TH9 immunity induces

type1 allergy including asthma.(34, 73-78)

TH17 cell is driven by IL-6 / IL-1 combining TGF beta(5, 79). Thus, TH17 cell is

closely related to TH22 immunological pathway. It is characterized by IL-17 secreting

CD4 T cell. TH17 cells are found to be important in chronic immune-complex

mediated disease such as rheumatic arthritis. Then, TH17 helper cell is the chronic T

helper cell related to TH22 immunity.(80)TGF beta with STAT5 can suppress the

acute IL-22 producing cells and enhance the chronic IL-17 producing cells(49, 81).

Because of the role of TGF beta in TH17 immunity, regulatory IL-17 producing cells

are noted.(82, 83) The effector cells of TH17 immunity include regulatory

neutrophils(N2), IL-17 producing CD4 T cells, and IgA2 producing B cells.(60,

84)TH17 immunity induces type3 immune-complex hypersensitivity including

ulcerative colitis.(85, 86)

TH3 cells are driven by IL-10 and TGF beta.(87, 88)Thus, TH3 cells are closely

related to THαβ immunological pathway. It also produces IL-10 as well as TGF beta.

Thus, TH3 helper cell is important to chronic antibody dependent cellular cytotoxic

hypersensitivity. TH3 cell is the chronic helper T cells corresponding to THαβ helper

cell. The TH3 immune effector cells include IL-13 producing regulatory NK

cells(CD56+CD16- NK2 cells), IL-10 and TGF beta secreting CD4 T cells,

suppressive CD8 T cells (CD8 Treg CD28-), and IgA1 producing B cells.(30, 38, 89,

90) IgA1 is produced in serum and is against viral protein antigens. TH3 immunity

induces type2antibody dependent cytotoxic hypersensitivity including chronic stage

of SLE.(91, 92)

Concluding remarks

This summary diagram: 4x2+2 immunological pathways are the whole pictures of

host immunological pathways. There are four eradicable immune responses:

TH1/TH2/TH22/THαβ which are corresponding to four tolerable immune responses:

TH1Like/TH9/TH17/TH3. It will match the four types of hypersensitivity. Then, we

can clearly understand the detailed immune response against different pathogens as

well as allergy/hypersensitivity.

Highlights

Recently, many new host immunological pathways other than traditional TH1/TH2

diagram were discovered including TH17, TH9, TH22, and Follicular helper T cells.

Recent studies suggest there is a link between the new immunological pathways and

the autoimmunities. However, a detailed framework is still not known.

Recent studies also suggest different pathogens are related to these newly discovered

host immunological pathways. However, we still need a detailed framework to

describe the functional classification of host immune responses.

Outstanding questions box

Further studies are needed to verify the accuracy of this new framework of host immunological pathways and their relationship to different pathogens and different types of autoimmune responses.

Acknowledgement

The author is very thankful for the instruction by Professors Alan Scott, Louis August Bourgeois, and Pien-Chien Huang in my PhD study in vaccine science track of Department of International Health of Johns Hopkins University Bloomberg School of Public Health. The author is also very grateful for the guidance of Professors Chi-Huey Wong and Alice, Lin-Tsing Yu in my postdoctorate research in Genomics Research Center of Academia Sinica, Taiwan.

Author's Information

The author, Wan-Chung Hu, graduated as a MD from National Taiwan University. Then, he went to Department of International Health, Johns Hopkins University Bloomberg School of Public Health for study and got a vaccine science PhD. His PhD thesis is to study host immune reaction against malarial infection. He did postdoctorate study for cancer immunotherapy in Genomics Research Center, Academia Sinica, Taiwan. He finished PGY training in Mackay Memorial Hospital and Shin-Kong Memorial Hospital both in Taiwan. Then, he is currently working as a chief resident in Department of Clinical Pathology, Far Eastern Memorial Hospital of Taiwan R.O.C. for resident training.

Please see the brief abstract in **BioRxiv doi:** https://doi.org/10.1101/006965.

References

1.   T. R. Mosmann, H. Cherwinski, M. W. Bond, M. A. Giedlin and R. L. Coffman: Two
types of murine helper T cell clone. I. Definition according to profiles of lymphokine
activities and secreted proteins. *J Immunol*, 136(7), 2348-57 (1986)

2.   W. C. Hu: Human immune responses to Plasmodium falciparum infection:
molecular evidence for a suboptimal THalphabeta and TH17 bias over ideal and
effective traditional TH1 immune response. *Malar J*, 12(1), 392 (2013)
doi:10.1186/1475-2875-12-392

3.   S. Kumar, R. A. Naqvi, N. Khanna, P. Pathak and D. N. Rao: Th3 immune
responses in the progression of leprosy via molecular cross-talks of TGF-beta, CTLA-4
and Cbl-b. *Clin Immunol*, 141(2), 133-42 (2011) doi:10.1016/j.clim.2011.06.007

4.   M. K. Levings, R. Sangregorio, F. Galbiati, S. Squadrone, R. de Waal Malefyt and
M. G. Roncarolo: IFN-alpha and IL-10 induce the differentiation of human type 1 T
regulatory cells. *J Immunol*, 166(9), 5530-9 (2001)

5.   L. E. Harrington, R. D. Hatton, P. R. Mangan, H. Turner, T. L. Murphy, K. M.
Murphy and C. T. Weaver: Interleukin 17-producing CD4+ effector T cells develop via
a lineage distinct from the T helper type 1 and 2 lineages. *Nat Immunol*, 6(11), 1123-
32 (2005) doi:10.1038/ni1254

6.   S. Eyerich, K. Eyerich, D. Pennino, T. Carbone, F. Nasorri, S. Pallotta, F. Cianfarani,
T. Odorisio, C. Traidl-Hoffmann, H. Behrendt, S. R. Durham, C. B. Schmidt-Weber and
A. Cavani: Th22 cells represent a distinct human T cell subset involved in epidermal
immunity and remodeling. *J Clin Invest*, 119(12), 3573-85 (2009)
doi:10.1172/JCI40202

7.   T. Chtanova, S. G. Tangye, R. Newton, N. Frank, M. R. Hodge, M. S. Rolph and C.
R. Mackay: T follicular helper cells express a distinctive transcriptional profile,
reflecting their role as non-Th1/Th2 effector cells that provide help for B cells. *J
Immunol*, 173(1), 68-78 (2004)

8.   M. Dominguez-Villar, C. M. Baecher-Allan and D. A. Hafler: Identification of T
helper type 1-like, Foxp3+ regulatory T cells in human autoimmune disease. *Nat
Med*, 17(6), 673-5 (2011) doi:10.1038/nm.2389

9.   S. Hori, T. Nomura and S. Sakaguchi: Control of regulatory T cell development by
the transcription factor Foxp3. *Science*, 299(5609), 1057-61 (2003)
doi:10.1126/science.1079490

10.   C. G. Vinuesa, M. A. Linterman, D. Yu and I. C. MacLennan: Follicular Helper T
Cells. *Annu Rev Immunol*, 34, 335-68 (2016) doi:10.1146/annurev-immunol-041015-
055605

11.   D. Fina, M. Sarra, R. Caruso, G. Del Vecchio Blanco, F. Pallone, T. T. MacDonald
and G. Monteleone: Interleukin 21 contributes to the mucosal T helper cell type 1
response in coeliac disease. *Gut*, 57(7), 887-92 (2008) doi:10.1136/gut.2007.129882

12.   K. Luthje, A. Kallies, Y. Shimohakamada, G. T. Belz, A. Light, D. M. Tarlinton and S.
L. Nutt: The development and fate of follicular helper T cells defined by an IL-21
reporter mouse. *Nat Immunol*, 13(5), 491-8 (2012) doi:10.1038/ni.2261

13.   Y. Liu, S. Yu, Z. Li, J. Ma, Y. Zhang, H. Wang, B. Yang and C. Wu: TGF-beta
enhanced IL-21-induced differentiation of human IL-21-producing CD4+ T cells via
Smad3. *PLoS One*, 8(5), e64612 (2013) doi:10.1371/journal.pone.0064612

14.   M. J. McCarron and J. C. Marie: TGF-beta prevents T follicular helper cell
accumulation and B cell autoreactivity. *J Clin Invest*, 124(10), 4375-86 (2014)
doi:10.1172/jci76179

15.   T. H. Tran, F. E. Utama, J. Lin, N. Yang, A. B. Sjolund, A. Ryder, K. J. Johnson, L. M.
Neilson, C. Liu, K. L. Brill, A. L. Rosenberg, A. K. Witkiewicz and H. Rui: Prolactin
inhibits BCL6 expression in breast cancer through a Stat5a-dependent mechanism.
Cancer Res, 70(4), 1711-21 (2010) doi:10.1158/0008-5472.can-09-2314

16.   R. J. Johnston, Y. S. Choi, J. A. Diamond, J. A. Yang and S. Crotty: STAT5 is a potent
negative regulator of TFH cell differentiation. *J Exp Med*, 209(2), 243-50 (2012)
doi:10.1084/jem.20111174

17.   F. A. Scheeren, M. Naspetti, S. Diehl, R. Schotte, M. Nagasawa, E. Wijnands, R.
Gimeno, F. A. Vyth-Dreese, B. Blom and H. Spits: STAT5 regulates the self-renewal
capacity and differentiation of human memory B cells and controls Bcl-6 expression.
Nat Immunol, 6(3), 303-13 (2005) doi:10.1038/ni1172

18.   F. Verdier, R. Rabionet, F. Gouilleux, C. Beisenherz-Huss, P. Varlet, O. Muller, P.
Mayeux, C. Lacombe, S. Gisselbrecht and S. Chretien: A sequence of the CIS gene
promoter interacts preferentially with two associated STAT5A dimers: a distinct
biochemical difference between STAT5A and STAT5B. *Mol Cell Biol*, 18(10), 5852-60
(1998)

19.   D. Baumjohann, T. Okada and K. M. Ansel: Cutting Edge: Distinct waves of BCL6
expression during T follicular helper cell development. *J Immunol*, 187(5), 2089-92
(2011) doi:10.4049/jimmunol.1101393

20.   M. A. Linterman, L. Beaton, D. Yu, R. R. Ramiscal, M. Srivastava, J. J. Hogan, N. K.
Verma, M. J. Smyth, R. J. Rigby and C. G. Vinuesa: IL-21 acts directly on B cells to
regulate Bcl-6 expression and germinal center responses. *J Exp Med*, 207(2), 353-63
(2010) doi:10.1084/jem.20091738

21.   R. I. Nurieva, Y. Chung, G. J. Martinez, X. O. Yang, S. Tanaka, T. D. Matskevitch, Y.

H. Wang and C. Dong: Bcl6 mediates the development of T follicular helper cells.

Science, 325(5943), 1001-5 (2009) doi:10.1126/science.1176676

22.   V. L. Bryant, C. S. Ma, D. T. Avery, Y. Li, K. L. Good, L. M. Corcoran, R. de Waal

Malefyt and S. G. Tangye: Cytokine-mediated regulation of human B cell

differentiation into Ig-secreting cells: predominant role of IL-21 produced by CXCR5+

T follicular helper cells. *J Immunol*, 179(12), 8180-90 (2007)

23.   Y. S. Choi, D. Eto, J. A. Yang, C. Lao and S. Crotty: Cutting edge: STAT1 is required

for IL-6-mediated Bcl6 induction for early follicular helper cell differentiation. *J*

Immunol, 190(7), 3049-53 (2013) doi:10.4049/jimmunol.1203032

24.   P. Schaerli, P. Loetscher and B. Moser: Cutting edge: induction of follicular

homing precedes effector Th cell development. *J Immunol*, 167(11), 6082-6 (2001)

25.   S. Nakayamada, Y. Kanno, H. Takahashi, D. Jankovic, K. T. Lu, T. A. Johnson, H. W.

Sun, G. Vahedi, O. Hakim, R. Handon, P. L. Schwartzberg, G. L. Hager and J. J. O'Shea:

Early Th1 cell differentiation is marked by a Tfh cell-like transition. *Immunity*, 35(6),

919-31 (2011) doi:10.1016/j.immuni.2011.11.012

26.   C. Schindler, D. E. Levy and T. Decker: JAK-STAT signaling: from interferons to

cytokines. *J Biol Chem*, 282(28), 20059-63 (2007) doi:10.1074/jbc.R700016200

27.   C. R. Mahon, D.C.Lehman and G. Manuselis: Textbook of Diagnostic

Microbiology. Elsevier

(2014)

28.   G. Iezzi, A. Boni, E. Degl'Innocenti, M. Grioni, M. T. Bertilaccio and M. Bellone:

Type 2 cytotoxic T lymphocytes modulate the activity of dendritic cells toward type 2

immune responses. *J Immunol*, 177(4), 2131-7 (2006)

29.   D. Gong, W. Shi, S. J. Yi, H. Chen, J. Groffen and N. Heisterkamp: TGFbeta

signaling plays a critical role in promoting alternative macrophage activation. *BMC*

Immunol, 13, 31 (2012) doi:10.1186/1471-2172-13-31

30.   P. Romero, A. Zippelius, I. Kurth, M. J. Pittet, C. Touvrey, E. M. Iancu, P. Corthesy,

E. Devevre, D. E. Speiser and N. Rufer: Four functionally distinct populations of

human effector-memory CD8+ T lymphocytes. *J Immunol*, 178(7), 4112-9 (2007)

31.   P. A. Baars, S. Sierro, R. Arens, K. Tesselaar, B. Hooibrink, P. Klenerman and R. A.

van Lier: Properties of murine (CD8+)CD27- T cells. *Eur J Immunol*, 35(11), 3131-41

(2005) doi:10.1002/eji.200425770

32.   D. G. Mordue, F. Monroy, M. La Regina, C. A. Dinarello and L. D. Sibley: Acute

toxoplasmosis leads to lethal overproduction of Th1 cytokines. *J Immunol*, 167(8),

4574-84 (2001)

33.   U. Jung, J. E. Foley, A. A. Erdmann, M. A. Eckhaus and D. H. Fowler: CD3/CD28-

costimulated T1 and T2 subsets: differential in vivo allosensitization generates

distinct GVT and GVHD effects. *Blood*, 102(9), 3439-46 (2003) doi:10.1182/blood-
2002-12-3936

34.   C. Mesnil, S. Raulier, G. Paulissen, X. Xiao, M. A. Birrell, D. Pirottin, T. Janss, P.
Starkl, E. Ramery, M. Henket, F. N. Schleich, M. Radermecker, K. Thielemans, L. Gillet,
M. Thiry, M. G. Belvisi, R. Louis, C. Desmet, T. Marichal and F. Bureau: Lung-resident
eosinophils represent a distinct regulatory eosinophil subset. *J Clin Invest*, 126(9),
3279-95 (2016) doi:10.1172/jci85664

35.   H. Toru, M. Eguchi, R. Matsumoto, M. Yanagida, J. Yata and T. Nakahata:
Interleukin-4 promotes the development of tryptase and chymase double-positive
human mast cells accompanied by cell maturation. *Blood*, 91(1), 187-95 (1998)

36.   S. Higuchi, M. Kobayashi, A. Yano, K. Tsuneyama, T. Fukami, M. Nakajima and T.
Yokoi: Involvement of Th2 cytokines in the mouse model of flutamide-induced acute
liver injury. *J Appl Toxicol*, 32(10), 815-22 (2012) doi:10.1002/jat.1706

37.   M. A. Cooper, T. A. Fehniger and M. A. Caligiuri: The biology of human natural
killer-cell subsets. *Trends Immunol*, 22(11), 633-40 (2001)

38.   D. Peritt, S. Robertson, G. Gri, L. Showe, M. Aste-Amezaga and G. Trinchieri:
Differentiation of human NK cells into NK1 and NK2 subsets. *J Immunol*, 161(11),
5821-4 (1998)

39.   H. Nagumo and K. Agematsu: Synergistic augmentative effect of interleukin-10
and CD27/CD70 interactions on B-cell immunoglobulin synthesis. *Immunology*, 94(3),
388-94 (1998)

40.   J. Hendriks, L. A. Gravestein, K. Tesselaar, R. A. van Lier, T. N. Schumacher and J.
Borst: CD27 is required for generation and long-term maintenance of T cell immunity.
Nat Immunol, 1(5), 433-40 (2000) doi:10.1038/80877

41.   K. B. Nguyen, L. P. Cousens, L. A. Doughty, G. C. Pien, J. E. Durbin and C. A. Biron:
Interferon alpha/beta-mediated inhibition and promotion of interferon gamma:
STAT1 resolves a paradox. *Nat Immunol*, 1(1), 70-6 (2000) doi:10.1038/76940

42.   M. Nagaki, H. Iwai, T. Naiki, H. Ohnishi, Y. Muto and H. Moriwaki: High levels of
serum interleukin-10 and tumor necrosis factor-alpha are associated with fatality in
fulminant hepatitis. *J Infect Dis*, 182(4), 1103-8 (2000) doi:10.1086/315826

43.   S. J. Aujla, Y. R. Chan, M. Zheng, M. Fei, D. J. Askew, D. A. Pociask, T. A. Reinhart,
F. McAllister, J. Edeal, K. Gaus, S. Husain, J. L. Kreindler, P. J. Dubin, J. M. Pilewski, M.
M. Myerburg, C. A. Mason, Y. Iwakura and J. K. Kolls: IL-22 mediates mucosal host
defense against Gram-negative bacterial pneumonia. *Nat Med*, 14(3), 275-81 (2008)
doi:10.1038/nm1710

44.   K. Wolk, S. Kunz, E. Witte, M. Friedrich, K. Asadullah and R. Sabat: IL-22
increases the innate immunity of tissues. *Immunity*, 21(2), 241-54 (2004)
doi:10.1016/j.immuni.2004.07.007

45. K. Ghoreschi, A. Laurence, X. P. Yang, C. M. Tato, M. J. McGeachy, J. E. Konkel, H.

L. Ramos, L. Wei, T. S. Davidson, N. Bouladoux, J. R. Grainger, Q. Chen, Y. Kanno, W. T.

Watford, H. W. Sun, G. Eberl, E. M. Shevach, Y. Belkaid, D. J. Cua, W. Chen and J. J.

O'Shea: Generation of pathogenic T(H)17 cells in the absence of TGF-beta signalling.

Nature, 467(7318), 967-71 (2010) doi:10.1038/nature09447

46. S. Trifari, C. D. Kaplan, E. H. Tran, N. K. Crellin and H. Spits: Identification of a

human helper T cell population that has abundant production of interleukin 22 and is

distinct from T(H)-17, T(H)1 and T(H)2 cells. *Nat Immunol*, 10(8), 864-71 (2009)

doi:10.1038/ni.1770

47. T. Duhen, R. Geiger, D. Jarrossay, A. Lanzavecchia and F. Sallusto: Production of

interleukin 22 but not interleukin 17 by a subset of human skin-homing memory T

cells. *Nat Immunol*, 10(8), 857-63 (2009) doi:10.1038/ni.1767

48. Y. Yoshida, A. Kumar, Y. Koyama, H. Peng, A. Arman, J. A. Boch and P. E. Auron:

Interleukin 1 activates STAT3/nuclear factor-kappaB cross-talk via a unique TRAF6-

and p65-dependent mechanism. *J Biol Chem*, 279(3), 1768-76 (2004)

doi:10.1074/jbc.M311498200

49. S. Rutz, R. Noubade, C. Eidenschenk, N. Ota, W. Zeng, Y. Zheng, J. Hackney, J.

Ding, H. Singh and W. Ouyang: Transcription factor c-Maf mediates the TGF-beta-

dependent suppression of IL-22 production in T(H)17 cells. *Nat Immunol*, 12(12),

1238-45 (2011) doi:10.1038/ni.2134

50. K. Zhao, D. Zhao, D. Huang, X. Song, C. Chen, B. Pan, Q. Wu, J. Cao, Y. Yao, L. Zeng

and K. Xu: The identification and characteristics of IL-22-producing T cells in acute

graft-versus-host disease following allogeneic bone marrow transplantation.

Immunobiology, 218(12), 1505-13 (2013) doi:10.1016/j.imbio.2013.05.005

51. R. Hussain, G. Dawood, N. Abrar, Z. Toossi, A. Minai, M. Dojki and J. J. Ellner:

Selective increases in antibody isotypes and immunoglobulin G subclass responses to

secreted antigens in tuberculosis patients and healthy household contacts of the

patients. *Clin Diagn Lab Immunol*, 2(6), 726-32 (1995)

52. Y. S. Goh, A. J. Grant, O. Restif, T. J. McKinley, K. L. Armour, M. R. Clark and P.

Mastroeni: Human IgG isotypes and activating Fcgamma receptors in the interaction

of Salmonella enterica serovar Typhimurium with phagocytic cells. *Immunology*,

133(1), 74-83 (2011) doi:10.1111/j.1365-2567.2011.03411.x

53. J. P. Coutelier, J. T. van der Logt, F. W. Heessen, A. Vink and J. van Snick: Virally

induced modulation of murine IgG antibody subclasses. *J Exp Med*, 168(6), 2373-8

(1988)

54. G. R. Siber, P. H. Schur, A. C. Aisenberg, S. A. Weitzman and G. Schiffman:

Correlation between serum IgG-2 concentrations and the antibody response to

bacterial polysaccharide antigens. *N Engl J Med*, 303(4), 178-82 (1980)

doi:10.1056/nejm198007243030402

55.   J. D. Turner, H. Faulkner, J. Kamgno, M. W. Kennedy, J. Behnke, M. Boussinesq

and J. E. Bradley: Allergen-specific IgE and IgG4 are markers of resistance and

susceptibility in a human intestinal nematode infection. *Microbes Infect*, 7(7-8), 990-

6 (2005) doi:10.1016/j.micinf.2005.03.036

56.   Z. Yao, Y. Kanno, M. Kerenyi, G. Stephens, L. Durant, W. T. Watford, A. Laurence,

G. W. Robinson, E. M. Shevach, R. Moriggl, L. Hennighausen, C. Wu and J. J. O'Shea:

Nonredundant roles for Stat5a/b in directly regulating Foxp3. *Blood*, 109(10), 4368-

75 (2007) doi:10.1182/blood-2006-11-055756

57.   H. X. Wei, Y. H. Chuang, B. Li, H. Wei, R. Sun, Y. Moritoki, M. E. Gershwin, Z. X.

Lian and Z. Tian: CD4+ CD25+ Foxp3+ regulatory T cells protect against T cell-

mediated fulminant hepatitis in a TGF-beta-dependent manner in mice. *J Immunol*,

181(10), 7221-9 (2008)

58.   H. Zan, A. Cerutti, P. Dramitinos, A. Schaffer and P. Casali: CD40 engagement

triggers switching to IgA1 and IgA2 in human B cells through induction of

endogenous TGF-beta: evidence for TGF-beta but not IL-10-dependent direct S mu--

>S alpha and sequential S mu-->S gamma, S gamma-->S alpha DNA recombination. *J*

Immunol, 161(10), 5217-25 (1998)

59.   C. Andre, F. Andre and C. Fargier: Distribution of IgA 1 and IgA 2 plasma cells in

various normal human tissues and in the jejunum of plasma IgA-deficient patients.

Clin Exp Immunol, 33(2), 327-31 (1978)

60.   B. He, W. Xu, P. A. Santini, A. D. Polydorides, A. Chiu, J. Estrella, M. Shan, A.

Chadburn, V. Villanacci, A. Plebani, D. M. Knowles, M. Rescigno and A. Cerutti:

Intestinal bacteria trigger T cell-independent immunoglobulin A(2) class switching by

inducing epithelial-cell secretion of the cytokine APRIL. *Immunity*, 26(6), 812-26

(2007) doi:10.1016/j.immuni.2007.04.014

61.   J. Ochoa-Reparaz, A. Rynda, M. A. Ascon, X. Yang, I. Kochetkova, C. Riccardi, G.

Callis, T. Trunkle and D. W. Pascual: IL-13 production by regulatory T cells protects

against experimental autoimmune encephalomyelitis independently of autoantigen. *J*

Immunol, 181(2), 954-68 (2008)

62.   D. A. Vignali, L. W. Collison and C. J. Workman: How regulatory T cells work. *Nat*

Rev Immunol, 8(7), 523-32 (2008) doi:10.1038/nri2343

63.   R. A. O'Connor, M. D. Leech, J. Suffner, G. J. Hammerling and S. M. Anderton:

Myelin-reactive, TGF-beta-induced regulatory T cells can be programmed to develop

Th1-like effector function but remain less proinflammatory than myelin-reactive Th1

effectors and can suppress pathogenic T cell clonal expansion in vivo. *J Immunol*,

185(12), 7235-43 (2010) doi:10.4049/jimmunol.1001551

64.   K. J. Oestreich, S. E. Mohn and A. S. Weinmann: Molecular mechanisms that

control the expression and activity of Bcl-6 in TH1 cells to regulate flexibility with a

TFH-like gene profile. *Nat Immunol*, 13(4), 405-11 (2012) doi:10.1038/ni.2242

65. J. Prochazkova, K. Pokorna and V. Holan: IL-12 inhibits the TGF-beta-dependent

T cell developmental programs and skews the TGF-beta-induced differentiation into a

Th1-like direction. *Immunobiology*, 217(1), 74-82 (2012)

doi:10.1016/j.imbio.2011.07.032

66. A. Gorter, P. S. Hiemstra, E. A. van der Voort, L. A. van Es and M. R. Daha:

Binding of human IgA1 to rat peritoneal macrophages. *Immunology*, 64(2), 207-12

(1988)

67. S. Balu, R. Reljic, M. J. Lewis, R. J. Pleass, R. McIntosh, C. van Kooten, M. van

Egmond, S. Challacombe, J. M. Woof and J. Ivanyi: A novel human IgA monoclonal

antibody protects against tuberculosis. *J Immunol*, 186(5), 3113-9 (2011)

doi:10.4049/jimmunol.1003189

68. L. Cosmi, F. Liotta, E. Maggi, S. Romagnani and F. Annunziato: Th17 and non-

classic Th1 cells in chronic inflammatory disorders: two sides of the same coin. *Int*

Arch Allergy Immunol, 164(3), 171-7 (2014) doi:10.1159/000363502

69. K. Gerlach, Y. Hwang, A. Nikolaev, R. Atreya, H. Dornhoff, S. Steiner, H. A. Lehr, S.

Wirtz, M. Vieth, A. Waisman, F. Rosenbauer, A. N. McKenzie, B. Weigmann and M. F.

Neurath: TH9 cells that express the transcription factor PU.1 drive T cell-mediated

colitis via IL-9 receptor signaling in intestinal epithelial cells. *Nat Immunol*, 15(7), 676-

86 (2014) doi:10.1038/ni.2920

70. V. Dardalhon, A. Awasthi, H. Kwon, G. Galileos, W. Gao, R. A. Sobel, M.

Mitsdoerffer, T. B. Strom, W. Elyaman, I. C. Ho, S. Khoury, M. Oukka and V. K. Kuchroo:

IL-4 inhibits TGF-beta-induced Foxp3+ T cells and, together with TGF-beta, generates

IL-9+ IL-10+ Foxp3(-) effector T cells. *Nat Immunol*, 9(12), 1347-55 (2008)

doi:10.1038/ni.1677

71. R. Goswami, R. Jabeen, R. Yagi, D. Pham, J. Zhu, S. Goenka and M. H. Kaplan:

STAT6-dependent regulation of Th9 development. *J Immunol*, 188(3), 968-75 (2012)

doi:10.4049/jimmunol.1102840

72. C. M. Westerberg, E. Ulleras and G. Nilsson: Differentiation of mast cell

subpopulations from mouse embryonic stem cells. *J Immunol Methods*, 382(1-2),

160-6 (2012) doi:10.1016/j.jim.2012.05.020

73. J. Kerzerho, H. Maazi, A. O. Speak, N. Szely, V. Lombardi, B. Khoo, S. Geryak, J.

Lam, P. Soroosh, J. Van Snick and O. Akbari: Programmed cell death ligand 2 regulates

TH9 differentiation and induction of chronic airway hyperreactivity. *J Allergy Clin*

Immunol, 131(4), 1048-57, 1057 e1-2 (2013) doi:10.1016/j.jaci.2012.09.027

74. C. Pilette, K. T. Nouri-Aria, M. R. Jacobson, L. K. Wilcock, B. Detry, S. M. Walker, J.

N. Francis and S. R. Durham: Grass pollen immunotherapy induces an allergen-

specific IgA2 antibody response associated with mucosal TGF-beta expression. *J*

Immunol, 178(7), 4658-66 (2007)

75.   M. Iikura, M. Yamaguchi, T. Fujisawa, M. Miyamasu, T. Takaishi, Y. Morita, T.

Iwase, I. Moro, K. Yamamoto and K. Hirai: Secretory IgA induces degranulation of IL-

3-primed basophils. *J Immunol*, 161(3), 1510-5 (1998)

76.   K. R. Bartemes, K. M. Cooper, K. L. Drain and H. Kita: Secretory IgA induces

antigen-independent eosinophil survival and cytokine production without inducing

effector functions. *J Allergy Clin Immunol*, 116(4), 827-35 (2005)

doi:10.1016/j.jaci.2005.07.014

77.   K. Obata, K. Mukai, Y. Tsujimura, K. Ishiwata, Y. Kawano, Y. Minegishi, N.

Watanabe and H. Karasuyama: Basophils are essential initiators of a novel type of

chronic allergic inflammation. *Blood*, 110(3), 913-20 (2007) doi:10.1182/blood-2007-

01-068718

78.   C. Sillaber, K. Geissler, R. Scherrer, R. Kaltenbrunner, P. Bettelheim, K. Lechner

and P. Valent: Type beta transforming growth factors promote interleukin-3 (IL-3)-

dependent differentiation of human basophils but inhibit IL-3-dependent

differentiation of human eosinophils. *Blood*, 80(3), 634-41 (1992)

79.   Y. Chung, S. H. Chang, G. J. Martinez, X. O. Yang, R. Nurieva, H. S. Kang, L. Ma, S.

S. Watowich, A. M. Jetten, Q. Tian and C. Dong: Critical regulation of early Th17 cell

differentiation by interleukin-1 signaling. *Immunity*, 30(4), 576-87 (2009)

doi:10.1016/j.immuni.2009.02.007

80.   S. C. Liang, X. Y. Tan, D. P. Luxenberg, R. Karim, K. Dunussi-Joannopoulos, M.

Collins and L. A. Fouser: Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells

and cooperatively enhance expression of antimicrobial peptides. *J Exp Med*, 203(10),

2271-9 (2006) doi:10.1084/jem.20061308

81.   X. Liu, S. Leung, C. Wang, Z. Tan, J. Wang, T. B. Guo, L. Fang, Y. Zhao, B. Wan, X.

Qin, L. Lu, R. Li, H. Pan, M. Song, A. Liu, J. Hong, H. Lu and J. Z. Zhang: Crucial role of

interleukin-7 in T helper type 17 survival and expansion in autoimmune disease. *Nat*

Med, 16(2), 191-7 (2010) doi:10.1038/nm.2077

82.   G. Beriou, C. M. Costantino, C. W. Ashley, L. Yang, V. K. Kuchroo, C. Baecher-Allan

and D. A. Hafler: IL-17-producing human peripheral regulatory T cells retain

suppressive function. *Blood*, 113(18), 4240-9 (2009) doi:10.1182/blood-2008-10-

183251

83.   K. S. Voo, Y. H. Wang, F. R. Santori, C. Boggiano, Y. H. Wang, K. Arima, L. Bover, S.

Hanabuchi, J. Khalili, E. Marinova, B. Zheng, D. R. Littman and Y. J. Liu: Identification

of IL-17-producing FOXP3+ regulatory T cells in humans. *Proc Natl Acad Sci U S A*,

106(12), 4793-8 (2009) doi:10.1073/pnas.0900408106

84.   Z. G. Fridlender, J. Sun, S. Kim, V. Kapoor, G. Cheng, L. Ling, G. S. Worthen and S.

M. Albelda: Polarization of tumor-associated neutrophil phenotype by TGF-beta: "N1" versus "N2" TAN. *Cancer Cell*, 16(3), 183-94 (2009) doi:10.1016/j.ccr.2009.06.017

85. P. Miossec and J. K. Kolls: Targeting IL-17 and TH17 cells in chronic inflammation. *Nat Rev Drug Discov*, 11(10), 763-76 (2012) doi:10.1038/nrd3794

86. I. Backert, S. B. Koralov, S. Wirtz, V. Kitowski, U. Billmeier, E. Martini, K. Hofmann, K. Hildner, N. Wittkopf, K. Brecht, M. Waldner, K. Rajewsky, M. F. Neurath, C. Becker and C. Neufert: STAT3 activation in Th17 and Th22 cells controls IL-22-mediated epithelial host defense during infectious colitis. *J Immunol*, 193(7), 3779-91 (2014) doi:10.4049/jimmunol.1303076

87. Z. M. Chen, M. J. O'Shaughnessy, I. Gramaglia, A. Panoskaltsis-Mortari, W. J. Murphy, S. Narula, M. G. Roncarolo and B. R. Blazar: IL-10 and TGF-beta induce alloreactive CD4+CD25- T cells to acquire regulatory cell function. *Blood*, 101(12), 5076-83 (2003) doi:10.1182/blood-2002-09-2798

88. A. Doetze, J. Satoguina, G. Burchard, T. Rau, C. Loliger, B. Fleischer and A. Hoerauf: Antigen-specific cellular hyporesponsiveness in a chronic human helminth infection is mediated by T(h)3/T(r)1-type cytokines IL-10 and transforming growth factor-beta but not by a T(h)1 to T(h)2 shift. *Int Immunol*, 12(5), 623-30 (2000)

89. A. Tulunay, S. Yavuz, H. Direskeneli and E. Eksioglu-Demiralp: CD8+CD28-, suppressive T cells in systemic lupus erythematosus. *Lupus*, 17(7), 630-7 (2008) doi:10.1177/0961203308089400

90. D. B. Keskin, D. S. Allan, B. Rybalov, M. M. Andzelm, J. N. Stern, H. D. Kopcow, L. A. Koopman and J. L. Strominger: TGFbeta promotes conversion of CD16+ peripheral blood NK cells into CD16- NK cells with similarities to decidual NK cells. *Proc Natl Acad Sci U S A*, 104(9), 3378-83 (2007) doi:10.1073/pnas.0611098104

91. B. Chaudhary and E. Elkord: Downregulation of immunosuppressive environment in patients with chronic HBV hepatitis on maintained remission. *Front Immunol*, 6, 52 (2015) doi:10.3389/fimmu.2015.00052

92. S. Ohga, A. Nomura, H. Takada, T. Tanaka, K. Furuno, Y. Takahata, N. Kinukawa, N. Fukushima, S. Imai and T. Hara: Dominant expression of interleukin-10 and transforming growth factor-beta genes in activated T-cells of chronic active Epstein-Barr virus infection. *J Med Virol*, 74(3), 449-58 (2004) doi:10.1002/jmv.20197

Figure legends

Figure 1. The summary diagram of host immunological pathways. In the middle, Tfh

side (follicular help T cell) begins the initiatory immunity; on the other hand, Treg

side (regulatory T cells) starts the regulatory immunity. Eradicable TH1 and tolerable

TH1-like(TH1k) are related in the diagonal line. Eradicable TH2 and tolerable TH9

are related in the diagonal line. Eradicable TH22 and tolerable TH17 are related in the

diagonal line. Eradicable THαβ and tolerable TH3 are related in the diagonal line.

Table 1. Summary of host immunological pathways

Immune pathways	Driven cytokine	Transcription factor	Effector cells	CD4 T cells	B cells	Pathogen	Hypersensitivities
Initiatory							
Tfh	IL-21	STAT1, STAT3, STAT5B		IL-21 CD4 T cells	IgM B cells	Acute infection	
Eradicable immunities							
TH1	IL-12	STAT4	Macrophage(M1), CTL(Tc1)	IFNg CD4 T cells	IgG3 B cells	Intracellular bacteria & protozoa	Type 4 DTH
TH2	IL-4	STAT6	Eosinophil(iEOS), basophil, mast cells(MCct)	IL-4 IL-5 CD4 T cells	IgG4/IgE B cells	Helminths & insects	Type 1 allergy
TH22	IL-1	STAT3	Neutrophils(N1)	IL-1, TNFα, IL-22 CD4 T cells	IgG2 B cells	Extracellular bacteria & fungi	Type3 Immune complex
TH$\alpha\beta$	IL-10 IFN$\alpha\beta$	STAT1 STAT2	NK cells(NK1), CTL(Tc2)	IL-10 CD4 T cells	IgG1 B cells	viruses	Type 2 ADCC

Immune pathways	Driven cytokine	Transcription factor	Effector cells	CD4 T cells	B cells	Pathogen	Hypersensitivities
Regulatory							
Treg	TGFβ	STAT5A, STAT5B		TGFβ CD4 T cells	IgA B cells	Chronic infection	
Tolerable immunities							
TH1like	IL-12, TGFβ	STAT4, STAT5	Macrophage(M2), CD8 Treg	IFNg/TGFβ CD4 T cells	IgA1 B cells	Intracellular bacteria & protozoa	Type 4 DTH
TH9	IL-4, TGFβ	STAT6, STAT5	Eosinophil(rEOS), basophil, mast cells(MCt)	IL-9 CD4 T cells	IgA2 B cells	Helminths & insects	Type 1 allergy
TH17	IL-6, TGFβ	STAT3, STAT5	Neutrophils(N2)	IL-17 CD4 T cells	IgA2 B cells	Extracellular bacteria & fungi	Type3 immune complex
TH3	IL-10, TGFβ	STAT1 STAT2 STAT5	NK cells(NK2), CD8 Treg	IL-10/TGFβ CD4 T cells	IgA1 B cells	viruses	Type 2 ADCC