A Framework of All Discovered Immunological Pathways and Their Roles for

Four Specific Types of Pathogens and Hypersensitivities

Running title: A Framework of Host Immunities

Wan-Chung Hu*

MD National Taiwan University

PhD Johns Hopkins University

Department of Clinical Pathology, Far Eastern Memorial Hospital, New Taipei City,

Taiwan

Department of Clinical Pathology, Taipei Tzu Chi Hospital, New Taipei City, Taiwan

*Correspondence: Wan-Chung Hu, Wanchung.Hu09@gmail.com (W. Hu)

Keywords: Th1/2, Th17, Th9, Th22, Treg, Tfh

Summary sentence: The summarized framework of host immunities to explain their

relations to specific pathogens and hypersensitivities

Abstract

Tfh initiates four eradicable immunities. Tfh includes FDC, LTi, IL21 CD4 T cell, and

IgG/M B cell. Treg initiates four tolerable immunities. Treg includes DCreg, ILCreg,

TGFβ CD4 T cell, and IgA B cell. TH1/TH1-like is immunity for intracellular

bacteria/protozoa and type 4 delayed type hypersensitivity. TH1 includes M1

macrophage, mDC2, Tc1 CD8 T cell, IFNg CD4 T cell, ILC1, iNKT1, and IgG3 B

cell. TH1-like includes M2 macrophage, ILC1, suppressive CD8 T cell, IFNg/TGFβ

CD4 T cell, regulatory iNKT cells, and IgA1 B cell. TH2/TH9 is immunity for

helminths and type1 IgE mediated hypersensitivity. TH2 includes iEOS eosinophil,

Langerhans cell, basophil/MCct mast cell, IL-4 CD4 T cell, ILC2, iNKT2, and

IgE/IgG4 B cell. TH9 includes rEOS eosinophil, basophils/mast cell MCt, IL-9 CD4

T cell, ILC2, regulatory iNKT cells, and IgA2 B cell. TH22/TH17 is immunity for

extracellular bacteria/fungi and type 3 immune complex hypersensitivity. TH22

includes N1 neutrophils, mDC1, IL-22 CD4 T cell, ILC3(NCR+), iNKT17, and IgG2

B cell. TH17 includes N2 neutrophils, IL-17 CD4 T cell, regulatory iNKT cells,

ILC3(NCR-), and IgA2 B cell. THαβ/TH3 is immunity for viruses and type 2

antibody dependent cytotoxic hypersensitivity. THαβ includes NK1 natural killer cell,

pDC, Tc2 CD8 T cell, IL10 CD4 T cell, ILC10, iNKT10, and IgG1 B cell. TH3

includes NK2 natural killer cell, suppressive CD8 T cell, ILC10, IL-10/TGFβ CD4 T

cell, regulatory iNKT cells, and IgA1 B cell.

Background of host immunities

Many host immunological pathways have been discovered, such as traditional TH1/TH2, TH3, TH17, TH22, Tfh, Treg, TH9, and Tr1 (THαβ). These identified pathways are not logically organized. In this study, I propose a detailed and complete picture regarding host immunological pathways (Fig. 1).

The traditional TH1/TH2 paradigm was proposed by Dr. Mosmann in 1986 [1]. TH1 was thought to provide host immunity against viruses and intracellular bacteria. TH2 provides host immunity against multicellular parasites (helminthes). In my PhD thesis, I proposed a new THαβ immunological pathway against viruses derived from traditional TH1 immunity [2]. TH1 immunity focuses on intracellular bacteria and protozoa. TH3 immunity and Tr1 immunological pathways were identified after TH1 and TH2 [3,4]. Recently, additional immune responses have been discovered, including TH17, TH22, Tfh, Treg, and TH1-like immunological pathways [5-7].

Initiatory immune response

Follicular helper T cells (Tfh) are believed to be key helper cells for B-cell germinal centers in lymph nodes. Tfh cells are characterized by IL-21-producing T cells [8]. Follicular dendritic cells (CD14+) are the antigen presenting cells [9]. Lymphoid tissue inducer cells (LTi) are the innate lymphoid cells for Tfh [10]. BCL6 is a key

transcription factor for Tfh. TGFβ with an STAT5 signal can constrain the

differentiation of IL-21-producing helper T cells [11]. IL-21 production is related to

STAT1and STAT3 activation. IL-21 production is also related to STAT5 activation

because immunosuppressive prolactin can cause STAT5a to suppress BCL6

expression [12]. However, STAT5b can upregulate BCL6[13]. STAT5a and STAT5b have

distinct target genes in immune responses [14]. The transcription factor that induces Tfh

should be STAT5b. BCL6 is important in Tfh development [15]. Tfh can induce B-cells

to start producing the IgM antibody and IL-21 also helps to B cell isotype switch to

IgG[16,17]. Besides protein antigen recognized by B cells and T cells, lipid antigens also

are recognized by NKT cells. The subtype iNKTfh plays a role in immunity of

follicular helper T cells[18]. Thus, T lymphocytes are the first to initiate adaptive host

immunity [19-21]. Different STAT proteins regulate different immunological pathways. If

the infection tends to be eradicable, the host immunological pathways mentioned in

the following text are generated with other cytokines.

Eradicable immune responses

TH1 immunity is driven by IL-12 and is the host immunity against intracellular

bacteria or protozoa. The antigen presenting cells for TH1 immunity are type 2

myeloid dendritic cells (CD141+ mDC2)[9]. The main effector cells of TH1 immunity

are stimulatory macrophages (M1), IFNg-secreting cytotoxic CD8 T cells (CD28+ Tc1), IFNg-secreting CD4 T cells, iNKT1 cells, and IgG3-producing B-cells [22,23]. The initiation of eradicable immunity also needs the help of innate lymphoid cells to produce initial cytokines to drive to different immunological pathway. For TH1 immunity, the key innate lymphoid cells are ILC1[24]. The key transcription factor for TH1 immunity is STAT4. T-bet also plays a vital role in the TH1 immunological pathway. TH1 immunity against self-antigen is type 4 delayed-type hypersensitivity, such as type 1 diabetes mellitus.

TH2 immunity is driven by IL-4. TH2 provides immunity against extracellular parasites (helminths). The antigen presenting cells for TH2 immunity are Langerhans cells (CD1a+)[9,25]. The main effector cells of TH2 immunity are eosinophils (iEOS), basophils/connective tissue mast cells (MCtc, mast cell tryptase, and chymase), IL-4/IL-5-secreting CD4 T cells, iNKT2 cells, ILC2, and IgG4/IgE-producing B-cells [26]. IgG4 activates eosinophils, and IgE activates mast cells, such as in acute anaphylaxis [27]. The function of IgG4–eosinophil is to activate eosinophil-mediated cellular immunity against parasites or insects. The function of IgE–mast cells is to expel helminths or insects through a physiological mechanism. Mast cells activated by IgE can release histamine, which causes bronchoconstriction, vomiting/nausea, rhinorrhea, skin itchiness, stomach acidification, increased local vascular

permeability, or increased bowel movement. These actions can all help to expel

helminths or insects physiologically. The key transcription factor for TH2 immunity is

STAT6. GATA3 also plays a vital role in the TH2 immunological pathway. TH2

immunity against self-antigen is a type1 immediate allergy, such as food/drug allergy

or urticaria.

THαβ is distinguished from traditional TH1 immunity [2]. THαβ immunity is

against viruses and was called Tr1 cell by a few previous researchers[4]. THαβ

immunity is driven by IFNa/b or IL-10. The antigen presenting cells for THαβ

immunity are plasmacytoid dendritic cells (pDC)[9]. The main effector cells of THαβ

immunity are IL-10-producing stimulatory NK cells (CD56−CD16 + NK1 cells), IL-

10/IL-27-secreting CD4 T cells, IL-10-secreting cytotoxic CD8 T cells (CD28+ Tc2),

iNKT10 cells, ILC10, and IgG1-producing B-cells [22,28-30]. The CD27 molecule is

important for virus immunity. The key transcription factors for THαβ immunity are

STAT1and STAT2[31]. THαβ immunity against self-antigen is a type 2 antibody-

dependent cytotoxic hypersensitivity, such as an acute stage of myasthenia gravis. IL-

10 is not merely an immunosuppressive cytokine; it can also have potent stimulatory

effects on NK cells, CTLs, and B-cells.

TH22 is the host innate immunity against extracellular bacteria and fungi. TH22

is driven by IL-6 or TNFa [32]. The antigen presenting cells for TH22 immunity are type

1 myeloid dendritic cells (CD1c+ mDC1)[9]. The main effector cells for TH22

immunity are PMNs (N1), IL-22-secreting CD4 T cells, complements, pentraxins,

iNKT17 cells, ILC3(NCR+), and IgG2-producing B-cells [6,33]. The key transcription

factor for TH22 is STAT3. AP1 and CEBP are also important transcription factors.

TGFβ can suppress IL-22 to skew to TH17 immunity [34]. TH22 against self-antigen is

a type 3 immune-complex and complement-mediated hypersensitivity, such as the

Arthus reaction. The extracellular or intracellular locations of protozoa or fungi

mainly decide the host immunological pathways.

Four IgG subtypes fit the four types of acute immunological pathways. Murine

IgG antibodies also have four subclasses. The following correlation exists between

murine and human IgG subtypes: Human IgG1<->Murine IgG2a; Human IgG2<-

>Murine IgG3; Human IgG3<->Murine IgG2b; and Human IgG4<->Murine IgG1 [35].

hIgG1/mIgG2a works against viral antigens; hIgG2/mIgG3 works against bacterial

antigen, especially polysaccharides; hIgG3/mIgG2b works against intracellular

bacteria; and hIgG4/mIgG1 works against parasite antigens [36-38]. It is also worth

noting that the immune response against fungus or protozoa is mainly based on

intracellular or extracellular. Extracellular fungus such as Candida spp or Aspergillus

spp usually trigger TH22 immunity, but intracellular fungus such as Histoplasma spp

will trigger TH1 immunity.

Regulatory immune responses

Treg is the host immune inhibitory mechanism [39] and is driven by IL-2 and

TGFβ. Regulatory dendritic cells (DCreg) are the antigen presenting cells for Treg[40].

Regulatory innate lymphoid cells (ILCreg)are the initial helpers for Treg production[41].

The main effector cells for Treg are the TGFβ-producing CD4 T cells, FOXP3

regulatory iNKT cells, and IgA-producing B-cells[42]. The key transcription factor for

the Treg pathway is STAT5, especially STAT5a. However, both STAT5a and STAT5b

play nonredundant roles in Treg generation [43]. They may act sequentially with

STAT5b activation first in Tfh signaling. STAT5b and STAT5a combined signaling

induces Treg generation. The combination of Treg and the aforementioned four

immunological pathways is important to shift adaptive immunity to tolerable

immunity. During initial infection, acute-stage fierce cytokines can rapidly kill

pathogens and infected cells or tissues. However, if the pathogen infects numerous

cells in a tissue, such as the liver, killing the infected cells would completely destroy

the organ [44]. Thus, a regulatory T cell STAT5 signal combined with

TH1/TH2/TH22/THαβ would allow the generation of CD4 T cells with less fierce

cytokines [43]. TH1-like/TH9/TH17/TH3 immunological pathways are generated during

chronic infection. IgA1 and IgA2 are the two types of IgA antibodies. IgA1 is the

dominant IgA antibody in serum, whereas IgA2 is the dominant IgA in mucosa. TGFβ

can induce IgA1 or IgA2 depending on the lymphoid follicle location [45]. In GULTs or

Peyer's patch, IgA2 is the dominant IgA antibody produced in the GI mucosa. In the

lymph nodes of other body locations, IgA1 is the dominant IgA antibody produced.

However, IgA1 is especially related to viral protein antigens, whereas IgA2 is

especially related to bacterial antigens, such as LPS. The heavy-chain locus sequence

of B-cell antibodies is IgM, IgD, IgG3, IgG1, IgA1, IgG2, IgG4, IgE, and IgA2. B-

cells double-express IgM and IgD. IgG3, IgG1, and IgA1 comprise the first group for

cellular immunity. IgG2, IgG4, IgE, and IgA2 can be viewed as the second group for

humoral immunity. The gene sequence order is important, and it affects the time

sequence of the isotype switch. IL-13 is a profibrogenic Treg-related cytokine related

to TGFβ signaling.

Tolerable immune responses

TH1-like cells (non-classic TH1) are initiated by TGFβ (STAT5 signaling) and

IFNg (STAT4 signaling). TH1-like cells with Foxp3 + regulatory character have been

identified [7]. TH1 helper cells and TH1-like cells are closely related [46]. TH1-like cells

are chronic host immunity of the TH1 immune response. Thus, these cells could be

related to chronic inflammation, such as long-term tuberculosis infection. The effector

cells of TH1-like immunity include suppressive macrophages (M2), ILC1,

suppressive CD8 T cells (CD28-CD8+Treg), IgA1-producing B-cells, regulatory

iNKT cells, and IFNg/TGFβ-producing CD4 T cells [23,33,47]. TH1-like immunity

induces type 4 delayed-type hypersensitivity, such as Crohn's disease.

TH9 cells are driven by IL-4 (STAT6 signaling) combined with TGFβ (STAT5

signaling) [48-50]. Thus, TH9 cells are closely related to the TH2 immunological

pathway. The cells are characterized by the IL-9-secreting CD4 T cell. TH9 cells are

important under a chronic allergic condition, such as asthma. Thus, TH9 helper cells

are chronic T helper cells related to TH2 immunity. The effector cells of TH9

immunity include regulatory eosinophils, basophils/mucosal mast cells (MCt, mast

cell tryptase), ILC2, IL-9-producing CD4 T cells, regulatory iNKT cells, and IgA2-

producing B-cells [33,51]. TH9 immunity induces type1 allergy, including asthma [26].

TH17 cells are driven by IL-6/IL-1 combined with TGFβ [5]. Thus, TH17 cells are

closely related to the TH22 immunological pathway. TH17 cells are characterized by

the IL-17-secreting CD4 T cell. TH17 cells are important in chronic immune-

complex-mediated diseases, such as rheumatic arthritis. The TH17 helper cell is the

chronic T helper cell related to TH22 immunity. TGFβ with STAT5 can suppress the

acute IL-22-producing cells and enhance the chronic IL-17-producing cells [34].

Because of the role of TGFβ in TH17 immunity, regulatory IL-17-producing cells are

noted. The effector cells of TH17 immunity include regulatory neutrophils (N2), ILC3(NCR-), IL-17-producing CD4 T cells, regulatory iNKT cells, and IgA2-producing B-cells [33,52]. TH17 immunity induces type 3 immune-complex hypersensitivity, including ulcerative colitis.

TH3 cells are driven by IL-10 and TGFβ [53,54]. Thus, TH3 cells are closely related to the THαβ immunological pathway. These cells also produce IL-10 and TGFβ. Thus, TH3 helper cells are important to chronic antibody-dependent cellular cytotoxic hypersensitivity. TH3 cells are the chronic helper T cells corresponding to the THαβ helper cells. The TH3 immune effector cells include IL-13-producing regulatory NK cells (CD56 + CD16−NK2 cells), ILC10, IL-10- and TGFβ-secreting CD4 T cells, suppressive CD8 T cells (CD28-CD8+ Treg), regulatory iNKT cells, and IgA1-producing B-cells [29,30,47,55,56]. IgA1 is produced in serum and works against viral protein antigens. TH3 immunity induces type 2 antibody-dependent cytotoxic hypersensitivity, including a chronic stage of SLE.

Concluding remarks

The summary diagram includes complete pictures of the $4 \times 2 + 2$ immunological pathways (Table 1). The TH1, TH2, TH22, and THαβ eradicable immune responses correspond to the TH1-like, TH9, TH17, and TH3 tolerable

immune responses, respectively, and match the four types of hypersensitivity. The detailed immune response against different pathogens and allergy/hypersensitivity can be clearly understood.

Authorship

The author, Wan-Chung Hu, fully contributes to the concept, literate search, writing, and final approval of this manuscript.

Acknowledgment

The author is very thankful for the instructions provided by Professors Alan Scott, Louis August Bourgeois, and Pien-Chien Huang during his PhD study on vaccine science from the Department of International Health, Johns Hopkins University, Bloomberg School of Public Health. The author is also very grateful for the guidance of Professors Chi-Huey Wong and Alice Lin-Tsing Yu in his postdoctorate research in the Genomics Research Center of Academia Sinica, Taiwan.

Conflict of Interest Disclosure

The author declare there is no conflict of interest

Author's information

The author, Wan-Chung Hu, graduated as a MD from National Taiwan University. He then completed his PhD in vaccine science from the Department of International Health, Johns Hopkins University, Bloomberg School of Public Health. The author's PhD thesis was on host immune reaction against malarial infection. He conducted a postdoctorate study on cancer immunotherapy at the Genomics Research Center, Academia Sinica, Taiwan. The author completed his PGY training from Mackay Memorial Hospital and Shin-Kong Memorial Hospital in Taiwan. He is currently working as a chief resident in the Department of Clinical Pathology, Far Eastern Memorial Hospital, Taiwan (R.O.C.), for resident training.

References

1.	Mosmann TR, Cherwinski H, Bond MW, Giedlin MA, Coffman RL. Two types of murine helper T cell clone. I. Definition according to profiles of lymphokine activities and secreted proteins. *J Immunol* 1986; 136: 2348-57

2.	Hu WC. Human immune responses to Plasmodium falciparum infection: molecular evidence for a suboptimal THalphabeta and TH17 bias over ideal and effective traditional TH1 immune response. *Malar J* 2013; 12: 392

3.	Kumar S, Naqvi RA, Khanna N, Pathak P, Rao DN. Th3 immune responses in the progression of leprosy via molecular cross-talks of TGF-beta, CTLA-4 and Cbl-b. *Clin Immunol* 2011; 141: 133-42

4.	Levings MK, Sangregorio R, Galbiati F, Squadrone S, de Waal Malefyt R, Roncarolo MG. IFN-alpha and IL-10 induce the differentiation of human type 1 T regulatory cells. *J Immunol* 2001; 166: 5530-9

5.	Harrington LE, Hatton RD, Mangan PR, Turner H, Murphy TL, Murphy KM, et al. Interleukin 17-producing CD4+ effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nat Immunol* 2005; 6: 1123-32

6.	Eyerich S, Eyerich K, Pennino D, Carbone T, Nasorri F, Pallotta S, et al. Th22 cells represent a distinct human T cell subset involved in epidermal immunity and remodeling. *J Clin Invest* 2009; 119: 3573-85

7.	Dominguez-Villar M, Baecher-Allan CM, Hafler DA. Identification of T helper type 1-like, Foxp3+ regulatory T cells in human autoimmune disease. *Nat Med* 2011; 17: 673-5

8.	Luthje K, Kallies A, Shimohakamada Y, Belz GT, Light A, Tarlinton DM, et al. The development and fate of follicular helper T cells defined by an IL-21 reporter mouse. *Nat Immunol* 2012; 13: 491-8

9.	Collin M, McGovern N, Haniffa M. Human dendritic cell subsets. *Immunology* 2013; 140: 22-30

10.	Lane PJ, McConnell FM, Withers D, Gaspal F, Saini M, Anderson G. Lymphoid tissue inducer cells: bridges between the ancient innate and the modern adaptive immune systems. *Mucosal Immunol* 2009; 2: 472-7

11.	McCarron MJ, Marie JC. TGF-beta prevents T follicular helper cell accumulation and B cell autoreactivity. *J Clin Invest* 2014; 124: 4375-86

12.	Tran TH, Utama FE, Lin J, Yang N, Sjolund AB, Ryder A, et al. Prolactin inhibits BCL6 expression in breast cancer through a Stat5a-dependent mechanism. *Cancer Res* 2010; 70: 1711-21

13.   Scheeren FA, Naspetti M, Diehl S, Schotte R, Nagasawa M, Wijnands E, et al. STAT5 regulates the self-renewal capacity and differentiation of human memory B cells and controls Bcl-6 expression. *Nat Immunol* 2005; 6: 303-13

14.   Verdier F, Rabionet R, Gouilleux F, Beisenherz-Huss C, Varlet P, Muller O, et al. A sequence of the CIS gene promoter interacts preferentially with two associated STAT5A dimers: a distinct biochemical difference between STAT5A and STAT5B. *Mol Cell Biol* 1998; 18: 5852-60

15.   Nurieva RI, Chung Y, Martinez GJ, Yang XO, Tanaka S, Matskevitch TD, et al. Bcl6 mediates the development of T follicular helper cells. *Science* 2009; 325: 1001-5

16.   Ettinger R, Sims GP, Fairhurst AM, Robbins R, da Silva YS, Spolski R, et al. IL-21 induces differentiation of human naive and memory B cells into antibody-secreting plasma cells. *J Immunol* 2005; 175: 7867-79

17.   Pene J, Gauchat JF, Lecart S, Drouet E, Guglielmi P, Boulay V, et al. Cutting edge: IL-21 is a switch factor for the production of IgG1 and IgG3 by human B cells. *J Immunol* 2004; 172: 5154-7

18.   Gapin L. Development of invariant natural killer T cells. *Curr Opin Immunol* 2016; 39: 68-74

19.   Choi YS, Eto D, Yang JA, Lao C, Crotty S. Cutting edge: STAT1 is required for IL-6-mediated Bcl6 induction for early follicular helper cell differentiation. *J Immunol* 2013; 190: 3049-53

20.   Schaerli P, Loetscher P, Moser B. Cutting edge: induction of follicular homing precedes effector Th cell development. *J Immunol* 2001; 167: 6082-6

21.   Nakayamada S, Kanno Y, Takahashi H, Jankovic D, Lu KT, Johnson TA, et al. Early Th1 cell differentiation is marked by a Tfh cell-like transition. *Immunity* 2011; 35: 919-31

22.   Iezzi G, Boni A, Degl'Innocenti E, Grioni M, Bertilaccio MT, Bellone M. Type 2 cytotoxic T lymphocytes modulate the activity of dendritic cells toward type 2 immune responses. *J Immunol* 2006; 177: 2131-7

23.   Gong D, Shi W, Yi SJ, Chen H, Groffen J, Heisterkamp N. TGFbeta signaling plays a critical role in promoting alternative macrophage activation. *BMC Immunol* 2012; 13: 31

24.   Eberl G, Colonna M, Di Santo JP, McKenzie AN. Innate lymphoid cells. Innate lymphoid cells: a new paradigm in immunology. *Science* 2015; 348: aaa6566

25.   Klechevsky E, Morita R, Liu M, Cao Y, Coquery S, Thompson-Snipes L, et al. Functional specializations of human epidermal Langerhans cells and CD14+ dermal dendritic cells. *Immunity* 2008; 29: 497-510

26.   Mesnil C, Raulier S, Paulissen G, Xiao X, Birrell MA, Pirottin D, et al. Lung-resident eosinophils represent a distinct regulatory eosinophil subset. *J Clin Invest*

2016; 126: 3279-95

27. Toru H, Eguchi M, Matsumoto R, Yanagida M, Yata J, Nakahata T. Interleukin-4 promotes the development of tryptase and chymase double-positive human mast cells accompanied by cell maturation. *Blood* 1998; 91: 187-95

28. Jung U, Foley JE, Erdmann AA, Eckhaus MA, Fowler DH. CD3/CD28-costimulated T1 and T2 subsets: differential in vivo allosensitization generates distinct GVT and GVHD effects. *Blood* 2003; 102: 3439-46

29. Peritt D, Robertson S, Gri G, Showe L, Aste-Amezaga M, Trinchieri G. Differentiation of human NK cells into NK1 and NK2 subsets. *J Immunol* 1998; 161: 5821-4

30. Kim HS, Jang JH, Lee MB, Jung ID, Kim YM, Park YM, et al. A novel IL-10-producing innate lymphoid cells (ILC10) in a contact hypersensitivity mouse model. *BMB Rep* 2016; 49: 293-6

31. Nguyen KB, Cousens LP, Doughty LA, Pien GC, Durbin JE, Biron CA. Interferon alpha/beta-mediated inhibition and promotion of interferon gamma: STAT1 resolves a paradox. *Nat Immunol* 2000; 1: 70-6

32. Trifari S, Kaplan CD, Tran EH, Crellin NK, Spits H. Identification of a human helper T cell population that has abundant production of interleukin 22 and is distinct from T(H)-17, T(H)1 and T(H)2 cells. *Nat Immunol* 2009; 10: 864-71

33. Walker JA, Barlow JL, McKenzie AN. Innate lymphoid cells--how did we miss them? *Nat Rev Immunol* 2013; 13: 75-87

34. Rutz S, Noubade R, Eidenschenk C, Ota N, Zeng W, Zheng Y, et al. Transcription factor c-Maf mediates the TGF-beta-dependent suppression of IL-22 production in T(H)17 cells. *Nat Immunol* 2011; 12: 1238-45

35. Hussain R, Dawood G, Abrar N, Toossi Z, Minai A, Dojki M, et al. Selective increases in antibody isotypes and immunoglobulin G subclass responses to secreted antigens in tuberculosis patients and healthy household contacts of the patients. *Clin Diagn Lab Immunol* 1995; 2: 726-32

36. Coutelier JP, van der Logt JT, Heessen FW, Vink A, van Snick J. Virally induced modulation of murine IgG antibody subclasses. *J Exp Med* 1988; 168: 2373-8

37. Siber GR, Schur PH, Aisenberg AC, Weitzman SA, Schiffman G. Correlation between serum IgG-2 concentrations and the antibody response to bacterial polysaccharide antigens. *N Engl J Med* 1980; 303: 178-82

38. Turner JD, Faulkner H, Kamgno J, Kennedy MW, Behnke J, Boussinesq M, et al. Allergen-specific IgE and IgG4 are markers of resistance and susceptibility in a human intestinal nematode infection. *Microbes Infect* 2005; 7: 990-6

39. Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science* 2003; 299: 1057-61

40.   Schmidt SV, Nino-Castro AC, Schultze JL. Regulatory dendritic cells: there is more than just immune activation. *Front Immunol* 2012; 3: 274

41.   Wang S, Xia P, Chen Y, Qu Y, Xiong Z, Ye B, et al. Regulatory Innate Lymphoid Cells Control Innate Intestinal Inflammation. *Cell* 2017; 171: 201-216.e18

42.   Monteiro M, Almeida CF, Caridade M, Ribot JC, Duarte J, Agua-Doce A, et al. Identification of regulatory Foxp3+ invariant NKT cells induced by TGF-beta. *J Immunol* 2010; 185: 2157-63

43.   Yao Z, Kanno Y, Kerenyi M, Stephens G, Durant L, Watford WT, et al. Nonredundant roles for Stat5a/b in directly regulating Foxp3. *Blood* 2007; 109: 4368-75

44.   Wei HX, Chuang YH, Li B, Wei H, Sun R, Moritoki Y, et al. CD4+ CD25+ Foxp3+ regulatory T cells protect against T cell-mediated fulminant hepatitis in a TGF-beta-dependent manner in mice. *J Immunol* 2008; 181: 7221-9

45.   Zan H, Cerutti A, Dramitinos P, Schaffer A, Casali P. CD40 engagement triggers switching to IgA1 and IgA2 in human B cells through induction of endogenous TGF-beta: evidence for TGF-beta but not IL-10-dependent direct S mu-->S alpha and sequential S mu-->S gamma, S gamma-->S alpha DNA recombination. *J Immunol* 1998; 161: 5217-25

46.   Prochazkova J, Pokorna K, Holan V. IL-12 inhibits the TGF-beta-dependent T cell developmental programs and skews the TGF-beta-induced differentiation into a Th1-like direction. *Immunobiology* 2012; 217: 74-82

47.   Romero P, Zippelius A, Kurth I, Pittet MJ, Touvrey C, Iancu EM, et al. Four functionally distinct populations of human effector-memory CD8+ T lymphocytes. *J Immunol* 2007; 178: 4112-9

48.   Gerlach K, Hwang Y, Nikolaev A, Atreya R, Dornhoff H, Steiner S, et al. TH9 cells that express the transcription factor PU.1 drive T cell-mediated colitis via IL-9 receptor signaling in intestinal epithelial cells. *Nat Immunol* 2014; 15: 676-86

49.   Dardalhon V, Awasthi A, Kwon H, Galileos G, Gao W, Sobel RA, et al. IL-4 inhibits TGF-beta-induced Foxp3+ T cells and, together with TGF-beta, generates IL-9+ IL-10+ Foxp3(-) effector T cells. *Nat Immunol* 2008; 9: 1347-55

50.   Goswami R, Jabeen R, Yagi R, Pham D, Zhu J, Goenka S, et al. STAT6-dependent regulation of Th9 development. *J Immunol* 2012; 188: 968-75

51.   Westerberg CM, Ulleras E, Nilsson G. Differentiation of mast cell subpopulations from mouse embryonic stem cells. *J Immunol Methods* 2012; 382: 160-6

52.   Fridlender ZG, Sun J, Kim S, Kapoor V, Cheng G, Ling L, et al. Polarization of tumor-associated neutrophil phenotype by TGF-beta: "N1" versus "N2" TAN. *Cancer Cell* 2009; 16: 183-94

53.   Chen ZM, O'Shaughnessy MJ, Gramaglia I, Panoskaltsis-Mortari A, Murphy WJ,

Narula S, et al. IL-10 and TGF-beta induce alloreactive CD4+CD25- T cells to acquire regulatory cell function. *Blood* 2003; 101: 5076-83

54.   Doetze A, Satoguina J, Burchard G, Rau T, Loliger C, Fleischer B, et al. Antigen-specific cellular hyporesponsiveness in a chronic human helminth infection is mediated by T(h)3/T(r)1-type cytokines IL-10 and transforming growth factor-beta but not by a T(h)1 to T(h)2 shift. *Int Immunol* 2000; 12: 623-30

55.   Tulunay A, Yavuz S, Direskeneli H, Eksioglu-Demiralp E. CD8+CD28-, suppressive T cells in systemic lupus erythematosus. *Lupus* 2008; 17: 630-7

56.   Keskin DB, Allan DS, Rybalov B, Andzelm MM, Stern JN, Kopcow HD, et al. TGFbeta promotes conversion of CD16+ peripheral blood NK cells into CD16- NK cells with similarities to decidual NK cells. *Proc Natl Acad Sci U S A* 2007; 104: 3378-83

Figure legends

Fig 1. Summary diagram of host immunological pathways. In the middle, the Tfh side (follicular help T cell) initiates initiatory immunity and the Treg side (regulatory T cells) initiates regulatory immunity. Eradicable TH1 and tolerable TH1-like (Th1L), eradicable TH2 and tolerable TH9, eradicable TH22 and tolerable TH17, and eradicable THαβ and tolerable TH3 are related along diagonal lines.

Table 1. Summary of host immunological pathways

Immune pathways	Driven cytokine, ILCs, DC	Transcription factor	Effector cells	CD4 T cells	B cells	NKT cells	Pathogen	Autoimmune
Initiatory								
Tfh	IL-21, FDC, LTi	STAT1, STAT3, STAT5B		IL-21 CD4 T cells	IgM/G B cells	iNKTfh	Acute infection	
Eradicable immunities								
TH1	IL-12, ILC1, mDC2	STAT4	Macrophage(M1), CTL(Tc1)	IFNg CD4 T cells	IgG3 B cells	iNKT1	Intracellular bacteria & protozoa	Type 4 DTH
TH2	IL-4, ILC2, LC	STAT6	Eosinophil(iEOS), basophil, mast cells(MCct)	IL-4 IL-5 CD4 T cells	IgG4/IgE B cells	iNKT2	Helminths & insects	Type 1 allergy
TH22	IL-1, mDC1, ILC3 NCR+	STAT3	Neutrophils(N1)	IL-1, TNFα , IL-22 CD4 T cells	IgG2 B cells	iNKT17	Extracellular bacteria & fungi	Type3 Immune complex
THα β	IL-10, pDC, IFNα ,ILC10	STAT1 STAT2	NK cells(NK1), CTL(Tc2)	IL-10 CD4 T cells	IgG1 B cells	iNKT10	viruses	Type 2 ADCC

Immune pathways	Driven cytokine, ILCs	Transcription factor	Effector cells	CD4 T cells	B cells	NKT cells	Pathogen	Hypersensitivities
Regulatory								
Treg	TGFβ , DCreg, ILCreg	STAT5A, STAT5B		TGFβ CD4 T cells	IgA B cells	Regulatory iNKT	Chronic infection	
Tolerable immunities								
TH1like	IL-12, TGFβ ,ILC1	STAT4, STAT5	Macrophage(M2), CD8 Treg	IFNg/TGFβ CD4 T cells	IgA1 B cells	Regulatory iNKT	Intracellular bacteria & protozoa	Type 4 DTH
TH9	IL-4, TGFβ , ILC2	STAT6, STAT5	Eosinophil(rEOS), basophil, mast cells(MCt)	IL-9 CD4 T cells	IgA2 B cells	Regulatory iNKT	Helminths & insects	Type 1 allergy
TH17	IL-6, TGFβ , ILC3 NCR-	STAT3, STAT5	Neutrophils(N2)	IL-17 CD4 T cells	IgA2 B cells	Regulatory iNKT	Extracellular bacteria & fungi	Type3 immune complex
TH3	IL-10, TGFβ , ILC10	STAT1 STAT2 STAT5	NK cells(NK2), CD8 Treg	IL-10/TGFβ CD4 T cells	IgA1 B cells	Regulatory iNKT	viruses	Type 2 ADCC