Chromosomal distribution of cyto-nuclear genes in a dioecious plant with sex

chromosomes

Josh Hough[*†], J. Arvid Ågren[*§], Spencer C.H. Barrett, and Stephen I. Wright

Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON,

Canada, M5S 3B2

[*] These authors contributed equally to this work; corresponding authors

[†] josh.hough@utoronto.ca

[§] arvid.agren@utoronto.ca

Running title: Cyto-nuclear interactions in a dioecious plant

Key words: Cyto-nuclear interactions, co-adaptation, sexual conflict, gene transfer

Word count: 2421

Abstract word count: 245

Number of figures: 1

Abstract

The coordination between nuclear and organellar genes is essential to many aspects of

eukaryotic life, including basic metabolism, energy production, and ultimately,

organismal fitness. Whereas nuclear genes are bi-parentally inherited, mitochondrial and

chloroplast genes are almost exclusively maternally inherited, and this asymmetry may

lead to a bias in the chromosomal distribution of nuclear genes whose products act in the

mitochondria or chloroplasts. In particular, because X-linked genes have a higher

probability of co-transmission with organellar genes (2/3) compared to autosomal genes

(1/2), selection for co-adaptation has been predicted to lead to an over-representation of

nuclear-mitochondrial (N-mt) or nuclear-chloroplast (N-cp) genes on the X chromosome

relative to autosomes. In contrast, the occurrence of sexually antagonistic organellar

mutations might lead to selection for movement of cyto-nuclear genes from the X

chromosome to autosomes to reduce male mutation load. Recent broad-scale comparative

studies of N-mt distributions in animals have found evidence for these hypotheses in

some species, but not others. Here, we use transcriptome sequences to conduct the first

study of the chromosomal distribution of cyto-nuclear interacting genes in a plant species

with sex chromosomes (*Rumex hastatulus*; Polygonaceae). We found no evidence of

under- or over-representation of either N-mt or N-cp genes on the X chromosome, and

thus no support for either the co-adaptation or the sexual-conflict hypothesis. We discuss

how our results from a species with recently evolved sex chromosomes fit into an

emerging picture of the evolutionary forces governing the chromosomal distribution of

N-mt and N-cp genes.

Introduction

The intimate relationships between nuclear and organellar genomes in eukaryotes

represent some of the most striking examples of co-evolved mutualisms (Gillham 1994;

Lane 2005; Aanen et al. 2014). The long co-evolutionary history of nuclear and

mitochondrial genomes is perhaps best illustrated by the finding that the vast majority of

mitochondrial genes in animals have been transferred to the nuclear genome (Adams and

Palmer 2003; Rand et al. 2004; Burt and Trivers 2006). Indeed, animal mitochondria now

encode only a few proteins after having lost the majority of their original genes (Berg and

Kurland 2000; Ridley 2000; Bar-Yaacov et al. 2012). Moreover, almost one fifth of the

Arabidopsis thaliana nuclear genome is of chloroplast origin (Martin 2003), suggesting

that organellar-to-nuclear gene movement has played a crucial role in the evolution of

plant genetic systems.

The evolution of cyto-nuclear interactions and the chromosomal distribution of

the genes involved should be influenced by the contrasting modes of inheritance of

organellar genes (maternal inheritance) and autosomal genes (bi-parental inheritance).

This difference may, for example, result in conflict between nuclear and organellar genes

over sex determination and sex ratio (Cosmides and Tooby 1981; Werren and

Beukeboom 1998), and several mitochondrial genes in plants are known to cause male

sterility (Burt and Trivers 2006; Touzet and Meyer 2014). In systems with XY sex

determination, where males are the heterogametic (XY) and females the homogametic

sex (XX), genes on the X chromosome spend 2/3 of their time in females (Rand et al.

2001) and therefore share a female-biased inheritance pattern relative to Y-linked or

autosomal genes, which may result in inter-genomic co-adaptation and or conflict.

A potential consequence of inter-genomic conflict and co-adaptation between

cyto-nuclear genes and other regions of the genome is a shift in the chromosomal location

of cyto-nuclear genes, either becoming more or less abundant on the X chromosome. This

idea has been explored by several recent studies (Drown et al. 2012; Hill and Johnson

2013; Dean et al. 2014; Rogell et al. 2014), and two main processes have been proposed

to account for the movement of genes to or from the X chromosome. The co-adaptation

hypothesis predicts that the co-transmission of X-linked and organellar genes should

result in selection for the co-adaptation of such genes, thus leading to an over-

representation of N-mt and N-cp genes on the X chromosome relative to autosomes

(Rand et al. 2004; Drown et al. 2012). In contrast, the sexual conflict hypothesis predicts

the opposite chromosomal distribution, with more cyto-nuclear genes occurring on

autosomes to alleviate mutation load in males. To date, empirical evidence for the above

hypotheses are mixed. Drown et al. (2012) used previously published reference genomes

to examine the chromosomal distribution of N-mt genes in 16 vertebrates and found a

strong under-representation of such genes on the X chromosomes relative to autosomes in

14 mammal species, but not in two avian species with ZW sex determining systems.

Dean et al. (2014) included seven additional species in their analysis with independently

derived sex chromosomes, and found that the under-representation of N-mt genes on the

X chromosome was restricted to therian mammals and *Caenorhabditis elegans*.

Here, we use sex-linked and autosomal transcriptome sequences to investigate the

chromosomal distributions of cyto-nuclear interactions in the dioecious annual plant

Rumex hastatulus (Polygonaceae). Examining cyto-nuclear interactions within a plant

species is of interest for several reasons (see Sloan 2014). First, plants carry an additional

maternally inherited organellar genome that is absent in animals, the chloroplast genome.

This provides an opportunity to compare the chromosomal distribution of two

independent kinds of cyto-nuclear interacting genes: nuclear-mitochondrial (N-mt) and

nuclear-chloroplast (N-cp). Second, whereas animal sex chromosomes evolved hundreds

of millions of years ago (180 MYA in mammals and 140 MYA in birds; Cortez et al.

2014), the origin of plant sex chromosomes is a more recent event (Charlesworth 2013).

In *R. hastatulus,* sex chromosomes are thought to have evolved approximately 15-16

MYA (Navajas-Perez et al. 2005) and genes on the Y chromosome show evidence of

degeneration, resulting in a considerable proportion of genes that are hemizygous on the

X chromosome (Hough et al. 2014). *Rumex hastatulus* therefore provides an opportunity

to test whether the early changes involved in sex chromosome evolution have been

associated with a concomitant shift in the chromosomal location of N-mt or N-cp genes.

Moreover, the presence in this system of X-linked genes that have recently become

hemizygous provides an opportunity to compare the chromosomal distributions of X-

linked genes that are hemizygous versus those that have retained Y-linked alleles (X/Y

genes). Hemizygous genes are particularly good candidates for evaluating evidence for

co-adaptation and/or sexual conflict because of their relatively older age (Hough et al.

2014), and because beneficial mutations in such genes are exposed to positive selection

regardless of dominance and may therefore spread more rapidly.

Methods

Gene identification and functional annotation

We used sex-linked and autosomal transcriptome sequence data for *R. hastatulus* reported

in Hough et al. (2014; GenBank Sequence Read Archive accession no. SRP041588), and

obtained three sets of genes with which to test for an over- or under-representation of

nuclear-mitochondrial (N-mt) or nuclear-chloroplast (N-cp) genes. In total our analyses

included 1167 autosomal genes, 624 X-linked genes, and 107 hemizygous X-linked genes

(see Hough et al. 2014; *Methods* and *SI Appendix* for full details regarding the

identification of such genes from transcriptome sequence data). For each gene set, we

queried the sequences translated in all reading frames against the *A. thaliana* protein

database using the BLASTx homology search implemented in Blast2GO (Conesa et al.

2005), with a significance threshold (BLAST ExpectValue) of $1x10^{-3}$, above which

matches were not reported. We limited our searches to the *A. thaliana* protein database

because sequence matches to this database returned more detailed functional information

than is available for most other species in the NCBI plant database. We obtained

BLASTx results for 1073 autosomal genes (90%), 567 X-linked genes (90%), and 95

hemizygous genes (89%). Gene Ontology (GO) terms associated with the hits from

BLASTx queries were then retrieved using the 'Mapping' function in Blast2GO, which

used BLAST accessions to link the queried sequences to functional information stored in

the GO database (The Gene Onotology Consortium 2008). Gene names were retrieved

using NCBI mapping files 'gene info' and 'gene2accession', and GO terms were assigned

to query sequences using the 'Annotation' function with an E-Value-Hit-Filter of $1x10^{-6}$

and an annotation cut off of 55 (default parameters). Finally, we ran InterProScan

(Quevillon et al. 2005) to retrieve sequence domain/motif information and merged the

corresponding annotations with previously identified GO terms. This procedure generated

output files containing GO ID's and functional descriptions for each gene in our data set

(files will be uploaded to GitHub).

Statistical analyses

We used a similar approach to Drown et al. (2012) and Dean et al. (2014) and estimated

the number of N-mt and N-cp genes on the X chromosome and autosomes, and then

compared each of these estimates to an expected number. The expected number of N-mt

genes was obtained by calculating the product of the proportion of all genes in the data

set with mitochondrial annotations (matching GO:0005739) and the number of annotated

genes in a given gene set. The expected numbers of N-cp genes were calculated similarly,

using GO:0009507. We then calculated the ratios of the observed-to-expected numbers

for both N-mt and N-cp genes in each gene set. The observed-to-expected ratio is

expected to equal one when there is no under- or over-representation, and greater than

one when there is an over-representation. We note that, unlike for X-linked genes, we did

not have information regarding the particular chromosome locations for autosomal genes,

and therefore could not obtain the expected numbers of N-mt and N-cp genes per-

autosome as in previous studies (Drown et al. 2012; Dean et al. 2014). The expected

numbers were thus calculated assuming that the set of autosomal genes represented a

random sample of the autosomal chromosomes in this species, which is likely a valid

assumption given that the sequences were obtained using whole transcriptome shotgun

sequencing (Hough et al. 2014). Calculating the expected-to-observed ratios across X-

linked, autosomal, and X-hemizygous genes thus allowed us to determine whether any of

these gene sets contained an under- or over-representation of N-mt and N-cp genes

compared to the expectation based on the proportion of such genes in the full data set. We

tested the significance of over- or under-representation using Fisher's exact tests, and

calculated 95% confidence intervals for the numbers of N-mt or N-cp genes using 10,000

replicate bootstrapped samples. All data analysis was done in R (R Development Core

Team 2013; scripts will be available for download from GitHub).

Results and Discussion

It has been suggested that cyto-nuclear genes may be either over- or under-represented on

the X chromosome compared to autosomes, depending on whether their interactions are

driven by co-adaptation or sexual conflict (Rand et al. 2001; Drown et al. 2012; Hill and

Johnson 2013; Dean et al. 2014; Rogell et al. 2014). We annotated sex-linked and

autosomal transcriptome sequences to test these predictions in the dioecious plant *R.*

hastatulus. We found that neither mitochondria- or chloroplast-interacting nuclear genes

were under- or over-represented on the X chromosome (Fisher's exact test, $P = 0.4947$

and $P = 0.3074$, respectively; Figure 1). This pattern indicates that neither the co-

adaptation nor the sexual conflict hypothesis alone is sufficient to explain the

chromosomal distribution of cyto-nuclear genes in *R. hastatulus*.

There are several factors that are expected to be important in determining cyto-

nuclear gene distributions, and these may explain the lack of bias in *R. hastatulus*. For

example, under both the co-adaptation and sexual conflict hypotheses, the age of the sex

chromosomes will determine the extent to which selection (either for co-adaptation, or

sexual antagonism) has had time to operate, which depends on the rate of gene movement

onto and off of the sex chromosomes. Whereas previous studies of cyto-nuclear genes in

animals have focused almost exclusively on ancient sex chromosome systems (Drown et

al. 2012; Dean et al. 2014; Rogell et al. 2014), our study focused on a dioecious plant

species in which sex chromosomes evolved more recently (~15 MYA; Navajas-Perez et

al. 2005), and many genes likely stopped recombining much more recently (Hough et al.,

2014). The lack of bias in the chromosomal distribution of cyto-nuclear genes may

therefore reflect the recent time scale of sex chromosome evolution rather than the

absence of biased gene movement. The relatively young age of sex chromosomes may

also have played a role in the lack of bias reported in the sex and neo-sex chromosomes

in three-spined stickleback, which evolved ~10 MYA (Kondo et al. 2004) and ~2 MYA,

respectively (Natri et al. 2013). Comparative studies of sex chromosomes of different age

will be central for understanding the rate at which organellar gene movement occurs.

In addition to being evolutionarily older, X-linked hemizygous genes are expected

to show a greater effect of over-or under-representation than genes with both X- and Y-

alleles because recessive mutations (involved in either co-adaptation or sexual conflict)

will be exposed to selection instead of masked by an alternate allele in a heterozygous

genotype. We detected a slightly greater under-representation of X-hemizygous N-mt

genes compared to autosomes or X-genes with retained Y-alleles, but the effect was not

statistically significant ($P = 0.4947$). The opposite pattern was evident for N-cp genes,

which were slightly over-represented on hemizygous genes, but again this effect was not

significant ($P = 0.3074$). A larger sample of hemizygous genes would be required to more

confidently assess whether such genes are in fact more often involved in cyto-nuclear

interactions than other genes on the X chromosome, and to test whether the opposite

pattern for N-mt and N-cp hemizygous genes is a result of a different rate of nuclear gene

transfer between mitochondrial and chloroplast genomes.

Another factor that will affect the chromosomal distribution of cyto-nuclear genes

is the number of N-mt and N-cp genes that were located on the autosome from which the

sex chromosomes evolved. Since the origins of mitochondria and chloroplasts both vastly

predate that of sex chromosomes (1.5-2 BYA compared to < 200 MYA; Dyall et al.

2004; Timmis et al. 2004; Cortez et al. 2014) gene transfer from organellar genomes to

the nuclear genome began long before the evolution of sex chromosomes. A bias in the

chromosomal distribution of cyto-nuclear genes in either direction may therefore arise if

the ancestral autosome was particularly rich or poor in cyto-nuclear genes. Indeed, it is

striking that autosomes in the animal species previously examined exhibited extensive

variation in the relative number of N-mt genes (see Drown et al. 2012 Figure 1 and Dean

et al. 2014 Figure 1 and Figure 2). That the ancestral number of N-mt and N-cp genes is

likely to be important is highlighted by the fact that the majority of genes involved in

mitochondrial DNA and RNA metabolism in *A. thaliana* are found on chromosome III

(Elo et al. 2003). If such a biased autosomal distribution of organellar variation is

representative of the ancestral sex chromosomes, the X chromosome could carry

significantly more N-mt or N-cp genes because of this ancestral gene number rather than

a biased rate of gene movement. Genetic mapping and comparative genomic studies of

genes that have transferred from organellar genomes after the origin of sex chromosomes

may provide a means to control for ancestral differences in gene number and provide a

better test of biases in organellar-nuclear gene movement.

To conclude, we conducted the first investigation of the extent to which co-

adaptation and sexual conflict have shaped the chromosomal distribution cyto-nuclear

genes in a plant species with sex chromosomes. We found no sign of under- or over-

representation of either N-mt or N-cp genes on the X chromosome, implying that neither

co-adaptation nor sexual conflict alone can explain the chromosomal distributions of

these genes. Instead, we suggest that additional factors, including the age of sex

chromosomes and the time since X-Y recombination became suppressed, are likely to

have been important determinants of the patterns we observed. To determine whether the

under-representation of mito-nuclear genes on the X chromosome previously reported in

therian mammals and *C. elegans* (Drown et al. 2012; Dean et al. 2014) is indeed due to

sexual conflict, rather than neutral processes such as the number of mito-nuclear genes on

ancestral sex chromosomes, future studies should focus on quantifying rates of gene

movement after sex chromosome origination. Cyto-nuclear conflict and co-evolution has

undoubtedly played a major role in many aspects of genome evolution in both plant and

animal systems; however, it remains unclear whether these processes have also shaped

the chromosomal distribution cyto-nuclear genes.

Acknowledgements

We thank Rebecca Dean and Devin M Drown for comments on the manuscript. This

research was supported by Discovery Grants to SCHB and SIW from the Natural

Sciences and Engineering Council of Canada. JH was supported by an Ontario Graduate

Fellowship and JAÅ by a Junior Fellowship from Massey College.

References

Aanen DK, Spelbrink JN, Beekman M. 2014. What cost mitochondria? The maintenance

of functional mitochondrial DNA within and across generations. Philos Trans R

Soc Lond B Biol Sci. 369.

Adams KL, Palmer JD. 2003. Evolution of mitochondrial gene content: gene loss and

transfer to the nucleus. Mol Phylogenet Evol. 29:380–395.

Bar-Yaacov D, Blumberg A, Mishmar D. 2012. Mitochondrial-nuclear co-evolution and

its effects on OXPHOS activity and regulation. Biochim Biophys Acta 1819:1107–

1111.

Berg OG, Kurland CG. 2000. Why mitochondrial genes are most often found in nuclei.

Mol Biol Evol. 17:951–961.

Burt A, Trivers R. 2006. Genes in conflict: the biology of selfish genetic elements.

Cambridge, MA: Belknap Press of Harvard University Press.

Charlesworth D. 2013. Plant sex chromosome evolution. J Exp Bot. 64:405–420.

Conesa A, et al. 2005. Blast2GO: a universal tool for annotation, visualization and

analysis in functional genomics research. Bioinformatics 21:3674–3676.

Cortez D, et al. 2014. Origins and functional evolution of Y chromosomes across

mammals. Nature 508:488–493.

Cosmides LM, Tooby J. 1981. Cytoplasmic inheritance and intragenomic conflict. J

Theor Biol. 89:83–129.

Dean R, Zimmer F, Mank JE. 2014. The potential role of sexual conflict and sexual

selection in shaping the genomic distribution of mito-nuclear genes. Genome Biol

Evol. 6:1096–1104.

Drown DM, Preuss KM, Wade MJ. 2012. Evidence of a paucity of genes that interact

with the mitochondrion on the X in mammals. Genome Biol Evol. 4:763–768.

Dyall SD, Brown MT, Johnson PJ. 2004. Ancient invasions: from endosymbionts to

organelles. Science 304:253–257.

Elo A, Lyznik A, Gonzalez DO, Kachman SD, Mackenzie SA. 2003. Nuclear genes that

encode mitochondrial proteins for DNA and RNA metabolism are clustered in the

Arabidopsis genome. Plant Cell 15:1619–1631.

Gillham NW. 1994. Organelle genes and genomes. New York: Oxford University Press.

Hill GE, Johnson JD. 2013. The mitonuclear compatibility hypothesis of sexual selection.

Proc R Soc B. 280:20131314.

Hough J, Hollister JD, Wang W, Barrett SCH, Wright SI. 2014. Genetic degeneration of

old and young Y chromosomes in the flowering plant *Rumex hastatulus*. Proc Natl

Acad Sci U S A. 111:7713–7718

Kondo M, Nanda I, Hornung U, Schmid M, Schartl M. 2004. Evolutionary origin of the

medaka Y chromosome. Curr Biol. 14:1664–1669.

Lane N. 2005. Power, sex, suicide: mitochondria and the meaning of life. Oxford, UK:

Oxford University Press.

Martin W. 2003. Gene transfer from organelles to the nucleus: frequent and in big chunks.

Proc Natl Acad Sci U S A. 100:8612–8614.

Natri HM, Shikano T, Merilä J. 2013. Progressive recombination suppression and

differentiation in recently evolved neo-sex chromosomes. Mol Biol Evol. 30:1131–

1144.

Navajas-Perez R, et al. 2005. The evolution of reproductive systems and sex-determining

mechanisms within *Rumex* (Polygonaceae) inferred from nuclear and

chloroplastidial sequence data. Mol Biol Evol. 22:1929–1939.

Quevillon E, et al. 2005. InterProScan: protein domains identifier. Nucleic Acids Res. 33:

W116–120.

R Development Core Team. 2013. R: A language and environment for statistical

computing. Vienna, Austria: R Foundation for Statistical Computing.

Rand DM, Clark AG, Kann LM. 2001. Sexually antagonistic cytonuclear fitness

interactions in *Drosophila melanogaster*. Genetics 159:173–187.

Rand DM, Haney RA, Fry AJ. 2004. Cytonuclear coevolution: the genomics of

cooperation. Trends Ecol Evol. 19:645–653.

Ridley M. 2000. Mendel's demon: gene justice and the complexity of life. London:

Weidenfeld & Nicolson.

Rogell B, Dean R, Lemos B, Dowling DK. 2014. Mito-nuclear interactions as drivers of

gene movement on and off the X chromosome. BMC Genomics 15:330.

Sloan DB. 2014. Using plants to elucidate the mechanisms of cytonuclear co-evolution.

New Phytol. doi: 10.1111/nph.12835

The Gene Ontology Consortiun. 2008. The Gene Ontology project in 2008. Nucleic

Acids Res. 36:D440–444.

Timmis JN, Ayliffe MA, Huang CY, Martin W. 2004. Endosymbiotic gene transfer:

organelle genomes forge eukaryotic chromosomes. Nat Rev Genet. 5:123–135.

Touzet P, Meyer EH. 2014. Cytoplasmic male sterility and mitochondrial metabolism in

plants. Mitochondrion doi: 10.1016/j.mito.2014.04.009

Werren JH, Beukeboom LW. 1998. Sex determination, sex ratios, and genetic conflict.

Annu Rev Ecol Syst. 29:233–261.

Figure 1. Representation of the chromosomal location of cyto-nuclear genes in *Rumex*

hastatulus. Dots represent the observed to expected ratio of mito-nuclear (N-mt) and

chloro-nuclear (N-cp) genes on autosomes, the X chromosome, and hemizygous X genes,

with the 95% confidence intervals estimated by bootstrapping (10,000 replicates). The

vertical dotted line at 1 represents no over- or under-representation.