THE IMPACTS OF HISTORICAL BARRIERS ON FLORISTIC PATTERNS OF

PLANT GROUPS WITH DIFFERENT DISPERSAL ABILITIES IN THE RYUKYU

ARCHIPELAGO, JAPAN

Koh Nakamura

21 COE Program, Graduate School of Engineering and Science, University of the Ryukyus,

Senbaru 1, Nishihara, Okinawa 903-0213, Japan

Email: h081743@sci.u-ryukyu.ac.jp (Corresponding author)

Rempei Suwa, Tetsuo Denda, and **Masatsugu Yokota**

Laboratory of Ecology and Systematics, Faculty of Science, University of the Ryukyus, Senbaru

1, Nishihara, Okinawa 903-0213, Japan

ABSTRACT. - The effects of historical barriers in biogeographical patterns are expected to

persist differently depending on dispersal abilities of organisms. We tested two hypotheses that

plant groups with different dispersal abilities display different floristic patterns, and that

historical barriers can explain floristic differentiation patterns in plants with low dispersal

ability but not in plants with higher dispersal ability, in the seed plant flora of the Ryukyu

Archipelago. This area is biogeographically interesting because several similar floristic

differentiation patterns have been proposed, all of which are primarily explained by two

historical barriers, the Tokara Tectonic Strait (Tokara Gap) and the Kerama Gap, which arose

during the formation of the islands. We calculated floristic dissimilarity distance among 26

islands based on data sets for three dispersal-ability classes. Clustering analyses based on the

floristic dissimilarity distance generated similar floristic patterns regardless of dispersal-ability

class. We propose that because the landscape resistance is so strong that migration of plants is

severely restricted regardless of their dispersal abilities, the similar floristic differentiation

patterns are generated. Multivariate regression analyses using Mantel's randomization test

indicated that floristic differentiations among islands were explained by the both effects of the

historical barriers and geographic distance in all dispersal-ability classes. Significance of the

historical barriers is not determined by the plant dispersal abilities but presumably by the spatial

distribution of the islands, stochastic dispersals, and time since the formation of the barriers.

INTRODUCTION

The interactions between historical and current processes that shape biogeographic patterns are

a key component of biogeographical studies (e.g., Candolle, 1820; Good, 1974; Brown &

Gibson, 1983; Cox & Moore, 1993; Crisci et al., 2003). In historical biogeography,

biogeographic patterns are explained based on past processes that have persisted over very long

periods of time, such as geological barriers formed by tectonic movements over millions of

years (Crisci et al., 2003). However, current processes such as ongoing dispersals can affect or

obscure evidence of historical patterns with migration across historical barriers (Wolf et al.,

2001; Giller et al., 2004). Therefore, historical barriers and current dispersals compete in

shaping biogeographic patterns; however, their relative importance is expected to vary

depending on abiotic and biotic factors such as the spatial configuration of habitats and the

dispersal ability of organisms. Increased knowledge of the shifting relative importance in

relation to these factors would enhance our understanding of processes that structure

biogeographic patterns. The Ryukyu Archipelago is a suitable study area for such studies.

The Ryukyu Archipelago is an assemblage of continental islands located between Kyushu

Island of Japan and Taiwan (Fig. 1). This area is biogeographically interesting because several

similar floristic differentiation patterns have been proposed, all of which are primarily

explained by two historical barriers, the Tokara Tectonic Strait (Tokara Gap) and the Kerama

Gap, which arose during the formation of the islands (Hara, 1959; Good, 1974; Maekawa,

1974; Shimabuku, 1984; Takhtajan, 1986; Kitamura et al., 1994). However, it is expected that

plant groups whose seeds/fruits exhibit differing dispersal abilities display different floristic

patterns because the effects of the historical barriers are dependent on dispersal ability. That is,

a lower dispersal ability results in a greater amount of time that the historical effects are able to

persist, whereas a higher dispersal ability results in fewer historical effects (Nekola & White,

1999). Therefore, it is thought that historical barriers can explain floristic differentiation

patterns in plants with low dispersal ability, but not in plants with higher dispersal ability. In the

latter case, it is expected that without an effective barrier to dispersal, the floristic

differentiation would be correlated to the geographic distance between the islands, and that the

floristic pattern is explained by this spatial effect.

To test this hypothesis, we compiled the distribution records of seed plants and grouped them

into three dispersal-ability classes for each island of the Ryukyu Archipelago to examine the

floristic patterns for each dispersal-ability class. We then used a multivariate approach to

examine how the significance of the effects of the two historical barriers and the geographic

distance differ with various dispersal-ability classes.

MATERIAL AND METHODS

Study area

The Ryukyu Archipelago is thought to have undergone extensive changes in land configuration

during the Cenozoic, and there was more than one period of landbridge connection between the

islands and surrounding landmasses (Kyushu to the north, and southeastern China via Taiwan to

the south). This enhanced the range expansion of various lineages of terrestrial organisms in

this region (Hatusima, 1975; Ota, 1998; Chiang & Schaal, 2006). It is proposed that by the

Pliocene or early Pleistocene, the landbridge was segmented at two areas called the Tokara Gap

and the Kerama Gap, where the sea is currently more than 1000 m deep (Kawana, 2002), as a

result of tectonic movements and eustatic sea-level rise. It is thought that the islands have not

since been connected across the gaps, even during the Quaternary glacial sea-level minima (Ota,

1998). Given this geohistory, many reports have demarcated the flora at one or both of the gaps

(e.g., Hara, 1959; Good, 1974; Maekawa, 1974; Shimabuku, 1984; Takhtajan, 1986; Kitamura

et al., 1994). For example, Takhtajan (1986) placed the demarcation line of the

Japanese–Korean and Ryukyu provinces and that of the Ryukyu and Taiwanian provinces at the

Tokara and Kerama Gaps, respectively. In addition, Good (1974) and Kitamura et al. (1994)

regarded the Tokara Gap as the demarcation between the Sino–Japanese Region of the Boreal

Kingdom and the Continental Southeast Asiatic Region of the Paleotropical Kingdom.

The spatial arrangement of the 26 islands examined in this study is shown in Fig. 1. These

islands were selected because of the availability of reliable lists of seed plant flora. Based on the

two gaps, the islands of the Ryukyu Archipelago are divided into three groups: the islands 1 to

12, 13 to 22, and 23 to 26 belong to the northern, central, and southern Ryukyus, respectively.

The Ryukyu Archipelago is in the subtropics, or the transition from tropical to warm temperate

zones (Good, 1974). The climate is moderate throughout the year with a mean temperature of

approximately 15°C during the winter and 28°C in summer, and annual precipitation exceeds

2000 mm with no dry season. Therefore, the islands are covered in well-developed

broad-leaved evergreen forests (Maekawa, 1974).

10   **Floristic data editing**

The vascular plant flora of the Ryukyu Archipelago comprises approximately 1800 species

(Hatusima & Nackejima, 1979); however, detailed records of each species for each island have

not been compiled in one source. We referred to six literature sources for distribution records of

native plants on the 26 islands (Hatusima & Amano, 1974, 1994; Hatusima et al., 1975; Niiro &

Shinjo, 1989; Hatusima, 1991; Kawakubo & Tagawa, 1991). We compiled data sets for three

dispersal-ability classes: high (269 wind-dispersed species with dust seeds and fruits having

hairy pappi), low (200 gravity-dispersed species mostly with acorns and dehiscent fruits), and

intermediate (224 bird-dispersed species mostly with sap-fruits). Each of the three classes

included a similar number of species. Estimation of dispersal abilities was based on dispersal

vectors estimated on the basis of seed/fruit morphology (e.g., Brown & Gibson, 1983).

Infraspecific taxa were not separated to avoid taxonomic incongruence among the literature

sources and to avoid giving intra- and inter-specific differences equal weighting.

For each dispersal-ability class, the pairwise floristic similarity between islands was calculated

using Simpson's similarity index (SI); SI $= \gamma \beta^{-1}$, $(\alpha > \beta)$, where γ is the number of species

shared between two islands, and α and β are the numbers of species on each of the two islands

(Simpson, 1943). SI is one of the most frequently used similarity indices in biogeographic

analyses (Brown & Gibson, 1983), and we selected this index because it does not give an

improperly low similarity value when the areas (and hence the numbers of species) being

compared differ greatly (Balgooy, 1971). Floristic dissimilarity distance was calculated by

subtracting the value of SI from 1.

9   **Historical and spatial variables**

The historical effect of the Tokara Gap was expressed as a categorical matrix using dummy

variables; a value of 1 was given to matrix elements comparing two islands crossing over the

Tokara Gap, and a value of 0 was given to matrix elements comparing two islands on either side

of the Tokara Gap. The matrix of the Kerama Gap was constructed in the same manner. This is

because the proposed dates of the formation of the gaps are very approximate (Ota, 1998),

meaning that we cannot use these dates as accurate and reliable variables. Pairwise geographic

distance (km) among the islands was calculated based on the geological coordinates at the

highest points of each island (range = 11.3–1042.0 km; average distance = 353.2 km; SD =

284.0 km).

21   **Data analyses**

To illustrate the floristic biogeographic pattern in each dispersal-ability class, clustering

analyses of the islands were conducted based on floristic dissimilarity distances using the

unweighted pair group method with arithmetic mean (UPGMA; Sneath & Sokal, 1973) with

PAUP* 4.0b10 (Swofford, 2002).

To examine the significance of the effects of the Tokara and Kerama Gaps and the inter-island

geographic distance on the variation in the floristic dissimilarity distance, we conducted partial

Mantel tests. A Mantel test is used to test the statistical significance of correlations of distance

matrices, where matrix elements are not independent, by randomization (Mantel, 1967; Smouse

et al., 1986; Legendre et al., 1994). Partial Mantel tests are used to conduct partial linear

regression analyses using a response matrix and two explanatory matrices to test the correlation

between the response variable and one of the explanatory variables while controlling for the

effect of the other explanatory variable to remove spurious correlation (Legendre, 2000; Bonnet

& Peer, 2002; Tuomisto et al., 2003; Legendre et al., 2005; Lichstein, 2007). We conducted

partial Mantel tests of the floristic dissimilarity distance with two pairs of explanatory variables,

Tokara Gap and geographic distance, and Kerama Gap and geographic distance, using the

software program 'zt' with 10000 randomizations (Bonnet & Peer, 2002).

To estimate the geographic extent of the significant correlation between the floristic

dissimilarity distance and the geographic distance, we calculated a correlation coefficient r for

given distance classes using GeneAlEx 6 software (Peakall & Smouse, 2006). Distance class

sizes were determined to include a similar number of pairwise comparisons (32 or 33 island

pairs) within each distance class. The calculated correlation coefficients were plotted as a

function of geographic distance. Statistical significance of r was tested by 1000 randomizations

for the null hypothesis of no correlation.

25   **RESULTS**

Floristic distance and clustering of islands

Mean values of the pairwise floristic dissimilarity distances were 0.240 (range = 0–0.576) in

high, 0.187 (0–0.485) in intermediate, and 0.260 (0–0.641) in low dispersal-ability classes.

UPGMA clusterings in three dispersal-ability classes are shown in Fig. 2. Seen in a broad

perspective, the clustering patterns were similar and approximately congruent with the

geographic arrangement of the islands in the archipelago. In all three classes, the islands of each

of the northern, central, and southern Ryukyus were separated, with some exceptional small

islands (islands 13–15, 26). Among the three regions, the islands of the central and southern

Ryukyus were more closely connected to each other than to the islands of the northern Ryukyus.

Regression analyses

The results of the partial Mantel tests are shown in Table 1. In the high-dispersal ability class,

the partial regression of the floristic distance matrix on the matrix of the Tokara Gap,

controlling the effect of the geographic distance, was positive and significant ($r = 0.31$, $p =$

0.0001; in partial Mantel tests of this study, significance level 5% with Bonferroni adjustment

was $p = 0.05/12 = 0.0042$), and so was the partial regression of the floristic distance matrix on

the geographic distance, controlling the effect of the Tokara Gap ($r = 0.31$, $p = 0.0002$). The

partial regression of the floristic distance matrix on the matrix of the Kerama Gap, controlling

the effect of the geographic distance, was negative but not significant ($r = - 0.07$, $p = 0.1876$),

whereas the effect of the geographic distance, controlling the effect of the Kerama Gap, was

significantly positive ($r = 0.29$, $p = 0.0002$). The results for the intermediate and low

dispersal-ability classes were generally in accordance with one another. Specifically, the partial

regression of the floristic distance matrix on the matrix of the Tokara Gap, controlling the effect

of the geographic distance, was significantly positive (intermediate: $r = 0.44$, $p = 0.0002$; low: r

$= 0.18$, $p = 0.0037$) and vice versa (intermediate: $r = 0.38$, $p = 0.0001$; low: $r = 0.49$, $p = 0.0001$).

The partial regression of the floristic distance matrix on the matrix of the Kerama Gap,

controlling the effect of the geographic distance, was significantly negative (intermediate: $r = -$

0.28, $p = 0.0003$; low: $r = - 0.31$, $p = 0.0006$) but the effect of the geographic distance,

controlling the effect of the Kerama Gap, was significantly positive (intermediate: $r = 0.49$, $p =$

0.0001; low: $r = 0.55$, $p = 0.0001$).

Geographic extent of spatial correlation

As shown in Figure 3, the analyses examining the geographic extent of the significant

correlation between the floristic dissimilarity distance and the geographic distance generated

very similar patterns in the three dispersal-ability classes. Specifically, the correlation

coefficients were positive and significant in the first four distance classes (56, 104, 139, 186

km), with an x-intercept of approximately 250 km. The geographic extent of positive

correlation can be influenced by the chosen distance class size (Double et al., 2005); however

changing distance class sizes generated similar patterns (data not shown).

DISCUSSION

Dispersal ability and floristic patterns

The UPGMA analyses generated similar clustering patterns regardless of dispersal-ability class.

This result rejects the hypothesis that plant groups with different dispersal abilities exhibit

different floristic patterns. Biogeographic patterning results from each organism's migration,

which is a function of its dispersal ability and landscape resistance to the movement of the

organism (Nekola & White, 1999). Therefore, the similar floristic patterns observed here are

likely explained from these two viewpoints.

One explanation for our results is that the classification of dispersal ability was not adequate

and there was no significant difference in the potential dispersal distances among the three

classes. It is difficult to accurately estimate the dispersal ability of plants because evidence of

dispersal, especially such a long-distance dispersal as to cross over sea, is not available for most

plant species (Cain, 1974). Also, with respect to long distance dispersal, estimating dispersal

abilities based on seed/fruit morphology might be misleading because the morphological

adaptations of seeds/fruits that are typically used to identify the standard dispersal vector tend

to govern short-distance dispersal but do not necessarily constitute the main mechanism

responsible for long-distance dispersal (Nathan, 2006). Long-distance dispersal can occur by

means of multiple stochastic dispersal vectors, even in the absence of specific adaptations for

each vector (Bullock & Clarke, 2000; Myers et al., 2004). Conversely, some analyses

examining the relationships between plant dispersal abilities and decrease-rate of floristic

similarity with geographic distance have indirectly illustrated that plants with seeds/fruits

adapted to bird-dispersal are more dispersal-limited than plants adapted to wind-dispersal

(Nekola & White, 1999; Tuomisto et al., 2003). Therefore, classification of the plant dispersal

abilities based on the seed/fruit morphology is not necessarily unreasonable. Assuming that our

dispersal-ability classification was accurate, a second explanation for our results must consider

the effect of landscape resistance.

Given that the landscape resistance is common to the analyses of the three dispersal-ability

classes, it is likely responsible for the similar results in our analyses. Landscape resistance is

caused by spatial configuration, particularly size and isolation of habitats. The more isolated

and/or the smaller the habitat, the less effective dispersal will be (Nekola & White, 1999;

Sklenář & Jørgensen, 1999). For example, in North American spruce-fir forests, isolated

habitats restricted to the highest peaks, which are analogous to the habitats of islands, exhibit

much stronger landscape resistance than more contiguous habitats (Nekola & White, 1999). In

the Ryukyu Archipelago, it is very plausible that the isolated distribution of relatively small

islands is causing strong landscape resistance. As shown in Fig. 3, the distance at which the

correlation between floristic dissimilarity distance and geographic distance becomes effectively

zero was approximately 200 km, regardless of the dispersal-ability class. This means that the

geographic extent of the correlation is confined to a comparatively small area while the

archipelago extends for more than 1300 km. We propose that because the landscape resistance

is so strong that migration of plants is severely restricted regardless of their dispersal abilities,

the plant groups whose potential dispersal distances in a contiguous landscape are different

generate similar differentiation patterns.

18   **Significance/insignificance of historical effects**

The partial Mantel tests indicated that the floristic differentiations among islands were

explained by both historical effect and geographic distance in all three dispersal-ability classes.

This result rejects the hypothesis that historical barriers can not explain floristic differentiation

patterns in plants with higher dispersal ability. The UPGMA analyses indicated that the

northern, central, and southern Ryukyus were essentially separated. This separation pattern

seems to be consistent with the hypotheses of the preceding studies, which explain the floristic

differentiation pattern primarily by the historical gaps (Hara, 1959; Good, 1974; Maekawa,

1974; Shimabuku, 1984; Takhtajan, 1986; Kitamura et al., 1994). However, our partial Mantel

tests indicated that geographic distance and the Tokara Gap have exhibited significantly

positive effects but that the effects of the Kerama Gap were insignificant or significantly

negative. The significantly positive effect of the Kerama Gap was spurious. Note that the

geographic distances between islands across the Kerama Gap are much larger than those of the

Tokara Gap (the shortest distances across each gap are 227.6 km between islands 22 and 23 and

36.6 km between islands 12 and 13; Fig. 1). However, the UPGMA analyses indicated that the

islands of the central and southern Ryukyus were more closely connected to each other than to

the islands of the northern Ryukyus. Therefore, the explanation for the insiginificant effect of

the Kerama Gap in high dispersal-ability class is that the floristic dissimilarity distance across

the Kerama Gap was no more that what was explained by the effect of the accompanying large

geographic distance. Similarly, the explanation for the negative effects in the intermediate and

low dispersal-ability classes is that the floristic dissimilarity distance across the gap was less

than what was expected from the large geographic distance, for which the effect of the Kerama

Gap was adjusted to be negative. This is because that although the significantly positive effect

of the Kerama Gap was spurious, the negative effects in intermediate and low dispersal-ability

classes could also be spurious because it is unlikely that the formation of the Kerama Gap

increased the floristic similarity among the islands across the gap, considering that the

landbridge had played a significant role in the migration of the plants (Hatusima, 1975; Chiang

& Schaal, 2006). Thus, the effect of the Kerama Gap would actually be positive but

indistinguishable compared to the effect of the large geographic distance.

When we consider that the Tokara Gap is a very narrow strait (the two closest islands across the

gap are only 36.6 km apart today), the significant effect of the Tokara Gap requires an

explanation as to why ongoing inter-island dispersal has not completely obscured the effect of

the past formation of the gap. The duration of the effects of a historical barrier is determined by

the amount of time required for organisms to cross it (Nekola & White, 1999). The Tokara Gap

is thought to have formed during the Miocene or at the latest in the early Pleistocene, roughly

more than two million years ago (Ota, 1998). A recent quantitatively derived understanding of

the long-distance dispersal of plants may help us understand the persistence of the historical

effects of the Tokara Gap. The expected time for an effective dispersal, i.e., the successful

establishment of one reproductive individual is calculated as the inverse of the product of seed

arrival probability and seed-to-adult survival probability (Clark et al., 1999). For example, with

a hypothetical source population (10^6 individuals, each with an annual fecundity of 10^4 seeds

and with mean dispersal distance of 50 m), the expected time for a single effective dispersal

event to occur beyond 150 km is longer than 100 billion years under the mean trend. Conversely,

an effective long-distance dispersal event beyond 415 km, expected to occur once in almost

10^{13} years under the mean trend, may occur once in 10 years as a result of events that break the

rules, such as hurricanes, tornadoes, and tropical cyclones. Similarly, an effective long-distance

dispersal event beyond 34 km may occur once in almost 10^8 years under the mean trend but

once a year under exceptional conditions (Clark et al., 1999). Long-distance dispersal is an

extremely rare phenomenon even at the scale of a few tens of km or a few hundred km, although

nonstandard mechanisms may help realize it (Nathan, 2006). Although these are rough

estimates for certain types of plants, they still serve as useful references. The time required for

long-distance dispersal to cross the Tokara Gap under the mean trend would be much longer

than the time since the formation of the gap (> 2 million years), although this time would have

allowed some species to cross the gap with nonstandard mechanisms such as summer typhoons.

Thus, it is likely that the historical effects of the Tokara Gap have been preserved because of the

rarity of long-distance dispersal across the gap beyond 36.6 km under the mean trend with the

time being limited. Therefore, the significance of the historical barriers is not determined by the

plant dispersal abilities but presumably by the spatial distribution of the islands, stochastic

dispersals, and time since the gap formation.

ACKNOWLEDGEMENTS

We thank Drs. A. J. Davis (Max Planck Institute for Chemical Ecology), N. Kawakubo (Gifu

University), S. Matsumura (University of the Ryukyus) and H. Ota (University of the Ryukyus)

for their valuable comments. This study was supported by a Research Fellowship for Young

Scientists from the Japan Society for the Promotion of Science (JSPS) to K.N., a Grant-in-Aid

for Scientific Research (C) (no. 17570083) to M.Y., and a grant for the 21st Century COE

program of the University of the Ryukyus.

LITERATURE CITED

Balgooy, M. M. J. van, 1971. Plant geography of the Pacific, based on a census of Phanerogam

  genera. *Blumea*, Supplement, **6**: 1-222.

Bonnet, E. & Y. V. de Peer, 2002. zt: A Software Tool for Simple and Partial Mantel Tests.

  Journal of Statistical Software, **7**: 1-12.

Brown, J. H. & A. C. Gibson, 1983. *Biogeography*. CV Mosby, St. Louis. 643 pp.

Bullock, J. M. & R. T. Clarke, 2000. Long distance seed dispersal by wind: measuring and

  modelling the tail of the curve. *Oecologia*, **124**: 506-521.

Cain, S. A., 1974. *Foundations of plant geography*. Hafner, New York. 556 pp.

Candolle, A. P. de, 1820. Géographie botanique. *Dictionnaire des Sciences Naturelles*: 359-422.

Chiang, T. Y. & B. A. Schaal, 2006. Phylogeography of plants in Taiwan and the Ryukyu Archipelago. *Taxon*, **55**: 31-41.

Clark, J. S., M. Silman, R. Kern, E. Macklin & J. HilleRisLambers, 1999. Seed dispersal near and far: patterns across temperate and tropical forests. *Ecology*, **80**: 1475-1494.

Cox, C. B. & P. D. Moore, 1993. *Biogeography*. Blackwell Science, Cambridge. 326 pp.

Crisci, J. V., L. Katinas & P. Posadas, 2003. *Historical biogeography*. Harvard University Press, Cambridge. 250 pp.

Double, M. C., R. Peakall, N. R. Beck & A. Cockburn, 2005. Dispersal, philopatry, and infidelity: dissecting local genetic structure in superb fairy-wrens (*Malurus cyaneus*). *Evolution*, **59**: 625-635.

Giller, P. S., A. A. Myers & B. R. Riddle, 2004. Earth history, vicariance, and dispersal. In: Lomolino M. V., D. F. Sax & J. H. Brown (eds.), *Foundations of biogeography*. The University of Chicago Press. Pp. 267-276.

Good, R., 1974. *The geography of the flowering plants*. Longman, London. 557 pp.

Hara, H., 1959. An outline of phytogeography of Japan. In: Hara, H. (ed.), *Distribution maps of flowering plants in Japan 2*. Inoue book company. Pp. 1-96.

Hatusima, S., 1975. *Flora of the Ryukyus* (added and corrected). Okinawa association of biology education, Naha. 1002 pp.

Hatusima, S., 1991. *Flora of the northern Ryukyus*. Asahi press publishing co., Kagoshima. 257pp.

Hatusima, S. & T. Amano, 1974. Plants of Kumejima Island. In: Okinawa nature study society (ed.), *Okinawa nature study society research report. Academic research report on the site proposed for Kumejima prefectural Natural Park*. Okinawa Prefecture. Pp. 11-39.

Hatusima, S. & T. Amano, 1994. *Flora of the Ryukyus, south of Amami Island*, 2nd edn. The

biological society of Okinawa, Nishihara. 393 pp.

Hatusima, S. & K. Nackejima, 1979. *Flower of the Ryukyu Islands*. Kohdansha, Tokyo. 368 pp.

Hatusima, S., T. Amano & Y. Miyagi, 1975. Flora of the Miyako Islands, Ryukyus. In: Okinawa

nature study society (ed.), *Okinawa nature study society research report. Academic*

research report on the site proposed for Okinawa prefectural natural park, Miyako Islands.

Okinawa Prefecture. Pp. 31-70.

Kawakubo, N. & H. Tagawa, 1991. Geographic distribution of vascular plants in the Satsunan

Islands, the Nansei Islands. In: Environmental Agency of Japan (ed.), *Study of essential*

factors for preservation of wildlife in Nansei Islands. WWF Japan. Pp. 51-109.

Kawana, T., 2002. Neotectonics of the Ryukyu Arc. In: Kimura, M. (ed.), *The formation of the*

Ryukyu Arc and migration of biota to the arc. Okinawa Times. Pp. 59-83.

Kitamura, S., G. Murata, & M. Hori, 1994. *Colored illustrations of herbaceous plants of Japan*

(Sympetalae). Hoikusha, Osaka. 297 pp.

Legendre, P., 2000. Comparison of permutation methods for the partial correlation and partial

mantel tests. *Journal of statistical computation and simulation*, **67**: 37-73.

Legendre, P., D. Borcard & P. R. Peres-Neto, 2005. Analyzing beta diversity: partitioning the

spatial variation of community composition data. *Ecological Monographs*, **75**: 435-450.

Legendre, P., F. J. Lapointe & P. Casgrain, 1994. Modeling brain evolution from behavior: a

permutational regression approach. *Evolution*, **48**: 1487-1499.

Lichstein, J. W., 2007. Multiple regression on distance matrices: a multivariate spatial analysis

tool. *Plant Ecology*, **188**: 117-131.

Maekawa, F., 1974. Origin and characteristics of Japan's Flora. In: Numata, M. (ed.), *The flora*

and vegetation of Japan. Kodansha scientific. Pp. 33-86.

Mantel, N., 1967. The detection of disease clustering and a generalized regression approach.

Cancer Research, **27**: 209-220.

Myers, J. A., M. Vellend, S. Gardescu & P. L. Marks, 2004. Seed dispersal by white-tailed deer: implications for long-distance dispersal, invasion, and migration of plants in eastern North America. *Oecologia* **139**: 35-44.

Nathan, R., 2006. Long-distance dispersal of plants. *Science* **313**: 786-788.

Nekola, J. C. & P. S. White, 1999. The distance decay of similarity in biogeography and ecology. *Journal of Biogeography*, **26**: 867-878.

Niiro, Y. & K. Shinjo, 1989. Vegetation and flora of Yonagunijima Islands. In: Yonaguni silkworm preservation association (ed.), *Project report on preservation and propagation of Yonaguni silkworm*. Yonaguni silkworm preservation association. Pp. 57-154.

Ota, H., 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications. *Researches on Population Ecology*, **40**: 189-204.

Peakall, R. & P. E. Smouse, 2006. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*, **6**: 288-295.

Shimabuku, K., 1984. Vascular plants. In: Japan association of biology education (ed.), *Natural life forms in Okinawa*. Okinawa association of biology education. Pp. 23-32.

Simpson, G. G., 1943. Mammals and the nature of continents. *American Journal of Science*, **241**: 1-31.

Sklenář, P., & P. M. Jørgensen, 1999. Distribution patterns of páramo plants in Ecuador. *Journal of Biogeography*, **26**: 681-691.

Smouse, P. E., J. C. Long & R. R. Sokal, 1986. Multiple regression and correlation extensions of the Mantel test of matrix correspondence. *Systematic Zoology*, **35**: 627-632.

Sneath, P. H. A. & R. R. Sokal, 1973. *Numerical taxonomy*. W.H. Freeman, San Francisco. 573 pp.

Swofford, D. L., 2002. *PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods)*,

Version 4.0b10. Sinauer Associates, Sunderland.

Takhtajan, A., 1986. Floristic regions of the world. University of California Press, Berkeley.

522 pp.

Tuomisto, H., K. Ruokolainen & M. Yli-Halla, 2003. Dispersal, environment, and floristic

variation of western Amazonian forests. *Science*, **299**: 241-244.

Wolf P. G., K. Schneider & T. A. Ranker, 2001. Geographic distributions of homosporous ferns:

does dispersal obscure evidence of vicariance?. *Journal of Biogeography*, **28**: 263-270.

1   **Table 1.** Summary of partial Mantel tests of floristic dissimilarity distance in three

2   dispersal-ability classes. The tests were conducted with two pairs of explanatory variables,

3   Tokara Gap-geographic distance, and Kerama Gap-geographic distance. The significance was

4   tested with 10000 randomizations. Significance level 5% with Bonferroni adjustment for

5   multiple comparisons was p = 0.05/12 = 0.0042. * = significant, n.s. = not significant

Dispersal ability	Explanatory variable	Controlled variable	r	p
High	Tokara	Distance	0.31	0.0001*
	Distance	Tokara	0.31	0.0002*
	Kerama	Distance	-0.07	0.1876n.s.
	Distance	Kerama	0.29	0.0002*
Intermediate	Tokara	Distance	0.44	0.0002*
	Distance	Tokara	0.38	0.0001*
	Kerama	Distance	-0.28	0.0003*
	Distance	Kerama	0.49	0.0001*
Low	Tokara	Distance	0.18	0.0037*
	Distance	Tokara	0.49	0.0001*
	Kerama	Distance	-0.31	0.0006*
	Distance	Kerama	0.55	0.0001*

FIGURE LEGENDS

Fig. 1. Location of the Ryukyu Archipelago in East Asia and distribution of the 26 islands

analyzed in the archipelago. Segregation of the archipelago into the northern, central, and

southern Ryukyus by the Tokara and Kerama Gaps is shown. Shaded areas indicate land

configuration in the early Pleistocene after segmentation of the landbridge by the formation of

the gaps, following Ota (1998). The names and the coordinates at the highest peaks (N, E) of the

islands are as follows; 1, Takeshima (30° 48′, 130° 25′); 2, Ioujima (30° 47′, 130° 18′); 3,

Kuroshima (30° 49′, 129° 56′); 4, Tanegashima (30° 38′, 130° 58′); 5, Yakushima (30° 20′, 130°

30′); 6, Kuchinoerabujima (30° 26′, 130° 13′); 7, Kuchinoshima (29° 58′, 129° 55′); 8,

Nakanoshima (29° 51′, 129° 51′); 9, Gajajima (29° 54′, 129° 32′); 10, Tairajima (29° 41′, 129°

32′); 11, Suwanosejima (29° 38′, 129° 42′); 12, Akusekijima (29° 27′, 129° 35′); 13,

Kodakarajima (29° 13′, 129° 19′); 14, Takarajima (29° 08′, 129° 12′); 15, Yokoatejima (28° 48′,

128° 59′); 16, Amamioshima (28° 17′, 129° 19′); 17, Kikaijima (28° 18′, 129° 58′); 18,

Tokunoshima (27° 46′, 128° 58′); 19, Okinoerabujima (27° 22′, 128° 34′); 20, Yoronjima (27°

02′, 128° 25′); 21, Okinawajima (26° 43′, 128° 13′); 22, Kumejima (26° 22′, 126° 46′); 23,

Miyakojima (24° 45′, 125° 22′); 24, Ishigakijima (24° 25′, 124° 11′); 25, Iriomotejima (24° 21′,

123° 53′); 26, Yonagunijima (24° 27′, 123° 00′).

Fig. 2. UPGMA clusterings of the 26 islands based on the floristic dissimilarity distance (1 −

Simpson's similarity index) in high, intermediate, and low dispersal-ability classes. Island

numbers of the northern, central, and southern Ryukyus are colored blue, red and green,

respectively. Island names are given in Fig. 1. The scale bars represent floristic dissimilarity

distance of 0.1.

Fig. 3. Plots of the floristic correlation coefficient (r) as a function of geographic distance in

high, intermediate and low dispersal-ability classes. Each distance class includes the similar

number of pairwise comparisons (32 or 33 island pairs). Dashed lines show the 95% confidence

interval for the null hypothesis of no correlation, based on 1000 randomizations.

cm

Fig. 1

Fig. 2

5 cm

Fig. 3