COMPARATIVE STUDIES ON ARCHITECTURAL STRATIFICATION AND

WOODY SPECIES DIVERSITY IN SUBTROPICAL EVERGREEN BROADLEAF

FORESTS ALONG A LATITUDINAL THERMAL GRADIENT OF THE RYUKYU

ARCHIPELAGO, JAPAN

S. M. Feroz, Rempei Suwa, Koh Nakamura, Akio Hagihara and Masatsugu Yokota

Laboratory of Ecology and Systematics, Faculty of Science, University of the Ryukyus,

Okinawa 903-0213, Japan

ABSTRACT: In order to compare stand structure and woody species diversity of subtropical

evergreen broadleaf forests along a latitudinal thermal gradient of the Ryukyu Archipelago,

tree censuses in a 750 m^2 plot in Okinawa Island and a 400 m^2 plot in Ishigaki Island were

performed. The number of layers increased along a latitudinal thermal gradient from four in

the forest of Okinawa Island to five in the forest of Ishigaki Island. The values of Shannon's

index H' and Pielou's index J' tended to increase from the top layer downward in the forest

of Okinawa Island, whereas in the forest of Ishigaki Island, these values tended to increase

from the bottom layer upward. High woody species diversity depended on small-sized trees in

the Okinawa forest, whereas it depended on large-sized trees in the Ishigaki forest. The

woody species diversity is higher in the Okinawa forest ($H' = 4.83$ bit) than in the Ishigaki

forest ($H' = 4.36$ bit). According to successively decreasing height of layers from the top

downward, the value of H' increased continuously from the top layer downward in the Okinawa forest. This increasing trend was different from the Ishigaki forest, where the value of H' increased up to the second layer and then decreased downward. In the Okinawa forest, the expected number of species increased continuously from the top toward the bottom layer, i.e. the bottom layer contained the highest potential number of species (65). However, in the Ishigaki forest, it increased from the top to the fourth layer and then decreased to the bottom layer, i.e. the fourth layer contained the highest potential number of species (90). The floristic composition in the Okinawa forest was different from that in the Ishigaki forest in terms of similarity index, though approximately half of the species were common between them. The highest degree of similarity in floristic composition was between the second and third layers in the Okinawa forest, whereas it was between the third and bottom layers in the Ishigaki forest. The degree of similarity in floristic composition among layers was higher in the Okinawa forest than in the Ishigaki forest. Except the top and the bottom layer respectively in the forests of Okinawa Island and Ishigaki Island, the spatial distribution of trees was random in each layer. The degree of overlapping in the spatial distribution of trees among layers in these two forests suggested that trees in the upper two layers in the Ishigaki forest can catch sufficient light, while light can not penetrate easily to the lower three layers in both of the forests. As a result, almost species in the lower layers might be shade-tolerant in both of the forests. For both of the forests, mean tree weight of each layer decreased from the top downward, whereas the corresponding tree density increased from the top downward. This trend resembled the mean weight–density trajectory of self-thinning plant populations.

INTRODUCTION

The coastal areas of the Western Pacific from subarctic eastern Siberia to equatorial SE Asia have forest climates with sufficient rainfall, which develops a sequence of five forest formations: subarctic evergreen conifer forests, cool-temperate deciduous broadleaf forests, warm-temperate lucidophyll forests, subtropical forests and tropical rain forests (Kira, 1991). Within the Western Pacific sequence of thermal vegetation zones, the subtropical zone whose major part is covered by dry area, only a small part including a chain of islands from Okinawa to Taiwan and South China, is sufficiently moist to allow the development of subtropical forests. Therefore, the subtropical forests in the Ryukyu Archipelago are precious from a phytogeographical viewpoint.

A well-developed evergreen broadleaf forest exists in the northern part of Okinawa Island and in the central part of Ishigaki Island of the Ryukyu Archipelago. It is ecologically important to know how floristic composition, spatial distribution of trees, multi-layering structure and woody species diversity among layers change along a latitudinal thermal gradient of the Ryukyu Archipelago.

The diversity of tree species is fundamental to total forest biodiversity, because trees provide resources and habitats for almost all other forest species (Hall & Swaine, 1976; Huston, 1994; Whitmore, 1998; Huang et al., 2003). Measures of species diversity play a central role in ecology and conservation biology. The most commonly employed measures of species diversity are the Shannon function, species richness (number of species), and evenness (the

distribution of abundance among the species, sometimes known as equitability). In addition,

the spatial distribution of trees has been a major source of interest for plant ecologists because

of its potential role in explaining the coexistence of tree species in species-rich forests

(Bunyavejchewin et al., 2003).

The degree of canopy multi-layering and the woody species diversity increase along a

latitudinal thermal gradient from higher latitudes to the tropics (Hozumi, 1975; Yamakura,

1987; Kira, 1991; Ohsawa, 1995; Kimmins, 2004). The canopy multi-layering structure, i.e.

architectural stratification, is an important factor in maintaining higher woody species

diversity (Roberts & Gilliam, 1995; Lindgren & Sullivan, 2001). However, there is a dearth of

studies reporting the effect of the architectural stratification on floristic composition, woody

species diversity and spatial distribution of trees in the subtropical evergreen broadleaf forest

in Okinawa Island. It is evident that there have apparently been no studies from such a point

of view in the subtropical evergreen broadleaf forest in Ishigaki Island. The purposes of this

study were, therefore, to distinguish architectural stratification, to quantify woody species

diversity, floristic composition and spatial distributions of trees on the basis of the

architectural stratification and to compare the above parameters between the forests in the

northern part (Okinawa Island) and the southern part (Ishigaki Island) of the Ryukyu

Archipelago.

22   **MATERIALS AND METHODS**

24   *Study sites and sampling*

1 Two study sites were respectively selected in Okinawa Island and Ishigaki Island of the

2 Ryukyus Archipelago. One is located in a subtropical evergreen broadleaf forest at Mt.

3 Yonaha (26º45´N and 128º10´E), the northern part of Okinawa Island. The bedrock is

4 composed of silicate having soil pH of 4.35 (Alhamd et al., 2004; Feroz et al., 2006). The

5 other is located in a subtropical evergreen broadleaf forest at Mt. Omoto (24º25′03″N and

6 124º11′17″E), the central part of Ishigaki Island. The bedrock is composed of silicate having

7 soil pH of 4.55.

9 For the former site, a sampling plot of area 750 m^2 (25 m × 30 m) was established and divided

10 into 120 quadrats (2.5 m × 2.5 m). The slope, altitude and direction of the plot were 24.5º, 250

11 m above sea level and NWW, respectively. For the latter site, a sampling plot of area 400 m^2

12 (20 m × 20 m) was established and divided into 64 quadrats (2.5 m × 2.5 m). The slope and

13 altitude and direction of the plot were 21.4º, 200 m above sea level and SW, respectively.

14 Woody plants over 0.10 m in height for the former site and all woody plants for the latter site

15 were numbered. They were identified to species according to the nomenclature of Walker

16 (1976). Tree height H (m) and stem diameter at a height of H/10 $D_{0.1H}$ (cm) were measured.

19 *Climate*

21 The climatic data for the year of 2003~2007 were collected from Nago Meteorological Station

22 for the northern part of Okinawa Island and Maezato Meteorological Observatory for the

23 central part of Ishigaki Island. Mean monthly minimum temperature and mean monthly

24 maximum temperature are respectively 16.2 ± 0.5 (SE) ºC in January and 29.1 ± 0.2 (SE) ºC

25 in July in Okinawa Island, and 18.2 ± 0.4 (SE) ºC in January and 29.2 ± 0.3 (SE) ºC in July in

26 Ishigaki Island. Mean annual temperature is 22.9 ± 0.3 (SE) ºC in Okinawa Island and 23.9 ±

1 0.3 (SE) ºC in Ishigaki Island. The warmth index is 214.2 ± 0.5 (SE) ºC month in Okinawa

2 Island and 227.2 ± 1.3 (SE) ºC month in Ishigaki Island, which values are within the range of

3 180 to 240 ºC month of the subtropical region defined by Kira (1977). Mean monthly rainfall

4 in both of the islands is over 100 mm throughout the year except for 84 ± 22 (SE) mm in

5 February in Okinawa Island and 93 ± 19 (SE) mm in December in Ishigaki Island. Mean

6 annual rainfall is 2050 ± 182 (SE) mm yr^{-1} in Okinawa Island and 1942 ± 159 (SE) mm yr^{-1} in

7 Ishigaki Island. Typhoons with strong winds and rains frequently strike the islands between

8 July and October.

11 *Architectural stratification*

13 The M–w diagram proposed by Hozumi (1975) was used to identify the multi-layering

14 structure of the forest stands. Tree weight w was assumed to be proportional to $D_{0.1H}^{2}H$ and it

15 was arranged in descending order. Average tree weight M_n from the maximum tree weight w_1

16 to the nth tree weight was calculated using the form of $M_n = \sum_{i=1}^{n} w_i \Big/ n$ ($n = 1,\ldots.$, total

17 number of trees N). If the M–w diagram is constructed by plotting the values of M against the

18 corresponding values of w on logarithmic coordinates, then some segments on the M–w

19 diagram are formed. Each segment is related to the layer with the specific characteristics of

20 the beta–type distribution designated by Hozumi (1971, 1975). Hozumi (1975) pointed out

21 that the segments on the M–w diagram can be written by either of the following equations:

22 $M = A\,w + B$                        (1)

23 $M = C\,w^{b}$                          (2)

24 where A, B, C and b are coefficients. These functions reflect some aspect of the manners of

25 packing trees into the three-dimensional space as realized by a forest stand.

A boundary between layers was distinguished by the relationships of the first

derivative S_1 and the second derivative S_2 to tree weight w on the M–w diagram. The S_1 and

S_2 are defined as follows:

$$S_1 = \frac{d \ln M}{d \ln w} \cong \frac{\Delta \ln M}{\Delta \ln w} \quad (3)$$

$$S_2 = \frac{d}{d \ln w}\left(\frac{d \ln M}{d \ln w}\right) = \frac{dS_1}{d \ln w} \cong \frac{\Delta S_1}{\Delta \ln w} \quad (4)$$

When the moving value of S_1 is plotted against the corresponding value of w, points of

inflection probably appear instead of the points of intersection between the segments on the

M–w diagram. Then, the second derivative S_2 has extrema, whose x-coordinates can be

regarded as the points of boundary between layers.

Species dominance

Dominance of a species was defined by importance value IV (%) of the species:

$$IV = \left(\frac{n_i}{\sum\limits_{i=1}^{Q} n_i} \times 100 + \frac{a_i}{\sum\limits_{i=1}^{Q} a_i} \times 100 + \frac{f_i}{\sum\limits_{i=1}^{Q} f_i} \times 100\right) \Big/ 3 \quad (5)$$

where n_i is the number of individuals of the ith species, a_i is the basal area at a height of

$H/10$ of the ith species, f_i is the number of quadrats in which the ith species appeared and Q

is the total number of quadrats.

Species-area relationship

The expected number of species S_q appeared within the number of quadrats q selected at

random from the total number of quadrats Q was calculated from the equation proposed by

Shinozaki (1963) (cf. Hurlbert, 1971):

$$S_q = \sum_{i=1}^{s}\left[1-\left(\begin{array}{c}Q-q_i\\q\end{array}\right)\bigg/\left(\begin{array}{c}Q\\q\end{array}\right)\right]$$ (6)

where q_i is the number of quadrats in which the ith species occurred and S is the total number

of species. The S_q-values were obtained for q-values of 1, 2, 4, 8, 16, 32, 64 and 120 for the

forest in Okinawa Island, and 1, 2, 4, 8, 16, 32 and 64 for the forest in Ishigaki Island.

Floristic similarity

The similarity of floristic composition between layers was calculated using the following

index C_Π (Horn, 1966; cf. Morishita, 1959; Kimoto, 1967):

$$C_\Pi = \frac{2\sum_{i=1}^{s}n_{Ai}n_{Bi}}{(\Pi_A+\Pi_B)N_A N_B}$$ (7)

$$\left(\Pi_A = \sum_{i=1}^{s}n_{Ai}^2\bigg/N_A^2,\ \Pi_B = \sum_{i=1}^{s}n_{Bi}^2\bigg/N_B^2\right)$$

where S is the total number of species, n_{Ai} and n_{Bi} are the number of individuals of the ith

species respectively belonging to Layer A and Layer B. The value of C_Π is 1.0 when the

number of individuals belonging to a species is the same between the two layers for all

species, i.e., floristic composition is completely the same between the layers, and is 0.0 when

no common species is found between them.

Equation (7) was also applied for measuring the degree of similarity in floristic

composition between the Okinawa and Ishigaki forests. In this case, S is the total number of

species in the two forests, n_{Ai} and n_{Bi} are the number of individuals of the ith species

respectively belonging to the Okinawa and Ishigaki forests.

Species diversity

The following two indices of Shannon's index (MacArthur & MacArthur, 1961) H' (bit) and

Pielou's (1969) index J' were used to measure woody species diversity or equitability

(evenness):

$$H' = \sum_{i=1}^{s} \frac{n_i}{N} \log_2 \frac{N}{n_i} \tag{8}$$

$$J' = \frac{H'}{H'_{max}} \quad (H'_{max} = \log_2 S) \tag{9}$$

where N is the total number of individuals.

Spatial distributions of trees

The unit-size $\overset{*}{m_u} - m_u$ method and the ρ–index with successive changes of quadrat sizes (Iwao,

1972) were used to analyze the spatial distribution of trees. Mean density m_u is defined as:

$$m_u = \frac{\sum_{j=1}^{q_u} n_j}{q_u} \tag{10}$$

where n_j is the number of individuals in the jth quadrat and q_u is the total number of quadrats

when the quadrat size is u. The q_u-values were 120, 60, 30, 15, 7, 4 and 2 respectively for the

u-values of 1 (2.5 m × 2.5 m), 2, 4, 8, 16, 30 and 60 in the Okinawa forest, and 64, 32, 16, 8, 4

and 2 respectively for the u-values of 1 (2.5 m × 2.5 m), 2, 4, 8, 16 and 32 in the Ishigaki

forest. On the other hand, mean crowding $\overset{*}{m}_u$ is defined by Lloyd (1967) as:

$$\overset{*}{m}_u = \frac{\sum_{j=1}^{q_u} n_j (n_j - 1)}{\sum_{j=1}^{q_u} n_j}$$ (11)

If the basic component of the spatial distribution is a single individual tree, individual

trees are considered to be randomly distributed when $\overset{*}{m}_u = m_u$, aggregately distributed when

$\overset{*}{m}_u > m_u$ and uniformly distributed when $\overset{*}{m}_u < m_u$ for any quadrat size. In order to provide

knowledge on the distribution pattern of clumps, Iwao (1972) proposed the ρ–index.

$$\rho_u = \frac{\overset{*}{m}_u - \overset{*}{m}_{u-1}}{m_u - m_{u-1}}$$ (12)

where for the smallest quadrat size ($u = 1$), $\rho_1 = \overset{*}{m}_1 / m_1$. When the values of ρ_u are plotted

against the quadrat sizes, a peak of the curve may suggest the clump area.

Overlapping in spatial distributions of trees between layers

The ω-index, which is proposed by Iwao (1977) for analyzing spatial association between

species, was applied to measure the degree of overlapping in spatial distributions of trees

among layers with successive changes of quadrat size u.

$$\omega_{(+)} = \frac{\gamma - \gamma_{(ind)}}{1 - \gamma_{(ind)}} \qquad (\gamma \geq \gamma_{(ind)})$$ (13)

2
$$\omega_{(-)} = \frac{\gamma - \gamma_{(ind)}}{\gamma_{(ind)}} \qquad (\gamma \le \gamma_{(ind)})$$

4 where γ and $\gamma_{(ind)}$ are respectively given in the forms:

6
$$\gamma = \frac{\sum\limits_{j=1}^{Q_u} n_{Aj} n_{Bj}}{\sqrt{\sum\limits_{j=1}^{Q_u} n_{Aj}^{\,2} \sum\limits_{j=1}^{Q_u} n_{Bj}^{\,2}}}$$

9
$$\gamma_{(ind)} = \frac{1}{Q_u} \frac{\sum\limits_{j=1}^{Q_u} n_{Aj} \sum\limits_{j=1}^{Q_u} n_{Bj}}{\sqrt{\sum\limits_{j=1}^{Q_u} n_{Aj}^{\,2} \sum\limits_{j=1}^{Q_u} n_{Bj}^{\,2}}}$$

Here, Q_u is the total number of quadrats taken as 120, 60, 30, 15 and 7 respectively for the u-

values of 1 (2.5 m $\times$ 2.5 m), 2, 4, 8 and 16 in the Okinawa forest, and 64, 32, 16, 8, 4 and 2

respectively for the u-values of 1 (2.5 m $\times$ 2.5 m), 2, 4, 8 and 16 in the Ishigaki forest, n_{Aj} and

n_{Bj} are the number of individuals of the jth quadrat respectively belonging to Layer A and

Layer B. The value of ω changes from the maximum of +1.0 for complete overlapping,

through 0.0 for independent occurrence, to the minimum of –1.0 for complete exclusion.

Dendrogram

The dendrograms for analyzing the degrees of floristic similarity among layers were

constructed following Mountford's (1962) method, or unweighted pair-group method using

arithmetic averages (Sneath & Sokal, 1973).

Regression analysis

The coefficients for nonlinear equations were determined with statistical analysis software

(KaleidaGraph V. 4.0, Synergy Software, USA). On the other hand, the coefficients for linear

and curvilinear equations were determined with the least-squares method.

9   **RESULTS**

Architectural stratification

The $M–w$ diagram in the Okinawa forest is illustrated in Fig. 1a. As is clear from Fig. 1b, the

$M–w$ diagram shows four phases, the first and fourth of which phases possess a property of

Eq. (1), i.e. Type I of the C–D curve tribe that contains eight types (Shinozaki & Kira 1961),

whereas the second and third phases possess a property of Eq. (2), i.e. a power function. The b

values in Eq. (2) were significantly different between the second and the third phases ($t = 212$,

$P \cong 0$). As a result, it was confirmed that the forest consists of four architectural layers.

Figure 1c shows that the extrema, represented by arrows, emerged apparently in the $S_2 - \ln w$

relationship. Their tree weights at boundaries between layers were estimated to be 508, 3.18

and 0.00953 cm^2 m.

On the other hand, the $M–w$ in the Ishigaki forest is illustrated in Fig. 2a. As is clear from Fig.

2b, the $M–w$ diagram shows five phases, all of which have the property of Eq. (1). As a result,

it was confirmed that the forest consists of five architectural layers. Figure 2c shows that the

extrema, represented by arrows, apparently emerged in the $S_2 - \ln w$ relationship. Their tree

weights at the boundaries between layers were estimated to be 706.3, 18.74, 0.6871 and

0.01236 cm^2 m.

Figure 3 shows the relationships between tree height H and weight w respectively for the

Okinawa and Ishigaki forests, which were respectively formulated (cf. Kira & Ogawa, 1971)

as follows:

$$\frac{1}{H} = \frac{1}{1.37w^{0.361}} + \frac{1}{21.0} \qquad (14)$$

$$\frac{1}{H} = \frac{1}{1.06w^{0.368}} + \frac{1}{21.0} \qquad (15)$$

The estimated maximum tree height was 21.0 m, which was the same in both of the forests.

The heights of the boundaries were determined as 8.0, 1.9 and 0.25 m in the Okinawa forest

and 7.6, 2.7, 0.87 and 0.21 m in the Ishigaki forest by substituting the tree weights at

boundaries obtained above for w respectively in Eqs. 14 and 15. Therefore, in the Okinawa

forest, the height range was 8.0 m $< H \leq 16.5$ m for the top layer, 1.9 m $< H \leq 8.0$ m for the

second layer, 0.25 m $< H \leq 1.9$ m for the third layer and 0.10 m $\leq H \leq 0.25$ m for the bottom

layer. In the Ishigaki forest, the height range was 7.6 $< H \leq 14.1$ m for the top layer, 2.7 $< H \leq$

7.6 m for the second layer, 0.87 $< H \leq 2.7$ m for the third layer, 0.21 $< H \leq 0.87$ m for the

fourth layer and 0.0 $< H \leq 0.21$ m for the bottom layer.

Species dominance

Table 1 lists importance value IV of five woody species for each layer in each of the forests in

order of species rank, which was determined from IV in the total stand. In the subtropical

evergreen broadleaf forest in Okinawa Island, a total of 26 families, 43 genera, 60 species and

4684 woody individuals were recorded. The most species rich family was Rubiaceae, which

contained 12 species. *Symplocos*, *Lasianthus* and *Ilex* were the species rich genera, each of

which contained five species. Out of the 60 species, only three species (5%) consisted of a

single individual. *Castanopsis sieboldii* (Mak.) Hatusima appeared in all layers with the

highest importance value, especially with a tremendously high value of 44% in the top layer.

This phenomena indicate that *C. sieboldii* is the most dominant and climax species. *Schima*

wallichii (DC.) Korth. was the second dominant species in terms of *IV* of 9.30% in the total

stand. A quite high *IV* of 22.8% of *S. wallichi* in the top layer compared with the very low *IV*

(ranging from 0.62 to 1.69%) in the lower three layers indicate the heliophilic nature of the

species.

In the subtropical evergreen broadleaf forest in Ishigaki Island, a total of 33 families, 52

genera, 77 species and 4157 woody individuals were recorded. The most species-rich family

was Aquifoliaceae, which contained six species. Aquifoliaceae consisted of *Ilex* alone, which

is the most species-rich genus. Out of the 77 species, only four species (5%) consisted of a

single individual. *Ardisia quinquegona* Blume was the most dominant species in terms of the

highest importance value in the total stand and in all layers, except the top layer (Table 1).

This species may be a subcanopy species, i.e. not a climax species, because of its

disappearance in the top layer. *Castanopsis sieblodii* (Mak.) Hatusima was a low rank species

in terms of the low importance value in the total stand, though it is a climax species in the

Ishigaki subtropical evergreen broadleaf forest (Foster et al., 1960). However, this species

appeared in all layers with a considerably higher importance value, especially the second

highest importance value in the top layers (Data not shown).

Species–area relationship

As shown in Fig. 4, the expected number of species increased, and then tended to be saturated

with increasing number of quadrats for the Okinawa forest, whereas it increased with

increasing number of quadrats for the Ishigaki forest. The relationships of the expected

number of species S_q to the number of quadrats q in each layer and the total stand were well

approximated by the following equation (Ogawa, 1980; cf. Hagihara, 1995):

$$\frac{1}{S_q} = \frac{1}{cq^d} + \frac{1}{S_{max}}$$ (16)

where c and d are coefficients, and S_{max} is the expected maximum number of species.

In the Okinawa forest, the expected maximum number of species was estimated to be 62 in

the total stand. The S_{max} increased from 26 in the top layer, through 46 in the second layer

and 61 in the third layer, to 65 in the bottom layer. Therefore, the bottom layer contained the

highest potential number of species. This result was different from the Ishigaki forest, where

the expected maximum number of species was estimated to be 92 in the total stand. The

fourth layer contained the highest potential number of species as the S_{max} increased from 44 in

the top layer, through 55 in the second layer and 77 in the third layer, to 90 in the fourth layer,

but decreased to 88 in the bottom layer.

Floristic similarity

The floristic similarities among layers of each forest were classified using dendrograms of

similarity index C_Π, as shown in Fig. 5. In the Okinawa forest, the strongest similarity in

floristic composition was marked between the second and the third layers with a C_Π-value of

0.80. The next was between the second-third and the bottom layers with a C_Π-value of 0.64.

The lowest C_Π-value of 0.48 was between the top and the lower three layers. On the other

hand, in the Ishigaki forest, the third and the bottom layers showed the highest similarity in

floristic composition with a C_Π-value of 0.94. The second and third highest similarities were

respectively between the third-bottom and the fourth layers with a C_Π-value of 0.69, and

between the third-bottom-fourth and the second layers with a C_Π-value of 0.36. The lowest

C_Π-value of 0.15 was between the top and the lower four layers.

Woody species diversity in the stratified forest stands

In the Okinawa forest, the values of H' and J' were respectively 2.75 bit and 0.66 for the top

layer, 4.37 bit and 0.81 for the second layer, 4.73 bit and 0.80 for the third layer, 4.33 bit and

0.73 for the bottom layer. In the Ishigaki forest, on the other hand, the values of H' and J'

were respectively 3.84 bit and 0.90 for the top layer, 4.49 bit and 0.87 for the second layer,

4.29 bit and 0.71 for the third layer, 4.21 bit and 0.70 for the third layer, 3.73 bit and 0.65 for

the bottom layer (Table 2). These results indicate that the values of diversity indices H' and

J' tended to increase from the top layer downward except for the bottom layer in the Okinawa

forest, whereas in the Ishigaki forest they tended to increase from the bottom layer upward

except for the H'-value of the top layer.

When species diversity compared among layers, the highest value of H' was in the third

layer in the Okinawa forest. This is because of the second highest value of J' and the highest

species richness (56 species) in the third layer. However, in the Ishigaki forest, the highest

value of H' occurred in the second layer. This is mainly due to the second highest value of

J'¸ though species richness (35 species) was the fourth highest in the second layer. On the

other hand, the lowest value of H' was in the top layer in the Okinawa forest. This is ascribed

to the lowest J'-value and species richness (18 species) in the top layer. The lowest value of

H' was in the bottom layer. This is ascribed to the lowest value of J', though species

richness (52 species) was the third highest. In addition, in the Okinawa forest, the values of

H' in the second layer and in the bottom layer were nearly the same, while the value of

J' was higher in the second layer than in the bottom layer. This is because an increase of

species richness from the second layer (41 species) to the bottom layer (53 species) could

compensate for a decrease of evenness J' from the second layer to the bottom layer. On the

other hand, in the Ishigaki forest, the values of H' in the top layer and in the bottom layer

were nearly the same, while J' was the highest in the top layer, but was the lowest in the

bottom layer. This is because an increase of evenness J' from the bottom layer to the top

layer could compensate for a decrease of species richness from the bottom layer (52 species)

to the top layer (19 species).

Spatial distributions of trees

The spatial distribution patterns of trees for each layer are shown in Fig. 6 based on the unit-

size $\overset{*}{m}_{u} - m_{u}$ relation and the ρ-index against quadrat size u. The spatial distribution of the

basic component for all layers, except the top layer in the Okinawa forest and the bottom

layer in the Ishigaki forest, was a single individual and its distribution was random, because

the $\overset{*}{m_u} - m_u$ relation line mimicked the Poisson line at a significance level of 0.01. However,

in the top layer of the Okinawa forest, the spatial distribution of trees showed a special trend.

The ρ-index showed three peaks respectively at quadrat size 1, 4 and 30. The first peak at

quadrat size 1 may be occasional appearance due to the underdispersion of individual trees,

and the second and third peaks may perhaps be related to topography. As a result, the top

layer probably consists of a double clump, the small and large clumps of which were

respectively 25 m^2 and 187.5 m^2. In light of mean area occupied by individuals of the top

layer trees in the Okinawa forest (9.1 ± 0.6 (SE) m^2), the small clump may include three

individuals and the large clump may include around eight small clumps. On the other hand, in

the bottom layer of the Ishigaki forest, the spatial distribution of trees was aggregated ($P = 1.1$

$\times 10^{-4}$), though its basic component was a single individual.

Overlapping in spatial distributions of trees among layers

The degree of overlapping ω with successive changes of quadrat sizes in spatial

distributions of trees among layers combined successively from the top layer downward is

shown in Fig. 7. In the Okinawa forest, the spatial distributions of trees were overlapped

between the top and the second layers, between the top-second and the third layer, and also

between the top-second-third and the bottom layer. Similarly, in the Ishigaki forest, the spatial

distributions of trees were nearly overlapped between the top-second and the third layers,

between the top-second-third and the fourth layers, and also between the top-second-third-

fourth and the bottom layers. Only a different degree of overlapping occurred between the top

and the second layers, where the spatial distributions tended to be independent with each other.

These results may indicate that trees in the upper two layers in the Ishigaki forest can catch

sufficient light, while light can not penetrate easily to the lower three layers in both of the

forests. As a result, almost species in the lower layers might be shade-tolerant in both of the

forests.

Mean tree weight and density among layers

As shown in Fig. 8, mean tree weight $\overline{w_i}$ of the ith layer decreased from the top ($i = 1$) toward

the bottom layer ($i = 4$), whereas the opposite trend was observed for the tree density ρ_i of the

ith layer. This trend was successfully expressed for the forest in Okinawa Island in the form

(Feroz et al., 2006):

$$\overline{w_i} = K\rho_i^{-\alpha}\left(1 - \frac{\rho_i}{\rho_{\mathrm{o}}}\right)$$　　　　　　　　(17)

In the Okinawa forest, the values of coefficients K, α, and ρ_{o} were estimated as 2.23 x 10^8

$\mathrm{cm}^2\,\mathrm{m}\,\mathrm{ha}^{-\alpha}$, 1.49 and 24575 ha^{-1}. The relationship between mean tree weight $\overline{w_i}$ and tree

density ρ_i in the Ishigaki forest was well described by Eq. 15 with the same values of the

coefficients obtained in the Okinawa forest ($R^2 = 0.98$).

20　　　　　　　　　　　　　　　　**DISCUSSION**

All species in both of the forests in Okinawa Island and Ishigaki Island are evergreen

broadleaf. The number of species in the Okinawa forest was lower than that in the Ishigaki

forest, though the study area in Okinawa Island was approximately twice as large as that in

Ishigaki Island. Similarly, the potential number of species in the Okinawa forest (62) was

lower than that in the Ishigaki forest (92).

In both of the forests, *Castanopsis sieboldii* (Mak.) Hatusima, *Daphniphyllum teijmannii* Zoll.

ex Kurz, *Neolitsea aciculata* (Bl.) Koidz. and *Dendropanax trifidus* (Thunb.) Makino were

typically facultative shade species (light-tolerant under high light conditions and shade-

tolerant under low light conditions) as they appeared in all layers, especially *Castanopsis*

sieboldii was the most dominant species in the Okinawa forest with its highest importance

value in each layer (Table 1). Although *Ardisia quinquegona* Blume was the most dominant

species in the Ishigaki forest, it was not typically a facultative shade species. This is because

this species did not appear in the top layer, i.e. mainly shade-tolerant, growing from the

bottom layer to the subcanopy layer (or second layer). *Ardisia quinquegona* also did not

appear in the top layer in the Okinawa forest, though it was one of the dominant species.

Schima wallichii (DC.) Korth. in the Okinawa forest was typically a sun species as it appeared

in the top layer with a tremendously high importance value, while it had a very low

importance value in the lower three layers. In the Ishigaki forest, however, this species only

appeared in the lower three layers, indicating a shade-tolerant nature which is different from

the Okinawa forest.

The similarity in floristic composition between the top layer and the lower three layers in the

Okinawa forest was weak, i.e. approximately one-third of the species from the lower three

layers may be able to grow into the top layer. In the Ishigaki forest, however, the floristic

composition between the top layer and the lower four layers was almost exclusive, i.e.

approximately one-sixth of the species from the lower four layers may be able to grow into

the top layer. This is because the floristic composition was more similar among layers in the

Okinawa forest than in the Ishigaki forest (Fig. 5). Moreover, a low similarity in floristic

composition was observed between these two forests, which is concluded from the low value

of similarity index ($C_{\Pi} = 0.33$), though approximately half of the total number of species were

common. This is mainly due to the different number of individuals belonging to the same

species between these two forests. For example, the number of individuals of *Castanopsis*

sieboldii (Mak.) Hatusima in the Okinawa forest ($10480\ ha^{-1}$) was not the same as in the

Ishigaki forest ($2600\ ha^{-1}$).

It is known that the diversity of a community depends on two things: species richness and the

evenness with which the individuals are apportioned among them (Pielou, 1975). As the

lower layers contained many species relative to their smaller height ranges (Table 2),

obviously these layers support high species richness of the forests. For example, 88% of the

total species with 29% of the total individuals in the Okinawa forest and 67% of the total

species with 30% of the total individuals in the Ishigaki forest packed within such thin bottom

layers of 15 cm and 21 cm deep, respectively.

In the Okinawa forest, the value of H' for small-sized trees having $H \geq 0.10$ m was quite

high as compared to that of H' for large-sized trees having DBH ≥ 4.5 cm (Table 3). This is

mainly caused by a large number of species for small-sized trees, though higher J'-value has

a small influence on the high value of H'. As shown in Fig. 9, the trend of increasing

diversity with successively decreasing height of layers from the top downward represents that

high woody species diversity depended on small-sized trees. Since small-sized trees provide a

natural habitat for animals living on the forest floor, conservation of small-sized trees in the

lower layers is indispensable to sound maintenance of Okinawan evergreen broadleaf forests

in Okinawa Island.

On the other hand, in the Ishigaki forest, the value of H' for large-sized trees having DBH $\geq$

4.5 cm was quite high as compared to that of H' for the total stand (Table 3). This is mainly

caused by a high evenness for large-sized trees, though smaller number of species has an

influence on the low value of H'. The diversity tends to increase up to the second layer and

then decrease downward (Fig. 9), which was different from the Okinawa forest. These results

may represent that the large-sized trees have an important role in maintaining high woody

species diversity. Thus, it may be thought that the decreasing trend of H' from the second

layer downward is a characteristic of the subtropical evergreen broadleaf forest in Ishigaki

Island.

The values of the diversity indices (H' and J') for the total stand in the Okinawa forest were

higher than those in the Ishigaki forest (Table 3). This is because of a quite high evenness in

the Okinawa forest compared to that in the Ishigaki forest, though species richness was lower

in the Okinawa forest than in the Ishigaki forest. However, woody species diversity for large-

sized trees is lower in the Okinawa forest than in the Ishigaki forest. This is caused by a

higher species richness and evenness in the Okinawa forest than in the Ishigaki forest.

Therefore, the statement of increasing multi-layering structure and woody species diversity

along a latitudinal thermal gradient from higher latitudes to the tropics (Hozumi, 1975;

Yamakura, 1987; Kira, 1991; Ohsawa, 1995; Kimmins, 2004) supported the present study for

large-sized trees, but it did not support for total trees.

In Table 3, tree density of 62452 ha^{-1} for trees having $H \geq 0.10$ m in the Okinawa forest was

quite low compared to that of 103925 ha^{-1} for total trees in the Ishigaki forest, whereas basal

area of 50.3 m^2 ha^{-1} in the Okinawa forest was almost the same as that of 49.4 m^2 ha^{-1} in the

Ishigaki forest. However, tree density and basal area for trees having DBH ≥ 4.5 cm in the

Okinawa forest were respectively 3080 ha^{-1} and 43.2 m^2 ha^{-1}, whose values are comparable

with 2775 ha^{-1} and 42.7 m^2 ha^{-1} in the Ishigaki forest. Therefore, it follows that the Ishigaki

forest is densely populated with small-sized trees compared to Okinawa forest. However, the

biomasses of these forest stands are possibly similar, because the basal areas are very close to

each other for trees having DBH ≥ 4.5 cm.

The trend that mean tree weight decreased from the top toward the bottom layer, whereas tree

density increased from the top downward (Fig. 8) was successfully formulated with Eq. 15.

This trend resembles the mean plant weight–density trajectory of self-thinning even-aged

plant populations (Hagihara, 2000), which start growing from initial plant densities lower than

the initial plant density of the population obeying the -3/2 power law of self-thinning (Yoda et

al., 1963). The relationship of mean tree height to tree density for the upper two layers in the

Okinawa forest supported Yamakura's quasi -1/2 power law of tree height, because the α -

value in Eq. 15 was close to 3/2 (Feroz et al., 2006). Our formulated equation is more general

than Yamakura's quasi -1/2 power law of tree height, because the equation describes lower

layers as well.

20   **ACKNOWLEDGEMENTS**

22 We are grateful to our colleagues, Drs. L. Alhamd and M.N.I. Khan, Messrs. B. Tanaka, P. Wane,

23 S. Sharma, K. Analuddin and Haque A.T.M. Rafiqul, and Mses. M. Wu and Y. Li, for their

24 cooperation and active participation in the field works. Special thanks go to Prof. T. Shinzato,

25 Subtropical Field Science Center at Yona, University of the Ryukyus, Profs. H. Ota and M.

Izawa and Accos. Prof. T. Denda, Faculty of Science, University of the Ryukyus, and Prof. Z.-L. Huang, South China Institute of Botany, Chinese Academy of Science, for their kind cooperation and valuable suggestions. This study was financed in part by a Grant-in-Aid for Scientific Research (No.16201009) from the Ministry of Education, Culture, Sports, Science and Technology, and by the 21st Century COE program of the University of the Ryukyus.

LITERATURE CITED

Alhamd, L., S. Arakaki & A. Hagihara., 2004. Decomposition of leaf litter of four tree species in a subtropical evergreen broad-leaved forest, Okinawa Island, Japan. *Forest Ecology and Management*. **202**: 1–11.

Bunyavejchewin, S., J. V. LaFrankie, P. J. Baker, M. Kanzaki, P. S. Ashton & T. Yamakura., 2003. Spatial distribution patterns of the dominant canopy dipterocarp species in a seasonal dry evergreen forest in western Thailand. *Forest Ecology and Management*. **175**: 87–101.

Feroz, S. M., A. Hagihara & M. Yokota., 2006. Stand structure and woody species diversity in relation to the stand stratification in a subtropical evergreen broadleaf forest, Okinawa Island. *Journal Plant Research*. **119**: 293–301.

Foster, H. L., C. H. Stensland, H. G. May, F. R. Fosberg & R. J. Alvis, 1960. *Military geology of Ishigaki-Shima, Rȳukyū-Rettō*, The United States Geological Survery.

Hagihara, A., 1995 The Size-Dependence curve as a generalized Competition-Density curve. *Japanese Journal of Forest Planning*. **24**: 15–23.

Hagihara, A., 2000. Time-trajectory of mean phytomass and density in self-thinning plant populations. *Bulletin of theFaculty of Science*, *University of the Ryukyus*. **70**: 99–112.

Hall, J. B. & M. D. Swaine., 1976. Classification and ecology of closed-canopy forest in Ghana. *Journal of Ecology*. **64**: 913–951.

Horn, H. S., 1966. Measurement of "overlap" in comparative ecological studies. *The American Naturalist*. **100**: 419–424.

Hozumi, K., 1971. Studies on the frequency distribution of the weight of individual trees in a forest stand. III. A beta-type distribution. *Japanese Journal of Ecology*. **21**: 152–167.

Hozumi, K., 1975. Studies on the frequency distribution of the weight of individual trees in a forest stand. V. The *M-w* diagram for various types of forest stands. *Japanese Journal of Ecology*. **25**: 123–131.

Huang. W., V. Pohjonen, S. Johansson, M. Nashanda, M. I. L. Katigula & O. Luukkanen., 2003. Species diversity, forest structure and species composition in Tanzanian tropical forests. *Forest Ecology and Management*. **173**: 11–24.

Hurlbert, S. H., 1971. The nonconcept of species diversity: a critique and alternative parameters. *Ecology*. **133**: 125–133.

Huston, M. A., 1994. *Biological diversity*. Cambridge University Press, Cambridge.

Iwao, S., 1972. Application of the $\overset{*}{m}$ –m method to the analysis of spatial patterns by changing the quadrat size. *Researches on Population Ecology*. **14**: 97–128.

Iwao, S., 1977. Analysis of spatial association between two species based on the interspecies mean crowding. *Researches on Population Ecology*. **18**: 243–260.

Kimmins, J. P., 2004. *Forest ecology*. Pearson Education, Inc., New Jersey.

Kimoto, S., 1967. Some quantitative analysis on the Chrysomelid fauna of the Ryukyu Archipelago. *Esakia*. **6**: 27–54.

Kira, T., 1977. *Forest vegetation of Japan. Introduction*. Pages 1-9/14 In: Shidei, T. & T. Kira (eds.). Primary productivity of Japanese forests. Productivity of terrestrial communities. University of Tokyo Press, Tokyo.

Kira, T., 1991. Forest ecosystems of east and southest Aisa in a global perspective. *Ecological Research*. **6**: 185–200.

Kira, T. & H. Ogawa., 1971. *Assessment of primary production in tropical and equatorial forests*. Pages 309–321 In: Productivity of forest ecosystems. Unesco, Paris.

Lindgren, P. M. F. & T. P. Sullivan., 2001. Influence of alternative vegetation management treatments on conifer plantation attributes: abundance, species diversity, and structural diversity. *Forest Ecology and Management*. **142**: 163–182.

Lloyd, M., 1967. Mean crowding. *Journal of Animal Ecology*. **36**: 1–30.

MacArthur, R. H. & J. W. MacArthur., 1961. On bird species diversity. *Ecology*. **42**: 594–598.

Morishita, M., 1959. Measuring of interspecific association and similarity between communities. *Memoirs of the Faculty of Science*, *Kyushu University, Series E* (*Biolog*y). **3**: 65–80.

Mountford, M. D., 1962. *An index of similarity and its application to classificatory problems*. In: Murphy, P. W. (ed). Progress in soil zoology. Butterworths, London. pp 43–50.

Ogawa, H., 1980. *Structure and function of populations*. Asakura-shoten, Tokyo (in Japanese).

Ohsawa, M., 1995. Latitudinal comparison of altitudinal changes in forest structure, leaf-type, and species richness in humid monsoon Asia. *Vegetatio*. **121**: 3–10.

Pielou, E. C., 1969. *An introduction to mathematical ecology*. John Wiley & Sons, Inc., New York.

Pielou, E. C., 1975. *Ecological diversity*. Wiley, New York.

Roberts, M. R. & F. S. Gilliam., 1995. Patterns and mechanisms of plant diversity in forested ecosystems: implications for forest management. *Ecological Application*. **5**: 969–977.

Shinozaki, K., 1963. Note on the species-area curve. Proceedings of Annual Meeting for the Ecological Society of Japan, Tokyo. p 5 (in Japanese).

Shinozaki, K. & T. Kira, 1961. The C-D rule, its theory and practical uses (Intraspecific competition among higher plants X). *Journal of Biology*, *Osaka City University*. **12**: 69–82.

Sneath, P. H. A. & R. R. Sokal., 1973. *Numerical taxonomy*. W. H. Freeman and Company, San Francisco.

Walker, E. H., 1976. *Flora of Okinawa and the Southern Ryukyu Islands*. Smithsonian Institution Press, Washington, D.C.

Whitmore, T. C., 1998. *An introduction to tropical rain forests*. Oxford University Press, New York.

Yamakura, T., 1987. An empirical approach to the analysis of forest stratification. I. Proposed graphical method derived by using an empirical distribution function. *The Botanical Magazine*, *Tokyo*. **101**: 153–162.

Yamakura, T., 1988. An empirical approach to the analysis of forest stratification. II. Quasi - 1/2 power law of tree height in stratified forest communities. *The Botanical Magazine*, *Tokyo*. **101**: 153–162

Yoda, K., T. Kira, H. Ogawa & K. Hozumi., 1963. Self-thinning in overcrowded pure stands under cultivated and natural conditions (Intraspecific competition among higher plants XI). *Jouranl of Biology*, *Osaka City University*. **14**: 107–129.

Table 1. Five dominant species in the Okinawa forest **(a)** and the Ishigaki forest **(b)** in order of species rank determined by importance value *IV* in the total stand.

(a)

Species rank	Name of species	*IV* %				
		Top layer	Second layer	Third layer	Bottom layer	Total stand
1	*Castanopsis sieboldii* (Mak.) Hatusima	43.60	12.63	26.12	21.74	21.30
2	*Schima wallichii* (DC.) Korth.	22.34	1.78	0.62	1.69	9.30
3	*Elaeocarpus japonicus* Sieb. & Zucc.	10.16	7.56	1.73	2.78	4.61
4	*Myrsine seguinii* Lév'l.	–	4.14	5.46	7.01	3.92
5	*Ardisia quinquegona* Bl.	–	4.97	6.88	3.72	3.74

(b)

Species rank	Name of species	*IV* %				
		Top layer	Second layer	Third layer	Fourth layer	Bottom layer	Total stand
1	*Ardisia quinquegona* Bl.	–	9.23	24.66	10.52	24.71	15.39
2	*Lasianthus plagiophyllus* Hance	–	–	8.53	19.10	9.45	7.93
3	*Psychotria manillensis* Bartl.	–	–	9.34	6.89	6.88	6.54
4	*Sarcandra glabra* (Thunb.) Nakai	–	0.84	0.79	10.98	3.18	6.19
5	*Camellia sasanqua* Thunb.	–	7.55	1.65	6.82	6.73	5.54

1   Table 2. Diversity indices among layers in the Okinawa forest **(a)** and the Ishigaki forest **(b)**.

3   **(a)**

Layer	Height range (m)	No. of trees	No. of species	H' (bit)	J'
		per plot area			
Top	$8.0 < H \leq 16.5$	139	18	2.75	0.66
Second	$1.9 < H \leq 8.0$	953	42	4.37	0.81
Third	$0.25 < H \leq 1.9$	2226	56	4.73	0.80
Bottom	$0.10 \leq H \leq 0.25$	1366	51	4.33	0.73

5   **(b)**

Layer	Height range (m)	No. of trees	No. of species	H' (bit)	J'
Top	$7.6 < H \leq 14.1$	42	19	3.84	0.90
Second	$2.7 < H \leq 7.6$	130	35	4.49	0.87
Third	$0.87 < H \leq 2.7$	915	56	4.29	0.71
Fourth	$0.21 < H \leq 0.87$	1806	65	4.21	0.70
Bottom	$0.0 < H \leq 0.21$	1264	52	3.73	0.65

1 Table 3. Comparison in species diversity and tree density between the Okinawa and Ishigaki forests.

Study site	Sample area (ha)	Tree size	Basal area $(m^2\,ha^{-1})$	No. of trees (ha^{-1})	S	H' (bit)	J'	Source
Okinawa forest	0.075	$H \geq 0.10$ m	50.3	62452	60	4.83	0.82	Feroz et al. 2006
		DBH ≥ 4.5 cm	43.2	3080	32	3.80	0.77	
Ishigaki forest	0.04	$H > 0.0$ m	49.4	103925	77	4.36	0.69	
		DBH ≥ 4.5 cm	42.7	2775	36	4.61	0.89	

4 S, the number of species per sample area.

 FIGURE LEGENDS

Fig. 1. Relationships of mean tree weight M (**a**), the first derivative S_1 (**b**) and the second

derivative S_2 (**c**) to tree weight w on logarithmic coordinates in the Okinawa forest. In the

M–w diagram (a), the regression curves of the top and bottom layers are given by equation

(1) ($R^2 = 0.99$) for the top layer and ($R^2 = 0.97$) for the bottom layer. The regression

curves of the second and third layers are given by equation (2) ($R^2 = 0.91$) for the second

layer and ($R^2 = 0.94$) for the third layer. The S_1 and S_2 are respectively defined as

equations (3) and (4). The moving values of S_1 and S_2 are plotted against their

corresponding geometric means of w.

Fig. 2. Relationships of (**a**) mean tree weight M, (**b**) the first derivative S_1 and (**c**) the second

derivative S_2 to tree weight w on logarithmic coordinates in the Ishigaki forest. In the M–w

diagram (a), the regression curves for all layers are given by Eq. (1) ($R^2 = 0.98$) for the top

layer, ($R^2 = 0.92$) for the second layer, ($R^2 = 0.90$) for the third layer, ($R^2 = 0.88$) for the

fourth layer and ($R^2 = 0.96$) for the bottom layer. The application of S_1 and S_2 is the same

as in Fig. 1.

Fig. 3. Relationships between tree height H and weight w. In the Okinawa forest (**a**), the

regression curve is given by Eq. (14) ($R^2 = 0.97$). Filled circles, top layer (8.0 m $< H \le$

16.5 m); open circles, second layer $(1.9$ m $< H \le 8.0$ m); Filled squares, third layer $(0.25$

m $< H \le 1.9$ m); open squares, bottom layer $(0.10$ m $\le H \le 0.25$ m). In the Ishigaki forest

(**b**), the regression curve is given by Eq. (15) ($R^2 = 0.96$). Filled circles, top layer (7.6 m $<$

$H \le 14.1$ m); open circles, second layer $(2.7$ m $< H \le 7.6$ m); Filled squares, third layer

$(0.87$ m $< H \le 2.7$ m); open diamond, fourth layer $(0.21$ m $< H \le 0.87$ m); open squares,

bottom layer $(0.0$ m $< H \le 0.21$ m).

Fig. 4. Species–area curves (area of one quadrat: 2.5 m $\times$ 2.5 m).

Filled diamond, total stand and other symbols are the same as in Fig. 3. In the Okinawa

forest **(a)**, the curves are given by equation (16), where $S_{max} = 62.4$ ($R^2 \cong 1.0$) for the total

stand, $S_{max} = 32.8$ ($R^2 \cong 1.0$) for the top layer, $S_{max} = 47.0$ ($R^2 \cong 1.0$) for the second layer,

$S_{max} = 60.30$ ($R^2 \cong 1.0$) for the third layer, $S_{max} = 61.1$ ($R^2 \cong 1.0$) for the bottom layer. In

the Ishigaki forest **(b)**, the curves are given by the same equation, where $S_{max} = 92.6$ ($R^2 \cong$

1.0) for the total stand, $S_{max} = 43.7$ ($R^2 \cong 1.0$) for the top layer, $S_{max} = 54.7$ ($R^2 \cong 1.0$) for

the second layer, $S_{max} = 71.5$ ($R^2 \cong 1.0$) for the third layer, $S_{max} = 90.2$ ($R^2 \cong 1.0$) for the

fourth layer, $S_{max} = 77.9$ ($R^2 \cong 1.0$) for the bottom layer.

Fig. 5. Dendrograms of the degree of similarity C_Π in floristic composition among layers in

the Okinawa forest **(a)** and in the Ishigaki forest **(b).**

Fig. 6. A schematic representation showing the relationships between mean crowding $\overset{*}{m}_u$ and

mean density m_u , and ρ -index with successive changes of quadrat size u in the Okinawa

forest **(a)** and in the Ishigaki forest **(b)**.

Fig. 7. Degree of overlapping ω with successive changes of quadrat size u in spatial

distributions of trees in the Okinawa forest **(a)** and in the Ishigaki forest **(b)**. The smallest

quadrat size ($u = 1$) is 2.5 m $\times$ 2.5 m.

Fig. 8. Relationship between mean tree weight $\overline{w}_i$ and tree density ρ_i among the layers. The

curves are given by Eq. (17) in the Okinawa and Ishigaki forests ($R^2 = 0.98$). Filled circle,

Okinawa forest; open circle, Ishigaki forest.

Fig. 9. Trends of H' along the layers combined from the top downward in the Okinawa forest

(a) and in the Ishigaki forest **(b)**. Symbols are the same as in Fig. 8.

Figure 1

Feroz et al.

Figure 2

Feroz et al.

Figure 3

Feroz et al.

Number of quadrat

Figure 4

Feroz et al.

Figure 5

Feroz et al.

Figure 6

Feroz et al.

(a)

(Top+2nd) & 3rd layers
Top & 2nd layers
(Top+2nd+3rd) &
Bottom layers

(b)

(Top+2nd) & 3rd layers
(Top+2nd+3rd+4th) & Bottom layers
(Top+2nd+3rd) & 4th layers
Top & 2nd layers

Quadrat size *u*

ω

Figure 7

Feroz et al.

The numbers 1 through 25 appear down the left margin of the page.

Figure 8

Feroz et al.

Top layer to downward successively

Figure 9

Feroz et al.

43