Title: A Genome-Wide Association Analysis Reveals Epistatic Cancellation of Additive

Genetic Variance for Root Length in *Arabidopsis thaliana*

Short Titie: Epistatic Cancellation of Additive Genetic Variance

Authors

Jennifer Lachowiec[1,2+], Xia Shen[3,4,5], Christine Queitsch[1*], Örjan Carlborg[3*]

Affiliations

[1]Department of Genome Sciences, University of Washington, USA

[2]Molecular and Cellular Biology Program, University of Washington, USA

[3]Department of Clinical Sciences, Swedish University of Agricultural Sciences, Uppsala,

Sweden

[4]Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm,

Sweden

[5]MRC Human Genetics Unit, MRC Institute of Genetics and Molecular Medicine, University of

Edinburgh, Edinburgh, UK

* Co-corresponding authors, organ.carlborg@slu.se, queitsch@uw.edu

+Present address: Department of Ecology and Evolutionary Biology, University of Michigan,

USA

Abstract

Efforts to identify loci underlying complex traits generally assume that most genetic variance is

additive. Here, we examined the genetics of *Arabidopsis thaliana* root length and found that the

narrow-sense heritability for this trait was statistically zero. This low additive genetic variance

likely explains why no associations to root length could be found using standard additive-model-

based genome-wide association (GWA) approaches. However, the broad-sense heritability for

root length was significantly larger, and we therefore also performed an epistatic GWA analysis

to map loci contributing to the epistatic genetic variance. This analysis revealed four interacting

pairs involving seven chromosomal loci that passed a standard multiple-testing corrected

significance threshold. Explorations of the genotype-phenotype maps for these pairs revealed

that the detected epistasis cancelled out the additive genetic variance, explaining why these loci

were not detected in the additive GWA analysis. Small population sizes, such as in our

experiment, increase the risk of identifying false epistatic interactions due to testing for

associations with very large numbers of multi-marker genotypes in few phenotyped individuals.

Therefore, we estimated the false-positive risk using a new statistical approach that suggested

half of the associated pairs to be true positive associations. Our experimental evaluation of

candidate genes within the seven associated loci suggests that this estimate is conservative; we

identified functional candidate genes that affected root development in four loci that were part of

three of the pairs. In summary, statistical epistatic analyses were found to be indispensable for

confirming known, and identifying several new, functional candidate genes for root length using

a population of wild-collected *A. thaliana* accessions. We also illustrated how epistatic

cancellation of the additive genetic variance resulted in an insignificant narrow-sense, but

significant broad-sense heritability that could be dissected into the contributions of several

individual loci using a combination of careful statistical epistatic analyses and functional genetic

experiments.

Author summary

Complex traits, such as many human diseases or climate adaptation and production traits in

crops, arise through the action and interaction of many genes and environmental factors. Classic

approaches to identify contributing genes generally assume that these factors contribute mainly

additive genetic variance. Recent methods, such as genome-wide association studies, often

adhere to this additive genetics paradigm. However, additive models of complex traits do not

reflect that genes can also contribute with non-additive genetic variance. In this study, we use

Arabidopsis thaliana to determine the additive and non-additive genetic contributions to the

phenotypic variation in root length. Surprisingly, much of the observed phenotypic variation in

root length across genetically divergent strains was explained by epistasis. We mapped seven

loci contributing to the epistatic genetic variance and validated four genes in these loci with

mutant analysis. For three of these genes, this is their first implication in root development.

Together, our results emphasize the importance of considering both non-additive and additive

genetic variance when dissecting complex trait variation, in order not to lose sensitivity in

genetic analyses.

Introduction

Identifying the loci underlying quantitative phenotypes is among the central challenges in

genetics. The current quantitative genetics paradigm is based on the assumption of additive gene

action, despite an increasing body of evidence showing that non-additive effects are crucial to

many, if not most, biological systems [1–4]. The key arguments for remaining within the

additive paradigm are that many genetic architectures with non-additive gene action display

considerable additive genetic variance in populations [5] and that additive model-based

approaches have facilitated detection of thousands of loci associated with many complex traits

[6]. However, the amount of additive variance contributed by epistasis for example, will vary

with the allele-frequencies in populations and therefore be a dynamic property of studied

populations [7–13 and references therein]. Therefore, the focus on additive variation alone can

leave a considerable amount of genetic variance unexplored [3,14], and for a full dissection of a

complex trait, it is often necessary to explore alternative approaches that capture non-additive

variation. Statistical epistasis has been shown to be pervasive in both humans and model

organisms [2,14]. Epistatic contributions to complex traits have primarily been identified in

experimental populations using candidate gene approaches [1,2,4,15–19] and QTL mapping, for

example [1,20]. Genome-wide association (GWA) analyses have more recently emerged as an

effective method for mapping loci underlying phenotypic variation for complex traits also in

natural populations. However, GWA analyses to detect epistatic interactions are still under-

utilized in efforts to identify loci contributing to the genetic architecture of complex traits. To

obtain a more complete compilation of the loci contributing to the variation of a complex trait, it

would therefore be valuable to also account for epistatic interactions in addition to additive

effects in existing and future datasets.

The drawbacks of studying epistasis in natural populations of higher organisms include the cost

and time to collect large datasets, the lack of power due to noisy phenotypes, and the difficulties

in performing the necessary follow-up studies to replicate the epistatic associations, identify the

polymorphic genes, and dissect the underlying mechanisms leading to statistical epistatic

associations. Efforts to utilize epistatic interactions for detection of novel loci, however, are of

importance for enhancing sensitivity in statistical analyses [21], for selecting the most promising

analytical approaches and experimental designs for future work, and for providing insights as to

what type of genetic and biological mechanisms can lead to non-additive genetic variance and

genotype-phenotype relationships. Public resources have recently been developed for studying

genotype-phenotype associations with GWA analyses in model organisms such as *Arabidopsis*

thaliana [22] and *Drosophila melanogaster* [23]. Many powerful experimental approaches are

available in these species, making them attractive for studies aiming to functionally dissect more

complex genetic mechanisms such as those underlying statistical epistatic associations. For these

reasons, these resources have the potential to contribute to the fundamental work of clarifying

how statistical epistatic analyses should more generally be interpreted and used in efforts to

dissect the genetic architectures of complex traits.

Two major challenges have often been discussed in relation with GWA analyses based on

statistical epistatic models: the lack of sufficiently large experimental datasets to obtain

reasonable statistical power and the computational complexity when screening the genome for

multi-locus epistasis. One of the causes of low power is the need to use stringent, multiple-

testing corrected significance-thresholds to account for the large number of statistical tests

performed when exploring epistasis between all possible combinations of loci [24]. As allele-

frequencies are often skewed in the populations used for GWA analyses, power is decreased

further due to the small number of observations in the multi-locus, minor-allele genotype classes.

To some extent these challenges can be overcome by increasing the sample size and making

efficient use of high-performance computing [25]. Leveraging genotyped populations of model

organisms can also overcome statistical limitations. The genomes of model organisms are often

smaller, which reduces the search-space during genome-wide scans. The existence of inbred

populations, such the *A. thaliana* RegMap [26] or 1001 Genomes Project [14] or the *D.*

melanogaster Drosophila Genetic Reference Panel [15] collections, provide advantages in power

compared to analyses in general outbred populations with heterozygotes. This is particularly true

in analyses of epistasis due to the simpler genotype-phenotype maps. Several standard additive

GWA analyses have been previously performed for many traits using the publically available *A.*

thaliana collections and data sets [27,28], including several studies of *A. thaliana* root

phenotypes [29–31]. *A. thaliana* has a small genome (~125 Mb), large numbers of readily

available inbred accessions, a large knowledge base on how to efficiently score quantitative

phenotypes in large numbers of individuals, and powerful experimental resources, such as

collections of knock-out lines. This makes such populations into particularly well-suited models

for performing exhaustive scans for epistasis and for evaluating whether non-additive genetic

analyses are able to uncover novel contributions of genes to the studied phenotypes. Further,

these advantages lead to a smaller theoretical decrease in power than expected due to sample size

alone.

Model organism GWA studies, however, are often based on a hundred or fewer individuals, and

valid concerns have been raised regarding the challenges of testing for associations with

hundreds of thousands of markers, or millions of combinations of epistatic markers in such small

samples. In particular, the chance of detecting a random association will increase especially

when the number of individuals in the minor two-locus genotype class is small. This situation is

not accounted for by common statistical multiple-testing correction. Further, there is also a risk

of statistical epistatic associations arising due to "apparent epistasis" [32] in which the multi-

locus genotype tags an unobserved, single polymorphism in the genome. Here, the genetic

variance used to detect the association originates from a true genetic effect in the genome, but

this effect is hidden in a standard GWA due to its low correlation with the individual genotyped

markers. Instead the high-order correlation between the hidden genetic variant and a multi-

marker genotype leads to an epistatic variance that can be detectable in the epistatic GWA

analysis. Although this scenario has only previously been discussed for linked variants [32], it

could occur for unlinked loci if the analyzed population is small and the number of evaluated

combinations of loci is large. Also, this scenario goes beyond standard statistics used to account

for multiple testing in GWA analyses, as the increased risk for false positives is not due to the

strength of the statistical association. Instead, it results from the inability to properly compare

alternative explanatory genetic models for the association using the available genotype data.

Unless particular care is taken, both of these scenarios could lead to an increased false-positive

rate among the reported loci, as well as incorrect interpretations of the presence of biological

interactions if significant pairwise statistical epistasis is interpreted as evidence of an underlying

genetic architecture with genetic interactions. These potential risks associated with epistatic

GWA analyses in small experimental populations should therefore be considered when such

analyses are used.

Here, we conduct a GWA study to dissect the genetics underlying *A. thaliana* root length in a

population of wild-collected accessions. Root length had a statistically zero narrow-sense

heritability in this population, but a larger significant broad-sense heritability. The genetic

contribution to this trait was therefore unlikely to be detected using standard additive genetic

models, and it was not surprising that no locus was associated with root-length in the standard

additive-model based GWA analyses. An exhaustive GWA scan for two-locus interactions,

however, revealed four genome-wide significant interacting pairs. All of these displayed

genotype-phenotype relationships in which epistasis cancelled the marginal additive genetic

effects of the interacting loci, explaining the low levels of additive genetic variance in the

population. A new statistical method was used to evaluate the robustness of the GWA

associations in the small population, and it was estimated that half of these pairs are at risk of

being false positives. We also found low risk that the statistical epistatic interactions were caused

by "apparent epistasis" [32]. Our experimental evaluations of genes in the associated regions

using T-DNA insertion mutants identified functional candidates that affected root length in four

regions that were part of three pairs, from which three genes were newly implicated in root

development [33]. Together, our results illustrate the value of epistatic GWA analyses for

revealing novel loci contributing to complex traits in small, inbred populations. Further, we

noticed an increased risk for false-positive epistatic associations and "apparent epistasis" due to

the small sample-size but the top-ranked associations proved, despite this, to be valuable in

further experimental validation, as several functional candidate genes were identified in loci that

would have been discarded based on stringent statistics alone. We also report several promising

functional candidate genes for future studies to dissect the molecular mechanisms underlying

root development in *natural* A. thaliana populations that would have remained undetected

without an epistatic GWA analysis.

Results

Epistasis is the primary source of genetic variance for root length

To study the genetic regulation of mean root length, we generated a comprehensive data set

containing the *A. thaliana* root mean lengths across 93 wild-collected accessions (Fig S1, Table

S1). The 93 accessions used here were previously genotyped at ~215,000 single nucleotide

polymorphisms (SNPs) and used in a standard GWA analysis for many plant phenotypes [27].

We grew seedlings in a randomized design on standard plant medium in temperature-controlled

chambers to reduce environmental variation among individuals [34]. This experimental design

allows for parsing of genetic and environmental factors underlying root length [35,36].

In our dataset, mean root length varied significantly across accessions ($p < 1 \times 10^{-10}$, ANOVA;

Fig1). To evaluate whether the observed variation in mean root length was under genetic control,

we estimated both the narrow- and broad-sense heritability. This analysis used a linear mixed

model that simultaneously estimated the polygenic additive and epistatic variance components

using the R/hglm package [37]. We estimated the total broad sense heritability to be 0.24 (95%

CI 0.10 – 0.38) for mean root length (Table 1), showing that the trait is under genetic control. We

also partitioned the genetic variance into the individual additive and epistatic variance

components and found that almost all the variance was epistatic (0.22; $p = 1.32 \times 10^{-6}$) and only

a small, insignificant fraction was additive (0.01; $p = 0.63$). These heritability patterns suggest

that an epistatic GWA analysis might be needed to identify the loci explaining the heritable

phenotypic variation in root length mean.

Table 1. Estimated narrow (h^2) and broad sense (H^2) heritabilities of root length mean and

variance in a population of 93 natural *A. thaliana* accessions.

The broad-sense heritability was intermediate for root length mean, with a negligible additive

contribution.

Trait	H^2		h^2
	ANOVA[1]	hglm[2]	hglm[2]
Mean root length (mm)	0.25	0.24	0.015
	(p = 0)	(s.e. = 0.09, p = 7.2 × 10^{-3})[3]	(s.e. = 0.056, p = 0.79)[3]

[1]Estimated using ANOVA with accession as a fixed effect, [2]Estimated using the R/hglm package (hglm) by fitting a

linear mixed model including both additive and epistatic kinship matrices as random effects, [3]p-values from Wald

tests for the heritability being larger than zero, and s.e. are the standard errors estimated via jackknife resampling.

No additive genome-wide associations were detected for mean root length

Two GWA analysis methods were used to screen for additive loci contributing to the phenotypic

variation in mean root length. The first method was a standard single-locus additive mixed-

model based approach as implemented in the R-package GenABEL [38] that accounted for

population-structure by modelling the genomic kinships between the accessions as a random

effect. Significance testing was done using a Bonferroni-corrected significance threshold. The

second method was based on a whole-genome generalized ridge regression heteroscedastic

effects model (HEM), in which all SNPs were included simultaneously and their contributions

estimated as random effects using the R/bigRR package [39]. Here, a 5% genome-wide

significance threshold was determined by permutation testing. None of the tested SNPs were

significantly associated with mean root length in either of these analyses (Fig S1a, b). Given the

low estimate for the narrow-sense heritability for mean root length in this population ($h^2 = 0.01$),

this outcome was not surprising.

An epistatic genome-wide association analysis uncovered multiple interacting loci that

contributed to mean root length

To identify loci that contributed to the epistatic genetic variance for mean root length, we

performed an exhaustive two-locus SNP-by-SNP epistasis analysis (PLINKv1.07) [40]. Despite

the limited population size, several factors increased our chances of finding novel loci using this

approach: the small additive and large epistatic genetic variance (Table 1), the high precision in

estimating the phenotypic values due to extensive replication, the presence of only four two-

locus genotypes in the population of inbred accessions, and a reduced number of pairwise

combinations to test in the small genome. To further enhance power and decrease the risk of

false-positives, we excluded pairs of SNPs where the minor two-locus genotype class contained

few observations. This was achieved by applying two data filters: first, only SNPs with a minor

allele frequency greater than 0.25 were included, and second all SNP-by-SNP combinations with

fewer than four accessions in the minor genotype class were removed. In the epistatic analyses,

we accounted for population structure by performing the GWA analysis using a linear mixed

model including the genomic kinship matrix. Significance for the pairs were determined using a

standard Bonferroni-correction for the number of independent pair-wise tests across the

independent linkage blocks across the *A. thaliana* genome [41]. In total, the epistatic analysis

included approximately 78 million such independent tests. Six SNP-pairs, representing seven

genomic locations and four unique combinations of loci, passed this threshold (Table 2). Two

pairs were detected twice by associations to tightly linked SNPs located 1.5 kb and 505 bp apart

(Chromosome 3 at 9,272,294 and 9,273,674 bp - SNPs 3_9272294/ 3_9273674 - and

Chromosome at 15,862,026 and 15,862,525 bp - SNPs 5_15862026/5_15862525, respectively),

and one SNP (Chromosome 3 at 66,596 bp - SNP 3_66596) was part of two unique interacting

pairs (Table 2). The other two SNP-pairs contained independent SNPs ($R^2 < 0.8$).

Table 2. Exhaustive two-dimensional GWAS-scan for epistasis identifies six significant

interactions, representing four unique pairs, associated with the root length mean.

Pair	SNP1[1]	SNP2[1]	Interaction (mm)	p-value[2] (GWA)	p-value[3] (SS)
1	rs346846662 (3_66596)	rs347034954/rs347048583 (3_9273674/3_9272294)	0.55	$1.5×10^{-10}$	0.003
2	rs347005428 (1_17257526)	rs346595296/rs346511785 (5_15862026/5_15862525)	0.63	$1.7×10^{-10}$	0.28
3	rs346846662 (3_66596)	rs347177728 (5_18241640)	0.63	$3.0×10^{-10}$	0.37
4	rs347374065 (3_10891195)	rs347377866 (5_1027939)	0.63	$2.3×10^{-10}$	0.95

11 [1]Name of associated SNP(s) at the epistatic locus as on the Affymetrix single nucleotide polymorphism (SNP) chip

12 (our abbreviation as Chromosome_PositionInBp); [2]p-value for the interaction effect of the two loci in the epistatic

13 GWA analysis [40]; [3]Probability that the pair is a false-positive when accounting for the small sample-size (SS)

14 [42].

Explorations of epistatic genotype-phenotype maps revealed an epistatic cancellation of the

additive genetic variance

The four SNP pairs that were significant in the epistatic GWA (Table 2; Fig 2a) explained no

additive variance, but together 51% of the phenotypic variance of mean root length (95% CI:

31.2% - 72.4% from bootstrap). To explore the origin of the low additive variance in the

presence of the revealed epistasis, we estimated the multi-locus genotype-phenotype (G-P) maps

and their respective genotype frequencies in the analyzed population. The G-P maps were highly

related for the four pairs: the minor allele double-homozygote was always associated with the

longest mean root length and the shortest mean root lengths were associated with the genotypes

combining one major- and one minor-allele homozygote (Fig 2b; Fig S2). Although this type of

epistasis will lead to marginal additive genetic effects for the two loci for many allele-

frequencies, there are also combinations of allele-frequencies where the epistasis will cancel the

additive genetic variance completely. We illustrate this epistatic cancellation of the additive

genetic variance for the G-P map and allele-frequencies of the most significant pair in Fig 2c.

For this pair, there are minimal differences in the average (marginal) mean root lengths for the

two genotypes at the individual loci, and hence a minor contribution to the additive genetic

variance. Their joint contribution to the phenotypic variance, however, is large due to the large

differences in root lengths among the four two-locus genotype classes. The epistatic GWA

analysis is able to reveal these pairs in this population as the differences between these genotypes

contribute to the epistatic genetic variance instead.

Evaluating the robustness of statistical epistatic associations to a small population size

We next applied a conservative genome-wide test to estimate the risk that the epistatic GWA

associations were false positives due to the small population size. A pseudo-marker was created

for each pair of SNPs in the epistatic scan, resulting in an epistatic pseudo-genome with 78

million markers. Given the epistatic effect estimated for the four pairs that reached genome-wide

significance in the epistatic GWA, we computed the genome-wide probability of these pairs

being false-positives using the R/p.exact package [42], assuming conservatively that all the

pseudo-markers are independent [42] (Table 3). The pair 3_10891195/5_1027939 was found to

be sensitive to population size (p = 0.94). The pairs 1_17257526/5_15862026 and

3_66596/5_18241640 were less sensitive (p = 0.28 and p = 0.37, respectively), and hence there is

a 90% probability that at least one of these two signals is not false. The last pair

3_66596/3_9273674 was not sensitive to population-size (p = 0.003). We conclude that it is

essential to consider the effect of population-size when interpreting the results from GWA

analyses in small populations. In our particular case, half of the original associations were at risk

of being false-positives due to small populations size.

Table 3. Estimation of the risk that epistatic pairs identified in the GWA analysis are due

to high-order LD to unobserved functional variants ("apparent epistasis").

Using the whole-genome re-sequencing data from the reference 1001-genomes *A. thaliana*

collection, we estimated the high-order LD (r^2) between the four pseudo-markers representing

our epistatic pairs and all sequencing variants that were not genotyped in our GWA analysis. A

bootstrap approach was used to estimate the mean and max r^2 between the epistatic pseudo-

markers and all the genome-wide sequencing-variants. The risk that an association might be due

to "apparent epistasis" was calculated as the risk of observing an $r^2_{max} > 0.8$ in this analysis.

	Significant pairs in two-dimensional epistatic GWA analysis			
Locus 1[1]	3_66596		1_17257526	3_10891195
Locus 2[1]	3_9273674	5_18241640	5_15862026	5_1027939
r^2_{mean}	0.01 ± 0.001	0.01 ± 0.001	0.01 ± 0.003	0.01 ± 0.001
r^2_{max}	0.52 ± 0.10	0.66 ± 0.15	0.78 ± 0.11	0.78 ± 0.09
$P(r^2_{max} > 0.8)$	0.002	0.18	0.42	0.42
MAF_{PM}	0.065	0.043	0.043	0.043

[1]Locus detected as part of epistatic pair named as Chromosome_PositionInBp ; r^2_{mean}: mean high-order LD between

epistatic pseudo-marker and genome-wide sequence variants ± standard error; r^2_{max}: maximum high-order LD

between epistatic pseudo-marker and $_{genome}$-wide sequence variants ± standard error; $P(r^2_{max} > 0.8)$: probability of

observing a high-order LD larger then 0.8 in a random sample of 93 accessions with MAF_{PM} for the epistatic

pseudo-marker; MAF_{PM}: minor-allele frequency for the epistatic pseudo-marker, i.e. the frequency of the minor-

allele double-homozygote for the epistatic pair.

Quantifying the risk of "apparent epistasis" for significant statistical epistatic associations

For "apparent epistasis" to be present, the hidden functional variant needs to be in high linkage

disequilibrium (measured as r^2) with one of the "epistatic pseudo-markers" described above.

Therefore, "apparent epistasis" involving physically unlinked markers is only expected for very

strong mutations. However, if any such hidden variant exists, by definition neither its genetic

effect nor its linkage disequilibrium to the "epistatic pseudo-markers" can be estimated from the

GWA datasets. However, if a GWA study is performed in a species with sequenced reference

populations, the genomic data from that population can be used to estimate the risk of observing

a high r^2 between the tested markers and other polymorphisms in the genome. Here, we used the

1001 Genomes Project *A. thaliana* genomes [22] to perform this analysis for the four significant

pairs in the epistatic GWA analysis. We defined bi-allelic "epistatic pseudo-markers" by

assigning individuals with the two-locus minor-allele homozygote genotype the minor pseudo-

marker genotype and all others the major pseudo-marker genotype. By using a bootstrap analysis

based on samples of 93 accessions from the 728 whole-genome re-sequenced accessions from

the 1001 Genomes Project data, we estimated the probability of observing a high r^2 ($r^2 > 0.8$)

between any individual SNP in the re-sequencing data and the "epistatic pseudo-markers" for the

four pairs. This analysis revealed that the "epistatic pseudo-markers" corresponding to the

identified pairs have very low r^2 to most of the hidden variants (mean $r^2 = 0.01$). There is a very

low risk to observe a high r^2 ($r^2 > 0.8$) to any hidden variant in our population for the most

strongly associated pair ($p < 0.005$). For the other three pairs, this risk is still low ($p = 0.18$ -

0.42) but sufficiently large to indicate that the results should be interpreted with caution (Table

3).

Functional exploration of candidate genes in the associated regions using T-DNA insertion

lines

Our statistical analyses suggest that half of the pairs with genome-wide significant statistical

epistasis might be false-positives due to the small population-size. As only four pairs were found,

we decided to also perform an initial experimental and bioinformatics analysis that included all

these pairs to evaluate whether the statistical risk of false-positives was reflected also by the

presence of known, or novel, functional candidate genes for root-length in the associated regions.

We therefore first extracted the genotypes for 13 million genome-wide SNPs from the whole-

genome re-sequencing data from the 1001 Genomes Project [43]. SNPs that were in LD ($r^2 >$

0.8) and located within 5kb upstream or downstream of the leading SNP in the epistatic GWA

were considered further as candidate functional mutations. In Table 4, we list the predicted

functional effects of the SNPs that fulfill these criteria, together with their locations, for each

pair. In total, thirteen genes and two transposons were found in LD with the leading SNPs in the

seven regions detected in the epistatic GWA analysis to root length.

Table 4. Summary of identified polymorphisms in LD with the leading SNPs from the

epistatic GWA analysis.

Four unique interacting pairs of loci were significantly associated with root length (Table 2).

Here, the genes located within 1000bp of the leading SNPs, together with the mutational effects

of the identified polymorphisms, are listed.

Chr	SNP position (bp)	In pair	Locus	# Mutations (# in LD)[1]	Mutation Effects
3	66596	1, 4	At3g01180	1 (0)	1 upstream
			At3g01185	1 (0)	1 syn
3	9,272,294	1	At3g25520	4 (21)	1 intron, 3 upstream
	9,273,674		At3g25530	14 (21)	1 mis, 1 splice, 1 syn, 2 intron, 9 upstream
			At3g25540	17 (21)	1 mis, 4 syn, 2 intron, 5 5'UTR, 5 upstream
			At3g25545	3 (21)	3 upstream
5	18,241,640	4	At5g45120	6 (5)	2 mis, 4 syn
1	17,257,526	2	At1g46624	1 (0)	1 non-coding exon
5	15,862,026	2	At5g39610	2 (4)	6 upstream[3]
	15,862,525		At5g39620	2 (4)	6 downstream[3]
3	10,891,195	3	At3g28865	4 (7)	4 non-coding exon
			At3g28870	2 (7)	1 mis, 1 syn
			At3g28880	3 (7)	3 upstream
5	1,027,939	3	At5g03840	1 (1)	1 upstream
			At5g03850	1 (1)	1 downstream

[1]The number of polymorphisms associated with each listed gene, with the number of these polymorphisms that are

in LD with the leading SNP in parenthesis. Some mutations are associated with more than one gene. [2]Abbreviations

for mutational effects: syn = synonymous variant; mis = missense variant; splice = splice site variant. [3]SNPs more

than 1000bp away from the nearest gene

Further, we identified T-DNA insertion lines for 12/15 genes and transposons comprising the

seven statistically epistatic loci and experimentally evaluated them for effects on root length. In

Table 5, we present the genes for which the T-DNA insertions significantly altered the root

length. Results for all evaluated T-DNA lines are presented in the Supplementary Text. For three

of these loci (*NAC6*, *TFL1*, *At2g28880*), we are the first to show that these underlying genes

contribute to root growth, and for one locus (*ATL5*), our results confirm previous findings.

T-DNA mutant analysis reveals four functional candidate genes for root development in

regions detected using an epistatic genome-wide association analysis

Decreased root length in ATL5/OLI5/PGY3 *T-DNA insertion mutants*

Two linked SNPs on chromosome 3 (3_9272294 and 3_9273674; Table 2) were associated with

A. thaliana root length through an epistatic interaction with a distal SNP on the same

chromosome (3_66596). Twenty-one SNPs located in four genes were in high LD with the two

linked SNPs (Table 4; Table 5). For three of these (At3g25520, At3g25530 and At3g25540), we

were able to obtain insertion mutants and scored their mean root length compared to wild-type

(*Col-0*) controls. The insertion mutant for At3g25520 had significantly shorter roots than wild

type ($p = 1.2 \times 10^{-4}$; Tukey's post-hoc test); the insertion mutants for At3g25530 and At3g25540

did not differ significantly ($p = 1.00$ and $p = 0.23$, respectively) from wild type (Fig 3).

Table 5. Function and expression patterns of genes in LD with leading SNP from epistatic GWAS analysis. The gene names or proposed function is listed for all genes harboring polymorphisms in LD with the leading epistatic SNPs in the whole genome interaction analysis.

Chr	Position (bp)	In pair	Locus	Gene/Putative function	Highest expression tissue[1]
3	66596	1, 4	At3g01180	*SS2*	cauline leaves
			At3g01185	unknown	NA
3	9,272,294/	1	At3g25520	*ATL5*	apical meristem, seedling root
	9,273,674		At3g25530	*LOH1*	rosette leaves
			At3g25540	*GR1*	seedling root, imbibed seed
			At3g25545	unknown	seedling root (vasculature), imbibed seed
5	18,241,640	4	At5g45120	eukaryotic aspartyl protease family protein	adult root (meristem), imbibed seed
1	17,257,526	2	At1g46624	gypsy-like retrotransposon	NA
5	15,862,026/	2	At5g39610	*NAC6*	seedling root (vasculature)
	15,862,525		At5g39620	RAB GTPase homolog G1	adult root vasculature, mature seed
3	10,891,195	3	At3g28865	LINE retrotransposon	NA
			At3g28870	histone deacetylase	mature seed
			At3g28880	Ankyrin repeat family protein	NA
5	1,027,939	3	At5g03840	*TFL1*	adult root vasculature (mature zone)
			At5g03850	nucleic acid-binding, OB-fold-like protein	seedling root, apical meristem

[1] Locations of high expression were obtained from the BAR eFP *Arabidopsis* Browser (Winter et al. 2007).

At3g25520 encodes ATL5/OLI5/PGY3, a 5S rRNA binding protein whose promoter is highly accessible in seedling roots [44], corresponding to expression in roots [45]. Our finding agrees with earlier studies, in which the loss-of-function mutant *oli5-1* exhibits shorter primary roots

than wild-type [33]. In the tested accessions, three SNPs were found in the *ATL5* promoter and a

fourth in an intron, suggesting variable transcriptional regulation across accessions.

A novel role for NAC6 in the regulation of root length

Two linked SNPs on chromosome 5 (5_15862026 and 5_15862525; Table 2) were detected by

their statistical epistatic interaction with an SNP on chromosome 1 (17,257,526 bp - SNP

1_17257526; Table 2). Both of these linked SNPs, as well as the other SNPs in high LD, were

intergenic with the nearest gene, *NAC6* (At5g39610; Table 5), which is located 1.9 kb away.

NAC6 is a transcription factor regulating leaf senescence [46] and is highly expressed in

senescing leaves and in maturing seeds [45]. We phenotyped the available At5g39610 insertion

mutant for *NAC6* and found a significant decrease in mean root length ($p = 5.82 \times 10^{-10}$; Tukey's

post-hoc test; Fig 4), strongly implicating *NAC6* in the regulation of root length.

The previously uncharacterized gene At3g28880 contributes to root length

A locus on chromosome 3 (10,891,195 - SNP 3_10891195; Table 2) was identified due to its

statistical epistatic interaction with a locus on chromosome 5 (1,027,939 bp - SNP 5_1027939;

Table 2). SNPs in LD with 3_10891195 were found in three genes: At3g28865, At3g28870, and

At3g28880 (Table 5). None of these genes previously had been associated with root length. By

examining available insertion mutants, we established a novel role for the gene At3g28880 in

root development; its mutant showed significantly decreased mean root length compared to wild-

type controls ($p = 4.5 \times 10^{-10}$; Fig 4), whereas At3g28865 did not. At3g28880 encodes a

previously uncharacterized ankyrin family protein. There is no information on developmental

expression patterns for this gene; although the gene's putative regulatory region is accessible in

seeds and seedling tissue [44]. Across the tested accessions, a number of promoter-proximal

SNPs were found in At3g28880, possibly altering gene regulation.

Mutant TFL1 increases mean root length

The SNP 5_1027939 was detected via its epistatic interaction with the At3g28880 locus on

chromosome 3 (Table 2). This polymorphism and the only other polymorphism in LD are both

intergenic between the genes At5g03840 and At5g03850 (Table 5). Phenotyping the insertion

mutants for these genes revealed that the mutant for At5g03840, which encodes *TFL1*, showed

significantly longer roots than wild-type controls ($p = 5.7 \times 10^{-10}$; Tukey post-hoc test; Fig 4),

whereas no significant phenotypic effect was found for the At5g03850 mutant. Interestingly,

these results implicate *TFL1* as a repressor of root growth.

TFL1's role in floral initiation and morphology are well-established in many plant species [47].

It also plays a role in regulating protein storage in vegetative tissues, such as roots or seeds [48].

TFL1 is highly expressed in both adult and seedling roots [45]. Our finding that *TFL1* controls

root length is particularly poignant as a previous study reported no root length defects for a

different *tfl1* mutant with multiple other pleiotropic phenotypes [49]. Our finding extends the

functional reach of this multi-functional regulator.

Discussion

This study was designed to identify genes regulating seedling root length. The narrow-sense

heritability for this trait was not significantly different from zero in our population, whereas there

was significant broad-sense heritability (Table 1). Notably, we observed a very large contribution

of epistatic variance to the phenotypic variation in mean root length. With these findings in hand,

we completed a comprehensive set of GWA analyses to identify additive and epistatic loci

affecting this trait.

Consistent with the low narrow-sense heritability for mean root length, we were unable to

identify any additive loci that reached the Bonferroni-corrected significance threshold in our

analyses. By accounting for the contribution of pairwise epistatic interactions to the genetic

variance of root length, however, we identified seven loci involved in four unique epistatically

interacting pairs of loci that together explained a large fraction of the phenotypic variance in the

population (Table 2). One of these pairs is highly significant also when accounting for the small

population-size, whereas there is a risk that two of the three remaining pairs are false-positives

due to this. We also showed that it is unlikely that the statistical epistatic interaction for the most

significant pair would result from "apparent epistasis" due to a major, hidden variant in the

genome. This risk is, however, not negligible for the three less significant pairs. Based on this,

we conclude that despite the fact that much of the genetic variance present in this population can

be explained by the genome-wide significant pairwise epistasis detected amongst these loci,

additional evidence is needed before any firm conclusions can be drawn based on the other three

epistatic pairs detected in the GWA analysis.

All four pairs detected in the initial epistatic GWA display an epistatic cancellation of additive

genetic variance (Fig 2c). The presence of this type of genetic architecture for root-length in this

population explains the low narrow-sense heritability in the population and why no loci could be

found in the standard GWA analyses. Although further studies are needed to explore how

general this phenomenon is, our result contrasts the common view in quantitative genetics that

most genetic variance for complex traits is expected to be additive [5]. It further illustrates the

importance of taking a richer modeling approach in GWA analyses to improve the sensitivity as

a too strict focus on additive effects in the initial statistical analyses might lead to important

findings being missed and lead to an underestimation of the full adaptive potential due to the

epistatic suppression of selectable additive genetic variance [10–12].

Our finding of extensive statistical epistasis in this study should not be interpreted as evidence

for root development being a biological trait with exceptionally high levels of biological

interactions. Rather it is important to notice that, although the levels of statistical epistasis

detected here originate in the epistatic genotype-phenotype maps for these pairs, the

exceptionally high levels of epistatic variance in the population is even more due to the allele-

frequencies for the involved loci in the population (Fig 2c). Hence if the loci displaying the G-P

map in Fig 2c instead, for example, would be present at intermediary and equal frequencies, they

would contribute high levels of additive genetic variance instead. Consequently, our results is an

important reinforcement of the fact that the levels of epistatic variance for a trait in a population

will be a dynamic outcome of changes in genotype-frequencies over time even though the

underlying genotype-phenotype map might be constant [7–13].

In practice, however, neither allele frequencies nor genotype-phenotype maps will be known for

the loci contributing to a particular trait. As illustrated by our findings for root length in this

population, it is important to utilize GWA analysis methods that account for both additive and

non-additive genetic variance to detect the contributing loci. Once this is done, further theoretical

and experimental work is needed to identify the underlying genotype-phenotype maps and

dissect the molecular underpinnings of the statistical associations regardless of whether these

were detected mainly via their contributions to the additive and/or epistatic genetic variance. We

here propose how to statistically quantify the risks of each inferred epistatic GWA association to

guide further experimental work based on these findings. We find that there were statistical

reasons to interpret three of the detected epistatic pairs with caution, both due to an increased

risk of false-positives due to the small population-size and an inflated risk for "apparent

epistasis". Three novel functional candidate genes were, however, identified in loci detected as

part of these associations. This illustrates clearly that the ultimate contribution from a study will

rely on the combined use of appropriate statistical analyses to quantify the confidence in a

particular locus and an in-depth experimental dissection of the detected associations before a

conclusion can be reached about which loci contribute to a particular trait.

Functional dissection of epistasis, however, is a daunting task in natural populations and

therefore model organisms are an indispensable resource in this work. A first step in taking on

this challenge is to reduce the list of potential candidate genes and mutations for further, in-depth

functional explorations. Here, we use mutational analysis of the genes that harbor SNPs in LD

with the associated SNPs in the epistatic GWA analysis to identify the most likely candidate

genes for the inferred associations. We tested six of the seven epistatic regions with available T-

DNA insertion mutants. For four of the loci, we succeeded in identifying candidate genes

affecting root length, three of which were newly implicated in root development. The majority of

single-gene mutations affects do not affect root length [4,35,50]. Our results demonstrate the

benefits of using statistical epistatic analyses in combination with experimental analysis of T-

DNA insertion mutants to identify novel functional candidate genes involved in root

development and provide a basis for future detailed studies of the complex gene networks

involved in this trait (Fig 4).

It will be a major endeavor to functionally explore the biological mechanisms contributing to the

statistical epistatic associations. None of the genes affecting root development contain non-

synonymous mutations with likely phenotypic effects. Also, none of these genes have previously

been described to interact with their respective epistatic locus. Future functional studies will have

to explore the effects of the observed regulatory SNPs in transgenic plants in several

backgrounds to study the whether these statistical interactions are the result of an underlying

biological interaction or not. This endeavor will require significant time and resources, yet we

believe that *A. thaliana* root length represents an excellent model trait for future in-depth studies

of this type of complex signals arising from GWA studies.

Materials and Methods

Phenotyping

In order to reduce environmental variation among accessions, eighteen individuals from each of

the 93 accessions studied in Atwell et al (2010) (Table S1) were vernalized at 4° as seedlings for

six weeks to synchronize growth and flowering. The five most developmentally advanced

seedlings from each accession were then transferred to soil in a randomized design. Plants were

grown in long-day conditions at 22°. Flats were rotated three times per week to reduce position

effects on plant development. Seeds were collected over a period of three months as the plants

dried. Equal numbers of seeds were pooled from 3-5 parent plants to reduce the environmental

contribution of plant placement across flats of plants. Ethanol sterilized seeds were planted on $1\times$

Murashige and Skoog (MS) basal salt medium supplemented with $1\times$ MS vitamins, 0.05% MES

(wt/vol), and 0.3% (wt/vol) phytagel in a semi-randomized design with n = 70 per accession.

Four sets of twenty-three accessions plus a control accession (Col-0) made up a set. Each set was

replicated three times providing the standard error for variance, with a total n = 210 planted for

each accession. The seeds were stratified at 4° for three days and then grown for seven days in

darkness with the plates in a vertical position. A photograph of each plate was taken, and root

length was measured manually using the 'freehand' function in ImageJ1.46r [51]. Non-

germination, missing organs, and delayed development were noted.

Data standardization

Thirty-one seedlings with hypocotyls less than 5mm were removed because it is likely that

germination was severely delayed [34]. Between-set differences were corrected by subtracting

the difference between Col-0 in each replicate and set and the global mean for Col-0. Systematic

differences between replicates were still present and corrected within each set by using the

residuals from a model in which root length was explained by the within-set replicates.

Heritability estimation

Heritability of root length variance and mean were estimated using the repeated measures with

genotype as a fixed effect using ANOVA. To parse the contribution of additive and epistatic

effects, a linear mixed model including both additive and epistatic effects as random effects was

fitted using the R/hglm package [37], i.e.

$$y = \mu + a + b + e,$$

where

$$a \sim N(0, G\sigma_a^2)$$

3 are the individual-level additive genetic effects, and G is the genomic kinship matrix;

$$b \sim N(0, G \circ G\sigma_b^2)$$

are the individual-level epistatic effects, and G is the genomic kinship matrix; *e* are independent

and normally distributed residuals. The narrow sense heritability was estimated as the ratio of the

additive genetic variance to the total phenotypic variance, and the broad sense heritability was

estimated as the ratio of the sum of the additive and epistatic variance to the total phenotypic

variance.

Genome-wide association analysis

Using the transformed mean and the genotypes for the 93 accessions from Atwell et al. (2010),

we performed a series of analyses to detect marker-trait associations. For the standard additive

GWA analysis with population structure controlled, we used the function egscore() from

R/GenABEL [38]. The functions bigRR() and bigRR_update() in the R/bigRR package [39]

were used to perform a GWA analysis, in which all genome-wide SNP effects are modeled

simultaneously as random effects, and the effects were estimated via the generalized ridge

regression method HEM. The maximum absolute effect sizes in 1000 permutations were

extracted to determine an empirical 5% genome-wide significance threshold (p = 0.0020).

To screen the genome for pairs of epistatic SNPs, we used the PLINK –epistasis procedure [40]

that is based on the model:

1
$$Y \sim b_0 + b_{1.A} + b_{2.B} + b_{3.AB} + e,$$

which considers allele by allele epistasis in $b_{3.AB}$. In the analysis, all possible pairs of SNPs with

a minor allele frequency > 25% were tested.

We filtered out the pairs where there were fewer than four accessions in the minor two-locus

genotype-class. Further, all combinations in which the p-value was > 1×10^{-8} were also removed

as it was considered unlikely that they would be significant after correction for population

structure and applying a multiple-testing corrected significance threshold. For the remaining

pairs, a linear mixed model including fixed, additive and epistatic effects, as in the PLINK based

initial scan, were fitted together with a kinship correction for population stratification, as in the

single locus analyses, using the package R/hglm. We derived a multiple-testing corrected

significance threshold for this epistatic analysis (p = 3.2×10^{-10}) by estimating the number of

independent tests based on the number of estimated LD blocks in the genome. For this, we used

the fact that the *A. thaliana* genome is approximately 125Mb with average LD-blocks of about

10kb (12,500; [41]), and then applied a Bonferroni correction for the 78 million independent

tests performed. Seventy-eight million double-recessive pseudo markers were created, and the

R/p.exact package [42] was used to double-assess the exact false discovery rate of each detected

epistatic pair based on its estimated epistatic effect size and allele frequency. Using data on the

sixty-three accessions from the GWAs that were unambiguously identified in the 1001 Genomes

data [22], we then identified additional SNPs in LD with the leading SNP using the function *LD*

in the *genetics* package in R across a 10kb region around the marker.

High-order LD between epistatic pseudo markers and whole-genome sequencing variants

From 728 whole-genome sequences in *A. thaliana* 1001 Genomes Project

(http://www.1001genomes.org), we extracted all the sequencing variants. All four significant

pseudo-markers had at least four accessions with the double-recessive genotype (i.e. the minor

allele for the epistatic pseudo-marker) in the 728 sequenced accessions. For each of the pseudo-

markers, we therefore sampled 93 accessions from the 728 sequenced ones, keeping the same

allele frequency as for the pseudo-marker representing the epistatic pair detected in the GWA

dataset, and obtained the genome-wide, high-order LD r^2 distribution across all the sequenced

variants. This procedure was repeated 50 times for each pseudo-marker, and we calculated the

estimates and standard errors of the r^2_{mean}, r^2_{max} and estimated the risk for "apparent epistasis" as

the probability of observing an $r^2_{max} > 0.8$ in a particular random sample.

Examination of SNP-associated genes

The Ensembl Variant Effect Predictor based on the TAIR10 release of the *A. thaliana* genome

was used to determine the effects of the leading SNPs and the SNPs in high LD with them.

Genes were considered as candidates if they were within 1kb of a variant. Expression of the

candidate genes was determined using the BAR eFP Arabidopsis browser [45].

Validation of interacting loci

T-DNA lines were obtained for the candidate genes (Table S2). Root length was ascertained as

described above (n = 20). Tukey's HSD post hoc test was used to compare root lengths between

the T-DNA lines and the wild-type accession (*Col-0*).

Acknowledgments

We thank Mats Pettersson and Ronnie Nelson for useful discussions and Karla Schultz and

members of the Queitsch lab for experimental assistance.

References

1. Huang W, Richards S, Carbone MA, Zhu D, Anholt RRH, Ayroles JF, et al. Epistasis

dominates the genetic architecture of Drosophila quantitative traits. Proc Natl Acad Sci U

S A. 2012;109: 15553–9. doi:10.1073/pnas.1213423109

2. Mackay TFC. Epistasis and quantitative traits: using model organisms to study gene-gene

interactions. Nat Rev Genet. Nature Publishing Group, a division of Macmillan Publishers

Limited. All Rights Reserved.; 2014;15: 22–33. Available:

http://dx.doi.org/10.1038/nrg3627

3. Nelson RM, Pettersson ME, Li X, Carlborg Ö. Variance Heterogeneity in Saccharomyces

cerevisiae Expression Data: Trans-Regulation and Epistasis. PLoS One. Public Library of

Science; 2013;8: e79507.

4. Queitsch C, Carlson KD, Girirajan S. Lessons from model organisms: phenotypic

robustness and missing heritability in complex disease. PLoS Genet. 2012/11/21 ed.

Public Library of Science; 2012;8: e1003041. doi:10.1371/journal.pgen.1003041

PGENETICS-D-12-00749 [pii]

5. Hill WG, Goddard ME, Visscher PM. Data and theory point to mainly additive genetic

variance for complex traits. Mackay TFC, editor. PLoS Genet. Public Library of Science;

2008;4: e1000008. doi:10.1371/journal.pgen.1000008

6. Visscher PM, Brown MA, McCarthy MI, Yang J. Five years of GWAS discovery. Am J

Hum Genet. 2012;90: 7–24. doi:10.1016/j.ajhg.2011.11.029

7.   Goodnight CJ. Population Differentiation and the Transmission of Density Effects between Generations. Evolution (N Y). Society for the Study of Evolution; 1988;42: 399–403 CR – Copyright © 1988 Society for th. doi:10.2307/2409244

8.   Goodnight CJ. Quantitative trait loci and gene interaction: the quantitative genetics of metapopulations. Heredity (Edinb). Nature Publishing Group; 2000;84: 587–598.

9.   Whitlock MC, Phillips PC, Moore FB-G, Tonsor SJ. Multiple fitness peaks and epistasis. Annu Rev Ecol Syst. JSTOR; 1995; 601–629.

10. Eitan Y, Soller M. Selection induced genetic variation. Evolutionary theory and processes: Modern horizons. Springer; 2004. pp. 153–176.

11. Le Rouzic A, Siegel PB, Carlborg Ö. Phenotypic evolution from genetic polymorphisms in a radial network architecture. BMC Biol. BioMed Central Ltd; 2007;5: 50.

12. Le Rouzic A, Carlborg Ö. Evolutionary potential of hidden genetic variation. Trends Ecol Evol. Elsevier; 2008;23: 33–37.

13. Monnahan PJ, Kelly JK. Epistasis Is a Major Determinant of the Additive Genetic Variance in Mimulus guttatus. Carlborg Ö, editor. PLOS Genet. 2015;11: e1005201. doi:10.1371/journal.pgen.1005201

14. Carlborg O, Haley CS. Epistasis: too often neglected in complex trait studies? Nat Rev Genet. Nature Publishing Group; 2004;5: 618–25. doi:10.1038/nrg1407

15. Mackay TF. The genetic architecture of quantitative traits. Annu Rev Genet. Annual Reviews 4139 El Camino Way, P.O. Box 10139, Palo Alto, CA 94303-0139, USA; 2001;35: 303–39. doi:10.1146/annurev.genet.35.102401.090633

16. Rowe HC, Hansen BG, Halkier BA, Kliebenstein DJ. Biochemical networks and epistasis shape the Arabidopsis thaliana metabolome. Plant Cell. 2008;20: 1199–216. doi:10.1105/tpc.108.058131

17. Pyun J-A, Kim S, Cha DH, Kwack K. Epistasis between polymorphisms in TSHB and ADAMTS16 is associated with premature ovarian failure. Menopause (New York, NY). 2013;

18. Sapkota Y, Mackey JR, Lai R, Franco-Villalobos C, Lupichuk S, Robson PJ, et al. Assessing SNP-SNP interactions among DNA repair, modification and metabolism related pathway genes in breast cancer susceptibility. PLoS One. Public Library of Science; 2013;8: e64896.

19. Vanhaeren H, Gonzalez N, Coppens F, De Milde L, Van Daele T, Vermeersch M, et al. Combining growth-promoting genes leads to positive epistasis in Arabidopsis thaliana. Elife. eLife Sciences Publications Limited; 2014;3: e02252. doi:10.7554/eLife.02252

20. Carlborg Ö, Jacobsson L, Åhgren P, Siegel P, Andersson L. Epistasis and the release of genetic variation during long-term selection. Nat Genet. Nature Publishing Group; 2006;38: 418–420.

21.   Carlborg Ö, Andersson L, Kinghorn B. The use of a genetic algorithm for simultaneous

mapping of multiple interacting quantitative trait loci. Genetics. Genetics Soc America;

2000;155: 2003–2010.

22.   Cao J, Schneeberger K, Ossowski S, Günther T, Bender S, Fitz J, et al. Whole-genome

sequencing of multiple Arabidopsis thaliana populations. Nat Genet. Nature Publishing

Group; 2011;43: 956–963.

23.   Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al. The

Drosophila melanogaster genetic reference panel. Nature. Nature Publishing Group;

2012;482: 173–178.

24.   Gibson G. Rare and common variants: twenty arguments. Nat Rev Genet. Nature

Publishing Group; 2012;13: 135–145. doi:10.1038/nrg3118

25.   Hemani G, Shakhbazov K, Westra H-J, Esko T, Henders AK, McRae AF, et al. Detection

and replication of epistasis influencing transcription in humans. Nature. Nature Publishing

Group; 2014;

26.   Horton MW, Hancock AM, Huang YS, Toomajian C, Atwell S, Auton A, et al. Genome-

wide patterns of genetic variation in worldwide Arabidopsis thaliana accessions from the

RegMap panel. Nat Genet. Nature Publishing Group; 2012;44: 212–216.

27.   Atwell S, Huang YS, Vilhjalmsson BJ, Willems G, Horton M, Li Y, et al. Genome-wide

association study of 107 phenotypes in Arabidopsis thaliana inbred lines. Nature.

2010/03/26 ed. 2010;465: 627–631. doi:10.1038/nature08800 nature08800 [pii]

28. Filiault DL, Maloof JN. A genome-wide association study identifies variants underlying

the Arabidopsis thaliana shade avoidance response. PLoS Genet. Public Library of

Science; 2012;8: e1002589. doi:10.1371/journal.pgen.1002589

29. Meijón M, Satbhai SB, Tsuchimatsu T, Busch W. Genome-wide association study using

cellular traits identifies a new regulator of root development in Arabidopsis. Nat Genet.

Nature Publishing Group, a division of Macmillan Publishers Limited. All Rights

Reserved.; 2014;46: 77–81. doi:10.1038/ng.2824

30. Slovak R, Göschl C, Su X, Shimotani K, Shiina T, Busch W. A Scalable Open-Source

Pipeline for Large-Scale Root Phenotyping of Arabidopsis. Plant Cell. 2014;26: 2390–

2403. doi:10.1105/tpc.114.124032

31. Rosas U, Cibrian-Jaramillo A, Ristova D, Banta JA, Gifford ML, Fan AH, et al.

Integration of responses within and across Arabidopsis natural accessions uncovers loci

controlling root systems architecture. Proc Natl Acad Sci U S A. 2013;110: 15133–8.

doi:10.1073/pnas.1305883110

32. Wood AR, Tuke MA, Nalls MA, Hernandez DG, Bandinelli S, Singleton AB, et al.

Another explanation for apparent epistasis. Nature. Nature Publishing Group; 2014;514:

E3–E5.

33. Fujikura U, Horiguchi G, Ponce MR, Micol JL, Tsukaya H. Coordination of cell

proliferation and cell expansion mediated by ribosome-related processes in the leaves of

Arabidopsis thaliana. Plant J. Wiley Online Library; 2009;59: 499–508.

34. Sangster TA, Salathia N, Undurraga, Milo R, Schellenberg K, Lindquist SL, et al. HSP90

affects the expression of genetic variation and developmental stability in quantitative

traits. Proc Natl Acad Sci U S A. 2008/02/22 ed. 2008;105: 2963–2968.

doi:10.1073/pnas.0712200105

35. Lempe J, Lachowiec J, Sullivan AM, Queitsch C. Molecular mechanisms of robustness in

plants. Curr Opin Plant Biol. 2013/01/03 ed. 2012; doi:S1369-5266(12)00172-0 [pii]

10.1016/j.pbi.2012.12.002

36. Geiler-Samerotte K, Bauer C, Li S, Ziv N, Gresham D, Siegal M. The details in the

distributions: why and how to study phenotypic variability. Curr Opin Biotechnol.

2013;24: 752–9. doi:10.1016/j.copbio.2013.03.010

37. Rönnegård L, Shen X, Alam M. hglm: A Package for Fitting Hierarchical Generalized

Linear Models. R J. 2010;2.

38. Aulchenko YS, Ripke S, Isaacs A, Van Duijn CM. GenABEL: an R library for genome-

wide association analysis. Bioinformatics. Oxford Univ Press; 2007;23: 1294–1296.

39. Shen X, Alam M, Fikse F, Rönnegård L. A novel generalized ridge regression method for

quantitative genetics. Genetics. Genetics Soc America; 2013;193: 1255–1268.

40. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: a

tool set for whole-genome association and population-based linkage analyses. Am J Hum

Genet. Elsevier; 2007;81: 559–575.

1   41.   Kim S, Plagnol V, Hu TT, Toomajian C, Clark RM, Ossowski S, et al. Recombination and

2   linkage disequilibrium in Arabidopsis thaliana. Nat Genet. Nature Publishing Group;

3   2007;39: 1151–1155.

4   42.   Shen X. Flaw or discovery? Calculating exact p-values for genome-wide association

5   studies in inbred populations. bioRxiv. 2015; Available:

6   http://biorxiv.org/content/early/2015/02/17/015339.abstract

7   43.   Ossowski S, Schneeberger K, Clark RM, Lanz C, Warthmann N, Weigel D. Sequencing

8   of natural strains of Arabidopsis thaliana with short reads. Genome Res. Cold Spring

9   Harbor Lab; 2008;18: 2024–2033.

44.   Sullivan AM, Arsovski AA, Lempe J, Bubb KL, Weirauch MT, Sabo PJ, et al. Mapping

and Dynamics of Regulatory DNA and Transcription Factor Networks in A. thaliana. Cell

Rep. 2014;8: 2015–2030. doi:10.1016/j.celrep.2014.08.019

45.   Winter D, Vinegar B, Nahal H, Ammar R, Wilson G V, Provart NJ. An "Electronic

Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data

sets. PLoS One. Public Library of Science; 2007;2: e718.

46.   Aeong Oh S, Park J-H, In Lee G, Hee Paek K, Ki Park S, Gil Nam H. Identification of

three genetic loci controlling leaf senescence in Arabidopsis thaliana. Plant J. Blackwell

Science Ltd; 1997;12: 527–535. doi:10.1046/j.1365-313X.1997.00489.x

47.   Jack T. Molecular and genetic mechanisms of floral control. Plant Cell. 2004;16 Suppl:

S1–17. doi:10.1105/tpc.017038

48.   Sohn EJ, Rojas-Pierce M, Pan S, Carter C, Serrano-Mislata A, Madueño F, et al. The

shoot meristem identity gene TFL1 is involved in flower development and trafficking to

the protein storage vacuole. Proc Natl Acad Sci U S A. 2007;104: 18801–6.

doi:10.1073/pnas.0708236104

49.   Larsson AS, Landberg K, Meeks-Wagner DR. The TERMINAL FLOWER2 (TFL2) Gene

Controls the Reproductive Transition and Meristem Identity in Arabidopsis thaliana.

Genetics. 1998;149: 597–605. Available: http://www.genetics.org/content/149/2/597.full

50.   Bolle C, Huep G, Kleinbölting N, Haberer G, Mayer K, Leister D, et al. GABI-DUPLO: a

collection of double mutants to overcome genetic redundancy in Arabidopsis thaliana.

Plant J. 2013;75: 157–71. doi:10.1111/tpj.12197

51.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image

analysis. Nat Methods. 2012;9: 671–675.

1 **Fig 1. The distributions of within-accession root length mean and variance in a sample of**

2 **93 natural *A. thaliana* accessions.**

3 There is a large dispersion of the root length mean at day seven among the accessions (*n* = 3 x 70

4 dark grown seedlings / genotype).

Fig 2. Four statistical epistatic interactions associated with mean root length.

(A) The x- and y-axes represent the five *A. thaliana* chromosomes. The positions of the seven

SNPs that are part of the four significant interacting pairs (Table 2) are indicated by a black dot.

Solid lines indicate support for an interaction by more than one linked SNP and dotted lines

indicate support by a single SNP. (B) Genotype-phenotype (G-P) map of the root means for the

four genotype combinations constituting the interaction between the SNPs on chromosome 1

(17,257,526 bp) and chromosome 5 (15,862,026 bp). The major allele is indicated by -1, and the

minor allele is indicated by 1. This G-P-map is a representative example for the other pairs in

which the accessions with a minor and major allele have lower phenotypic values than those with

either both major or both minor alleles (Fig S2). (C) G-P map for the most significant epistatic

pair illustrates how this type of epistasis cancels the additive genetic variance at the allele-

frequencies observed for the two loci in the analyzed population of wild-collected *A. thaliana*

accessions.

1   **Fig 3. Four candidate genes exhibit altered root growth in T-DNA insertion lines.** T-DNA

2   lines for candidate genes in six of the seven epistatic regions were tested for root length

3   differences to wild-type (*Col-0*). The gene models for the tested T-DNA insertion lines are

4   pictured. For four of the six loci, a T-DNA line exhibited significantly different root length from

5   *Col-0* (significantly different lines are marked with a * in the figure).

Fig 4. Summary of combined statistical and functional support for loci underlying root

length.

Four epistatic pairs, involving seven unique loci (connected boxes), were identified at a genome-

wide significance-threshold in a two-dimensional genome-wide GWA scan. Green, yellow, and

red lines connect pairs of loci with very low ($p = 0.003$), intermediate ($0.28 < p < 0.37$), and high

($p = 0.95$) risk for the interaction being a false-positive when accounting for population size. The

risk of the statistical epistatic association resulting from high-order LD to an unobserved

functional variant in the genome (i.e. "apparent epistasis") is illustrated by arrow color, in which

yellow indicates an intermediate risk ($0.18 < p < 0.42$) and green a very low risk ($p = 0.0021$).

Green boxes indicate loci that the T-DNA insertion line analyses suggest the named genes to be

involved in root development. When considering the joint statistical and functional results, two

pairs emerge as highly likely true positive two-locus associations: 3_66596/3_9273674 due to

very strong statistical support and one identified functional candidate gene, and

3_10891195/5_1027939 where the identification of functional candidate genes at both loci

suggest that the two-locus association in the original genome-wide scan is true despite the lower

statistical support in after the conservative statistical correction for sample-size. For the other

two pairs, the results are inconclusive. There is strong support for one of the two associated loci

(3_66596 from its statistical interaction with 3_9273674 and 1_17257526 by the detection of the

a functional candidate gene in the T-DNA analysis), but weaker support for the second locus.

Further work is thus needed to conclude whether these pairs represent true positive two-locus

associations, or whether they are false-positives due to the small population-size or high-order

LD ("apparent epistasis") to unknown functional variants.

Supporting information captions

Table S1. *A. thaliana* accessions phenotyped for GWA.

Table S2. T-DNA lines tested for root phenotypes.

Fig S1. No significant associations are found in GWAS for root length using an additive model.

(A) In the GWAS based on an additive model no SNPs reached the Bonferroni threshold. (B) Using a whole-genome generalized ridge regression model, in which SNPs were modeled as random effects, no SNPs reached the Bonferroni threshold.

Fig S2. Similar genotype-phenotype maps for all significantly interacting pairs of loci.

(A-C) Mean root length is displayed for all four two-locus genotype-classes for the four pairs of significantly interacting loci. The major allele is indicated by -1 and the minor allele is indicated by 1. The combination of two minor alleles (1,1) always has the longest root length compared to the other three-allele combinations, which are more similar to one another. Pictured are mean root lengths for the epistatic pairs for SNPs (A) 3_66596 and 3_9272294/3_9273674, (B) 3_66596 and 5_18241640, and (C) 3_10891195 and 5_1027939.