

Title: Secondary contact and local adaptation contribute to genome-wide patterns of clinal variation in *Drosophila melanogaster*

Authors: Alan O. Bergland¹, Ray Tobler^{1,2}, Josefa González^{1,3}, Paul Schmidt⁴, Dmitri Petrov¹

1. Department of Biology, Stanford University, Stanford, CA 94305-5020

2. Institut für Populationsgenetik, Vetmeduni Vienna, Veterinärplatz 1, Vienna A-1210, Austria

3. Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra). Passeig Maritim de la Barceloneta 37-49. 08003 Barcelona, Spain.

4. Department of Biology, The University of Pennsylvania, Philadelphia, PA 19104

Running title: Latitudinal clines in *Drosophila*

Keywords: *Drosophila melanogaster*, adaptation, secondary contact, latitudinal clines, parallelism

Contact information for corresponding author:

Alan O. Bergland

Department of Biology

Stanford University

Stanford, CA 94305-5020

email: bergland@stanford.edu

phone: (650) 736 2249

fax: (650) 723 6132

Abstract

Populations arrayed along broad latitudinal gradients often show patterns of clinal variation in phenotype and genotype. Such population differentiation can be generated and maintained by historical demographic events and local adaptation. These evolutionary forces are not mutually exclusive and, moreover, can in some cases produce nearly identical patterns of genetic differentiation among populations. Here, we investigate the evolutionary forces that generated and maintain clinal variation genome-wide among populations of *Drosophila melanogaster* sampled in North America and Australia. We contrast patterns of clinal variation in these continents with patterns of differentiation among ancestral European and African populations. Using established and novel methods we derive here, we show that recently derived North America and Australia populations were likely founded by both European and African lineages and that this admixture event contributed to genome-wide patterns of parallel clinal variation. The pervasive effects of admixture meant that only a handful of loci could be attributed to the operation of spatially varying selection using an F_{ST} outlier approach. Our results provide novel insight into the well-studied system of clinal differentiation in *D. melanogaster* and provide a context for future studies seeking to identify loci contributing to local adaptation in a wide variety of organisms, including other invasive species as well as some temperate endemics.

Introduction

All species live in environments that vary through time and space. In many circumstances, such environmental heterogeneity can act as a strong selective force driving adaptive differentiation among populations. Thus, a major goal of evolutionary and ecological genetics has been to quantify the magnitude of adaptive differentiation among populations and to identify loci underlying adaptive differentiation in response to ecologically relevant environmental variation.

Phenotypic and genetic differentiation between populations has been examined in a variety of species. In some cases, patterns of differentiation are directly interpretable in the context of circumscribed environmental differences that occur over short spatial scales (Richardson et al. 2014) For instance, differences in salinity experienced by freshwater and marine populations of sticklebacks has led to the identification of key morphological, physiological, and genetic differences between replicate pairs of populations (Kitano et al. 2010; Jones et al. 2012). Similarly, pigmentation morph frequency closely tracks variation in substrate color for a variety of species (reviewed in Gray & McKinnon 2007) thereby providing an excellent opportunity to directly relate environmental variation to phenotypic and genetic differentiation.

Patterns of genetic and phenotypic variation have also been examined in species arrayed along broad geographical transects such as latitudinal clines (Endler 1993). In this paradigm, the goal has often been to identify the phenotypic and genetic basis for adaptation to temperate environments. In certain cases it has been possible to directly relate latitudinal variation in specific environmental variables to aspects of phenotypic and genetic differentiation (e.g., photoperiod and critical photoperiod or flowering time; Bradshaw & Lounibos 1977; Stinchcombe *et al.* 2004). In general, the collinearity of multiple ecological and environmental variables along latitudinal clines often complicates the direct relation of environmental variation to specific phenotypic and genetic

differences. Nonetheless, because many genetically based phenotypic clines within species often mirror deeper phylogenetic differentiation between endemic temperate and tropical species, it is clear that populations distributed along latitudinal clines have adapted to aspects of temperate environments (Gibert et al. 2001).

Latitudinal clines have been extensively studied in various drosophilid species, most notably *Drosophila melanogaster*. Parallel clines in morphological (Rajpurohit et al. 2008; Telonis Scott et al. 2011), stress tolerance (Hoffmann et al. 2002; Schmidt et al. 2005), and life-history traits (Schmidt et al. 2005; Lee et al. 2011) have been identified in *D. melanogaster* populations distributed along multiple continents. These phenotypic clines demonstrate that flies from poleward locales are generally more hardy albeit less fecund, reflecting a classic trade-off between somatic maintenance and reproductive output (Schmidt et al. 2005) that would be alternately favored between populations exposed to harsh winters versus more benign tropical environments. Extensive clinal variation in various genetic markers has also been identified (Sezgin et al. 2004; Hoffmann & Weeks 2007). In some cases clinal genetic variants have been directly linked to clinally varying phenotypes (Schmidt et al. 2008; Paaby et al. 2010; Lee et al. 2013; Paaby et al. 2014a), whereas in other cases parallel clinal variation at genetic markers has been documented across multiple continents (Turner et al. 2008; González et al. 2010; Reinhardt et al. 2014). Taken as a whole, there is abundant evidence that local adaptation to spatially varying selection pressures associated with temperate environments has shaped clinal patterns of phenotypic and genetic variation in *D. melanogaster*.

Demographic forces can also heavily shape patterns of clinal variation (Endler 1993) and the recent demographic history of *D. melanogaster* may be particularly relevant to our understanding of clinal patterns of genetic variation in this species. *D. melanogaster* is an Afro-tropical species (David & Capy 1988) that has colonized the world in the wake of human migration. Population genetic inference suggests that *D. melanogaster* first migrated out of Africa to Eurasia approximately 15,000 years ago (Li & Stephan 2006) and eventually migrated eastward across Asia, arriving to South East Asia approximately 2.5Kya (Laurent et al. 2011). *D. melanogaster* invaded the Americas and Australia within the last several hundred years and likely colonized these continents in their entirety quickly (Bock & Parsons 1981; Keller 2007). Historical records suggest that *D. melanogaster* colonized North America and Australia each in a single event (Bock & Parsons 1981; Keller 2007).

In contrast, however, population genetic (Caracristi & Schlötterer 2003; Duchon et al. 2013; Kao et al. 2014) and morphological evidence (Capy et al. 1986; Ferveur et al. 1996; Coyne et al. 1999; Takahashi et al. 2001; Rouault et al. 2004) suggest that, for the Americas at least, there were multiple colonization events with some migrants coming from Africa and some from Europe. While there is less evidence that Australia experienced multiple waves of colonization by *D. melanogaster* such a scenario is plausible given the high rates of human migration and inter-continental travel during the 19th century. If North America and Australia experienced multiple waves of immigration from highly differentiated ancestral populations (Pool et al. 2012), many genetic variants would appear clinal, even in the absence of spatially varying selection pressures (Endler 1993; Caracristi & Schlötterer 2003).

Investigating whether North America and Australia represent secondary contact zones is, therefore, crucial for our understanding of the extent of spatially varying selection operating on this species. We note that the models of dual colonization and adaptive differentiation as evolutionary forces that generate and maintain clinal variation in North America and Australia are not mutually exclusive. Notably, one plausible model is that dual colonization of these continents generated patterns of clinal variation and spatially varying selection has subsequently slowed the rate of genetic homogenization among populations. Accordingly, we sought to investigate whether genome-wide patterns of clinal genetic variation in North America and Australia show signals of dual colonization and local adaptation.

We find that both North American and Australian populations show several genomic signatures consistent with secondary contact and suggest that this demographic process is likely to have generated patterns of clinal variation at a large fraction of the genome in both continents. Despite this genome-wide signal of recent admixture, we find evidence that spatially varying selection has shaped patterns of allele frequencies at some loci along latitudinal clines using an F_{ST} outlier approach. Some of these F_{ST} outliers are in or near genes previously identified to underlie life-history and stress tolerance traits. However, in other cases, previously identified functional polymorphisms were not identified as F_{ST} outliers demonstrating the inherent challenge of identifying the genetic targets of spatially varying selection. We discuss these findings in relation to the well-documented evidence of spatially varying selection acting on this species as well as the interpretation of patterns of genomic variation along broad latitudinal clines in general.

Materials and Methods

Genome-wide allele frequency estimates. We utilized novel and publically available genome-wide estimates of allele frequencies of *D. melanogaster* populations sampled world-wide (Figure 1, Table S1). Allele frequency estimates of six North American populations are described in Bergland (2014). Allele frequency estimates of three European populations are described in Bastide et al. (2013) and Tobler et al. (2014). Allele frequency estimates from 20 African populations with full genome sequence that are described in Pool et al. (2012). Allele frequency estimates of two Australian populations are described in Kolaczkowski et al. (2011). Allele frequency estimates from an additional two Australian populations are reported here for the first time. Allele frequency estimates from these additional Australian populations were made by pooling ten individuals from each of 22 isofemale lines originating from Innisfail (17°S) or Yering Station (37°S), Australia (isofemale lines kindly provided by A. Hoffmann). Sequencing library preparation and mapping followed methods outlined in Bergland et al. (2014). Because Australian data were low coverage (~10X per sample, on average), we combined the two northern populations and two southern populations into two new, synthetic populations which we refer to as ‘tropical’ and ‘temperate,’ respectively. In several analyses, *D. simulans* was used as an outgroup. In these analyses, we used genome-wide allele frequency estimates from a world-wide collection *D. simulans* (Begun et al. 2007), using genome-wide *lift-over* files reported in Bergland et al. (2014).

We performed SNP quality filtering similar to the methods presented in Bergland et al. (2014). Briefly, we excluded SNPs within 5bp of polymorphic indels, SNPs within repetitive regions, SNPs with average minor allele frequency less than 15% in both North

America and Australia, SNPs with low (<5) or excessively high read depth (>2 times median read depth) and SNPs not present in the *Drosophila* Genetic Reference Panel (Mackay et al. 2012). African samples were not quality filtered for read depth because allele frequency estimates from these samples were derived from sequenced haplotypes and not pooled samples. Regions of inferred admixture (Pool et al. 2012) in African samples (i.e., introgression of European haplotypes back to African populations) were removed from analysis.

Estimation of the population tree. We calculated Nei's genetic distance (Nei 1972) between each pair of populations as,

$$-\log\left(\frac{\sum p_1 \cdot p_2}{\sqrt{\sum p_1^2} \cdot \sqrt{\sum p_2^2}}\right)$$

where p_1 and p_2 are allele frequency estimates in population 1 and 2, and generated a population tree using the neighbor-joining algorithm implemented in the R (R Core Team 2014) package *ape* (Paradis et al. 2004). To generate bootstrap values for each node, we randomly sampled 10,000 SNPs 100 times. We generated estimates of the population tree using the whole genome and for each chromosome focusing on SNPs that occur within the large, cosmopolitan inversions (Corbett-Detig & Hartl 2012), outside these inversions, or both.

Estimation of the proportion African and European ancestry. We used a simple linear regression method to estimate the proportion of African and European ancestry for each newly derived North American and Australian population. This method follows an approach outlined in Alkorta-Aranburu et al. (2012) where each newly derived population is modeled as a linear combination one African and one European population using an intercept-free regression model. Ancestry coefficients were calculated averaging over 100 bootstrap replicates of 5000 SNPs per derived population per pair of ancestral populations. We generated ancestry estimates using the whole genome and for each chromosome separately focusing on SNPs that occur within the large, cosmopolitan inversions, outside these inversions, or both.

To verify that our regression based method of ancestry proportion estimation is accurate and robust to various demographic scenarios, we simulated several demographic models using coalescent simulations as implemented in *ms* (Hudson 2002). To assess the accuracy of estimated ancestry proportion, we first simulated a three population model where with one ancestral population, one ancient population that diverged from the ancestral population $0.1N_e$ generations ago, and one derived population that results from admixture between the ancestral and ancient populations $0.001N_e$ generations ago. In these simulations, we varied the proportion of lineages in the derived population that originated from the ancient and ancestral populations. A graphical model of this demographic history is presented in Supplemental Figure 2 and *ms* code is available from DataDryad under accession doi:10.5061/dryad.gg5nv.

Next, we sought to verify that single- and dual-colonization scenarios produce distinctive ancestry proportion clines. In these models, we simulated an ancient

population that diverged from an ancestral population $0.1Ne$ generations ago. In the single-colonization scenario a newly derived population was founded from the ancestral population and a series of four additional populations were derived from this initial newly derived population through a series of serial founder events. In the dual-colonization scenario, two newly derived populations were founded by either the ancestral or the ancient population. These newly derived populations each gave rise to another newly derived population through a serial founder event, and a fifth newly derived population resulted from the merger of these later populations. In both single- and dual-colonization models, migration was only allowed between neighboring newly derived populations. A graphical representations of these models can be found in Supplemental Figures 3 and *ms* code is available from DataDryad under accession doi:10.5061/dryad.gg5nv. In the single- and dual-colonization models, we varied the age of initial colonization of the newly derived populations and the extent of the population bottleneck during the initial and serial founder event(s).

Formal tests of admixture. To further assess whether newly derived North America and Australian populations result from admixture of European and African lineages of flies, we performed formal tests of admixture using the f_3 and D statistics (Reich et al. 2009; Patterson et al. 2012). Briefly, these statistics assess whether a proposed tree topology is consistent with the data.

The f_3 statistic, denoted (C; A, B) assesses whether the data are consistent with the topology ((A, C), B) or ((B,C), A). A significantly negative f_3 statistic demonstrates that the data are consistent with both topologies, thus indicating that population C is derived from an admixture event from populations A and B or populations closely related to A and B. However, non-significant or positive f_3 statistic does not preclude the possibility of admixture (Patterson et al. 2012).

The D statistic, which is derived and conceptually similar to the ABBA-BABA statistic (Patterson et al. 2012), assesses whether the data are consistent with the proposed topology ((W, X), (Y, Z)). In general, a significantly positive D statistic indicates that population W results from admixture between population X and Y, or populations closely related to X and Y. A significantly negative D statistic can indicate that population X results from admixture between population W and Y, or populations closely related to W and Y. While, in general, the sign of significant D statistics can be difficult to interpret the most general interpretation is that the data do not conform to the proposed ((W, X), (Y, Z)) topology.

There are three possible D statistics any four populations: ((W, X), (Y, Z)), ((W, Y), (X, Z)), and ((W, Z), (X, Y)). In our analysis, we used *D. simulans* as an outgroup (population Z in this notation). Accordingly, the first two D statistics are most easily interpretable in our analysis. In this context, population W represents the newly derived North American or Australian population and population X and Y represent the putative source African and European populations, respectively. To simplify the interpretation of these D statistics, and to provide a conservative analysis of admixture, we report the D statistic corresponding to the minimum absolute D of ((W, X), (Y, Z)) and ((W, Y), (X, Z))

For each North American and Australian population, we calculated f_3 and D using each European population as one putative donor population and each African population

as the other putative donor population. f_3 and D statistics were calculated using the *threepop* and *fourpop* programs included in *TreeMix* version 1.13 (Pickrell & Pritchard 2012) with 500 bootstrap replicates using a block size of 500 SNPs. Multiple testing correction was performed using the Bonferroni correction applied across all p -values from f_3 or D statistics.

F_{ST} outlier identification. We used the T_{F-LK} statistic (Bonhomme et al. 2010) to identify F_{ST} outliers in North America and Australia. This statistic is related to classic Lewontin-Krakauer test for F_{ST} outliers, under the assumption that the distribution of F_{ST} is proportional to a χ^2 distribution with degrees of freedom equal to one less the number of populations examined (Lewontin & Krakauer 1973). Various assumption underlying Lewontin-Krakauer test have been criticized (Robertson 1975) and the T_{F-LK} statistic attempts to correct for these by conditioning the distribution of F_{ST} values on the inferred underlying population tree. Under a variety of demographic scenarios (Bonhomme et al. 2010; Mita et al. 2013), including secondary contact (Lotterhos & Whitlock 2014), the T_{F-LK} statistic has generally been found to have low false positive- and high true positive-rates in F_{ST} outlier detection. Multiple testing correction for the T_{F-LK} statistic was performed using the false discovery rate methods implemented in the q -value package (Storey & Tibshirani 2003).

Differentiation and rates of parallelism at various SNP classes. To assess rates of co-differentiation we calculated the odds that SNPs fell above one of three F_{ST} quantile thresholds (85, 90, 95%) in both North America and Australia. We compared this value to the odds of co-differentiation from 500 sets of randomly selected SNPs that were matched to the focal SNPs by recombination rate (Comeron et al. 2012), chromosome, inversion status (at the large, cosmopolitan inversions *In(2L)t*, *In(2R)NS*, *In(3L)Payne*, *In(3R)K*, *In(3R)Payne*, *In(3R)Mo*, *In(X)A*, and *In(X)Be*), average read depth in North America and Australia, and heterozygosity in both continents. To control for the possible autocorrelation in signal along the chromosome, we divided the genome into non-overlapping 50Kb blocks and randomly sampled, with replacement, one SNP per block 500 times.

Next, we tested if SNPs at various annotation classes (e.g., short-introns, synonymous, non-synonymous, UTR; Figure 7) were more likely than expected by chance to be co-differentiated or show parallel changes in allele frequency between temperate and tropical locales in both North America and Australia conditional on them being co-differentiated. To assess rates of parallelism, we calculated the fraction of SNPs that were significantly co-differentiated and varied in a parallel fashion between North America and Australia for each SNP class and their matched, genomic controls, again controlling for the spatial distribution of SNPs along the chromosome. We report the difference in rates of parallelism. Standard deviations of the $\log_2(\text{odds-ratio})$ of co-differentiation and for differences in the rates of parallelism are calculated as in Bergland et al. (2014).

Results

Data. We examined genome-wide estimates of allele frequencies from ~30 populations of *D. melanogaster* sampled throughout North America, Australia, Europe and Africa

(Fig. 1). Our analyses largely focused on patterns of variation in North American and Australian populations and, consequently, we primarily focus on two sets of SNP markers. First, we utilized allele frequency estimates at ~500,000 high quality SNPs that segregate at intermediate frequency (MAF > 15%) in North America. The second set was composed of ~300,000 SNPs that segregate at intermediate frequency in Australia. For analyses that examine patterns of polymorphism in both North America and Australia, we examined SNPs that were at intermediate frequency in both continents, yielding a dataset of ~190,000 SNPs. Because of the low sequencing coverage in the Australian populations, it is unclear if the reduced polymorphism in that continent reflects the demographic history of those populations or experimental artifact. Although our analysis primarily focused on patterns of polymorphism in North America and Australia, we also examined allele frequency estimates at both sets of polymorphic SNPs in populations sampled in Europe and Africa.

Genomic signals of secondary contact. We performed a series of independent analyses to examine whether North America and Australia represent secondary contact zones of European and African populations of *D. melanogaster*. First, we constructed a neighbor-joining tree based on genome-wide allele frequency estimates from populations sampled world-wide using 100 sets of 10,000 randomly sampled SNPs. Neighbor joining trees were generated for SNPs residing within inversions, outside inversions, or both genome-wide (Figure 2) and for each chromosome separately (Supplemental Text 1). As expected, African populations exhibited the greatest diversity (Pool et al. 2012) and clustered at the base of the tree while European populations clustered at the tip. North American and Australian populations generally clustered between African and European populations (Figure 2), a pattern that supports the model (Kopelman et al. 2013) that both North American and Australian populations result from secondary contact of European and African ones.

In general, the topology of these population trees was independent of which SNPs were included in the analysis (i.e., compare Figure 2 and Supplemental Figure 1). One notable exception was the placement of the Portuguese population which falls within the North American samples when examining SNPs within inversions. In addition, the topology of the African populations varied depending on the set of SNPs under consideration. However the bootstrap support for the nodes among African populations are generally low, likely stemming from low coverage and imprecise estimates of allele frequencies in these populations. In chromosome specific analyses (Supplemental Text 1), nodes that do not agree with the genome-wide analysis generally had low bootstrap support.

Next, we calculated the proportion of African ancestry in North American and Australian populations by modeling these populations as a linear combination of African and European ancestry. Neutral coalescent simulations using *ms* found that this method accurately estimates ancestry proportions (Supplemental Figure 2). These simulations also revealed that a cline in ancestry proportion is unlikely under a single colonization scenario, even with extensive drift (Supplemental Figure 3), yet is stable and persistent under a dual-colonization scenario for long periods of time following initial colonization (Supplemental Figure 3).

In general, the proportion of African ancestry in North American and Australian populations is negatively correlated with latitude (Figure 3). Conversely, the proportion of European ancestry is positively correlated with latitude (Figure 3). One notable exception to these general ancestry patterns is a single population from Zambia (population ZI from Pool et al. 2012). This population is notable in that it has an exceptionally long branch in our neighbor joining analysis (Figure 2), likely indicating that it may be highly diverged from other African populations. In addition, the X-chromosome lacks a significant pattern of clinal variation in ancestry (Supplemental Figure 4), a pattern consistent with analyses presented in Kao et al. (2014).

Nonetheless, the general pattern of decreasing African ancestry with increasing latitude remains when using nearly any combination of African and European populations and for SNPs inside inversions, outside inversions, or both sampled genome-wide or on any particular autosome. Moreover, our estimate of ancestry proportion of the North Carolina population is in agreement with estimates made using other methods and data-sets (Duchen et al. 2013; Kao et al. 2014).

Finally, we calculated f_3 and D statistics (Reich et al. 2009; Patterson et al. 2012) – commonly referred to as formal tests of admixture – for each North American and Australian population using each sampled European and African population as a putative source population. We observe significantly negative f_3 statistics (Figure 4A) for each North American population when using the Italian or Austrian populations as European source populations and various African populations as the alternate source population (Supplemental Table 1). Significant admixture from Portuguese population into North America was only observed in southern populations using the f_3 statistic. In general, f_3 statistics were not significantly negative for the Australian populations after correcting for multiple testing. Note, however, that temperate and tropical populations in Australia show evidence of admixture using the f_3 statistic before stringent Bonferroni multiple testing correction and remain marginally significant following multiple testing correction (Supplemental Table 1).

We observe significantly positive D statistics for each newly derived North American and Australian population using various combinations of European and African populations as putative source populations (Figure 4B, Supplemental Table 1). Similar to the f_3 analysis, the Portuguese population shows little evidence of admixture into populations sampled in northern North America. It is worth noting that evidence of admixture using the f_3 and D statistics does not conclusively demonstrate that the sampled donor populations (i.e., the European and African populations) are the actual donor populations. Rather, evidence of admixture using these statistics implies that the sampled donor populations, or other unsampled yet closely related populations, are likely the donor populations.

Taken together, these results support the view that both North America and Australia represent secondary contact zones between European and African lineages of *D. melanogaster*. Our results confirm an earlier model (David & Capy 1988) and recent genomic evidence (Kao et al. 2014) that European *D. melanogaster* colonized high latitude locales in North American and Australia whereas African flies colonized low latitude locales in these regions. Genome-wide, low-latitude populations are more similar to African ones whereas high-latitude populations are more similar to European ones.

Under this dual-colonization scenario, we would expect that a large fraction of the genome varies clinally. Indeed, among North American populations of *D. melanogaster* approximately one third of all common SNPs, on the order of 10^5 , are clinal (following the analysis in Bergland et al. 2014). The vast extent of clinal variation in North America, then, is consistent with a dual colonization scenario which would generate patterns of clinal variation at a large fraction of the genome. However, these results do not preclude the existence of spatially varying selection that could also be acting among these populations which could explain patterns of differentiation reported for some loci (e.g., (Sezgin et al. 2004; Hoffmann & Weeks 2007; Paaby et al. 2010; 2014)) and could slow the rate of homogenization of allele frequencies at neutral polymorphisms throughout the genome among clinally distributed populations. We note that a similar analysis of the extent of clinality in Australia is not possible because we lack genome-wide allele frequency estimates from intermediate latitude populations in that continent.

Genomic signals of parallel local adaptation along latitudinal gradients. Our previous analysis supports the model that the demographic history of *D. melanogaster* has contributed to genome-wide patterns of differentiation among temperate and tropical populations of *D. melanogaster* living in North America and Australia. Regardless of this putative demographic history, multiple lines of evidence suggest that populations of flies living along broad latitudinal gradients have adapted to local environmental conditions that may be associated with aspects of temperate environments (see *Discussion*). Accordingly, we performed several tests to assess whether there is a strong, observable genomic signal of local adaptation.

First, we sought to identify F_{ST} outliers using the T_{F-LK} method that attempts to identify SNPs subject to spatially varying selection while maintaining a high power and low false positive rate (Bonhomme et al. 2010; Mita et al. 2013; Lotterhos & Whitlock 2014). This method models the distribution of F_{ST} values after conditioning on the observed population tree among the sampled populations. We identified several hundred significantly differentiated SNPs in North America (Figure 5; Supplemental Table 2), some of which are in or near genes that have been previously implicated in adaptation to spatially varying selection pressures (e.g., *Abd-B*; Fabian et al. 2012) or likely affect life-history traits and correlates (e.g., *AlstR*, *TyR*, *DopR*, *sNPF*) via modulation of endocrine signaling (Bergland 2011). Intriguingly however, the amino acid polymorphism in *cpo* (3R: 13793588) previously implicated in clinal variation in diapause propensity (Schmidt et al. 2008) is significantly differentiated prior to, but not following, multiple testing correction in our F_{ST} outlier analysis ($T_{F-LK} = 16.2$, p -value = 0.006, q -value = 0.52). Similarly, the extensively studied threonine/lysine polymorphism (Kreitman 1983; Powell 1997) that encodes the Fast and Slow allozyme variants at Alcohol dehydrogenase (*Adh*, 2L:14617051) is not significantly differentiated following multiple testing correction in our F_{ST} outlier analysis ($T_{F-LK} = 17.3$, p -value = 0.004, q -value = 0.45).

While a limited number of polymorphisms were identified as significantly differentiated among North American populations, no significantly differentiated SNPs were observed among the Australian populations after correcting for multiple testing (Figure 5). Note that the genome-wide average F_{ST} among North American populations is lower than among Australian populations (0.025 vs. 0.08 respectively), suggesting that

the lack of significantly elevated F_{ST} values in Australia is not due to a lack of population differentiation but rather a high genome-wide differentiation likely caused by recent secondary contact.

The exact number of SNPs with significantly elevated F_{ST} in any particular continent will be subject to a various of considerations including the number of sampled populations, the precision of allele frequency estimates, and the power of particular analytic methods to detect outlier F_{ST} . Some of these factors vary between our North American and Australian samples and thus our power to detect significant elevation of F_{ST} will vary between continents. Therefore, we investigated the general patterns of differentiation and parallelism between the sets of populations sampled in North America and Australia. In addition, we also examined patterns of differentiation and parallelism between these continents and populations sampled from the Old-World (i.e., Europe and Africa).

For these analyses, we first examined whether SNPs that were highly differentiated among one set of populations were also differentiated in another set (hereafter, ‘co-differentiated’). To perform this analysis, we calculated the odds ratio (see *Materials and Methods*) that SNPs fell above a particular quantile threshold of the F_{ST} distribution in any two sets of populations (Figure 6A). We performed this analysis for SNPs that fell either within or outside of the large cosmopolitan inversions. We find that SNPs that are highly differentiated in North America are also highly differentiated in Australia. In addition, we find that SNPs that are highly differentiated in either North America or Australia are also highly differentiated between Europe and Africa. Although patterns of co-differentiation are higher among SNPs within the large, cosmopolitan inversion than for SNPs outside the inversions, the qualitative patterns remain the same for either SNP class suggesting that clinal variation in inversions *per se* does not drive the observed high levels of co-differentiation.

SNPs that are co-differentiated among temperate and tropical populations in North America, Australia, or the Old-World can be differentiated in a parallel way or at random among each geographic region. We show here that there is a high degree of parallelism at the SNP level, genome-wide, among polymorphisms that are highly differentiated in any two sets of populations (Figure 6B). Patterns of parallelism at highly co-differentiated SNPs are similar among SNPs within or outside the large cosmopolitan inversions again suggesting that clinal variation in inversions are not driving genome-wide patterns of parallelism.

High rates of co-differentiation and parallelism among temperate and tropical populations sampled throughout the world can be interpreted in two ways. On the one hand, these patterns could be taken as evidence of parallel adaptation to aspects of temperate environments. On the other hand, these patterns are consistent with the model presented above that North American and Australian populations are the result of recent secondary contact between European and African lineages of flies (see *Results: Genomic signals of secondary contact*).

To differentiate these alternative interpretations, we estimated rates enrichment of highly co-differentiated SNPs and rates of parallelism at highly co-differentiated SNPs among classes of polymorphisms that that we expect, *a priori*, to be more or less likely to contribute to local adaptation. We reasoned that SNPs falling in short-introns, which have been previously shown to evolve neutrally (Lawrie et al. 2013), would be the least likely

to contribute to local adaptation. In contrast, SNPs in other functional classes (e.g., coding, UTR, intron) might be more likely to contribute to local adaptation along latitudinal clines (Reinhardt et al. 2014). We contrasted rates of co-differentiation and parallelism at these putatively functional SNP classes with rates at the short-intron (hereafter ‘neutral’) SNPs and at control SNPs matched to each class by several important biological and experimental features. These comparisons also take into account the spatial distribution of SNPs along the chromosome (see *Materials and Methods*). We reasoned that if parallel adaptive processes have contributed to genome-wide signals of co-differentiation and parallelism in Australia and North America, (1) some functional SNP classes would show a higher rate of co-differentiation and parallelism than neutral SNPs, (2) functional SNPs would show a higher rate of co-differentiation and parallelism than their control SNPs, and (3) neutral SNPs would show a lower rate of co-differentiation and parallelism than their control SNPs.

We find little evidence that various functional classes show differences in rates of co-differentiation or parallelism than either neutral SNPs or their matched controls (Fig. 7AB). Moreover, neutral SNPs show similar rates of co-differentiation and parallelism as their matched controls (Fig. 7AB). There is suggestive evidence that SNPs falling in 5’ UTRs show greater of co-differentiation than expected by chance, but this comparison is not significant after correcting for multiple tests (see $F_{ST} > 95\%$ Fig. 7A; $p_{naive} = 0.01$; $p_{corrected} = 0.24$). Moreover, highly co-differentiated SNPs in 5’UTR are not more likely to be parallel than expected by chance (Fig. 7B), suggesting that the observed excess of co-differentiation may be a statistical artifact. All other tests of excess co-differentiation or parallelism at different SNP classes were not significantly different from expectation ($p > 0.05$).

Taken together, the tests we performed to identify strong genomic signals of parallel adaptation along latitudinal clines were equivocal. We show that there were a modest number of F_{ST} outliers among North American populations sampled along a broad latitudinal cline and no observable F_{ST} outliers among Australian populations (Figure 5) suggesting that the bulk of the F_{ST} distribution is generated by the demographic history of this species. We show that SNPs with high F_{ST} among any one set of populations are likely to have high F_{ST} among other sets of population (Figure 6A). Furthermore, SNPs that are highly co-differentiated are likely to vary in a parallel fashion among geographic regions (Figure 6B). While this result could suggest parallel adaptation, it is also consistent with the dual colonization model we present above. Finally, we show that rates of co-differentiation and parallelism at highly co-differentiated SNPs are similar between functional SNPs, neutral SNPs, and their matched control SNPs (Figure 7) suggesting that the evolutionary forces shaping allele frequencies along latitudinal clines are similar across SNPs that are more- or less-likely to contribute to local adaptation.

Discussion. Herein we report results from a series of analyses that (1) examine whether populations of *D. melanogaster* sampled throughout North America and Australia show signatures of recent secondary contact between European and African lineages, and (2) examine whether there is a genomic signal of spatially varying selection acting along latitudinal gradients. We find that both North America and Australia show several signatures of secondary contact (Figures 2-4). Notably, high latitude populations are closely related to European populations, whereas low latitude populations are more

closely related to African ones. This result implies that a large portion of clinal variation within these continents could, in principal, be generated by the dual colonization of both North America and Australia (Figure 1). Consistent with this view, SNPs that are highly differentiated between temperate and tropical locales in either North America or Australia are also highly likely to be differentiated in a parallel way between Europe and Africa (Figures 6, 7). In addition, we report that genome-wide scans for significantly differentiated polymorphisms identified a limited number of outlier loci (Figure 5). Taken together, our results support the model that recent secondary contact in North America and Australia has generated clinal variation at a large fraction of polymorphisms genome-wide and that spatially varying selection acting at a moderate number of loci acts to slow the rate of genomic homogenization between geographically separated populations.

Secondary contact and the generation of clinal variation in allele frequencies. Recent secondary contact between formerly (semi-) isolated populations is a potent force that can generate clinal variation genome-wide (Endler 1993). In *D. melanogaster*, high levels of genetic differentiation have been observed between temperate and tropical populations sampled in North America and Australia (Turner et al. 2008; Kolaczkowski et al. 2011; Fabian et al. 2012; Reinhardt et al. 2014). In North America at least, most of these highly differentiated SNPs vary clinally (i.e., in a roughly monotonic fashion along latitudinal gradients at false-discovery rate < 10%; Bergland et al. 2014). Moreover, surveys of allele frequencies along latitudinal clines in both North America and Australia at allozymes (Sezgin et al. 2004), SNPs (Sezgin et al. 2004; Lavington et al. 2014; Bergland et al. 2014), microsatellites (Gockel et al. 2001), and transposable elements (González et al. 2010) have repeatedly demonstrated that approximately one third of all surveyed polymorphisms are clinal in either continent. At face value the high proportion of clinal polymorphisms throughout *D. melanogaster*'s genome suggests that demographic processes such as secondary contact have contributed to the generation of clinal variation in this species among recently colonized locales (Bock & Parsons 1981; Keller 2007).

Accordingly, we tested if newly derived populations of *D. melanogaster* show signatures of recent secondary contact. Using a variety of tests, we show that genome-wide patterns of genetic variation from populations sampled in North America and Australia are consistent with recent secondary contact (Figures 2-4). While historical records from North America (Keller 2007) and Australia (Bock & Parsons 1981) suggest a single point of colonization of *D. melanogaster*, results from morphological, behavioral, and genetic studies reported here and elsewhere (Caracristi & Schlötterer 2003; Rouault et al. 2004; Duchon et al. 2013; Kao et al. 2014) suggest that a dual colonization scenario is more likely. At least for the Americas active trade between Europe and western Africa supports the model that North America represents a secondary contact zone.

Australia did not experience the same types of trade with the Old World and throughout the 19th century intercontinental travel to Australia was primarily restricted to British ships. However, British ships traveling to Australia ported in South Africa and India then, after the opening of the Suez Canal in East Africa (Bach 1976). This raises the possibility that secondary contact between European and African fruit fly lineages could have occurred immediately prior to the successful colonization of Australia by *D. melanogaster* in the mid 19th century (Bock & Parsons 1981). Under this mixed-lineage,

single colonization scenario, rapid ecological sorting of colonizing lineages to temperate and tropical niches (Agosta & Klemens 2008) may have created a gradient where European flies were initially predominant at high latitudes and African flies predominant at low latitudes within Australia.

Although secondary contact is capable of generating patterns of clinal variation genome-wide, clines generated through this demographic process are transient. As admixed populations approach migration-selection equilibrium, clines at neutral loci should attenuate. Moreover, once at equilibrium, neutral differentiation should be minimal (Slatkin 1987) for species such as *D. melanogaster* where Nm has been estimated to be on the order of ~ 1 (Yamazaki *et al.* 1986; Singh & Rhomberg 1987) and long-distance dispersal is believed to be frequent (Coyne & Milstead 1987).

Thus, the critical question in determining whether the vast amount of clinal variation in North American and Australian flies has been generated by demography or selection is whether or not this species is at migration-selection equilibrium in these continents. There are several reasons why we suspect this species is not at equilibrium. First, *D. melanogaster* appeared in North America and Australia in the mid- to late 19th century (Bock & Parsons 1981; Keller 2007), or on the order of 1000 generations ago, assuming approximately 10 generations per year. Estimates of local, demic N are on the order of 10^4 (McKenzie 1980; McInnis *et al.* 1982; Powell 1997) implying that m is on the order of 10^{-4} (if $Nm \sim 1$). If these estimates are accurate to the order of magnitude, it would take approximately 2500 generations to get about half way to equilibrium (Whitlock 1992) or $\sim 10,000$ generations to fully approach equilibrium (Whitlock & McCauley 1999). Thus, from a simple demographic perspective, it would seem unlikely that *D. melanogaster* has reached migration-selection-drift equilibrium.

Others have suggested that non-African populations of *D. melanogaster* are not at equilibrium. In general, non-African populations of *D. melanogaster* show a reduction in diversity coupled with an excess of rare variants (Mackay *et al.* 2012). This genome-wide pattern is consistent with a population bottleneck during colonization followed by population expansion. Others have noted that non-African populations of *D. melanogaster* also have higher levels of linkage-disequilibrium (LD) than expected under the standard neutral model (Andolfatto & Przeworski 2000; Haddrill *et al.* 2005; Langley *et al.* 2012) whereas LD in African populations is more consistent with neutrality (Andolfatto & Wall 2003 *cf.* Langley *et al.* 2012). Although genome-wide elevation of LD could be caused by various factors including pervasive positive- or negative-selection, admixture would also possibly generate this signal.

Previous studies examining departure from equilibrium models in *D. melanogaster* have concluded that caution should be taken when conducting genome-wide scans for positive-selection given the non-equilibrium nature of this species (Andolfatto & Przeworski 2000). Notably, demographic forces such as population bottlenecks can, in principal, mimic many of the signatures left by some types of adaptive evolution. A complimentary approach to quantify the magnitude of adaptive evolution and to identify loci subject to selection is to identify polymorphisms that are differentiated between populations that are subject to divergent selection pressures. However, results presented here demonstrate that, for *D. melanogaster* at least, signatures of adaptive evolution from genome-wide patterns of differentiation along latitudinal clines in newly derived

populations in North America and Australia should be taken with a similar or even greater degree of caution as traditional scans for recent, positive selection.

Spatially varying selection and the maintenance of clinal variation in allele frequencies.

Whereas secondary contact is capable of generating clinal variation, spatially varying selection is required for its long-term maintenance. There is little doubt that populations of *D. melanogaster* living along broad latitudinal clines in temperate environments have adapted to spatially varying selection pressures. Support for the idea of local adaptation along latitudinal clines comes from three main lines of evidence.

First, certain phenotypes show repeatable clines along latitudinal and altitudinal gradients that mirror deeper phylogenetic variation among temperate and tropical species. For instance, aspects of body size vary clinally in North America (Coyne & Beecham 1987) and Australia (Kennington et al. 2003) as well as along altitudinal/latitudinal clines in India (Bhan et al. 2014) and altitudinal clines within Africa (Pitchers et al. 2013; Klepsatel et al. 2014). Given such patterns of parallelism within and among continents, including within the ancestral African range, the most plausible explanation is that parallel selection pressures have generated these patterns of latitudinal and altitudinal variation. These intraspecific clines mimic interspecific patterns among temperate and tropical endemic drosophilids following Bergmann's rule (Blanckenhorn & Demont 2004; Shelomi 2012) again implicating that natural selection has shaped these patterns of genetically based, phenotypic variation.

Second, certain genetic and phenotypic clines in *D. melanogaster* have shifted over decadal scales. Shifts in these clines are consistent with adaptation to aspects of global climate change wherein alleles common in low-latitude populations have become more prevalent in high-latitude ones over the last 20 years (Hoffmann & Weeks 2007).

Finally, here we identify several hundred polymorphisms in North America that are significantly differentiated (see *Results*, Figure 5, and Supplemental Table 3). Although the function of many of these polymorphisms is presently unknown, several are within the genes known to affect life-history traits and correlates that vary among temperate and tropical populations (see *Results*).

The identification of significantly differentiated SNPs within North America can be taken as evidence of local adaptation to spatially varying selection pressures. However, the observation that two SNPs (one in *cpo* and one in *Adh*) that each likely contribute to local adaptation fall in an upper, but not extreme, tail of the F_{ST} distribution suggests that there are many more ecologically relevant and functional polymorphisms that have contributed to local adaptation in *D. melanogaster*. However, the signal of high differentiation caused by spatially varying selection at these SNPs is likely masked by recent admixture that has contributed to a high level of differentiation genome-wide. In light of these results, we suggest that scans for local adaptation based on patterns of genetic differentiation in *D. melanogaster* are an important first step in identifying adaptively differentiated clinal polymorphisms but that additional evidence, such as functional validation (Schmidt et al. 2008; Paaby et al. 2010; 2014), should be gathered before concluding that differentiation is caused by adaptive processes.

Conclusions. It has long been recognized that genetic differentiation among populations can be caused by both adaptive and demographic (neutral) processes (Wright 1943). Due

to *D. melanogaster*'s large effective population size (Karasov et al. 2010), high migration rate (Coyne & Milstead 1987), and rapid decay of linkage disequilibrium (Mackay et al. 2012) others have concluded that differentiation among populations sampled along latitudinal gradients is primarily caused by spatially varying selection. Work presented here supports the notion that spatially variable selection does contribute to some differentiation among populations.

However, several genome-wide signatures presented here (Figures 2-4) and elsewhere (Caracristi & Schlötterer 2003; Duchon *et al.* 2013; Kao *et al.* 2014; Bergland *et al.* 2014) indicate that populations of flies in North America and Australia result from admixture of European and African lineages. High-latitude (temperate) populations in North America and Australia are more closely related to European populations whereas low-latitude (tropical) populations are more closely related to African ones (Figures 2, 3) suggesting that admixture occurred along a latitudinal gradient and that this demographic event generated clinal genetic variation at roughly 1/3 of all common SNPs (Bergland et al. 2014). These colonizing lineages of flies were likely already differentially adapted to the temperate and tropical conditions that they encountered in North America and Australia. Consequently the recent demographic history of this species in North America and Australia is collinear with both local adaptation within these newly colonized continents and among the ancestral ranges. One practical consequence of the collinearity of demography and adaptation is that the identification of clinality at any particular locus cannot be taken exclusively as evidence of spatially varying selection.

The collinearity of demography and adaptation may be a general feature of a variety of species. For instance, the successful colonization of novel locales by invasive species such as *D. melanogaster* is often facilitated by multiple waves of invasion. In addition to increasing propagule pressure (Simberloff 2009), multiple independent invasions are also thought to facilitate invasion success by buffering the loss of genetic diversity that accompanies bottlenecks associated with colonization. Moreover, some evidence suggests that rapid adaptation following invasion (reviewed in Dormontt et al. 2011), often results directly from hybridization between independently invading subpopulations (reviewed in Lee 2002). If invasive species are widely distributed in their native range (Bates et al. 2013), this raises the possibility that successful invasions may often result from admixture of populations that are already differentially adapted to selection pressures that vary along broad spatial gradients in the introduced ranges.

The collinearity of demography and adaptation may also occur in temperate endemic species. This phenomenon may be particularly true in marine taxa as a result of admixture between high-latitude refugial populations and low-latitude populations following the last glacial maxima ~20,000 years ago (Bernatchez & Wilson 1998; Maggs *et al.* 2008). For these species, secondary contact between high- and low-latitude populations would result in clinal variation at neutral loci as well as loci that contribute to adaptation along latitudinal gradients (e.g., Adams et al. 2006). In this scenario, neutral and adaptive genetic clines may be indistinguishable.

The ability to identify targets of spatially varying selection is now possible in many model and non-model species (Li et al. 2008; Savolainen et al. 2013). While these genome-wide approaches can be extremely powerful, great care must be taken to ensure that signals of local adaptation are uniquely identifiable and independent from pervasive (but often subtle) signals of demography. F_{ST} outlier approaches can in some cases enable

the detection of loci that underlie local adaptation following complex and confounding demographic scenarios. However, as we show here, loci known to underlie functionally relevant, phenotypic variation are not necessarily detected as statistically significant outliers, likely due to the pervasive signals of recent demographic events. To ameliorate these concerns, we suggest that proposed genetic targets of spatially varying selection be functionally verified, or that patterns of spatial variation at loci known *a priori* to underlie fitness related phenotypic variation be investigated. Alternatively, the genomic targets of local adaptation can be identified by examining population differentiation over small spatial scales (Richardson et al. 2014) or over short time periods (Bergland et al. 2014) that are likely to be orthogonal to the demographic history of the focal species.

Acknowledgements. We thank Joyce Kao, Heather Machado, and Annalise Paaby for insightful comments on earlier versions of this manuscript. We thank David Lawrie for providing lists of neutrally evolving SNPs and Ary Hoffmann for graciously providing isofemale lines from Australia. Finally, we thank Richard Hudson for help with *ms* and Nick Patterson and Peter Ralph for assistance interpreting *D* statistics. AOB was supported by an NIH National Service Research Award (F32 GM097837). JG is a Ramon y Cajal fellow (RYC-2010-07306) supported by grants from the European Commission (Marie Curie CIG PCIG-GA-2011-293860) and from the Spanish Government (Fundamental Research Projects Grant BFU-2011-24397). This work was supported by NSF DEB 0921307 (awarded to PS) and NIH R01GM089926 (awarded to PS and DP).

References

- Adams SM, Lindmeier JB, Duvernell DD (2006) Microsatellite analysis of the phylogeography, Pleistocene history and secondary contact hypotheses for the killifish, *Fundulus heteroclitus*. *Molecular Ecology*, **15**, 1109–1123.
- Agosta SJ, Klemens JA (2008) Ecological fitting by phenotypically flexible genotypes: implications for species associations, community assembly and evolution. *Ecology Letters*, **11**, 1123–1134.
- Alkorta-Aranburu G, Beall CM, Witonsky DB *et al.* (2012) The Genetic Architecture of Adaptations to High Altitude in Ethiopia. *PLoS genetics*, **8**, e1003110.
- Andolfatto P, Przeworski M (2000) A genome-wide departure from the standard neutral model in natural populations of *Drosophila*. *Genetics*, **156**, 257–268.
- Andolfatto P, Wall JD (2003) Linkage disequilibrium patterns across a recombination gradient in African *Drosophila melanogaster*. *Genetics*.
- Bach J (1976) *A Maritime History of Australia*. Thomas Nelson Limited, Melbourne.
- Bastide H, Betancourt A, Nolte V *et al.* (2013) A genome-wide, fine-scale map of natural pigmentation variation in *Drosophila melanogaster*. *PLoS genetics*, **9**, e1003534.
- Bates AE, McKelvie CM, Sorte CJB *et al.* (2013) Geographical range, heat tolerance and invasion success in aquatic species. *Proceedings of the Royal Society B: Biological Sciences*, **280**, 20131958.
- Begun DJ, Holloway AK, Stevens K *et al.* (2007) Population Genomics: Whole-Genome Analysis of Polymorphism and Divergence in *Drosophila simulans*. *PLoS biology*, **5**, e310.
- Bergland AO (2011) Mechanisms of nutrient-dependent reproduction in dipteran insects. In: *Mechanisms of life history evolution: the genetics and physiology of life history traits and trade-offs* (eds Flatt T, Heyland A). Oxford University Press, Oxford.

- Bergland AO, Behrman EL, O'Brien KR, Schmidt PS, Petrov DA (2014) Genomic evidence of rapid and stable adaptive oscillations over seasonal time scales in *Drosophila*. *PLoS genetics*, **10**, e1004775.
- Bernatchez L, Wilson CC (1998) Comparative phylogeography of Nearctic and Palearctic fishes. *Molecular Ecology*.
- Bhan V, Parkash R, Aggarwal D (2014) Effects of body-size variation on flight-related traits in latitudinal populations of *Drosophila melanogaster*. *Journal of genetics*.
- Blanckenhorn WU, Demont M (2004) Bergmann and converse Bergmann latitudinal clines in arthropods: two ends of a continuum? *Integrative and Comparative Biology*, **44**, 413–424.
- Bock IR, Parsons PA (1981) Species of Australia and New Zealand. In: (eds Ashburner M, Carson HL, Thompson JN), pp. 291–306. *Genetics and biology of Drosophila*, London.
- Bonhomme M, Chevalet C, Servin B *et al.* (2010) Detecting Selection in Population Trees: The Lewontin and Krakauer Test Extended. *Genetics*, **186**, 241–262.
- Bradshaw WE, Lounibos LP (1977) Evolution of dormancy and its photoperiodic control in pitcher-plant mosquitoes. *Evolution*, **31**, 546.
- Capy P, David JR, Allemand R *et al.* (1986) Genetic analysis of *Drosophila melanogaster* in the French West Indies and comparison with populations from other parts of the world. *Genetica*, **69**, 167–176.
- Caracristi G, Schlötterer C (2003) Genetic differentiation between American and European *Drosophila melanogaster* populations could be attributed to admixture of African alleles. *Molecular biology and evolution*, **20**, 792–799.
- Comeron JM, Ratnappan R, Bailin S (2012) The many landscapes of recombination in *Drosophila melanogaster*. *PLoS genetics*, **8**, e1002905.
- Corbett-Detig RB, Hartl DL (2012) Population genomics of inversion polymorphisms in *Drosophila melanogaster*. *PLoS genetics*, **8**, e1003056.
- Coyne JA, Beecham E (1987) Heritability of two morphological characters within and among natural populations of *Drosophila melanogaster*. *Genetics*.
- Coyne JA, Milstead B (1987) Long-distance migration of *Drosophila*. 3. Dispersal of *D. melanogaster* alleles from a Maryland orchard. *American Naturalist*, **130**, 70–82.
- Coyne JA, Wicker-Thomas C, Jallon J-M (1999) A gene responsible for a cuticular hydrocarbon polymorphism in *Drosophila melanogaster*. *Genetical research*, **73**, 189–203.
- David J, Capy P (1988) Genetic variation of *Drosophila melanogaster* natural populations. *Trends in Genetics*, **4**, 106–111.
- Dormontt EE, Lowe AJ, Prentis PJ (2011) Is rapid adaptive evolution important in successful invasions. In: *Fifty Years of Invasion Ecology* (ed Richardson DM), pp. 175–189. West Sussex.
- Duchen P, Živković D, Hutter S, Stephan W, Laurent S (2013) Demographic inference reveals African and European admixture in the North American *Drosophila melanogaster* population. *Genetics*, **193**, 291–301.
- Endler JA (1993) *Geographic Variation, Speciation, and Clines*. Princeton University Press, Princeton, NJ.
- Fabian DK, Kapun M, Nolte V *et al.* (2012) Genome-wide patterns of latitudinal differentiation among populations of *Drosophila melanogaster* from North America.

- Molecular Ecology*, **21**, 4748–4769.
- Ferveur J-F, Cobb M, Boukella H, Jallon J-M (1996) World-wide variation in *Drosophila melanogaster* sex pheromone: behavioural effects, genetic bases and potential evolutionary consequences. *Genetica*, **97**, 73–80.
- Gibert P, Moreteau B, Pétavy G, Karan D, David JR (2001) Chill-coma tolerance, a major climatic adaptation among drosophila species. *Evolution*, **55**, 1063–1068.
- Gockel J, Kennington WJ, Hoffmann A, Goldstein DB, Partridge L (2001) Nonclinality of molecular variation implicates selection in maintaining a morphological cline of *Drosophila melanogaster*. *Genetics*, **158**, 319–323.
- González J, Karasov TL, Messer PW, Petrov DA (2010) Genome-wide patterns of adaptation to temperate environments associated with transposable elements in *Drosophila*. *PLoS genetics*, **6**, e1000905.
- Gray SM, McKinnon JS (2007) Linking color polymorphism maintenance and speciation. *Trends in Ecology & Evolution*, **22**, 71–79.
- Haddrill PR, Thornton KR, Charlesworth B, Andolfatto P (2005) Multilocus patterns of nucleotide variability and the demographic and selection history of *Drosophila melanogaster* populations. *Genome Research*, **15**, 790–799.
- Hoffmann AA, Weeks AR (2007) Climatic selection on genes and traits after a 100 year-old invasion: a critical look at the temperate-tropical clines in *Drosophila melanogaster* from eastern Australia. *Genetica*, **129**, 133–147.
- Hoffmann AA, Anderson A, Hallas R (2002) Opposing clines for high and low temperature resistance in *Drosophila melanogaster*. *Ecology Letters*, **5**, 614–618.
- Hudson RR (2002) Generating sample under a Wright-Fisher neutral model of genetic variation. *Bioinformatics*, **18**, 337–338.
- Jones FC, Grabherr MG, Chan YF *et al.* (2012) The genomic basis of adaptive evolution in threespine sticklebacks. *Nature*, **484**, 55–61.
- Kao JY, Zubair A, Salomon MP, Nuzhdin SV, Campo D (2014) Population genomic analysis uncovers African and European admixture in *Drosophila melanogaster* populations from the southeastern United States and Caribbean Islands. *bioRxiv*, 009092.
- Karasov T, Messer PW, Petrov DA (2010) Evidence that adaptation in *Drosophila* is not limited by mutation at single sites. *PLoS genetics*, **6**, e1000924.
- Keller A (2007) *Drosophila melanogaster*'s history as a human commensal. *Current Biology*, **17**, R77–R81.
- Kennington WJ, Gockel J, Partridge L (2003) Testing for asymmetrical gene flow in a *Drosophila melanogaster* body-size cline. *Genetics*, **165**, 667–673.
- Kitano J, Lema SC, Luckenbach JA *et al.* (2010) Adaptive divergence in the thyroid hormone signaling pathway in the stickleback radiation. *Current Biology*, **20**, 2124–2130.
- Klepsatel P, Gáliková M, Huber CD, Flatt T (2014) Similarities and differences in altitudinal versus latitudinal variation for morphological traits in *Drosophila melanogaster*. *Evolution*, **68**, 1385–1398.
- Kolaczowski B, Kern AD, Holloway AK, Begun DJ (2011) Genomic differentiation between temperate and tropical Australian populations of *Drosophila melanogaster*. *Genetics*, **187**, 245–260.
- Kopelman NM, Stone L, Gascuel O (2013) The behavior of admixed populations in

- neighbor-joining inference of population trees. *Pacific Symposium on Biocomputing*, 273–284.
- Kreitman M (1983) Nucleotide polymorphism at the alcohol dehydrogenase locus of *Drosophila melanogaster*. *Nature*, **304**, 412–417.
- Langley CH, Stevens K, Cardeno C *et al.* (2012) Genomic variation in natural populations of *Drosophila melanogaster*. *Genetics*, **192**, genetics.112.142018–598.
- Laurent SJY, Werzner A, Excoffier L, Stephan W (2011) Approximate Bayesian Analysis of *Drosophila melanogaster* polymorphism data reveals a recent colonization of Southeast Asia. *Molecular biology and evolution*, **28**, 2041–2051.
- Lavington E, Cogni R, Kuczynski C *et al.* (2014) A small system—high-resolution study of metabolic adaptation in the central metabolic pathway to temperate climates in *Drosophila melanogaster*. *Molecular biology and evolution*, **31**, 2032–2041.
- Lawrie DS, Messer PW, Hershberg R, Petrov DA (2013) Strong purifying selection at synonymous sites in *D. melanogaster*. *PLoS genetics*, **9**, e1003527.
- Lee CE (2002) Evolutionary genetics of invasive species. *Trends in Ecology & Evolution*, **17**, 386–391.
- Lee SF, Eyre Walker YC, Rane RV *et al.* (2013) Polymorphism in the neurofibromin gene, *Nf1*, is associated with antagonistic selection on wing size and development time in *Drosophila melanogaster*. *Molecular Ecology*, **22**, 2716–2725.
- Lee SF, Sgró CM, Shirriffs J *et al.* (2011) Polymorphism in the couch potato gene clines in eastern Australia but is not associated with ovarian dormancy in *Drosophila melanogaster*. *Molecular Ecology*, **20**, 2973–2984.
- Lewontin RC, Krakauer J (1973) Distribution of gene frequency as a test of the theory of the selective neutrality of polymorphisms. *Genetics*, **74**, 175–195.
- Li H, Stephan W (2006) Inferring the demographic history and rate of adaptive substitution in *Drosophila*. *PLoS genetics*, **2**, e166.
- Li YF, Costello JC, Holloway AK, Hahn MW (2008) “Reverse Ecology” and the power of population genomics. *Evolution*, **62**, 2984–2994.
- Lotterhos KE, Whitlock MC (2014) Evaluation of demographic history and neutral parameterization on the performance of FST outlier tests. *Molecular Ecology*, **23**, 2178–2192.
- Mackay TFC, Richards S, Stone EA *et al.* (2012) The *Drosophila melanogaster* Genetic Reference Panel. *Nature*, **482**, 173–178.
- Maggs CA, Castilho R, Foltz D *et al.* (2008) Evaluating signatures of glacial refugia for North Atlantic benthic marine taxa. *dx.doi.org*, **89**, S108–S112.
- McInnis DO, Schaffer HE, Mettler LE (1982) Field dispersal and population sizes of native *Drosophila* from North Carolina. *American Naturalist*, **119**, 319–330.
- McKenzie JA (1980) An ecological study of the alcohol dehydrogenase (*Adh*) polymorphism of *Drosophila melanogaster*. *Australian Journal of Zoology*, **28**, 709–716.
- Mita S, Thuillet AC, Gay L *et al.* (2013) Detecting selection along environmental gradients: analysis of eight methods and their effectiveness for outbreeding and selfing populations. *Molecular Ecology*, **22**, 1383–1399.
- Nei M (1972) Genetic distance between populations. *American Naturalist*, **106**, 283–292.
- Paaby AB, Bergland AO, Behrman EL, Schmidt PS (2014) A highly pleiotropic amino acid polymorphism in the *Drosophila* insulin receptor contributes to life-history

- adaptation. *Evolution*, **68**, 3395–3409.
- Paaby AB, Blacket MJ, Hoffmann AA, Schmidt PS (2010) Identification of a candidate adaptive polymorphism for *Drosophila* life history by parallel independent clines on two continents. *Molecular Ecology*, **19**, 760–774.
- Paradis E, Claude J, Strimmer K (2004) APE: analyses of phylogenetics and evolution in R language. *Bioinformatics*, **20**, 289–290.
- Patterson N, Moorjani P, Luo Y *et al.* (2012) Ancient admixture in human history. *Genetics*, **192**, 1065–1093.
- Pickrell JK, Pritchard JK (2012) Inference of population splits and mixtures from genome-wide allele frequency data. *PLoS genetics*, **8**, e1002967.
- Pitchers W, Pool JE, Dworkin I (2013) Altitudinal clinal variation in wing size and shape in African *Drosophila melanogaster*: one cline or many? *Evolution*, **67**, 438–452.
- Pool JE, Corbett-Detig RB, Sugino RP *et al.* (2012) Population genomics of Sub-Saharan *Drosophila melanogaster*: African diversity and Non-African admixture. *PLoS genetics*, **8**, e1003080.
- Powell J (1997) *Progress and Prospects in Evolutionary Biology : The Drosophila Model*. Oxford University Press.
- R Core Team (2014) R: A Language and Environment for Statistical Computing.
- Rajpurohit S, Parkash R, Ramniwas S, Singh S (2008) Variations in body melanisation, ovariole number and fecundity in highland and lowland populations of *Drosophila melanogaster* from the Indian subcontinent. *Insect Science*, **15**, 553–561.
- Reich D, Thangaraj K, Patterson N, Price AL, Singh L (2009) Reconstructing Indian population history. *Nature*, **461**, 489–494.
- Reinhardt JA, Kolaczowski B, Jones CD, Begun DJ, Kern AD (2014) Parallel geographic variation in *Drosophila melanogaster*. *Genetics*, **197**, 361–373.
- Richardson JL, Urban MC, Bolnick DI, Skelly DK (2014) Microgeographic adaptation and the spatial scale of evolution. **29**, 165–176.
- Robertson A (1975) Remarks on the Lewontin-Krakauer test. *Genetics*, **80**, 396.
- Rouault J-D, Marican C, Wicker-Thomas C, Jallon J-M (2004) Relations between cuticular hydrocarbon (hc) polymorphism, resistance against desiccation and breeding temperature; a model for hc evolution in *D. melanogaster* and *D. simulans*. *Genetica*, **120**, 195–212.
- Savolainen O, Lascoux M, Merilä J (2013) Ecological genomics of local adaptation. *Nature Reviews Genetics*, **14**, 807–820.
- Schmidt PS, Matzkin L, Ippolito M, Eanes WF (2005) Geographic variation in diapause incidence, life-history traits, and climatic adaptation in *Drosophila melanogaster*. *Evolution*, **59**, 1721–1732.
- Schmidt PS, Zhu C-T, Das J *et al.* (2008) An amino acid polymorphism in the couch potato gene forms the basis for climatic adaptation in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 16207–16211.
- Sezgin E, Duvernell DD, Matzkin LM, Duan Y, Zhu CT (2004) Single locus latitudinal clines in metabolic genes, derived alleles, and their relationship to temperate adaptation in *Drosophila melanogaster*. *Genetics*, **168**, 923–931.
- Shelomi M (2012) Where are we now? Bergmann's rule *sensu lato* in insects. *The American Naturalist*, **180**, 511–519.

- Simberloff D (2009) The Role of Propagule Pressure in Biological Invasions. *Annual Review of Ecology, Evolution, and Systematics*, **40**, 81–102.
- Singh RS, Rhomberg LR (1987) A comprehensive study of genic variation in natural populations of *Drosophila melanogaster*. I. Estimates of gene flow from rare alleles. *Genetics*, **115**, 313–322.
- Slatkin M (1987) Gene flow and the geographic structure of natural populations. *Science*, **236**, 787–792.
- Stinchcombe JR, Weinig C, Ungerer M *et al.* (2004) A latitudinal cline in flowering time in *Arabidopsis thaliana* modulated by the flowering time gene FRIGIDA. *Proceedings of the National Academy of Sciences of the United States of America*, **101**, 4712–4717.
- Storey JD, Tibshirani R (2003) Statistical significance for genomewide studies. *Proceedings of the National Academy of Sciences of the United States of America*, **100**, 9440–9445.
- Takahashi A, Tsaur S-C, Coyne JA, Wu C-I (2001) The nucleotide changes governing cuticular hydrocarbon variation and their evolution in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences of the United States of America*, **98**, 3920–3925.
- Telonis Scott M, Hoffmann AA, Sgró CM (2011) The molecular genetics of clinal variation: a case study of ebony and thoracic trident pigmentation in *Drosophila melanogaster* from eastern Australia. *Molecular Ecology*, **20**, 2100–2110.
- Tobler R, Franssen SU, Kofler R *et al.* (2014) Massive habitat-specific genomic response in *D. melanogaster* populations during experimental evolution in hot and cold environments. *Molecular biology and evolution*, **31**, 364–375.
- Turner TL, Levine MT, Eckert ML, Begun DJ (2008) Genomic analysis of adaptive differentiation in *Drosophila melanogaster*. *Genetics*, **179**, 455–473.
- Whitlock MC (1992) Temporal fluctuations in demographic parameters and the genetic variance among populations. *Evolution*, **46**, 608–615.
- Whitlock MC, McCauley DE (1999) Indirect measures of gene flow and migration: $F_{ST} \approx 1/(4Nm+1)$. *Heredity*, **82**, 117–125.
- Wright S (1943) Isolation by distance. *Genetics*, **28**, 114–1398.
- Yamazaki T, Choo J-K, Watanabe TK, Takahata N (1986) Gene flow in natural populations of *Drosophila melanogaster* with special reference to lethal allelism rates and protein variation. *Genetics*, **113**, 73–89.

Data Accessibility:

- Raw DNA sequence data for Inisfail and Yering Station samples: NCBI SRA: BioProject 270869
- *ms* code to generate simulations of various colonization models: Data Dryad: doi:10.5061/dryad.gg5nv
- All other genomic data are previously published.

Author Contributions

AOB and RT analyzed the data. JG generated sequencing libraries. AOB, RT, JG, PS, and DP wrote the manuscript.

Figure legends.

Figure 1. Map of collection locales (squares) and proposed colonization routes of *D. melanogaster* (arrows).

Figure 2. Estimated population tree of sampled locales. Loci were sampled across all chromosomes, focusing on SNPs residing within (+), outside (-) large, cosmopolitan inversions, or both(\pm). See Supplemental Figure 1 for population trees based on each chromosome.

Figure 3. Proportion African and European ancestry in North American and Australian populations. Thin grey lines represent the ancestry estimate for any particular African or European population averaged over all European and African populations. Black line represents the average ancestry proportions from all African and European populations. The blue circle represents the spring Pennsylvania sample and the red circle represents the fall Pennsylvanian sample. See Supplemental Table X for ancestry estimates for each pairwise combination of African and European population, and for SNPs on each chromosome, and with locations inside and/or outside inversions.

Figure 4. Observed standardized Z scored of (A) f_3 and (B) D statistics for each North American and Australian population when considering every possible combination of African and European population and a putative donor population. The dashed line represents the Bonferroni significance threshold at $\alpha < 0.05$. Values below the threshold in (A) are significantly different from 0. Values above or below the threshold in (B) are significantly different from 0.

Figure 5. Number of significantly differentiated SNPs in North America and Australia at various FDR (q -value) thresholds.

Figure 6. Patterns of co-differentiation and parallelism between North American, Australian, and Old-world populations. (A) \log_2 odds-ratio that SNPs fall above the F_{ST} quantile cut-off (x -axis) in both sets of populations (NA: North America; AUS: Australia; OW: Old-World). (B) Proportion of SNPs that vary in a parallel way given that they fall above the F_{ST} quantile cut-off in both sets of populations. Confidence bands represent 95% confidence intervals.

Figure 7. Patterns of (A) co-differentiation and (B) parallelism among various classes of SNPs relative to their matched controls. Vertical lines represent 95% confidence intervals. Horizontal dotted lines represents the null expectations. See *Materials and Methods* for details.

Supplemental Text 1. Neighbor-joining trees for each chromosome in Newick format.


Supplemental Figure 1. Verification that the linear model method of ancestry proportion estimation is accurate.

Supplemental Figure 2. Verification that the linear model method of ancestry proportion estimation is robust under various demographic scenarios.

Supplemental Figure 3. Ancestry estimates for each chromosome partitioned by inversion status.

Supplemental Table 1. f_3 and D statistics for each pair-wise combination of African and European populations.

Supplemental Table 2. Annotated table of F_{ST} outlier SNPs at $q < 0.15$ in North America.



+/- inversions

+ inversions

- inversions

Figure 2





Figure 3






Fig. 6







A**B**

Fig. 7




A

S. Fig. 1


**B**Estimated ancestry coef.
(linear model)

S. Fig. 2


A




B




C




D




E



F





bioRxiv preprint doi: <https://doi.org/10.1101/009084>; this version posted January 6, 2015. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.

