How predator hunting-modes affect prey behaviour: Capture

deterrence in *Drosophila melanogaster*

Abhijna Parigi[1,2,3], Cody Porter[2,4], Megan Cermak[2], William R. Pitchers[1,3] and Ian Dworkin[1,2,3*]

Affiliations:

1 Program in Ecology, Evolutionary Biology, and Behaviour, Michigan State University, East

Lansing, MI 48824, USA

2 Department of Zoology, Michigan State University, East Lansing, MI 48824, USA

3 BEACON Centre for the Study of Evolution in Action, Michigan State University, East

Lansing, MI 48824, USA

4 Current address: Department of Zoology and Physiology, University of Wyoming. Laramie,

WY, USA.

* Correspondence to Ian Dworkin: idworkin@msu.edu

Links to videos (also available in supplement)

Abdominal Lifting http://dx.doi.org/10.6084/m9.figshare.1185638

Stopping Behaviour http://dx.doi.org/10.6084/m9.figshare.1185639

Retreat http://dx.doi.org/10.6084/m9.figshare.1185640

ABSTRACT

Hunting mode, the distinct set of behavioural strategies that a predator employs while

hunting, can be an important determinant of the prey organism's behavioural response.

However, few studies have considered how a predator's hunting mode influences anti-

predatory behaviours of a prey species. Here we document the influence of active hunters

(zebra jumping spiders, *Salticus scenicus*) and ambush predators (Chinese praying

mantids, *Tenodera aridifolia sinensis*) on the capture deterrence anti-predatory

behavioural repertoire of the model organism, *Drosophila melanogaster*. We

hypothesized that *D. melanogaster* would reduce overall locomotory activity in the

presence of ambush predators, and increase activity with active hunters. First we

observed and described the behavioural repertoire of *D. melanogaster* in the presence of

the predators. We documented three previously undescribed behaviours- abdominal

lifting, stopping and retreat- which were performed at higher frequency by *D.*

melanogaster in the presence of predators, and may aid in capture deterrence. Consistent

with our predictions, we observed an increase in the overall activity of *D. melanogaster*

in the presence of jumping spiders (active hunter). However, counter to our prediction,

mantids (ambush hunter) had only a modest influence on activity. We also observed

considerable intra and inter-individual variation in response to both predator types. Given

these new insights into *Drosophila* behaviour, and with the genetic tools available,

dissecting the molecular mechanisms of anti-predator behaviours may now be feasible in

this system.

INTRODUCTION

Predation, a ubiquitous selective force, gives rise to and determines the nature of defensive traits in prey populations (Edmunds, 1974; Juliano & Gravel 2002; Goslin & Rodd, 2007; Langerhans, 2007; Lima & Dill, 1990; Sansom, Lind & Cresswell, 2009). Predator hunting-modes, i.e., the set of behaviours that predators employ to pursue and capture their prey (Schoener, 1971; Huey & Pianka 1981; Preisser, Orrock & Schmitz 2007), have been shown to induce distinct prey responses (Schmitz, 2008) that in turn influence the productivity of ecological communities. In habitats dominated by active hunters there is lower species evenness and higher above-ground net primary productivity compared to habitats dominated by ambush hunters (Schmitz, 2008). The authors suggest the observed differences in prey productivity to be driven by hunting mode specific trade-offs between foraging and seeking refuge. Although studies often describe the effects of predators on prey traits (i.e. DeWitt, Robinson & Wilson, 2000; Reznick, Butler & Rodd, 2001; Relyea, 2001), it is rare for the role of predator hunting-mode to be explicitly considered.

Here we investigate segregating differences in the anti-predatory behavioural repertoire of the fruit fly, *Drosophila melanogaster*, in response to two predator species differing in hunting modes. Based on (Schmitz, 2008), we predicted that fruit flies, in the presence of a familiar predator, would exhibit hunting-mode specific modifications in activity levels. We used *D. melanogaster* because, although it is one of the most well-studied model organisms, there is a relative paucity of information regarding *D. melanogaster's* natural history, ecology and behaviour, including habitat, food resources, and natural enemies (but see Reaume & Sokolowski, 2006; David & Capy, 1988; Turelli

& Hoffmann, 1991; Schmidt, et al. 2005; Fleury et al. 2004; Wilfert & Jiggins, 2014;

Stephan & Li, 2006). While anti-predator behaviours are well studied as targets of

selection in prey (Juliano & Gravel, 2002; Stoks, McPeek & Mitchell, 2003; Magurran et

al. 1992), the genetic bases of such behaviours are seldom investigated. Given the range

of genetic and genomic tools available for *D. melanogaster,* along with its complex

behavioural repertoire and suitability for experimental evolution, understanding the anti-

predatory behaviours persisting in a natural population of the fruit fly brings us one step

closer to deciphering the molecular mechanisms for anti-predator behaviours.

Previous work has examined the effects of natural enemies on population and

community structures of *Drosophila spp*. Worthen (1989) studied the effects of predation

by rove beetles (staphylinids) on the coexistence of three mushroom-feeding *Drosophila*

species, and Escalante & Benado (1990) showed that ant predators regulate population

densities of wild *D. starmeri* (cactophillic fruit fly). In *D. melanogaster per se*, the role of

parasites in influencing larval and adult behaviours has been extensively studied (Milan,

Kacsoh & Schlenke, 2012; Kacsoh et al., 2013; Polak & Starmer, 1998). Despite this

literature, we know little about the predators of *D. melanogaster* adults in the wild, nor

the nature of anti-predatory behaviours segregating in natural populations.

We documented the influence of two predators, the zebra jumping spider (*Salticus*

scenicus) and juvenile Chinese praying mantids (*Tenodera aridifolia sinensis*) on the

capture-deterrence behaviours of *D. melanogaster* individuals derived from a wild-caught

population. The zebra spider is an active hunter, locating prey visually (with an extensive

visual field attained by antero-medially positioned simple eyes) (Dill, 1975; Horner,

Stangl & Fuller, 1988). Mantids are generally ambush predators, waiting for prey to enter

their attack range (Prete, Klimek & Grossman, 1990). Despite numerous differences,

zebra spiders and juvenile Chinese mantids are similar in two relevant ways. First, both

species primarily detect prey visually (Forster, 1979; Harland, Jackson, Macnab, 1999;

Jackson, & Blest, 1982; Prete 1999) and are likely incapable of depth perception when

their prey item is motionless (Prete, 1999; Freed, 1984). Second, small adult diptera

account form a substantial proportion of the diet of both predators in the wild (Iwasaki,

1998; Okuyama, 2007).

Based on the findings of Schmitz (2008), we predicted that fruit flies, in the

presence of a familiar predator, would exhibit hunting-mode specific modifications in

activity levels. To maximize distance from the actively hunting spider, our prediction was

that flies would increase their overall activity levels, whereas, to reduce the probability of

encountering a stationary threat (the mantid, an ambush predator), we expected flies to

decrease overall activity.

Under controlled laboratory conditions, we documented the behaviours of

individual adult *D. melanogaster* with and without the two predator species. Our results

suggest that in the presence of zebra spiders, *D. melanogaster* increases its overall

locomotory activity, performs a distinct "stopping" behaviour and increases the

performance of a newly described abdominal lifting behaviour (the function of which is

as of yet unknown). Counter to our prediction though, *D. melanogaste*r's locomotion, and

most other behaviours are not substantially altered in the presence of mantids. However,

upon direct encounter with a mantid, many individuals of *D. melanogaster* perform (a

previously undescribed) retreat behaviour- a response not generally elicited by jumping

spiders. Furthermore we observe considerable intra- and inter-individual variation in

response to predators. We discuss our results in terms of conditionally expressed

behaviours as they relate to predator hunting mode, co-evolutionary history of predators

and prey, and in terms of broadening our understanding of the behavioural ecology of *D.*

melanogaster.

METHODS

Drosophila Population and Culture Conditions

The *Drosophila melanogaster* population used in this study originated from a

natural population at Fenn Valley Vineyards in Fennville, Michigan (GPS coordinates:

42.57, -86.14) during the summer of 2010. A lab population (henceforth referred to as

FVW) was initiated from this collection using the progeny of over 500 single-pair

matings of field caught *D. melanogaster* as well as wild caught males. This design

allowed us to screen out the sympatric congener, *D. simulans,* which was present in our

collections at a frequency of about 5%. Screening involved setting up single pair mating

in vials and discarding all lines with *D. simulans*-like genital morphology. After

screening, ~1500 individuals were placed into cage ($32.5cm^3$, BugDorm BD43030F) to

establish the FVW population. The population is currently maintained in this cage at an

adult density ~ 3000 individuals in a room maintained at 23^0C (+/- 1^0C), and 40-70% RH.

Adults were allowed to lay eggs in 10 bottles with 50-60 ml of a standard yeast-cornmeal

food for 2-3 days. These bottles were then removed and kept in a Percival incubator

(Model: I41VLC8) at 24^0C and 65% RH throughout the larval stages. All flies and larvae

were maintained in a 12 hr light/dark cycle with lights on at 08:00 hours.

For the experiments, pupae were collected 24 hours before they emerged as

adults. Pupae were removed from bottles using forceps and individual pupae were placed

into 1.5 ml microcentrifuge tubes. Each tube was pre-filled with ~ 0.5 ml of yeast-

cornmeal food and its cap was punctured for gas exchange. Upon emergence, adult flies

were sexed visually without anesthesia and housed in these tubes in the incubator until

needed for behavioural assays. Age of flies used in behaviour analysis was 3-7 days. By

using socially naïve flies in our assays, we were able to establish a consistent baseline of

social experience among all individuals, allowing us to eliminate the potentially

confounding influence of variation in social experience on behaviour that is well-

documented in *Drosophila* (Yurkovic et al., 2006; Levine, 2004; Krupp et al., 2008;

Lefranc et al., 2001; Chabaud et al., 2009).

Spiders

S. scenicus individuals were collected throughout the spring/summer of 2012 on

the campus of Michigan State University. Spiders were housed individually in vials in a

room maintained at 23^0C (+/- 1^0C) and 30-50% RH and fed ~5 *D. melanogaster* a week.

Prior to use in behavioural assays, spiders were starved for at least 48 hours. Each spider

was used in only a single behavioural assay.

Mantids

Mantid egg cases were both collected near the campus of Michigan State

University as well as ordered from Nature's Control (Medford, Oregon). Mantid egg

cases were stored at 4^0C and transferred to 25^0C and 70% RH for hatching. Given the

substantial changes in mantid body size across moults (Iwasaki, 1990), only first instar

nymphs were used for experiments. Prior to behavioural assay, mantids were starved for

at least 24 hours and each mantid was used only once.

Behavioural Assays

All assays were performed 1-4 hours after the incubator lights came on in the

morning (08:00). Behavioural assays were recorded with an Aiptek AHD H23 digital

camcorder attached to a tripod under a combination of natural and fluorescent light that is

present in the room wherein the FVW population and spiders are maintained. For each

predator (spiders and mantids), we recorded the behaviour for each of 15 male and 15

female socially naïve, virgin flies (collected as described above). We used a chamber

constructed from the bottom of a 100 x15mm petri dish inverted on top of a glass plate

with a sheet of white paper beneath to maximize the visibility of flies and predators.

For each assay, an individual fly was aspirated into the chamber and allowed to

acclimate for 5 minutes. After this acclimation period, flies were recorded for 5 minutes.

A single spider or mantid was then introduced to the chamber and behaviours were

recorded for an additional 10 minutes or until capture. The chamber was washed with 10-

30% ethanol and rinsed with reverse osmosis water after each assay to remove olfactory

cues.

.

Behaviours Recorded

All *Drosophila* behaviours were categorized and analysed as either "states" or

"events". Behavioural states have measurable duration and are mutually exclusive with

other states (e.g. individuals cannot simultaneously walk and run). Behavioural events

are discrete behaviours that occur instantaneously and are also mutually exclusive with

each other (e.g. turning versus jumping) but not always mutually exclusive with

behavioural states. For example, an individual could perform a wing display (event) while

simultaneously walking (state), but it could not jump (event) while simultaneously

running (state). In this study we treated flying as an event because the structure of the

experimental chamber constrained flight duration. Attempted flight by *D. melanogaster*

could result in landing due to contact with a wall of the petri dish. We also recorded when

a fly was not visible (occluded) to the observers analysing video. We recorded a total of 6

discrete events and 5 behavioural states in *D. melanogaster* in response to predation by

spiders and mantids (Table 1). In order to interpret an individual fly's behaviour in the

context of predatory encounters, we designated two keys to describe the location of the

predator in regard to its interactions with the fly. As flies might alter their behaviour

when a predator is within striking distance, we recorded predator location based on

whether or not it was within striking distance of the fly (~ 5mm from the spider/mantid,

also see *Spider location/ Mantid location* in Figure 1).

Video Processing

Recorded behaviours were viewed with VLC media player (version 2.0.3) and

analysed by two observers using a manual event recorder, JWatcher V1.0 software

(Blumstein, 2006). One observer (A.P.) viewed each video and verbally announced the

occurrence of behaviours while the other observer (C.P./ M.C.) recorded the occurrence

of these behaviours with JWatcher. Because *Drosophila* anti-predatory behaviours are

often complex and occur rapidly, we analysed all videos at 0.5X speed.

Controlling for effects of season and disturbance

We conducted experiments with spiders between October and December 2012 and those

with mantids from March and May 2013. To confirm that predator species-specific

behavioural differences were not confounded with seasonal differences in behaviour, we

performed 6 additional assays (alternating between spider and mantid treatments) within

the span of one week. Following a spider assay, the plates were wiped down with 30%

ethanol followed by a rinse with RO water before a mantid assay was conducted.

Additionally, the process of adding a predator to the arena invariably resulted in a

disturbance that likely startled the fly (unrelated to the presence of a predator). To

confirm that behaviours induced by this disturbance were not confounded with predator

induced behavioural differences, we performed 3 control assays. Here, after 5 minutes of

acclimatization without a predator (see above for more details), the arena containing the

fruit fly was disturbed gently (~ magnitude of disturbance caused by the addition of a

predator). For all controls, video processing and behaviours recorded were identical to

mantid and spider treatments described above. See Supplement b, S1 for a detailed

description of these control experiments and their results.

Data processing and statistical analysis

A custom Python script was used to parse Jwatcher formatted data files into a

comma-separated-value (CSV) file for analysis in **R** (version 3.0.1).

To analyse the effects of predator state (i.e., presence or absence of predators) on

the time dedicated to locomotory behavioural states, and number of occurrence for

behavioural events, we fit generalized linear mixed effects models (using both `glmer`

function in the lme4 package version 1.0-5, and the `MCMCglmm` function in the

MCMCglmm package version 2.17) with predator state, total duration of assay with and

without a predator (duration), sex, temperature and recording time as fixed effects, and

individual by predator state and date as random effects . Formally, the model was:

231                 $y \sim \beta_0 + \beta_1 PS + \beta_2 D + \beta_3 Ag + \beta_4 T + \beta_5 ST + \beta_6 Sx + \beta_7 + \varepsilon$

Where y is a vector of time spent in a behavioural state. β_1 is the regression coefficient for

predator state, β_2 is for duration in each predator state, β_3 is for age of the fly, β_4 is for

temperature, β_5 is for time at which assay was started, β_6 is for sex of the fly and β_7 is for

date on which the assay was performed. We estimated random effects for individuals

including variation in response to predator state and duration of assay, and we fit an

independent random effect for date. Thus we fit a repeated effects (longitudinal) mixed

effects model allowing for variation among individuals for the influence of predator

presence and duration of assay where for the i^{th} individual

$$
\begin{pmatrix} \beta_{0i} \\ \beta_{1i} \\ \beta_{2i} \end{pmatrix} \sim MVN \left(\begin{bmatrix} \mu_{\beta_0} \\ \mu_{\beta_1} \\ \mu_{\beta_2} \end{bmatrix}, \begin{bmatrix} \sigma^2_{\beta_0} & \sigma_{\beta_0,\beta_1} & \sigma_{\beta_0,\beta_2} \\ \sigma_{\beta_0,\beta_1} & \sigma^2_{\beta_1} & \sigma_{\beta_1,\beta_2} \\ \sigma_{\beta_0,\beta_2} & \sigma_{\beta_1,\beta_2} & \sigma^2_{\beta_2} \end{bmatrix} \right)
$$

243   and (independent of the above)

244                         $\beta_7 \sim N(0, \sigma^2_j)$ where $j = 1 \dots$ date

Preliminary analyses were inconsistent with the need to fit higher order interactions

among fixed effects, so interaction terms were not considered further. The one exception

was for "stopping" behaviour where individuals almost exclusively performed this in the

presence of the predators. For the behavioural states (locomotion, grooming and

stopping) we assumed normally distributed variation. For the counts of events (abdominal

lift, jumping, etc) we used a log-link function and assumed the variation was poisson

distributed. Estimation using both maximum likelihood (`lmer`) and simulating the

posterior distribution (`MCMCglmm`) provided similar results for fixed effects, and

generally for random effect components as well.

Among individual coefficients of variation were calculated by dividing the square root of

the among individual variance component from the model by its respective fixed effect

estimate (i.e. its "mean"). While confidence intervals were consistent for fixed effects,

the intervals were more difficult to estimate given the complexities of the random effect

structure of the model, and some caution is warranted for their interpretation.

260   To test for non-random associations in the temporal structure of behavioural

patterns we constructed transition frequencies using the "msm" library (version 1.2)

(Jackson, 2011) in **R**. To test for both for first order Markov processes between

behaviours (transition probabilities), as well as the influence of predator presence on

these transition probabilities, we fit log-linear models (assuming poisson distributed data)

with the transition frequency matrices (Crawley, 2012) using `glm` in **R**. As advocated by

(Crawley, 2012; Bakeman & Gottman, 1997) we fit a saturated log-linear model (with

lag0, lag1 and predator state as the effects in the model) and tested the influence of

Table 1 Names and descriptions of all observed behaviours. Videos are provided at the end of Supplement b.

Behaviour	Description
Abdominal lift (ab)	Momentary rearing up on abdomen (see video 1)
Fly	Moving through space by wing use
Jump	Instantaneous movement between points without wing use
Pause	Noticeable period of inactivity; transitional
Turn	180 degree change in orientation without change in position
Wing display (wd)	Momentary lifting up of wings without singing or vibration
Groom	Running legs over any body part-often while otherwise stationary
Walk	Movement through space by ambulation
Run	Rapid movement through space by ambulation
Stop	Immobile (see video 2)
Retreat	Walking in reverse upon encounter with an object (like a predator) (see video 3)

deleting the terms (i.e. third order interaction) on change in deviance. We used modified

"Z-scores", adjusted using sequential Bonferroni to assess the deviation of particular cells

in the transition frequency matrix from expected values (assuming independence). For the

visual transition probability matrices, we combined the behavioural event "pause" with

the behavioural state "stop" because 1) we wanted to reduce the complexity of the matrix

and 2) the main difference between the two behaviours is that pause is instantaneous and

stop has duration. All transition diagrams were constructed in Inkscape (version 0.48.2,

Harrington, 2004-2005).

RESULTS

From pilot observations (not included in analysis), we (I.D., A.P. and C.P.)

catalogued and described *Drosophila melanogaster* behaviours observed in the presence

of a predator (Table 1). Among the behaviours listed in Table 1, abdominal lifting (ab,

supplement b, video 1) and retreat (supplement b, video 3), to our knowledge, have not

been previously described in *D. melanogaster* literature.

Flies perform a range of anti-predatory behaviours in response to a zebra spider

To visualize each individual fruit fly's response to the presence of a zebra

jumping spider, we generated ethograms (see Figure 1a and Supplement a). For the two

predator states (spider present and spider absent) we measured the mean proportion of

time dedicated to each behavioural state, as well as the number of occurrences per minute

for each behavioural event. When a spider was present, *D. melanogaster* increased the

proportion of time it spent walking and running by 50% (95% CI: 21-79% increase)

(a)

(b)

Figure 1

D. melanogaster used a greater proportion of its behavioural repertoire and performed each behaviour at a higher frequency in the presence of a jumping spider than in the presence of a juvenile mantid. a) Representative ethogram of a male, 4 day old *D. melanogaster* in response to a zebra jumping spider. b) Ethogram of a male, 5 day old *D. melanogaster* in response to a juvenile Chinese paying mantid. Light grey background represents time in the arena before the addition of a predator and dark grey background is when the predator was present in the chamber. Each black bar represents the occurrence of a behaviour during the observation period. Top half of the figure (separated by *Predator location*) consists of events and the bottom half consists of states. Because states have duration, the width of each black bar corresponds to the duration of a state. *Predator location* (i.e., *Spider location* in a and *Mantid location* in b) indicates whether or not the predator was within striking distance of the fruit fly at that time point. This information is relevant only after the predator was added to the chamber (~ 300 s into the assay). Dark grey bars in *Predator location* indicate that the spider was within striking distance and light grey regions indicate that the spider was out of striking distance. *Predator location* is white when the predator is absent from the arena or after successful capture. If capture did not occur, *predator location* remains light grey in colour.

Figure 2

Fruit flies increase overall activity levels in the presence of jumping spiders (a) and (b) but not in the presence of mantids (c) and (d). Plots (a) and (b) show change in mean number of occurrences per minute of each behavioural state as a result of the addition of a predator. Plots (c) and (d) show mean change in percentage of total time spent in a given behavioural state caused by the addition of a predator. On the left of the dotted line, behavioural changes correspond to the presence of a spider whereas on the right of the dotted line, behavioural differences are due to the presence of a juvenile praying mantid. Error bars are ± 95% CI.

while grooming 60% less (95% CI: 43-77% decrease). This is shown in Figure 2 and

Supplement b Figure S1 (treatment contrasts with 95% CI in figures are provided to

enable assessment of significance). While they were observed at low frequencies prior to

the addition of spiders, *D. melanogaster* substantially increased the frequency of pauses,

jumps and flights (per minute) in the presence of spiders (Figure 2, Supplement b Figure

S3). For instance the frequency of abdominal lifts increased from 1.51/minute to

4.0/minute (95% CI: 1.55-9.93), while jumping showed a 6.6X increase from 0.73/minute

to 4.82/minute (95% CI: 1.77-12.17). "Stopping"; a motionless state that likely aids in

capture deterrence (see Videos 2, Supplement b) was not performed by *D. melanogaster*

in the absence of spiders (Supplement b Figure S2). However in the presence of spiders,

the average total time spent "stopping" increased to ~25.8 seconds (95% CI: 10.1 - 41.7

seconds). When interacting with spiders, flies were only observed to perform the

"retreat" behaviour once (of 30 individuals). Interestingly, we did not see significant sex

specific differences in either frequencies of occurrence (Supplement b Figure S3) or

proportion of time allocate (Supplement b Figure S1) to the majority of measured

behaviours (But see S3 panels "pause" and "turn").

Given the design of our experiment, we were able to model the degree to which

individuals varied in their responses to the jumping spiders. Individuals varied greatly

both in their baseline activity levels as well as in their propensities to respond to jumping

spiders. The among-individual coefficient of variation for time spent grooming in the

absence of predators was 57.7% (40.1-74.2%). While most individuals reduced their

grooming activity in the presence of predators, the degree to which they did so varied

substantially, with the among individual coefficient of variation for the decrease being

67.8% (26.5-94.6%), as shown in Figure 3a. For walking, the among-individual

coefficient of variation was 80.3% (50-105%) in the absence of the spider, and 135% (1-

181%) for the magnitude of increase in the presence of the spider (Figure 3b).

Performance of the stopping behaviour by *D. melanogaster* in the presence of spiders

varied substantially among individuals, with the among-individual coefficient of variation

being 168% (95% CI: 123-214%). This is driven in part by the fact that 40% of

individual flies never performed stopping, even in the presence of the spider. There was

a negative correlation (-0.84), between the amount of time individuals spent grooming

before and after the addition of the spiders (Table 2). That is, on average, individuals who

were more active prior to the addition of the spider reduced their activity to a greater

extent in the presence of the spider. A similar negative correlation (-0.66) for among

individual activity for locomotion, was observed (Table 2).

To visualise the temporal associations among behavioural sequences, we

constructed transition matrices (Supplement b Tables S1, S2, S5 and S6) and transition

probability diagrams for all pairs of behaviours in the presence (Figure 4a) and absence

(Supplement b Figure S7) of spiders. In response to jumping spiders, transitions among

behaviours are somewhat more dispersed (with many connections between behaviours),

suggesting that there is weak temporal association between fruit fly behaviours. Indeed

these qualitative conclusions are supported based on the Z-scores. In the absence of

spiders 8 possible transitions were significant (after controlling for multiple comparison,

Supplement b Table S2), while 13 transitions were significant in the presence of the

spider (Supplement b Table S1). Most of these differences were due to the increase in

behaviours potentially involved with anti-predation activity (i.e. flight, abdominal

Figure 3

Inter individual behavioural variation in response to predators is present in natural populations. Reaction norms visualize how each individual fruit fly responded to the introduction of a spider (panels a and b) or a mantid (panels c and d) into the assay chamber. Measures are in seconds. Each line corresponds to response of one individual. Estimates are derived from the predicted values for each individual from the mixed models.

Table 2 Individual flies show negative correlations between behavioural states before and after the introduction of a predator. There is considerable variation among individuals in time spent performing specific behaviours (i.e. walking and grooming), with and without predators. However, there is a strong negative correlation within individuals for time spent before and after introduction of the predator. That is, individuals who spend more time performing a specific behaviour prior to the addition of a predator, reduce that behaviour to an even greater amount (than the average for the sample) once the predator is introduced. The one exception is for grooming for the mantid trials. Diagonals of the table contain the standard devation (mean of the posterior distribution) for individual behavioural responses (95% CIs in paratheses) from the random effects of the models. Above the diagonal are covariances between predictors (and CIs in parantheses). Below the diagonal are correlation coefficients for the covariances between the predictors.

Grooming, Spider

	Intercept	Pred.State	Time
Intercept	89.2 (62.0, 115.3)	-68.2 (-88.7, -34.1)	26.4 (14.0, 36.0)
Pred.State	-0.84	61.7 (18.8, 84.5)	-14 (-25.5, 17.1)
Time	0.75	-0.3	10.4 (4.4, 14.9)

Walking, Spider

	Intercept	Pred.State	Time
Intercept	81.9 (43.3, 109.8)	-60.2 (-89.5, 27.5)	25.6 (-8.8, 36.7)
Pred.State	-0.66	67.3 (0.36, 98.5)	11.5 (-22.2, 29.2)
Time	0.62	0.15	13 (5.6, 18.2)

Grooming, Mantid

	Intercept	Pred.State	Time
Intercept	122.8 (59.6, 175)	-20.2 (-117, 106)	46.0 (-16.9, 74.5)
Pred.State	-0.05	60.5 (0.13, 100.2)	-18.3 (-47.9, 34.0)
Time	0.8	-0.26	21.5 (2.6, 33.8)

Walking, Mantid

	Intercept	Pred.State	Time
Intercept	144.8 (86.2, 198)	-100.3 (-162.6, 38.8)	63.2 (31.6, 90.3)
Pred.State	-0.86	80.5 (0.21, 139.3)	-45.2 (-76.4, 19.6)
Time	0.94	-0.86	29.4 (11.3, 43.5)

lifting). However, while the results of the log-linear analysis (across the whole transition

frequency matrix) supported the dependence of current behavioural states on the previous

state (resid df=71, deviance=632, p < 0.001), the inclusion of predator status did not

influence this dependence (resid df = 71, deviance = 59, p = 0.8).

Flies perform a previously undescribed retreat behaviour in response to mantids

In contrast to their behaviour in the presence of jumping spiders, the presence of a

juvenile mantid had a minimal influence on *D. melanogaster*'s locomotory activity

(Figure 2c and 2d, also see ethograms in Figure 1b and Supplement a). Time spent

grooming, walking, running and stopping was highly variable (with estimates including

zero) in the presence of a juvenile praying mantid (Figure 2d, Supplement b Figures S1

and S2). Grooming decreased by 20% (95% CI: 83% decrease to 36% increase),

locomotion increased by 9% (13% decrease to 32% increase) and stopping increased by

40% (8% decrease to 85% increase). Similarly, the presence of a mantid did not influence

the frequency at which *D. melanogaster* tended to perform most instantaneous behaviours

(Figure 2c, Supplement b Figure S4). However, as was observed in the presence of

spiders, flies substantially increased the frequency of abdominal lifting, from 0.45/minute

to 5.6/min (95% CI: 2.56 - 12.17) in the presence of a juvenile mantid (Supplement b

Figure S4). Upon encounter with a mantid, half of the individuals (15/30) performed a

previously undescribed reversal behaviour (Supplement b video 3), which we term

"retreat". As with the zebra spiders, we saw no significant sex specific differences in

response to mantids.

Although the presence of a mantid had a small effect on fly behaviour, flies did

vary considerably in their grooming and walking activities. Indeed, the among-individual

variability in proportion of time spent grooming and walking is greater in magnitude in

the presence of the mantids than spiders (Figure 3). Evidence for negative co-variation

for intra-individual behaviour before and after the addition of the predator was not

strongly supported (i.e. 95% CIs for covariances included zero) (Table 2).

Transition matrices and transition probability diagrams (Supplement b Figure

S7b, Figure 4b and Tables S3, S4, S7 and S8) show patterns of temporal association

among behaviours. In response to juvenile mantids, the transitions diagram is less

dispersed than that in the presence of jumping spiders (Figure 4), suggesting that the

degree of association between behaviours in the presence of mantids is more extreme.

While most behaviours (abdominal lift, fly, groom, jump, run, stop, and turn) tend to

transition to walking, we also see stronger associations between other pairs of behaviours.

For example, after performing the retreat behaviour, flies often either performed the

abdominal lift or turn, while flight is often followed by stopping. These observations are

supported by the findings that in the absence of mantids, 12 transitions showed

significant deviations from expectations (Supplement b, Table S4). In comparison, in the

presence of mantids 23 transitions showed a significant deviation from expected values

(Supplement b, Table S3). Interestingly, as with the spiders the log-linear model supports

the non-independence of behavioural states (resid df=71, deviance=1054, p <0.001), but

not for the additional influence of predator state on this non-independence (resid df = 71,

deviance = 72, p=0.4).

(a)

(b)

Figure 4

Spiders and mantids had different effects on the temporal associations between pairs of *D. melanogaster* behaviours. a) A diagram representing probability of transitioning from one fly behavioural state to the other in the presence of a zebra jumping spider. b) A diagram representing probability of transitioning from one fly behavioural state to the other in the presence of a juvenile praying mantid. Thickness of arrows indicates transition probability between the two behaviours. The arrowhead points to the behaviour being transitioned to. Thickness of the box around behavioural state (groom, run, occl, retreat, stop and walk) indicate the mean proportion of total time spent in that behaviour, whereas thickness of the box around behavioural events (fly, jump, turn, wd, ab) indicates mean number of occurrences per minute of that behaviour. To reduce the complexity of the web we combined the behaviours "pause" with the behaviour "stop". Behavioural transitions that occurred less than 10 times have not been shown in the figure.

DISCUSSION

Prey organisms can alter their behaviour to reduce the likelihood of detection,

capture or encounter with a predator (Lima, 1998). For example, when predators are

present, ground squirrels dedicate more time to vigilance behaviours (like scanning for a

predator, see (Bachman, 1993) and some aquatic insects spend more time in refuges

(Kohler & McPeek, 1989). These changes in behaviour may alter the use of resources,

and potentially the fitness of an organism. However, the nature and intensity of non-

consumptive effects of a predator on its prey are a function of several predator specific

factors, one of which is the predator's hunting mode (Preisser, Orrock & Schmitz, 2007).

Predator hunting mode, i.e., the set of behavioural strategies that a predator employs to

pursue and capture its prey (Schoener, 1971; Huey & Pianka, 1981; Schmitz, 2008) can

be an important determinant of a prey organism's anti-predatory behavioural response

(Schmitz, 2008). In this study, we describe the anti-predatory behavioural repertoire of a

natural population of *Drosophila melanogaster* in response to predation by the zebra

jumping spider (*Salticus scenicus)* and juvenile Chinese praying mantids (*Tenodera*

aridifolia sinensis). Among other characteristics, zebra spiders and praying mantids differ

in their hunting mode. While we discuss our findings with respect to hunting mode

differences, we recognize that other attributes differing among the predators may

contribute to the observed differences in prey behavioural repertoires. However, as our

experimental design was meant to minimize the effects of many possible confounding

factors (e.g. time of day, temperature, humidity) it seems likely that, in part, our results

reflect hunting mode differences.

In response to active hunters (those that constantly patrol for prey), we predicted

that fruit flies would increase their overall activity levels (including flight) in order to

maintain maximum distance from the predator at all times; To reduce the likelihood of an

encounter with an ambush predator however (i.e., a predator that only attacks when a

prey organism wanders in to its strike zone), we predicted that *D. melanogaster* would

respond by decreasing locomotory activities (Schmitz, 2008). Our results, however, were

only partially in line with these predictions. While the actively hunting jumping spiders

induce a clear increase in overall activity, we found the presence of juvenile mantids- our

ambush predators- to have minimal influence on fruit fly activity levels (Figure 2,

Supplement b Figure S2). It has been previously argued that ambush predators might be a

predictable source of threat to prey organisms (Preisser, Orrock & Schmitz, 2007;

Schmitz, 2008) as opposed to the diffuse and variable threat imposed by active hunters

(Schmitz, 2008). Therefore, it is perhaps surprising that fruit flies show a stronger

behavioural response to the threat of active hunters (zebra jumping spiders). However,

our predictions are based on studies on a grasshopper and its two predatory spider species

that differ in hunting mode. Given that selection pressures faced by adult diptera are

different from those experienced by grasshoppers (orthoptera), such predictions may not

be generalizable. Several factors including body size and dispersal patterns may

contribute to this difference. Many species of jumping spiders, including *S. scenicus,* are

often seen in the natural habitat of *D. melanogaster* (personal observations of A.P., C.P.

and I.D.), and are likely to be ecologically relevant predators of *Drosophila*. Mantids

however, are rarely found in areas where fruit flies are abundant (personal observations

of A.P. and I.D.), at least in Eastern North America. Therefore, it is likely that fruit flies,

having experienced a longer evolutionary history with small jumping spiders, are better

able to recognize these spiders as a threat. In addition, the disturbance created by a

constantly patrolling zebra spider may be partly responsible for the increased activity

levels seen in *D. melanogaster* (either due to actual mechanical disturbance or because

flies are able to detect moving objects quicker than stationary ones). In this study, we are

unable to tease apart the effects of evolutionary recognition versus constant mechanical

disturbance on the differences in flies' activity levels. Further experimentation with

harmless but constantly moving heterospecifics (such as field crickets) or immobilized

active hunters might be useful in addressing these issues.

We also identified a number of (to our knowledge) undescribed behaviours of *D.*

melanogaster, potentially relating to its interactions with predators. The behaviour we

called "stopping" (Table 1) was observed numerous times after a direct (but failed) attack

by a spider (Supplement 3 video 1). While *D. melanogaster* will spend time without any

ambulatory activity (walking, running), they are almost always observed to be active

(generally grooming) during these periods. However, when fruit flies performed the

stopping behaviour, there was a complete lack of movement on the part of the fly, even

when video was viewed at a few frames/second. When a fruit fly was "stopped", the

spider had to search for the fly, irrespective of the physical proximity between the spider

and the fly. In salticids, while the principal eyes have high spatial acuity, secondary eyes

are primarily used to detect moving objects (Harland, Jackson & Macnab, 1999; Land,

1971). Because salticids are unable to accommodate by changing the shape of their lens,

they need to extensively sample their visual field to see details in object shape and form

(Harland, Jackson & Macnab, 1999; Land, 1971; Blest, Hardie & McIntyre, 1981).

Scanning for prey by such sampling is likely a slow process unless guided by the motion

sensing peripheral eyes, giving motionless prey the advantage of staying hidden (at least

for a few seconds) while in plain sight of their salticid predator. Thus, *D. melanogaster*

may be using the "stopping" behaviour as a potential mechanism to reduce the likelihood

of detection by the spider.

Additionally, in the presence of both predators, *D. melanogaster* substantially

increase the frequency at which it performed abdominal lifts. To our knowledge,

abdominal lifting has not been described in *D. melanogaster* literature before and may be

relevant in an anti-predatory context. While studying courtship behaviours in female *D.*

melanogaster, Lasbleiz, Ferveur & Everaerts (2006) described two behaviours perhaps

similar to the abdominal lifting described here: abdominal drumming and abdominal

extension. Abdominal drumming (described as "quickly repeated vertical movements of

the abdomen which is tapped on the substrate") was only seen in males during courtship

display, and abdominal extensions (described as " abdomen raised by 15-30 degrees")

were also closely associated with courtship. Because abdominal lifting was often directed

at a predator or followed a failed predatory encounter, we suspect abdominal lifting to be

different from abdominal extensions and abdominal drumming, and with a possibly anti-

predatory function. We speculate that if abdominal lifting is indeed anti-predatory, it

could function in one of several possible ways. First, abdominal lifting may be a signal of

prey condition directed at the predator as a form of pursuit deterrence, comparable to

stotting in the Thomson's gazelle (FitzGibbon & Fanshawe, 1988). Second, because *D.*

melanogaster are often surrounded by conspecifics, abdominal lifting may be a means

though which one fly warns its conspecifics of the presence of a potential threat (similar

in function to fin flicking in tetras, (Brown, Godin & Pedersen, 1999). Finally it may be

an indication of some sort of physiological priming of the fly in preparation for a fight-

or-flight response. Determining whether it is a specific anti-predator behaviour, as well as

the details of its function need to be a focus of future work.

In response to the juvenile praying mantids, half of the fruit flies we observed

(15/30) performed a reverse walking behaviour which we have called "retreat", where the

flies walked in reverse, away from the predator (supplement b, video 3). This was often

(but not always) interspersed with the abdominal lifting behaviour. Phenomenologically,

this behaviour may be similar to that described in Bidaye et al. (2014). Bidaye el al.

(2014) identified neurons that upon activation changed walking direction in *D.*

melanogaster. Bidaye et al's reverse walking behaviour appears to be a smooth and

continuous behaviour, whereas "retreat" was often discontinuous and accompanied by

abdominal lifting. If the two "retreat" behaviours are related, the observed disassociation

between retreat and abdominal lifting as well as its continuous nature (in Bidaye el al.,

2014) may be a function of how the neurons were perturbed.

We also investigated how the presence of the different predators may influence

non-random associations among behaviours. We observed that in the presence of both

predators there was an increase in the number of behavioural transitions that deviated

from expectations under independence (from 12 to 23 with the mantid, and 8 to 13 with

the spider). Despite this, the log-linear model (analysing the whole transition frequency

matrix) did not support the influence of predator state on the frequencies of transitions.

This may be partly due to the relatively modest sample sizes (in terms of both individuals

and transitions among behaviours). Further work is necessary to validate and extend this

sequential analysis.

While we show that there are some predator hunting-mode specific behavioural

differences in *D. melanogaster's* anti-predator response, we reiterate two important

caveats. First, although the primary distinction between the zebra jumping spider and

juvenile Chinese praying mantids as predators is their hunting-mode, other factors

between these species (for example, size, colour, odour) may also influence differences in

fruit fly behaviours. Replicating the observations with other predator pairs that differ in

hunting-mode is necessary to confirm hunting-mode's influence on anti-predatory

repertoires. Secondly, our assay chambers are an artificial environment and do not

resemble the conditions under which *D. melanogaster* face predators in the wild. Due to

the nature of our assay chamber, *D. melanogaster* were unable to employ behavioural

strategies that may reduce encounters with predators (e.g., utilizing a refuge). Therefore

we were only able to describe the capture-deterrence repertoire of *D. melanogaster*

behaviour. We believe that our study is a necessary first step to describing and

documenting the complete anti-predatory behavioural repertoire of *D. melanogaster* and

we foresee future work to be conducted in a modified chamber, under more "natural"

conditions. Doing so will allow us to take this premier model genetic system and make it

into an ecological model as well.

ACKNOWLEDGEMENTS

We thank Dr. Fred Dyer for loaning us video cameras. Thanks to Dr. Ben DeBivort for

discussions and suggestions and to Dr. Reuven Dukas and Dr. Marla Sokolowski and

members of the ███████ lab for comments on the manuscript. This material is based in

part upon work supported by the National Science Foundation under Cooperative

Agreement No. DBI-0939454. Any opinions, findings, and conclusions or

recommendations expressed in this material are those of the author(s) and do not

necessarily reflect the views of the National Science Foundation.

REFERENCES

Bachman, G. C. (1993) The effect of body condition on the trade-off between vigilance

and foraging in Belding's ground squirrels. Anim Behav 46: 233–244.

Bakeman, R. & Gottman, J. M. (1997) Observing interactions. New York: Cambridge

University Press.

Bidaye, S. S., Machacek, C., Wu, Y. & Dickson, B. J. (2014) Neuronal Control of

Drosophila Walking Direction. Science 344, 97–101. (doi:10.1126/science.1249964)

Blest, A. D., Hardie, R. C. & McIntyre, P. (1981) The spectral sensitivities of identified

receptors and the function of retinal tiering in the principal eyes of a jumping spider. J.

Comp. Physiol., A 145: 227-239.

Blumstein, D. T. (2006) The multi-predator hypothesis and the evolutionary persistence

of antipredator behavior. Ethology 112: 209–217. (doi:10.1111/j.1439-

0310.2006.01209.x)

Brown, G., Godin, J. & Pedersen, J. (1999) Fin-flicking behaviour: a visual antipredator

alarm signal in a characin fish, *Hemigrammus erythrozonus*. Anim Behav 58: 469–475.

(doi:10.1006/anbe.1999.1173)

Chabaud, M.A., Isabel, G., Kaiser, L. & Preat, T. (2009) Social facilitation of long-

lasting memory retrieval in *Drosophila*. Curr Biol 19: 1654–1659.

(doi:10.1016/j.cub.2009.08.017)

Crawley, M. J. (2012) The R book. John Wiley & Sons.

David, J. R. & Capy, P. (1988) Genetic variation of *Drosophila melanogaster* natural

populations. Trends Genet 4: 106–111.

DeWitt, T. J., Robinson, B. W. & Wilson, D. S. (2000) Functional diversity among

predators of a freshwater snail imposes an adaptive trade-off for shell morphology. Evol

Ecol Res 2: 129–148.

Dill, L. M. (1975) Predatory behavior of the zebra spider, *Salticus scenicus* (Araneae:

Salticidae). Can J Zool 53: 1284–1289.

Edmunds, M. (1974) Defence in animals: a survey of anti-predator defences. Longman

Harlow, New York.

Escalante, A. & Benado, M. (1990) Predation on the cactophilic fly, *Drosophila starmeri*,

in the columnar cactus, *Pilosocereus lanuginosus*. Biotropica, 48–50.

FitzGibbon, C. D. & Fanshawe, J. H. (1988) Stotting in Thomson's gazelles: an honest

signal of condition. Behav Ecol Sociobiol 23: 69–74.

Fleury, F., Ris, N., Allemand, R., Fouillet, P., Carton, Y. & Boulétreau, M. (2004)

Ecological and genetic interactions in Drosophila-parasitoids communities: a case study

with *D. melanogaster*, *D. simulans* and their common Leptopilina parasitoids in south-

eastern France. Genetica 120: 181–194.

Forster, L. M. (1979) Visual mechanisms of hunting behaviour in *Trite planiceps*, a

jumping spider (Araneae: Salticidae). NZ J Zool 6: 79–93.

(doi:10.1080/03014223.1979.10428351)

Freed, A. N. (1984) Foraging behaviour in the jumping spider *Phidippus audax*: bases for

selectivity. J Zool 203: 49–61.

Gosline, A. K. & Rodd, F. H. (2007) Predator-induced plasticity in guppy (*Poecilia*

reticulata) life history traits. Aquat Ecol 42, 693–699. (doi:10.1007/s10452-007-9138-7)

Harland, D. P., Jackson, R. R. & Macnab, A. M. (1999) Distances at which jumping

spiders (Araneae: Salticidae) distinguish between prey and conspecific rivals. J Zool 24:

357–364.

Harrington, B. et al (2004-2005) Inkscape. http://www.inkscape.org/

Horner, N. V., Stangl, F. B. & Fuller, G. K. (1988) Natural history observations of

Salticus austinensis in north-central Texas. J Arachnol 16: 260-262.

Huey, R. B. & Pianka, E. R. (1981) Ecological consequences of foraging mode. Ecology

4: 991-999.

Iwasaki, T. (1990) Predatory behavior of the praying mantis, *Tenodera aridifolia* I. Effect

of prey size on prey recognition. J Ethol 8: 75–79.

Iwasaki, T. 1998 Prey menus of two praying mantises, *Tenodera aridifolia* (Stoll) and

Tenodera angustipennis Saussure (Mantodea: Mantidae). Entomol Sci 4: 529-532.

Jackson, R. R. & Blest, A. D. (1982) Short Communication: The distances at which a

primitive jumping spider, *Portia fimbriata*, makes visual discriminations. J Exp Biol 97:

441–445.

Jackson, C. H. (2011) Multi-State Models for Panel Data: The msm Package for R. J Stat

Softw 38(8): 1-29. URL http://www.jstatsoft.org/v38/i08/'

Juliano, S. A. & Gravel, M. E. (2002) Predation and the evolution of prey behavior: an

experiment with tree hole mosquitoes. Behav Ecol 13, 301–311.

Kacsoh, B. Z., Lynch, Z. R., Mortimer, N. T. & Schlenke, T. A. (2013) Fruit Flies

Medicate Offspring After Seeing Parasites. Science 339: 947–950.

(doi:10.1126/science.1229625)

Kohler, S. L. & McPeek, M. A. (1989) Predation risk and the foraging behavior of

competing stream insects. Ecology 70: 1811–1825.

Krupp, J. J., Kent, C., Billeter, J.-C., Azanchi, R., So, A. K. C., Schonfeld, J. A., Smith,

B. P., Lucas, C. & Levine, J. D. (2008) Social experience modifies pheromone expression

and mating behavior in male *Drosophila melanogaster*. Curr Biol 18: 1373–1383.

(doi:10.1016/j.cub.2008.07.089)

Langerhans, R. B. (2007) Evolutionary consequences of predation: avoidance, escape,

reproduction, and diversification. Predation in organisms 177-220. Springer Berlin

Heidelberg. (doi: 10.1007/978-3-540-46046-6_10)

Land, M. F. (1971) Orientation by jumping spiders in the absence of visual feedback. J

Exp Biol 54: 119–139.

Lasbleiz, C., Ferveur, J.-F. & Everaerts, C. (2006) Courtship behaviour of *Drosophila*

melanogaster revisited. Anim Behav 72: 1001–1012.

(doi:10.1016/j.anbehav.2006.01.027)

Lefranc, A., Jeune, B., Thomas-Orillard, M. & Danchin, E. (2001) Non-independence of

individuals in a population of *Drosophila melanogaster*: effects on spatial distribution

and dispersal. C R Acad Sci III, Sci Vie 324: 219–227.

Levine, J. D. (2004) Sharing time on the fly. Curr Opin Cell Biol 16: 210–216.

(doi:10.1016/j.ceb.2004.02.009)

Lima, S. L. & Dill, L. M. (1990) Behavioral decisions made under the risk of predation: a

review and prospectus. Can J Zool 68, 619–640.

Lima, S. L. (1998) Nonlethal effects in the ecology of predator-prey interactions.

BioScience 48, 25–34.

Lima, S. L. & Bednekoff, P. A. (1999) Temporal variation in danger drives antipredator

behavior: the predation risk allocation hypothesis. Am Nat 153: 649-659

Magurran, A. E., Seghers, B. H., Carvalho, G. R. & Shaw, P. W. (1992) Behavioural

consequences of an artificial introduction of guppies (*Poecilia reticulata*) in N. Trinidad:

Evidence for the evolution of anti-predator behaviour in the wild. Proc R Soc B 248:

117–122. (doi:10.1098/rspb.1992.0050)

Milan, N. F., Kacsoh, B. Z. & Schlenke, T. A. (2012) Alcohol consumption as self-

medication against blood-borne parasitesin the fruit fly. Curr Biol 22: 488–493.

(doi:10.1016/j.cub.2012.01.045)

Okuyama, T. (2007) Prey of two species of jumping spiders in the field. Appl Entomol

Zool 42: 663–668. (doi:10.1303/aez.2007.663)

Polak, M. & Starmer, W. T. (1998) Parasite-induced risk of mortality elevates

reproductive effort in male *Drosophila*. Proc Royal Soc B 265: 2197–2201.

Preisser, E. L., Orrock, J. L. & Schmitz, O. J. (2007) Predator hunting mode and habitat

domain alter nonconsumptive effects in predator-prey interactions. Ecology 88: 2744–

2751.

Prete, F. R., Klimek, C. A. & Grossman, S. P. (1990) The predatory strike of the praying

mantis, *Tenodera aridifolia sinensis*. J Insect Physiol 36: 561–565.

Prete, F. R. (1999) The Praying Mantids 93-141. Baltimore: The Johns Hopkins

University Press.

Reaume, C. J. & Sokolowski, M. B. (2006) The nature of *Drosophila melanogaster*. Curr

Biol 16: R623-R628.

Relyea, R. A. (2001) Morphological and behavioral plasticity of larval anurans in

response to different predators. Ecology 82: 523–540.

Reznick, D., Butler, M. J., IV & Rodd, H. (2001) Life-history evolution in guppies VII.

The comparative ecology of high and low predation environments. Am Nat 157: 126–

140. (doi:10.1086/318627)

Sansom, A., Lind, J. & Cresswell, W. (2009) Individual behavior and survival: the roles

of predator avoidance, foraging success, and vigilance. Behav Ecol 20, 1168–1174.

(doi:10.1093/beheco/arp110)

Schmidt, P. S., Matzkin, L., Ippolito, M. & Eanes, W. F. (2005) Geographic variation in

diapause incidence, life-history traits, and climatic adaptation in *Drosophila*

melanogaster. Evolution 59: 1721–1732.

Schmitz, O. J. (2008) Effects of predator hunting mode on grassland ecosystem function.

Science 319: 952–954. (doi:10.1126/science.1152355)

Schoener, T. W. (1971) Theory of feeding strategies. Annu Rev Ecol Syst 2: 369–404.

Stephan, W. & Li, H. (2006) The recent demographic and adaptive history of *Drosophila*

melanogaster. Heredity 98: 65–68. (doi:10.1038/sj.hdy.6800901)

Stoks, R., McPeek, M. A. & Mitchell, J. L. (2003) Evolution of prey behavior in response

to changes in predation regime: damselflies in fish and dragonfly lakes. Evolution 57:

574–585.

Turelli, M. & Hoffmann, A. A. (1991) Rapid spread of an inherited incompatibility factor

in California *Drosophila*. Nature 353: 440–442.

Wilfert, L. & Jiggins, F. M. (2014) Flies on the move: an inherited virus mirrors

Drosophila melanogaster's elusive ecology and demography. Mol Ecol 23: 2093–2104.

(doi:10.1111/mec.12709)

Worthen, W. B. (1989) Predator-mediated coexistence in laboratory communities of

mycophagous *Drosophila* (Diptera: Drosophilidae). Ecol Entomol 14: 117–126.

(doi:10.1098/rspb.1998.0559)

Yurkovic, A., Wang, O., Basu, A. C. & Kravitz, E. A. (2006) Learning and memory

associated with aggression in *Drosophila melanogaster*. Proc Natl Acad Sci U.S.A. 103:

17519–17524. (doi:10.1073/pnas.0608211103)

Supplemental material

Table of contents

Supplemental material type	Page numbers
Supporting information	1-2
Supplemental figures	2-7
Supplemental tables	9-14
Videos	14

Supporting information

Differential response to spiders versus mantids

Because spider and mantid population densities vary by season, we had to temporally

segregate the spider assays from the mantid assays. We conducted all spider observations

between October and December 2012 and all the mantid observations between March and May

2013. Comparing time allocation and frequencies of occurrences in the predator absent state

between the two predator treatments suggest that behavioural modifications were predator

induced, and not due to seasonal effects (Figure S5 and S6). Although the assays were carried

out under highly controlled conditions, to confirm that predator species-specific behavioural

differences were not confounded with seasonal differences in behaviour, we performed 6

additional assays (alternating between spider and mantid treatments) within the span of one

week. The control experiments show no evidence of confounding effects of season with *D.*

melanogaster's anti-predator behavioural repertoire (Table S9, S11 and S12 below). Ethograms

are shown in Supplement a. Furthermore, to confirm that the disturbance we caused (to the assay

chamber) during the addition of a predator did not confound behavioural responses to the

predator, we did 3 "no predator" control assays. For these "no predator" controls, instead of

adding a predator to the arena, we caused a mild disturbance (~ to intensity of disturbance caused

while adding the predator) without actually adding any predator. We found that disturbance

caused during predator addition was not responsible for observed behavioural modifications

(Table S10 and S13). Finally, "no predator" controls also ruled our temporal differences in fruit

fly activity levels (Table S10 and S13)

Supplemental figures

Figure S1

Presence of jumping spiders causes fruit flies to walk more and groom less, whereas the presence

of mantids causes weaker, more variable (and not significant) changes in fruit fly activity levels.

Here we show coefficient plots from the output of mixed effects models using the package

MCMCglmm to visualize duration of two behaviours (**grooming in black** and **locomotion in**

red) as a function of predator state (present vs absent of spiders, left panels and mantids, right

panels), time spent in the assay [Time (cent.)] total, sex of the fly, start time of the assay,

temperature in the room and age of the fly. The continuous covariate "Time" was centered

around the mean and therefore reflects the average increase in time spent per minute of the assay.

Estimates are in seconds. Error bars are ± 95% CI.

41
Figure S2

Fruit flies "stop" significantly longer in the presence of spiders (left panel), and to a much lesser

extent (and not significantly) in the presence of mantids (right panel). Here we have coefficient

plots made from the output of mixed effects models using the package MCMCglmm to visualize

duration of "stopping" as a function of predator state (present vs absent), time spent in the assay

[Time (cent.)], sex of the fly, start time of the assay, temperature in the room and age of the fly.

Estimates are in seconds. Error bars are ± 95% CI. The continuous covariate "Time" was

centered around the mean and therefore reflects the average increase in time spent per minute of

the assay. Although assays were performed between 9 am and 12 pm each day, start time for the

mantid assays significantly affected the total time that flies spent "stopping".

Figure S3

In the presence of spiders, fruit flies increased the frequency with which they performed flights,

pauses and jumps. Here we show coefficient plots made from the output of mixed effects models

where the events were modeled using a poisson generalized linear mixed model with a log-link

function fit using the `MCMCglmm` function, and estimates remain on a natural log scale.

Coefficient plots were used to visualize frequency of each individual behavioural event (ab, fly,

pause, wd, turn and jump) as a function of predator state (present vs absent of a spider), time

spent in the assay [Time (cent.)], sex of the fly, start time of the assay, temperature in the room

and age of the fly. All estimates are scaled to number of events per minute. Error bars are ± 95%

CI.

Figure S4

Fruit flies performed abdominal lifts are a higher frequency in the presence of a juvenile mantid.

Coefficient plots made from the output of mixed effects models where the events were modeled

using a poisson generalized linear mixed model with a log-link function fit using the `MCMCglmm`

function, and estimates remain on a natural log scale. Coefficient of the model output are used to

visualize frequency of each individual behavioural event (ab, fly, pause, wd, turn and jump) as a

function of predator state (present vs absent of a mantid), time spent in the assay [Time (cent.)],

sex of the fly, start time of the assay, temperature in the room and age of the fly. All estimates

are scaled to number of events per minute. Error bars are ± 95% CI. Although assays were

performed between 9 am and 12 pm each day, start time for the mantid assays significantly

affected the frequency at which *D. melanogaster* performed the "Fly", "Wd" and "Jump"

behaviours.

Figure S5

Hunting mode induced behavioural differences in fruit fly behaviours were not confounded with

seasonal effects. Here we show percentage time spent in each behavioural state (left) and number

of occurrences per minute for each behavioural event (right) as measured for individual fruit flies

before the addition of a spider (white circles) and before the introduction of a mantid (black

circles) into the chamber. Error bars are ± 2 * SEs. Overlapping error bars suggest that there was

minimal effect of season on the behavioural repertoire of fruit flies.

Figure S6

Seasonal differnces in fruit fly behaviours did not confound behavioural differences induced by

different hunting-modes. Flies measured before the addition of a spider did not differ in

behaviour from flies measured before the addition of a mantid.

Figure S7

a) A diagram representing probability of transitioning from one fly behaviour to the other

when individuals were measured before the addition of a spider **b)** A diagram representing

probability of transitioning from one fly behaviour to the other for individuals measured

before the addition of a juvenile mantid. Thickness of arrows indicates transition

probability between the two behaviours. The arrowhead points to the behaviour being

transitioned to. Thickness of the box around behavioural state (groom, run, occl, retreat,

stop and walk) indicate the mean proportion of total time spent in that behaviour, whereas

thickness of the box around behavioural events (fly, jump, turn, wd, ab) indicates mean

number of occurrences per minute of that behaviour. To reduce the complexity of the web

we combined the behaviours "pause" with the behaviour "stop". Behavioural transitions

that occurred less than 10 times have not been shown in the figure.

 Supplemental tables

 Table S1 Transition frequency matrix when a spider was present in the chamber. Each row

 represents the number of times one behaviour (row name) transitioned to another behaviour

 (column name). Numbers in blue represent transitions that occurred more often that expected

 under a model of independence, whereas numbers in red are transitions that occurred less often

 than expected (see methods).

Spider

Behavior	ab	capture	fly	groom	jump	retreat	run	stop	occl	turn	walk	wd	Total
ab	0	2	**8**	9	7	0	**45**	6	5	10	**112**	7	211
capture	0	0	0	0	0	0	0	0	0	0	0	0	0
fly	4	2	0	15	7	0	**34**	**25**	10	1	58	1	157
groom	11	3	16	0	10	0	8	11	5	7	**77**	2	150
jump	4	2	19	8	0	0	9	13	2	0	28	0	85
retreat	1	0	0	0	0	0	0	0	0	0	0	0	1
run	**43**	4	**33**	13	9	1	0	18	3	4	51	9	188
stop	9	3	32	10	16	0	19	0	1	9	**38**	0	137
occl	3	0	6	4	5	0	4	4	0	1	14	0	41
turn	8	1	0	4	3	0	14	2	0	0	**66**	2	100
walk	**122**	7	36	**87**	26	0	42	60	14	**63**	0	24	481
wd	5	0	4	0	0	0	9	0	0	1	29	0	48
Total	210	24	154	150	83	1	184	139	40	96	473	45	1599

 Table S2 Transition frequency matrix before a spider was added to the chamber. Each row

 represents the number of times one behaviour (row name) transitioned to another behaviour

 (column name). Numbers in blue represent transitions that occurred more often that expected

 under a model of independence, whereas numbers in red are transitions that occurred less often

 than expected (see methods)s.

Behavior	ab	capture	fly	groom	jump	retreat	run	stop	occl	turn	walk	wd	Total
ab	0	0	2	16	1	0	30	1	2	10	65	2	129
capture	0	0	0	0	0	0	0	0	0	0	0	0	0
fly	1	0	0	9	1	0	6	2	0	0	15	0	34
groom	7	0	11	0	8	0	18	1	5	10	122	0	182
jump	0	0	1	7	0	0	0	0	1	0	9	0	18
retreat	0	0	0	0	0	0	0	0	0	0	0	0	0
run	38	0	5	14	1	0	0	4	2	1	42	3	110
stop	1	0	4	0	2	0	2	0	0	1	10	0	20
occl	2	0	0	6	0	0	5	0	0	3	16	0	32
turn	4	0	1	5	0	0	2	0	3	0	72	1	88
walk	74	0	8	126	5	0	42	13	18	64	0	12	362
wd	1	0	0	0	0	0	4	0	0	0	13	0	18
Total	128	0	32	183	18	0	109	21	31	89	364	18	993

Table S3 Transition frequency in the presence of a juvenile praying mantid. Each row represents the number of times one behaviour (row name) transitioned to another behaviour (column name). Numbers in blue represent transitions that occurred more often that expected whereas numbers in red are transitions that occurred less often than expected.

Mantid

Behavior	ab	capture	fly	groom	jump	retreat	run	stop	occl	turn	walk	wd	Total
ab	0	1	9	2	1	9	10	5	4	6	127	0	174
capture	0	0	0	0	0	0	0	0	0	0	0	0	0
fly	0	0	0	7	0	1	5	60	2	0	50	0	125
groom	0	0	17	0	3	0	18	3	5	13	305	1	365
jump	0	0	0	1	0	0	0	13	0	0	18	1	33
retreat	9	0	2	0	0	0	0	1	1	7	3	0	23
run	11	0	3	18	0	0	0	17	1	12	68	0	130
stop	4	0	45	9	4	5	21	0	0	5	139	0	232
occl	1	0	3	7	1	0	5	1	0	0	73	0	91
turn	4	1	2	6	1	3	15	0	1	0	164	0	197
walk	145	2	43	326	23	5	51	132	73	150	0	49	999
wd	0	0	1	0	0	0	2	0	0	1	47	0	51
Total	174	4	125	376	33	23	127	232	87	194	994	51	2420

Table S4 Transition frequency matrix before a juvenile mantid was added to the chamber. Each row represents the number of times one behaviour (row name) transitioned to another behaviour

133 (column name). Numbers in blue represent transitions that occurred more often that expected

134 whereas numbers in red are transitions that occurred less often than expected.

No mantid

Behavior	ab	capture	fly	groom	jump	retreat	run	stop	occl	turn	walk	wd	Total
ab	0	0	0	0	0	1	1	0	0	1	**31**	0	34
capture	0	0	0	0	0	0	0	0	0	0	0	0	0
fly	0	0	0	1	0	0	0	**19**	1	0	11	0	32
groom	0	0	9	0	1	0	4	**1**	3	**12**	**212**	0	242
jump	0	0	0	3	0	0	1	4	0	0	6	0	14
retreat	1	0	0	0	0	0	0	1	1	0	0	0	3
run	0	0	0	9	1	0	0	1	0	0	15	0	26
stop	0	0	**14**	**1**	1	0	1	0	0	1	54	0	72
occl	0	0	1	2	0	0	0	0	0	1	55	0	59
turn	2	0	0	2	0	0	0	0	4	0	**79**	0	87
walk	31	0	8	**236**	10	1	19	**45**	48	**71**	0	21	490
wd	0	0	0	2	0	0	0	1	0	0	**18**	0	21
Total	34	0	32	256	13	2	26	72	57	86	481	21	1080

137 **Table S5** Transition probability from one behaviour (row name) to the other (column name) in

138 the presence of a zebra jumping spider. Transition probabilites are obtained by dividing each

139 transition frequeny (see table S1) between a pair of behaviours by the total number of times a

140 given behaviour was performed (row sums in table S1).

Spider

Behavior	ab	capture	fly	groom	jump	retreat	run	stop	occl	turn	walk	wd
ab	0.00	0.01	0.04	0.04	0.03	0.00	0.22	0.03	0.02	0.05	0.55	0.03
capture	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
fly	0.03	0.01	0.00	0.10	0.04	0.00	0.22	0.16	0.06	0.01	0.37	0.01
groom	0.07	0.02	0.11	0.00	0.07	0.00	0.05	0.07	0.03	0.05	0.52	0.01
jump	0.05	0.02	0.22	0.09	0.00	0.00	0.11	0.15	0.02	0.00	0.33	0.00
retreat	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
run	0.24	0.02	0.18	0.07	0.05	0.01	0.00	0.10	0.02	0.02	0.28	0.05
stop	0.07	0.02	0.23	0.07	0.12	0.00	0.14	0.00	0.01	0.07	0.28	0.00
occl	0.07	0.00	0.15	0.10	0.12	0.00	0.10	0.10	0.00	0.02	0.34	0.00
turn	0.08	0.01	0.00	0.04	0.03	0.00	0.14	0.02	0.00	0.00	0.67	0.02
walk	0.27	0.02	0.08	0.19	0.06	0.00	0.09	0.13	0.03	0.14	0.00	0.05
wd	0.10	0.00	0.08	0.00	0.00	0.00	0.19	0.00	0.00	0.02	0.60	0.00

143 **Table S6** Transition probability from one behaviour (row name) to the other (column name)

144 before a zebra jumping spider was introduced into the arena. Transition probabilites are obtained

145 by dividing each transition frequeny (see table S1) between a pair of behaviours by the total

146 number of times a given behaviour was performed (row sums in table S1).

No spider

Behavior	ab	capture	fly	groom	jump	retreat	run	stop	occl	turn	walk	wd
ab	0.00	0.00	0.02	0.13	0.01	0.00	0.24	0.01	0.02	0.08	0.51	0.02
capture	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
fly	0.03	0.00	0.00	0.26	0.03	0.00	0.18	0.06	0.00	0.00	0.44	0.00
groom	0.04	0.00	0.06	0.00	0.04	0.00	0.10	0.01	0.03	0.05	0.67	0.00
jump	0.00	0.00	0.06	0.39	0.00	0.00	0.00	0.00	0.06	0.00	0.50	0.00
retreat	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
run	0.36	0.00	0.05	0.13	0.01	0.00	0.00	0.04	0.02	0.01	0.39	0.03
stop	0.05	0.00	0.20	0.00	0.10	0.00	0.10	0.00	0.00	0.05	0.50	0.00
occl	0.06	0.00	0.00	0.19	0.00	0.00	0.16	0.00	0.00	0.09	0.50	0.00
turn	0.05	0.00	0.01	0.06	0.00	0.00	0.02	0.00	0.03	0.00	0.83	0.01
walk	0.21	0.00	0.02	0.36	0.01	0.00	0.12	0.04	0.05	0.18	0.00	0.03
wd	0.06	0.00	0.00	0.00	0.00	0.00	0.22	0.00	0.00	0.00	0.72	0.00

149 **Table S7** Transition probability from one behaviour (row name) to the other (column name) in

150 the presence of a juvenile praying mantid. Transition probabilites are obtained by dividing each

151 transition frequeny (see table S1) between a pair of behaviours by the total number of times a

152 given behaviour was performed (row sums in table S1).

Mantid

Behavior	ab	capture	fly	groom	jump	retreat	run	stop	occl	turn	walk	wd
ab	0.00	0.01	0.05	0.01	0.01	0.05	0.06	0.03	0.02	0.03	0.73	0.00
capture	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
fly	0.00	0.00	0.00	0.06	0.00	0.01	0.04	0.48	0.02	0.00	0.40	0.00
groom	0.00	0.00	0.05	0.00	0.01	0.00	0.05	0.01	0.01	0.04	0.84	0.00
jump	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.41	0.00	0.00	0.56	0.03
retreat	0.39	0.00	0.09	0.00	0.00	0.00	0.00	0.04	0.04	0.30	0.13	0.00
run	0.08	0.00	0.02	0.14	0.00	0.00	0.00	0.13	0.01	0.09	0.52	0.00
stop	0.02	0.00	0.19	0.04	0.02	0.02	0.09	0.00	0.00	0.02	0.60	0.00
occl	0.01	0.00	0.03	0.08	0.01	0.00	0.05	0.01	0.00	0.00	0.80	0.00
turn	0.02	0.01	0.01	0.03	0.01	0.02	0.08	0.00	0.01	0.00	0.83	0.00
walk	0.15	0.00	0.05	0.34	0.02	0.01	0.05	0.14	0.08	0.16	0.00	0.05
wd	0.00	0.00	0.02	0.00	0.00	0.00	0.04	0.00	0.00	0.02	0.92	0.00

Table S8 Transition probability from one behaviour (row name) to the other (column name) before a juvenile praying mantid was introduced into the arena. Transition probabilites are obtained by dividing each transition frequeny (see table S1) between a pair of behaviours by the total number of times a given behaviour was performed (row sums in table S1).

No mantid

Behavior	ab	capture	fly	groom	jump	retreat	run	stop	occl	turn	walk	wd
ab	0.00	0.00	0.00	0.00	0.00	0.03	0.03	0.00	0.00	0.03	0.91	0.00
capture	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
fly	0.00	0.00	0.00	0.03	0.00	0.00	0.00	0.59	0.03	0.00	0.34	0.00
groom	0.00	0.00	0.04	0.00	0.00	0.00	0.02	0.00	0.01	0.05	0.88	0.00
jump	0.00	0.00	0.00	0.21	0.00	0.00	0.07	0.29	0.00	0.00	0.43	0.00
retreat	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.33	0.33	0.00	0.00	0.00
run	0.00	0.00	0.00	0.35	0.04	0.00	0.00	0.04	0.00	0.00	0.58	0.00
stop	0.00	0.00	0.19	0.01	0.01	0.00	0.01	0.00	0.00	0.01	0.75	0.00
occl	0.00	0.00	0.02	0.03	0.00	0.00	0.00	0.00	0.00	0.02	0.93	0.00
turn	0.02	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.05	0.00	0.91	0.00
walk	0.07	0.00	0.02	0.50	0.02	0.00	0.04	0.10	0.10	0.15	0.00	0.04
wd	0.00	0.00	0.00	0.10	0.00	0.00	0.00	0.05	0.00	0.00	0.86	0.00

Table S9 Proportion of time spent in a given behavioural state by each individual fruit fly before and after introducing a treatment (i.e., a disturbance, spider or mantid) to the assay chamber.

Treatment	Individual	Grooming		Walking		Running		Stopping		Retreat	
		Absent	Present	Absent	Present	Absent	Present	Absent	Present	Absent	Present
Disturbance	1	0.820	0.728	0.180	0.265	0.000	0.005	0.000	0.000	0.000	0.000
	2	0.590	0.265	0.338	0.658	0.000	0.009	0.044	0.015	0.000	0.000
	3	0.264	0.219	0.658	0.685	0.048	0.036	0.013	0.052	0.000	0.000
Spider	1	0.528	0.490	0.359	0.394	0.022	0.011	0.000	0.003	0.000	0.000
	2	0.116	0.000	0.844	0.861	0.000	0.086	0.000	0.053	0.000	0.000
	3	0.706	0.181	0.262	0.368	0.010	0.076	0.000	0.347	0.005	0.002
Mantid	1	0.419	0.764	0.474	0.217	0.008	0.005	0.061	0.014	0.000	0.000
	2	0.673	0.075	0.327	0.785	0.000	0.014	0.000	0.093	0.000	0.006
	3	0.684	0.963	0.309	0.037	0.000	0.000	0.000	0.000	0.000	0.000

Table S10 Number of occurrences per minute of each behavioural event before and after the

166 introduction of a treatment (i.e., a disturbance, spider or mantid) to the assay chamber.

		Pause		Turn		Ab		Wd		Fly		Jump	
Treatment	Individual	Absent	Present	Absent	Present	Absent	Present	Absent	Present	Absent	Present	Absent	Present
Disturbance	1	0.000	0.006	0.000	0.006	0.000	0.003	0.000	0.000	0.000	0.002	0.000	0.003
	2	0.003	0.013	0.003	0.013	0.060	0.023	0.003	0.005	0.082	0.027	0.000	0.005
	3	0.000	0.002	0.000	0.002	0.027	0.052	0.003	0.010	0.045	0.090	0.018	0.016
Spider	1	0.009	0.013	0.000	0.005	0.043	0.046	0.006	0.000	0.006	0.003	0.000	0.000
	2	0.020	0.018	0.032	0.045	0.012	0.000	0.012	0.027	0.048	0.252	0.000	0.000
	3	0.015	0.011	0.003	0.011	0.026	0.090	0.000	0.002	0.018	0.041	0.003	0.000
Mantid	1	0.052	0.017	0.016	0.006	0.031	0.002	0.008	0.002	0.070	0.020	0.003	0.000
	2	0.005	0.060	0.005	0.012	0.023	0.111	0.000	0.003	0.000	0.076	0.000	0.006
	3	0.000	0.000	0.003	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

169 **Table S11** Coefficients from a linear model (lm) for control individuals meansured before and after

170 the addition of a spider. While estimate of posterior means are similar to those of the main spider

171 dataset, Due to low sample sizes, CIs are large.

Grooming			
Coefficient	*Posterior mean*	*Lower CI*	*Upper CI*
Intercept	181.88	-23.34	387.10
Predator state	-75.38	-377.82	227.05
Duration	0.45	-0.34	1.24
Locomotion			
Intercept	156.76	37.32	276.21
Predator state	17.27	-158.75	193.29
Duration	0.24	-0.22	0.70

174 **Table S12** Coefficients from a linear model (lm) for control individuals meansured before and

175 after the addition of a mantid. Estimate of posterior means are similar to those of the main mantid

176 dataset, but due to low sample sizes, CIs large.

Grooming			
Coefficient	**Posterior mean**	**Lower CI**	**Upper CI**
Intercept	134.79	-1399.38	1668.96
Predator state	341.56	-2661.10	3344.23
Duration	-0.80	-12.76	11.17
Locomotion			
Intercept	345.69	-919.99	1611.36
Predator state	-323.02	-2800.19	2154.15
Duration	1.64	-8.24	11.51

179 **Table S13** Coefficients from a linear model (lm) for control individuals meansured before and after

180 a disturbance. Despite low sample sizes, it is clear that disturbance had minimal effect on fruit fly

181 behaviours.

Grooming			
Coefficient	**Posterior mean**	**Lower CI**	**Upper CI**
Intercept	200.00	60.11	339.81
Disturbance	0.00	-197.78	197.78
Duration	2.38	-0.53	5.29
Locomotion			
Intercept	132.30	-0.28	264.84
Disturbance	0.00	-187.47	187.47
Duration	-1.05	-3.81	1.70

184 **Links to videos describing novel behaviours**

185 1) Abdominal Lifting http://dx.doi.org/10.6084/m9.figshare.1185638

186 2) Stopping Behaviour http://dx.doi.org/10.6084/m9.figshare.1185639

187 3) Retreat http://dx.doi.org/10.6084/m9.figshare.1185640