1   **Adaptation to larval crowding in *Drosophila ananassae* and *Drosophila nasuta***

2   ***nasuta*: increased larval competitive ability without increased larval feeding rate**

4   Archana Nagarajan[1*], Sharmila Bharathi Natarajan[2*], Mohan Jayaram[3], Ananda Thammanna[4],

5   Sudarshan Chari[5], Joy Bose, Shreyas V. Jois[6] and Amitabh Joshi[†]

6   *Evolutionary Biology Laboratory, Evolutionary & Organismal Biology Unit, Jawaharlal Nehru*

7   *Centre for Advanced Scientific Research, Jakkur P.O., Bangalore 560064, India.*

*: equal contribution; †: Corresponding author
9

1.   Present address: Centre for Organelle Research (CORE), Faculty of Science and

Technology, University of Stavanger, N-4036 Stavanger, Norway.

2.   Present address: Department of Pharmocology, School of Medicine, University of

Pennsylvania, Philadelphia 19104, USA.

3.   Present address: Department of Molecular Reproduction, Development and Genetics,

Indian Institute of Science, Bangalore 560012, India.

4.   Present address: Jubilant Biosys Ltd., 96 Yeshwantpur Industrial Suburb IInd Stage,

Tumkur Road, Bangalore 560 022, India.

5.   Present address: Department of Zoology, Michigan State University, East Lansing, MI

48824, USA.

6.   Present address: Department of Animal Physiology, Zoology Institute, Biocenter, Cologne

50674, Germany.

Email addresses:

archana0681@gmail.com

sharmilanatarajan79@gmail.com

jmohan80@gmail.com

ananda_thammanna@jubilantbiosys.com

charisud@msu.edu

caspase.joy@gmail.com

joisshreyasv@yahoo.co.in

ajoshi@jncasr.ac.in

Key words: life-history evolution; experimental evolution; development time; dry weight;

competition; *K*-selection.

Running title: Alternative routes to competitive ability

Word count: Abstract – 252; Main Text – 8167; Figures – 7; Tables – 6; References - 42

Abstract

The standard view of adaptation to larval crowding in fruitflies, built on results from 25 years of multiple experimental evolution studies on *D. melanogaster*, is that enhanced competitive ability evolves primarily through increased larval feeding and foraging rate, at the cost of efficiency of food conversion to biomass, and increased larval tolerance to nitrogenous wastes. These results, moreover, were at odds from the predictions of classical *K*-selection theory, notably the expectation that selection at high density should result in the increase of efficiency of conversion of food to biomass, and were better interpreted through the lens of α-selection. We show here that populations of *D. ananassae* and *D. n. nasuta* subjected to extreme larval crowding evolve greater competitive ability and pre-adult survivorship at high density primarily through a combination of reduced larval duration, faster attainment of minimum critical size for pupation, greater time efficiency of food conversion to biomass, increased pupation height with a relatively small role of increased urea/ammonia tolerance, if at all. This is a very different suite of traits than that seen to evolve under similar selection in *D. melanogaster* and seems to be closer to the expectations from the canonical theory of *K*-selection. We discuss possible reasons for these differences in results across the three species. Overall, the results reinforce the view that our understanding of the evolution of competitive ability in fruitflies needs to be more nuanced than before, with an appreciation that there may be multiple evolutionary routes through which higher competitive ability can be attained.

Introduction

Adaptation to crowding leading to greater competitive ability, first conceptualized as *r*-
and *K*-selection theory (MacArthur and Wilson 1967), is an important phenomenon in
ecology and evolution that was extensively studied in *Drosophila melanogaster* over the
past three decades, using experimental evolution approaches (reviewed in Joshi 1997;
Mueller 1997; Prasad and Joshi 2003). With lifespan, fecundity, starvation resistance and
dessication resistance, some of the other traits extensively examined via experimental
evolution in *D. melanogaster*, there were conflicting reports of differing patterns of
correlated responses to selection between laboratories (Ackermann et al. 2001; Prasad and
Joshi 2003; Rose et al. 1996), that led, in part, to a growing appreciation that evolution is
often "local" (Rose et al. 2005). In the case of adaptation to crowding, however, the
pattern of correlated responses to selection was largely consistent over two separate
selection experiments, involving *D. melanogaster* populations originating from different
geographical sources (*r*- and *K*-populations: Mueller and Ayala 1981; UU and CU
populations: Joshi and Mueller 1996), and the suite of traits through which selected
populations evolved greater competitive ability was very different from the canonical
version of *r*- and *K*-selection theory that emphasized the centrality of increased food to
biomass conversion efficiency as an adaptation to chronic crowding (Mueller 2009).

Results from the earlier selection studies on adaptation to crowding in *D.
melanogaster* have been reviewed extensively (Joshi 1997; Mueller 1997; Joshi et al.
2001; Prasad and Joshi 2003; Dey et al. 2012) and, consequently, we will just summarize
the major observed correlated responses of pre-adult traits to selection. Relative to the low
density *r*-populations, the crowding adapted *K*-populations exhibited greater (i) pre-adult

survivorship, when assayed at high larval density (Bierbaum et al. 1989), (ii) pre-adult

competitive ability (Mueller 1988), (iii) larval feeding rate (Joshi and Mueller 1988), (iv)

larval foraging path length (Sokolowski et al. 1997), (v) pupation height (Mueller and

Sweet 1986; Joshi and Mueller 1993), and (vi) minimum food requirement for pupation

(Mueller 1990). Pre-adult survivorship of the r- and K-populations did not differ

significantly when assayed at low larval density (Bierbaum et al. 1989). Pre-adult

development time in the K-populations was lower than that of the r-populations at low (30

larvae per vial) and moderately high (160 larvae per vial) density, but greater at very high

(320 larvae per vial) density (Bierbaum et al. 1989). The evolution of higher larval feeding

rate and pupation height in crowding adapted populations was subsequently verified with

a separate set of populations (rK and $r{\times}rK$ populations), by Guo et al (1991).

Relative to the control UU populations, the larval crowding adapted CU

populations exhibited greater (i) pre-adult survivorship, when assayed at high larval

density (Mueller et al. 1993; Shiotsugu et al. 1997), (ii) larval feeding rate (Joshi and

Mueller 1996), (iii) larval foraging path length (Sokolowski et al. 1997), (iv) minimum

food requirement for pupation (Joshi and Mueller 1996), and (v) tolerance to nitrogenous

wastes like urea (Shiotsugu et al. 1997; Borash et al. 1998) and ammonia (Borash et al.

1998). Pre-adult survivorship of the CU and UU populations did not differ significantly

when assayed at low larval density (Mueller et al. 1993; Shiotsugu et al. 1997). CU pre-

adult development time was similar to the UU populations when assayed at low larval

density, but lower than the UU populations when assayed at high larval density (Borash

and Ho 2001). Pupation height of the CU populations was greater than the UU controls

during early generations of CU selection (Mueller et al. 1993), but did not differ

significantly from the controls after about 60 generations of selection (Joshi and Mueller

1996); possible reasons for this are extensively discussed by Joshi et al. (2003). Urea and

ammonia tolerance were not assayed on the *r*- and *K*-populations, whereas competitive

ability was not assayed on the CU and UU populations.

The *K*-populations differed from the *r*-populations in experiencing higher larval as

well as adult density and, moreover, were kept on an overlapping generation regime

whereas the *r*-populations were kept on discrete generations (Mueller and Ayala 1981).

The CU populations, on the other hand, differed from the UU controls only in larval

density (Joshi and Mueller 1996). Given the congruence in the pre-adult traits that evolved

in the CU and *K*-populations, it is likely that these traits represented a response primarily

to levels of larval crowding. Thus, the overall picture that emerged from these two studies

was that long-term exposure to larval crowding in *Drosophila* selected for increased larval

feeding and foraging activity, at the cost of efficiency at converting food to biomass, a

greater tendency to pupate away from the food, and a higher tolerance to toxic levels of

metabolic waste. It makes sense that these traits would contribute to greater pre-adult

survivorship in a crowded larval culture characterized by diminishing food levels, greater

chance of pupal drowning on the increasingly mushy food surface, and a rapid buildup of

levels of nitrogenous waste. This has been the canonical view of adaptations to crowding

in *Drosophila* for the past fifteen years or so (Mueller 1997; Joshi et al. 2001; Prasad and

Joshi 2003; Mueller 2009; Mueller and Cabral 2012).

The notion that faster feeding is a strong correlate of pre-adult competitive ability

in *Drosophila* has a lot of support. Populations selected for faster feeding rate were also

found to be better competitors (Burnet et al. 1977)), and populations selected for rapid

pre-adult development evolved both reduced feeding rate and reduced competitive ability

(Prasad et al. 2001; Shakarad et al. 2005; Rajamani et al. 2006). Similarly, populations

selected for increased parasitoid resistance evolved both reduced feeding rate and reduced

competitive ability (Fellowes et al. 1998, 1999). The notion that there is a cost to faster

feeding (Joshi and Mueller 1996) was supported by the rapid return of feeding rates to

control levels when CU populations were maintained at moderate densities (Joshi et al.

2003). There is also evidence for a trade-off between urea/ammonia tolerance and larval

feeding rate. Populations selected for greater urea and ammonia tolerance, respectively,

also showed reduced larval feeding rate (Borash et al. 2000) and larval foraging path

length (Mueller et al. 2005), and populations selected for greater urea tolerance did not

show higher survivorship than controls at high larval density (Shiotsugu et al. 1997). In

general, larval feeding rate and forgaing path length appear to be positively correlated

(Joshi and Mueller 1988, 1996; Sokolowski et al. 1997; Borash et al. 2000; Prasad et al.

2001; Mueller et al. 2005). Thus, the evolution of competitive ability in *Drosophila* seems

to be the outcome of a balance between mutually antagonistic traits like increased larval

feeding and forgaing behaviour (and perhaps pupation height), greater tolerance to

nitrogenous wastes, and a reduced efficiency of conversion of food to biomass. Indeed, the

CU populations exhibited a temporal polymorphism for two of these traits: offspring of

early eclosing flies in a crowded culture showed higher feeding rates, whereas offspring of

late eclosing flies showed greater urea/ammonia tolerance than controls (Borash et al.

1998).

Given that the above view of adaptation to larval crowding in *Drosophila* was built

around studies on a single species (*D. melanogaster*), we wanted to investigate whether

other species of *Drosophila* would also respond to larval crowding by evolving essentially

the same set of traits. If the genetic architecture of traits relevant to fitness under larval

crowding is reasonably conserved across congeners, then we should see a similar pattern

of correlated responses to selection for adaptation to larval crowding in other *Drosophila*

species. We report results from two selection experiments, on *D. ananassae* and *D. nasuta*

nasuta, involving selection for adaptation to larval crowding. *D. ananassae* is ecologically

and phylogenetically closer to *D. melanogaster* (*Sophophora* Subgenus, Melanogaster

Group, Melanogaster Subgroup), being a cosmopolitan human commensal and belonging

to the Melanogaster Group, Ananassae Subgroup and Ananassae Species Complex,

whereas *D. n. nasuta* belongs to the *Drosophila* Subgenus, Immigrans Group, Nasuta

Subgroup and Frontal Sheen Complex, and is found primarily in orchards and open land.

Our results showed that selected populations of both *D. ananassae* and *D. n. nasuta*

exhibited similar patterns of correlated responses to selection for adaptation to larval

crowding, but that these populations evolved greater competitive ability through traits

very different from those seen earlier in *D. melanogaster* populations subjected to similar

selection.

Materials and methods

Experimental populations

This study used eight laboratory populations each of *D. ananassae* and *D. n. nasuta*. Four

control *D. ananassae* populations (AB$_{1-4}$: **A**nanassae **B**aseline) were derived from a single

population, initiated in May-June 2001 with about 300 wild-caught females from

Bangalore, India, and maintained as a single population on a 21-day discrete generation

cycle for 34 generations (Sharmila Bharathi *et al*. 2003). The four AB populations were

maintained on a 21-day discrete generation cycle at 25º ± 1 C, ~90% relative humidity,

constant light and on cornmeal medium. Larval density was regulated at 60-80 larvae per

vial (9 cm × 2.4 cm) with 6 mL food. Forty vials were set up per replicate population.

Twelve days after egg-collection, eclosed adults (1500-1800) from all 40 vials were

collected into Plexiglas cages ($25 \times 20 \times 15$ cm^3) containing a Petridish (9 cm diameter) of

food that was changed every alternate day, and a moistened ball of cotton. On day 18 after

egg-collection, the flies were given food along with a generous smear of live yeast-acetic

acid paste. On day 21 from the previous egg-collection, fresh food plates were put into the

cages and eggs collected from them after 18 h to initiate the next generation.

181   From the AB populations, four populations (ACU$_{1-4}$: **A**nanassae **C**rowded as

larvae and **U**ncrowded as adults) were derived, one each from each of AB$_{1-4}$, two

generations after the AB populations were established. The ACU populations were

selected for adaptation to larval crowding by subjecting them to a density of 550-600 eggs

per vial with 1.5 mL of food. In initial generations the density was lower; the final

densities were attained by generation 15 of ACU selection. The ACU populations were

otherwise maintained the same as the AB controls, except that only 20 vials of eggs were

collected each generation (to keep the number of breeding adults similar to controls: 1500-

1800), and the collection of eclosed adults into the cages continued until day 18 from

previous egg-collection, as eclosion in a crowded culture is staggered over several days.

191   Four control *D. n. nasuta* populations (NB$_{1-4}$: **N**asuta **B**aseline) were derived, after

24 generations as a single population on a 21 day discrete generation cycle, from a

laboratory population established using about 70 females collected from orchards and

domestic garbage dumps in different parts of Bangalore, India, during October-November

2001 (Sharmila Bharathi *et al*. 2003). The four NB populations were maintained on a 21-

day discrete generation cycle at $25^\circ \pm 1$ C, ~90% relative humidity, constant light and on

cornmeal medium. The larval density was regulated at 60-80 larvae per vial (9 cm $\times$ 2.4

cm) with 6 mL food. Fifty two vials were set up per replicate population to keep the

number of breeding adults at about 1500-1800. Twelve days after egg-collection, eclosed

adults from all 40 vials per replicate population were collected into Plexiglas cages (25 $\times$

20×15 cm^3) containing a Petridish of food that was changed every alternate day, and a

moistened ball of cotton. On day 18 after egg-collection, the flies were given food along

with a generous smear of live yeast-acetic acid paste. On day 21 from the previous egg-

collection, fresh food plates were put into the cages and eggs collected from them after 18

h to initiate the next generation.

From the NB populations, four populations (NCU$_{1-4}$: **N**asuta **C**rowded as larvae

and **U**ncrowded as adults) were derived, one each from each of NB$_{1-4}$, two generations

after the NB populations were established. The NCU populations were selected for

adaptation to larval crowding by subjecting them to a density of 350-400 eggs per vial

with 2 mL of food. In initial generations the density was lower; the final densities were

attained by generation 15 of NCU selection. This difference from ACU in larval density is

to compensate for the larger size at each stage of the *D. nasuta* larvae. The NCU

populations were otherwise maintained in the same way as the NB controls, with similar

adult population size, except that the collection of eclosed adults into the cages continued

until day 18 from previous egg-collection, and only 40 vials with eggs were set up.

Since each ACU or NCU population was derived from one control population,

selected and control populations of each species bearing identical numerical subscripts are

more related to each other than to other populations with which they share the selection

regime. Therefore, control and selected populations with identical subscripts were treated

as blocks, representing ancestry, in the statistical analyses.

Collection of flies for assays

All control and selected populations were maintained under common (control-type)

rearing conditions for one complete generation prior to assays, to eliminate non-genetic

parental effects. The progeny of these flies, hereafter 'standardized flies', were then used

for the various assays. To obtain progeny for assays, standardized flies in cages were

provided yeast-acetic acid paste on food for three days before egg collection. A fresh

Petridish with food was then placed in the cages and the flies were allowed to lay eggs for

~14 h, after which eggs were removed from the food with a moistened paintbrush and

placed into vials for setting up the various assays. All assays were conducted at 25 ± 1^{o}C,

under constant light.

Pre-adult survivorship

After 42 generations of ACU selection, eggs laid by standardized flies were placed into

vials at a density of either 70 or 600 per vial containing 1.5 mL of food. Eight such vials

were set up for each replicate AB and ACU population at each density in single-species

culture. Eight such vials at each density were also set up in two-species cultures, in

competition with a common competitor, a white eyed mutant population of *D.*

melanogaster, maintained for about 90 generations in the laboratory on a three week

discrete generation cycle. The white eyed population was derived from spontaneously

occurring mutant individuals in the JB populations (Sheeba et al. 1998) in our laboratory.

For the two-species cultures, eggs laid by standardized ACU, AB or white eyed flies were

collected and placed into vials at a density of either 70 (35 ACU or AB eggs and 35 eggs

from the white eyed mutant population) or 600 (300 ACU or AB eggs and 300 eggs from

the white eyed mutant population) per vial containing 1.5 mL of food. Eight such vials

were set up for each replicate AB and ACU population at each density. The number of

flies eclosing in each vial was recorded and used to calculate pre-adult survivorship.

At generation 76 of NCU selection, an assay similar to that described above was

set up using the NB and NCU populations and the white eyed *D. melanogaster*. The only

difference from the *D. ananassae* assay was that the low and high density treatments

comprised of 70 or 350 eggs in vials with 2 mL of food.

Duration of pre-adult life-stages and pre-adult development time

After 53 generations of ACU selection, pre-adult development time was assayed at a high

density of 600 eggs per vial with 1.5 mL of food. Eggs from standardized AB and ACU

flies were dispensed into each vial using a moistened paintbrush. Eight such vials were set

up per replicate population. After the pupae darkened, vials were checked every 6 h and

the number of eclosing flies recorded.

After 71 generations of ACU selection, pre-adult development time was assayed at

a low density of 30 eggs per vial with 6 mL of food (An earlier study at generation 41 of

ACU selection had shown that pre-adult development time did not differ significantly

between low densities of 30 or 70 eggs per vial with 6 mL of food: data not shown). Eggs

from standardized AB and ACU flies were dispensed into each vial using a moistened

paintbrush. Ten vials were set up per replicate population. After the pupae darkened, the

vials were monitored for eclosion at 2 h intervals, and the number of eclosing flies

recorded. As part of the same assay, egg hatching time and the duration of each larval

instar and the pupal stage were also determined. For assaying egg hatching time, 30 eggs

from the standardized flies were arranged on a small agar cube in a food vial in six rows

of five eggs each. Ten such vials were set up per population. Fifteen hours after egg

laying, the vials were checked for any hatched eggs once every hour, till no eggs hatched

for three consecutive hours. For assaying instar and pupal duration, eggs of approximately

identical age were harvested over a three hour period from the standardized flies and

dispensed into vials with 6 mL of food at a density of 30 eggs per vial. Sixty such vials

were set up per population. Forty three hours after the midpoint of the three hour egg

laying period, four vials per population were removed from the incubators and immersed

in hot water. The dead larvae were removed and kept in 70% ethanol for subsequent

examination. Every two hours, this process was repeated. The larval instars were

differentiated based on the number of 'teeth' in the larval mouth hooks. From these data,

the number of larvae of each instar present in each two-hourly sample was determined,

and the median time of each molt was obtained by interpolation. For pupal duration and

pre-adult development time, ten vials per population were set up with 30 eggs in 6 mL of

food. After the first pupa was seen, the vials were screened every two hours and any new

pupae that had formed were marked on the vial with different coloured marker pens.

Thereafter, the vials were monitored for eclosion and the number of eclosing males and

females in each vial was determined every two hours. These observations yielded data on

egg to pupa and egg to adult development time, from which the pupal duration could be

calculated.

After 84 generations of NCU selection, pre-adult development time was assayed at

a high density of 350 eggs per vial with 2 mL of food. Eggs from standardized NB and

NCU flies were dispensed into each vial using a moistened paintbrush. Eight such vials

were set up per population. After the pupae darkened, the vials were checked every 6 h

and the number of eclosing flies recorded.

After 62 generations of NCU selection, egg hatching time, larval instar duration,

pupal duration and pre-adult development time of the NB and NCU populations were

assayed exactly as described above for the ACU and AB populations.

Larval feeding rate

After 71 generations of ACU selection, the feeding rates of AB and ACU larvae were

measured at physiologically equalized ages, based on the difference in AB and ACU

development time. This was done by collecting eggs from the standardized ACU flies 5 h

later than the AB flies. Thus, at the time of assay, ACU larvae were 58 h old, whereas AB

larvae were 63 h old and, thus, approximately in the same relative stage of their larval

development. Following Joshi and Mueller (1996), about a hundred eggs laid over a four

hour period were collected from standardized flies and placed into two Petridishes with

non-nutritive agar each for AB and ACU populations. Twenty-four hours later, twenty-

five newly hatched larvae were transferred from these agar Petridishes to a Petridish

containing a thin layer of non-nutritive agar overlaid with 1.5 mL of 37.5% yeast

suspension. Four such Petridishes were set up per population. The larvae were then

allowed to feed for 58 (ACU) or 63 (AB) h, by which time they were in the early third

instar. At this point, 20 larvae from each population were assayed for feeding rate,

following the procedure of Joshi and Mueller (1996), by placing them individually in a

small Petridish (5 cm diameter) containing a thin layer of agar overlaid with a thin layer of

10% yeast suspension. After allowing for a 15 sec acclimation period, feeding rate was

measured under a stereozoom microscope as the number of cephalopharyngeal sclerite

retractions in a 1 min period. Selected and control populations, matched by the subscripted

indices, were assayed together, with one larva from the selected population and one from

the control population being assayed alternately. The same procedure was followed for

assaying larval feeding rate of 20 larvae from each NCU and NB population, after 77

generations of NCU selection.

Larval foraging path length

Twenty early third instar larvae for each population were assayed for larval foraging path

length. The collection of larvae for the assay was exactly as described above for larval

feeding rate assays. For assaying foraging path length, individual larvae were placed in a

small Petridish (5 cm diameter) containing a thin layer of agar overlaid with a thin layer of

10% yeast suspension. After allowing for a 15 sec acclimation period, the larvae were

allowed to move around on the Petridish for 1 min. The path traversed by the larvae on the

yeast surface was traced onto a transparency sheet and later measured with a thread and

ruler. Selected and control populations, matched by subscripted indices, were assayed

together at generations 52 and 81 of ACU and NCU selection, respectively.

Pupation height

Eggs from standardized flies were collected and placed into vials with 6 mL of food at a

density of 50 (ACU/AB) or 70 (NCU/NB) eggs per vial. Ten and eight such vials were set

up per population for *D. ananassae* and *D. n. nasuta*, respectively. Once all pupae had

formed, pupation height of each pupa was measured, following Joshi and Mueller (1993),

as the distance between the surface of the food medium to the point between the anterior

spiracles of the pupa. Any pupa on or touching the surface of the food was given a

pupation height of zero. Assays were conducted at generations 39 and 34 of ACU and

NCU selection, respectively.

Larval urea and ammonia tolerance

After 49 generations of ACU selection, eggs laid over a four hour period by standardized

flies were collected and exactly 30 eggs per 6 mL food were placed into vials at three

concentrations each of urea (0, 14 and 18 g/L) or ammonia (0, 15 and 30 g/L NH_4Cl) in

the food. These are values that allowed the detection of differences in urea/ammonia

tolerance between selected and control populations in previous studies on *D. melanogaster*

(Shiotsugu et al. 1997; Borash et al. 1998). Ten vials were set up per population at each

concentration of either urea or ammonia, and the number of eclosing flies in each vial was

recorded to calculate pre-adult survivorship. Both urea and ammonia were used as there is

some confusion about which compound actually increases in concentration in crowded

larval cultures, although it is likely that it is ammonia in the case of *D. melanogaster*

(Botella et al. 1985; Borash et al. 1998). The NCU and NB populations were assayed in an

identical manner after 76 generations of NCU selection, except that the egg-laying period

was 12 hours, and the concentrations used were 0, 9 and 11 g/L for urea and 0, 15 and 20

g/L for ammonia, as earlier studies had shown that *D. n. nasuta* larvae were more sensitive

to urea and ammonia than their *D. ananassae* counterparts, and *D. n. nasuta* females are

not as fecund as *D. ananassae* females (data not shown).

Minimum feeding time and dry weight after minimum feeding

These assays were conducted only on the ACU and AB populations. After 45 generations

of ACU selection, critical minimum feeding time for pupation was assayed by setting up

freshly hatched larvae, from eggs laid by standardized flies, onto Petridishes with agar

overlaid with 1.5 mL of 37.5% yeast suspension, as described above for the feeding rate

and foraging path length assays. Twenty five larvae were placed into each Petridish, and

sixty such Petridishes were set up per population. At 46, 49, 52 and 55 h after egg hatch, a

total of 150 larvae per population per time point were removed from the yeast, gently

washed in water to remove any yeast sticking to their bodies, and placed in ten vials

containing 5 mL of non-nutritive agar, at a density of 15 larvae per vial. These vials were

subsequently monitored for pupation and eclosion, and the pre-adult survivorship after

feeding for different periods of time noted. Using the information from this assay, a

second assay was carried out after 68 generations of ACU selection to measure the dry

weights at eclosion of ACU and AB flies that had fed as larvae for different durations of

time, roughly corresponding to pre-adult survivorship of 13, 25, 50 and 60%, respectively

(feeding for 54, 56, 59 and 62 h for AB, and 50, 55, 58 and 60 h for ACU populations,

respectively). Freshly hatched larvae were collected, shifted to yeast and then removed

and placed into agar vials after feeding for different durations corresponding to pre-adult

survivorship of 13, 25, 50 and 60% for ACU and AB populations, exactly as described

above. Once eclosions began in the agar vials, flies were collected every 4 h and frozen

for subsequent weighing. Frozen flies were sorted into batches of five males or five

females each and dried at 70^{o}C for 36 h before weighing in batches.

Statistical analyses

Since assays on ACU/AB and NCU/NB populations were conducted at different times,

data from the two species were analyzed separately. All traits were subjected to

completely randomized block analyses of variance (ANOVA). All ANOVAs treated

replicate population 1..4 (representing ancestry) as random blocks crossed with the fixed

factor selection regime. Additional fixed factors, crossed with both selection regime and

block, were included when relevant. These factors were larval density and type of culture

(single- or two-species) for pre-adult survivorship, pre-adult life-stage for life-stage

duration, urea or ammonia concentration in the food for urea and ammonia tolerance, and

larval feeding duration or survivorship for the assays on minimum critical feeding time

and of dry weight at eclosion after feeding for different time durations corresponding to

four different pre-adult survivorship levels. Development time data from low and high

density were analyzed separately, rather than incorporating larval density as a factor,

because the low and high density assays were conducted at different times for each

species. Data on pre-adult survivorship were arcsine-squareroot transformed before

ANOVA. For all traits, ANOVAs were done on replicate population mean values and,

therefore, only fixed factor effects and interactions could be tested for significance. All

analyses were implemented using Statistica for Windows rel.5.0 B, (Stat Soft 1995).

Multiple comparisons were done using Tukey's HSD test.

Results

Pre-adult survivorship

Pre-adult survivorship at high larval density is the primary trait expected to be under direct

selection in populations exposed to high larval crowding each generation. With regard to

survivorship at low versus high larval density, the broad pattern of results was very similar

in the two species (Figure 1). In both species, pre-adult survivorship was, on an average,

significantly higher at low rather than high larval density, and in selected compared to

control populations (Figure 1, Table 1). The trend was for survivorship to be higher, on

average, in single-species cultures than in two-species cultures, but the difference was

significant only in the case of *D. ananassae* (Table 1). Both species showed a significant

selection regime × larval density interaction (Table 1), with the survivorship of selected

and control populations not differing significantly at low density, and with selected

populations showing significantly higher survivorship than controls at high density

(Figure 1). Both species also showed a significant culture type × larval density interaction

(Table 1). In *D. ananassae*, averaged over selection regimes, survivorship at low density

in both types of cultures (single- or two-species) was similar, whereas survivorship at high

density was significantly greater in single-species than two-species cultures (Figure 1). In

D. n. nasuta, the opposite pattern was seen. Averaged over selection regime, survivorship

at low density in single-species cultures was higher than in two-species cultures (Figure

1), but not significantly so. However, at high density, survivorship in two-species cultures

was higher than in single-species cultures but, again, not significantly so. In fact, the only

clear significant difference in the multiple comparisons was that between single-species

cultures at low and high densities (Figure 1.). In the case of *D. ananassae*, there was also a

significant culture type × selection regime interaction (Table 1), but the only clearly

significant differences in multiple comparisons were between survivorship in two-species

cultures at high density on the one hand, and in the other three culture type × selection

regime combinations on the other (Figure 1).

Pre-adult development time

In general, pre-adult development time in *D. n. nasuta* was higher than that in *D.*

ananassae, as also noted earlier (Sharmila Bharathi et al. 2004). Pre-adult development

time was, in general, greater in males than in females, and greater at high rather than low

larval density, as is usually the case in *Drosophila* (Figures 2,3, Table 2).

Interestingly, in both species at both larval densities, there was a significant main

effect of selection regime, with selected populations consistently showing substantially

lower development time than control populations at both densities (Figures 2,3, Table 2).

On an average, AB development time was higher than that of ACU populations by about

14 and 29 h at low and high larval density, respectively (Figures 2A, 3A), whereas that of

NB populations was greater than that of NCU populations by about 17 and 29 h at low and

high larval density, respectively (Figures 2B,3B). In three of the four assays, there was no

significant selection regime × sex interaction (Table 2). In *D. n. nasuta* at high larval

density, control males and females had similar development times whereas NCU males

had significantly higher development time than NCU females, albeit by just about 1 h

(Figure 3B).

Pre-adult life-stage duration

The overall pattern of pre-adult life-stage durations in both species was similar to what is

seen in *Drosophila* spp. in general. Consistent with the pre-adult development time results,

both species showed a main effect of selection regime (Table 3), with life-stage durations

being greater in control rather than selected populations, on an average (Figure 4). In *D.*

ananassae, a 20 h egg stage was followed by two larval instars of about 24 h each, a third

larval instar of about 46 h and a pupal stage lasting about 87 h (Figure 4A); there was no

significant selection regime × life-stage duration interaction. In *D. n. nasuta*, all pre-adult

life-stages differed significantly from one another in duration, with a 23 h egg stage,

followed by two larval instars of about 27 and 29 h, respectively, a longer third larval

instar of about 63 h and a pupal stage lasting about 87 h (Figure 4B). The longer larval

instar durations in *D. n. nasuta* are consistent with their larger adult size compared to *D.*

ananassae (Sharmila Bharathi et al. 2004). *D. n. nasuta* also showed a significant

selection regime × life-stage duration interaction (Table 3), with a large (~12 h) reduction

in third instar duration in the NCU populations (Figure 4B). We note that, given the time

interval between subsequent observations (1 h for egg-hatch; 2 h for larval instars), our

screening might be too coarse to pick up small but consistent differences on the order of

30 min – 1 h. However, the differences between selected and control populations in the

durations of different pre-adult life-stages, though often not significant, did add up

roughly to the overall difference seen in pre-adult development time between selected and

control populations in both species.

Larval traits

Surprisingly, larval feeding rate did not differ significantly between selected and control

populations in either species (Table 4). Mean (± s.e.) larval feeding rates in sclerite

retractions per min were, in fact, extremely similar between selected and control

populations of both species (AB: 128.33 ± 3.13; ACU: 131.55 ± 3.05; NB: 112.76 ± 5.29;

NCU: 113.03 ± 4.34). In both species, selected populations showed significantly greater

mean (± s.e.) pupation height in cm than control populations (AB: 1.02 ± 0.28; ACU: 1.60

± 0.20; NB: 0.44 ± 0.12; NCU: 1.54 ± 0.11) (Table 4). In case of larval foraging path

length, the results varied between species, with *D. ananassae* showing a significant main

effect of selection regime while *D. n. nasuta* did not (Table 4). AB populations had mean

(± s.e.) foraging path length in cm of 5.52 ± 0.61, compared to 7.38 ± 0.60 in the ACU

populations. NB and NCU populations had mean (± s.e.) foraging path length in cm of

4.24 ± 0.53 and 4.08 ± 0.43, respectively.

Larval urea and ammonia tolerance

There was no clear pattern to the results of assays on pre-adult survivorship in the

presence of metabolic wastes like urea (Figure 5) and ammonia (Figure 6) in the food

medium, except for strong evidence for the toxic effects of these compounds, reflected in

significant main effects of concentration in all four ANOVAs (Table 5).

ACU populations had lower survivorship than AB controls across all levels of

urea, including 0 g/L, and thus showed no evidence of greater or lesser urea tolerance than

the AB controls (Figure 5A), with a significant main effect of selection regime but no

significant selection regime × concentration interaction (Table 5). There was suggestive

evidence for greater ammonia tolerance in the ACU compared to the AB populations

(Figure 6A). At 0 g/L of ammonia in the food, ACU populations had significantly lower

survivorship than AB controls, whereas at 30 g/L, the ACU survivorship was higher,

though not significantly so, than the AB survivorship (Figure 6A). The selection regime $\times$

concentration interaction was also significant for this assay (Table 5).

In the case of *D. n. nasuta*, there was no significant effect of selection regime or of

the selection regime $\times$ concentration interaction in the urea tolerance assay (Table 5).

However, there was a non-significant trend of NCU survivorship being lower than NB at 0

g/L and higher than NB at 11 g/L (Figure 5B). In case of ammonia, NCU populations had

consistently lower survivorship than NB controls at all concentrations (Figure 6B) and

there was no significant effect of selection regime or of the selection regime $\times$

concentration interaction, yielding no suggestion of increased ammonia tolerance in the

NCU populations.

Minimum critical feeding time

In the minimum critical feeding time assay, only significant main effects of selection

regime and feeding duration were seen (Table 6), with pre-adult survivorship increasing

from feeding durations from 46-55 h, and with ACU survivorship being higher than that of

the AB populations at every feeding duration (Figure 7A). At 55 h of feeding as larvae,

ACU survivorship was over 50% while that of the AB populations was less than 30%

(Figure 7A). Overall, ACU populations seemed to attain similar levels of survivorship

approximately 6 h before the AB controls (Figure 7A), which is commensurate with the

54.8 h difference between their first and second instar durations combined. These results

suggest that larvae in the ACU populations attain their minimum critical size for pupation

approximately 5-6 h before their AB controls.

In the ANOVA on data from the assay of dry weight at eclosion of individuals that

had fed for different durations of time as larvae, corresponding to four different levels of

pre-adult survivorship, only the main effects of sex and survivorship were significant

(Table 6). On an average, females were significantly heavier than males, and dry weight at

eclosion tended to increase with survivorship (Figure 7B). At no survivorship level was

there a significant difference between the dry weights of AB and ACU flies (Figure 7B),

suggesting that the minimum critical size for pupation in the selected and control

populations is probably not different, even though the ACU populations attain it faster

than the AB controls.

Discussion

It is clear that both the ACU and NCU populations did evolve adaptations to larval

crowding over the course of selection: in both species, pre-adult survivorship at high

density and competitive ability was greater in selected than in control populations (Figure

1, Table 1). The overall pattern of correlated responses to selection in the two species was

also similar, and, more importantly, different from that reported earlier in case of *D.*

melanogaster in that it did not involve the evolution of higher feeding rates (Table 4).

In clear contrast to what was seen earlier in *D. melanoagster*, both ACU and NCU

populations evolved faster pre-adult development time than their controls (Table 2), with

the difference being apparent at both low (Figure 2) and high (Figure 3) assay density. The

pre-adult development time difference between selected and control populations is entirely

due to reduction in the duration of larval instars (Figure 4). In absolute terms, the

reductions in pre-adult development time in the selected populations (~14 h in ACU, ~17

h in NCU) are quite large. For comparison, populations of *D. melanogaster* subjected to

strong directional selection for reduced pre-adult development time showed a reduction of

~16 h in the larval duration, and ~10 h in pupal duration, after 50 generations of selection

(Prasad et al. 2001). In the earlier studies on *D. melanogaster*, CU populations showed no

difference from controls in development time at low density, whereas they were faster

developing than controls when assayed at high density (Borash and Ho 2001). While the

K-populations did show faster development than *r*-populations when assayed at low

density (Bierbaum et al. 1989), this cannot be unequivocally ascribed to density-dependent

selection. The *K*-populations were on an overlapping generation maintenance regime and,

relative to a discrete generation regime like that of the *r*-populations, this itself would

impose direct selection for faster development. If we put the results on life-stage duration

(Figure 4) and development time (Figures 2,3) together with those from the minimum

feeding time assay (Figure 7), the ACU and NCU populations have evolved faster

development, especially in the first two larval instars, enabling them to reach the critical

minimum size for pupation about 6 h earlier than controls. This is particularly impressive

considering that *D. melanogaster* populations selected directly for rapid development

reached the critical size only about 2 h before controls after 50 generations of selection

(Prasad et al. 2001). At the same time, the weights of ACU and NCU flies are not different

from controls after feeding for different durations close to the minimum feeding time

(Figure 7, Table 6). Thus, in terms of time, ACU and NCU larvae are clearly more

efficient at converting food to biomass than controls during the first two (and an early part

of the third) larval instars, becoming equally heavy adults as controls after feeding for

about 6 h less. Of course, whether they are more efficient also in terms of food consumed

cannot be determined from these experiments. However, these results do suggest that

unlike the *K*- and CU populations, (Mueller 1990; Joshi and Mueller 1996), the ACU and

NCU populations have evolved to become more rather than less efficient than controls.

The faster development of the ACU and NCU populations, relative to controls, is also

interesting in the context of observations that faster development correlates with greater

competitive ability across species of *Drosophila* (Krijger et al. 2001), even though direct

selection for rapid development leads to the evolution of reduced competitive ability

(Shakarad et al. 2005) through reduced larval feeding rate, foraging path length and

pupation height (Prasad et al. 2001) as well as reduced larval urea tolerance (Joshi et al.

2001).

Unlike in *D. melanogaster*, larval feeding rates did not evolve to become greater in

the ACU and NCU populations (Table 4), although these populations clearly evolved

greater competitive ability than controls (Figure 1). In fact, we measured feeding rates at

various points during ACU and NCU selection, and also at various larval stages, and there

was never any difference between selected and control populations in either species (data

not shown). Interestingly, larval foraging path length was greater in ACU populations than

in AB populations, whereas NCU and NB populations had very similar foraging path

lengths (Table 4). Thus, at least in *D. ananassae*, the consistent positive relationship

between larval feeding rate and foraging path length seen in *D. melanogaster* (Joshi and

Mueller 1988, 1996; Sokolowski et al. 1997; Borash et al. 2000; Prasad et al. 2001;

Mueller et al. 2005) seems to be uncoupled. Pupation height evolved to become greater in

both ACU and NCU populations, relative to controls, mirroring results from the *K*-

populations (Mueller and Sweet 1986), but not the CU populations (Joshi and Mueller

1996; but see also Mueller et al. 1993 and Joshi et al. 2003). One possible reason for the

evolution of increased pupation height in the *K*-populations but not the CU populations

has been speculated to be the greater hardness and dryness of the cornmeal based food (*r*-

and *K*-populations) compared to banana based food (UU and CU populations). It was

suggested that banana food being much softer and more fluid compared to cornmeal food,

perhaps the UU populations were also under selection for increased pupation height due to

greater risk of pupal drowning on the food surface even at their relatively low rearing

density (Joshi et al. 2003). In this context, we note that our ACU/AB and NCU/NB

populations are maintained on a relatively hard and dry cornmeal food and this might be

the reason for the evolution of greater pupation height in selected populations, relative to

controls.

Compared to other traits, the results on urea and ammonia tolerance in the selected

and control populations were not very clear. While pre-adult survivorship clearly

decreased with increasing amounts of urea or ammonia in the food (Figures 5,6, Table 5),

there was no clear evidence for greater urea or ammonia tolerance in the selected

populations, except for greater ammonia tolerance in the ACU populations, relative to the

AB controls (Figure 6A). There was a slight suggestion of a trend towards increased urea

tolerance in the NCU populations, relative to NB controls in that NCU survivorship at 0

g/L was below that of NB populations and became slightly higher than NB populations at

11 g/L (Figure 5B). However, there was no significant selection regime × urea

concentration interaction (Table 5), indicating that this is at best a suggestive result. It

might be that crowded *D. ananassae* and *D. n. nasuta* cultures have experienced

differential levels of urea versus ammonia build-up to which they have adapted.

Moreover, we have seen, in other studies with *D. melanogaster*, that the detection of

between-population differences in urea/ammonia tolerance is often affected by an

interaction between the specific concentrations used and the larval density (Avani Mital,

Gitanjali P. Vaidya and Amitabh Joshi, *unpubl. data*). It is, therefore, possible that we

were unable to detect significant differences between selected and control populations due

to the specific concentrations and larval densities used in our assays of urea and ammonia

tolerance. It is also known that larval exposure to urea markedly reduces subsequent

fecundity in *D. melanogaster*, and that populations selected for increased larval urea

tolerance undergo smaller fecundity declines than controls when reared as larvae on food

with urea (Shiotsugu et al. 1997). Thus, it is also possible that the ACU and NCU

populations might have evolved greater tolerance to the detrimental effects of urea on

fecundity; our experiments did not explore this possibility.

Overall, the crowded *D. ananassae* and *D. n. nasuta* populations seem to have

evolved greater competitive ability and pre-adult survivorship at high density primarily

through a combination of reduced duration of the larval stage, faster attainment of

minimum critical size for pupation, greater efficiency of food conversion to biomass,

increased pupation height and, perhaps, greater urea/ammonia tolerance. This is in

contrast to *D. melanogaster*, in which crowding adapted populations evolve greater

competitive ability and pre-adult survivorship at high density primarily through a

combination of increased larval feeding rate and foraging path length, at the cost of

reduced efficiency of food conversion to biomass, and greater urea/ammonia tolerance

(Mueller 1997; Prasad and Joshi 2003; Mueller 2009). Thus, the ACU and NCU

populations appear to have responded to crowding in a manner closer to the canonical

notion of *K*-selection, in contrast to the primarily α-selection responses shown by the *K*-

and CU populations of *D. melanogaster* (see also Dey et al. 2012). Thus, the results

underscore the fact that there are, in principle, multiple routes to the evolution of greater

competitive ability (Joshi et al. 2001; Dey et al. 2012), something that has also been

experimentally demonstrated in the context of inter-specific competition in *Drosophila*

(Joshi and Thompson 1995).

At this point, we can only speculate about the reason(s) for why the evolution of

increased competitive ability in the ACU and NCU populations occurred in a manner so

different from that seen earlier in *D. melanogaster*. We believe there are three possible

reasons for the observed discrepancy in correlated responses to selection for adaptation to

larval crowding across the three species, and the three proposed explanations are not

mutually exclusive. First, it is possible that the pattern of genetic variances and

covariances among traits relevant to survival in a crowded culture is different from that in

D. melanogaster in the two species studied here (*D. ananassae* and *D. n. nasuta*). For

example, maybe the AB and NB populations do not harbour additive genetic variance for

larval feeding rate. Such differences of genetic architecture could simply reflect between-

species differences, being the result of historical selection pressures and contingencies.

However, if that were the case, we would have expected the results from *D. ananassae* to

be closer to *D. melanogaster*, as compared to *D. n. nasuta*, given the phylogenetic and

ecological relationships of these species (see Introduction, last paragraph). A second, more

likely, possibility is that these differences of genetic architecture between *D. ananassae*

and *D. n. nasuta* on the one hand, and *D. melanogaster* on the other, are because the ACU

and NCU populations represent selection on relatively recently wild-caught populations

(see Materials and methods) whereas the *K-* and CU populations involved selection on

populations that had already been in the laboratory for a large number of generations

(Mueller and Ayala 1981; Joshi and Mueller 1996). There might be significant differences

in the genetic architecture of traits related to fitness under larval crowding between wild

and laboratory populations. Finally, it is possible that the discrepancies between the results

of this study and those from earlier work on *D. melanogaster* are due to differences in the

specific details of the respective maintenance regimes of selected populations in the

various studies. Such small differences in selection regimes have been implicated in the

different trajectories of pupation height in the *K-* versus the CU populations (Joshi et al.

2003). The *K*-populations were maintained on a serial transfer system in half-pint milk

bottles with about 350 mL food per bottle (Mueller and Ayala 1981), while the CU

populations were maintained in 6 dram vials (2.2 cm diameter) at a density of about 1000-

1500 eggs per vial in about 5 mL of food. Unlike in our ACU and NCU populations,

neither food amount nor number of eggs per vial were exactly controlled in CU

maintenance. In the CU populations, eggs were laid by adult females onto thin films of

food on four watch glasses per cage. These films of food with eggs on them were then

sliced up and divided roughly equally among 40 vials that each contained about 3 mL of

food (A. Joshi, *pers. obs.*). Thus, our ACU and NCU populations had a smaller absolute

amount of food per rearing container (bottle/vial) than the *K-* and CU populations, and

also differed from the latter in specific densities of eggs per mL of food. It is, therefore,

possible that the time course of food depletion and nitrogenous waste build-up in the ACU

and NCU cultures is somewhat different from that in the *K*- and CU populations. It has

been shown theoretically that optimal feeding rates are likely to decline as the

concentration of nitrogenous waste in the food increases (Mueller et al. 1995). Thus, at

least in principle, it is possible that the optimal feeding rates in the ACU and NCU

populations are actually less than they were for the *K*- and CU populations, and that is

why increased feeding rates did not evolve in our experiments. Subsequent studies will

attempt to discriminate between these various explanations. However, for the time being,

we believe that these results underscore the need for a more nuanced understanding of

adaptations to larval crowding in *Drosophila*, with a greater appreciation for the fact that

increased competitive ability can be attained through the evolution of fairly different

suites of traits. We also need to be cognizant of the fact that seemingly small differences

of maintenance in otherwise similar selection regimes might mediate the evolution of very

different trajectories in phenotypic space.

Acknowledgements

We thank Larry Mueller for much helpful discussion, and N. Rajanna and M. Manjesh for

help in the laboratory. A. Nagarajan thanks the Council of Scientific and Industrial

Research, Government of India, for financial assistance in the form of Junior and Senior

Research Fellowships. S. B. Natarajan was supported by a doctoral fellowship from the

Jawaharlal Nehru Centre for Advanced Scientific Research. Shreyas Jois' stay in the

laboratory was supported by the joint Summer Research Fellowship programme of the

three Indian science academies. This work was supported by funds from the Department

of Science and Technology, Government of India, to A. Joshi. Preparation of the

manuscript was supported in part by a J. C. Bose National Fellowship from the

Department of Science and Technology, Government of India, to A. Joshi. The authors

declare no competing interests.

References

References

Ackermann M., Bijlsma R., James A. C., Partridge L., Zwaan B. J. and Stearns S. C. 2001

Effects of assay conditions in life-history experiments with *Drosophila melanogaster*. *J.*

Evol. Biol. **14,** 199–209.

Bierbaum T. J., Mueller L. D. and Ayala F. J. 1989 Density-dependent evolution of life-

history traits in *Drosophila melanogaster*. *Evolution* **43,** 382–392.

Borash D. J., Gibbs A. G., Joshi A. and Mueller L. D. 1998 A genetic polymorphism

maintained by natural selection in a temporally varying environment. *Am. Nat.* **151,** 148–

156.

Borash D. J., Teótonio H., Rose M. R. and Mueller L. D. 2000 Density-dependent natural

selection in *Drosophila*: correlations between feeding rate, development time and

viability. *J. Evol. Biol.* **13,** 181–187.

Borash D. J. and Ho G. T. 2001 Patterns of selection: stress resistance and energy storage

in density-dependent populations of *Drosophila melanogaster*. *J. Insect Physiol.* **47,**

1349–1356.

Botella L. M., Gonzalez A. C., Moya A. and Mensua J. L. 1985 Larval stop, delayed

development and survival in over-crowded cultures of *Drosophila melanogaster*: effect of

urea and uric acid. *J. Insect. Physiol.* **31,** 179–185.

Burnet B., Sewell D. and Bos M. 1977 Genetic analysis of larval feeding behaviour in

Drosophila melanogaster. II. Growth relations and competition between selected lines.

Genet. Res. **30,** 149–161.

Dey S., Bose J. and Joshi A. 2012 Adaptation to larval crowding in *Drosophila ananassae*

leads to the evolution of population stability. *Ecol. Evol.* **2,** 941–951.

Fellowes M. D. E., Kraaijeveld A. R. and Godfray H. C. J. 1998 Trade-off associated with

selection for increased ability to resist parasitoid attack in *Drosophila melanogaster*. *Proc.*

R. Soc. Lond. B **265,** 1553–1558.

Fellowes M. D. E., Kraaijeveld A. R. and Godfray H. C. J. 1999 Association between

feeding rate and parasitoid resistance in *Drosophila melanogaster*. *Evolution* **53,** 1302–

1305.

Guo P. Z., Mueller L. D. and Ayala F. J. 1991 Evolution of behavior by density-dependent

natural selection. *Proc. Natl. Acad. Sci. USA* **88,** 10905–10906.

Joshi A. and Mueller L. D. 1988 Evolution of higher feeding rate in *Drosophila* due to

density-dependent natural selection. *Evolution* **42,** 1090–1092.

Joshi A. and Mueller L. D. 1993 Directional and stabilizing density-dependent natural

selection for pupation height in *Drosophila melanogaster*. *Evolution* **47,** 176–184.

Joshi, A. and Thompson J. N. 1995 Alternative routes to the evolution of competitive

ability in two competing species of *Drosophila*. *Evolution* **49,** 616–625.

Joshi A. and Mueller L. D. 1996 Density-dependent natural selection in *Drosophila*: trade-

offs between resource acquisition and utilization. *Evol. Ecol.* **10,** 463–474.

Joshi A. 1997 Laboratory studies of density-dependent selection: adaptations to crowding

in *Drosophila melanogaster*. *Curr. Sci.* **72,** 555–562.

Joshi A., Prasad N. G. and Shakarad M. 2001 *K*-selection, α-selection, effectiveness and

tolerance in competition: density-dependent selection revisited. *J. Genet.* **80,** 63–75.

Joshi A., Castillo R. B. and Mueller L. D. 2003 The contribution of ancestry, chance and

past and ongoing selection to adaptive evolution. *J. Genet.* **82,** 147–162.

Krijger C. L., Peters Y. C. and Sevenster J. G. 2001 Competitive ability of neotropical

Drosophila predicted from larval development times. *Oikos* **92,** 325–332.

MacArthur R. H. and Wilson E. O. 1967 *The theory of island biogeography*. Princeton

University Press, Princeton, NJ, USA.

Mueller L. D. and Ayala A. J. 1981 Trade-off between *r*-selection and *K*-selection in

Drosophila populations. *Proc. Natl. Acad. Sci. USA* **78,** 1303–1305.

Mueller L. D. and Sweet V. F. 1986 Density-dependent natural selection in *Drosophila*:

evolution of pupation height. *Evolution* **40,** 1354–1356.

Mueller L. D. 1988 Evolution of competitive ability in *Drosophila* by density-dependent

natural selection. *Proc. Natl. Acad. Sci. USA* **85,** 4383–4386.

Mueller L. D. 1990 Density-dependent selection does not increase efficiency. *Evol. Ecol.*

4, 290–297.

Mueller L. D., Graves J. L. and Rose M. R. 1993 Interactions between density-dependent

and age-specific selection in *Drosophila melanogaster*. *Func. Ecol.* **7,** 469–479.

Mueller L. D. 1997 Theoretical and empirical examination of density-dependent selection.

Annu. Rev. Ecol. Syst. **28,** 269–288.

Mueller L. D., Folk D. G., Nguyen N., Nguyen P., Lam P., Rose M. R. and Bradley T. J.

2005 Evolution of larval foraging behavior in *Drosophila* and its effect on growth and

metabolic rates. *Physiol. Entomol.* **30,** 262–269.

Mueller L. D. 2009 Fitness, demography, and population dynamics in laboratory

experiments. In *Experimental evolution: concepts, methods and applications of selection*

experiments (Garland T. Jr., Rose M. R., eds.), pp. 197–216. University of California

Press, Berkeley, CA, USA.

Mueller L. D., Rauser C. L. and Rose M. R. 2007 Population dynamics, life history and

demography: lessons from *Drosophila*. *Adv. Ecol. Res.* **37,** 77–99.

Mueller L. D. and Cabral L. G. 2012 Does phenotypic plasticity for adult size versus food

level in *Drosophila melanogaster* evolve in response to adaptation to different rearing

densities? *Evolution* **66,** 263–271.

Prasad N. G., Shakarad M., Anitha D., Rajamani M. and Joshi A. 2001 Correlated

responses to selection for faster development and early reproduction in *Drosophila*: the

evolution of larval traits. *Evolution* **55,** 1363–1372.

Prasad N. G. and Joshi A. 2003 What have two decades of laboratory life-history

evolution studies on *Drosophila melanogaster* taught us? *J. Genet.* **82,** 45–76.

Rajamani M., Raghavendra N., Prasad N. G., Archana N., Joshi A. and Shakarad M. 2006

Reduced larval feeding rate is a strong evolutionary correlate of rapid development in

Drosophila melanogaster. *J. Genet.* **85,** 209–212.

Rose M. R., Nusbaum T. J. and Chippindale A. K. 1996 Laboratory evolution:

experimental wonderland and the Cheshire cat syndrome. In *Adaptation* (Rose M. R.,

Lauder G. V., eds.), pp. 221–241. Academic Press, San Diego, CA, USA.

Rose M. R., Passananti H. B., Chippindale A. K., Phelan J. P., Matos M., Teotónio H. and

Mueller L. D. 2005 The effects of evolution are local: evidence from experimental

evolution in *Drosophila. Integr. Comp. Biol.* **45,** 486–491.

Shakarad M., Prasad N. G., Gokhale K., Gadagkar V., Rajamani M. and Joshi A. 2005

Faster development does not lead to correlated evolution of greater competitive ability in

Drosophila melanogaster. Biol. Lett. **1,** 91–94.

Sharmila Bharathi N., Prasad N. G., Shakarad M. and Joshi A. 2003 Variation in adult life

history and stress resistance across five species of *Drosophila. J. Genet.* **82,** 191–205.

Sharmila Bharathi N., Prasad N. G., Shakarad M. and Joshi A. 2004 Correlates of sexual

dimorphism for dry weight and development time in five species of *Drosophila. J. Zool.*

Lond. **264,** 87–95.

Sheeba V., Madhyastha N. A. A. and Joshi A. 1998 Oviposition preference for normal

versus novel food resources in laboratory populations of *Drosophila melanogaster. J.*

Biosci. **23,** 93–100.

Shiotsugu J., Leroi A. M., Yashiro H., Rose M. R. and Mueller L. D. 1997 The symmetry

of correlated responses in adaptive evolution: an experimental study using *Drosophila.*

Evolution **51,** 163–172.

Sokolowski M. B., Pereira H. S. and Hughes K. 1997 Evolution of foraging behavior in

Drosophila by density-dependent selection. *Proc. Natl. Acad. Sci. USA* **94,** 7373–7377.

StatSoft. 1995 *Statistica Vol. I: general conventions and statistics 1*. StatSoft Inc., Tulsa,

OK, USA.

FIGURES

Figure 1. Mean pre-adult survivorship at low and high larval density in the selected and

control populations of the two species, when cultured alone or in the presence of equal

numbers of eggs of white-eyed *D. melanogaster* competitors. Error bars are standard

errors around the mean of the four replicate population mean values. Significant

differences between pairs of means are indicated by labeling them with different letters.

For both species, as the three-way interaction between type of culture, selection regime

and larval density was not significant, only the difference between the mean pre-adult

survivorships of selected and control populations at high and low larval densities,

averaged over type of culture (single- or two-species), is highlighted.

Figure 2. Mean male and female pre-adult development time at low larval density in the

selected and control populations of the two species. Error bars are standard errors around

the mean of the four replicate population mean values. For both species, significant

differences between pairs of means are indicated by labeling them with different letters.

For *D. ananassae*, as the interaction between sex and selection regime was not significant,

only the difference between the mean pre-adult development time of males and females,

averaged over selection regimes is highlighted.

Figure 3. Mean male and female pre-adult development time at high larval density in the

selected and control populations of the two species. Error bars are standard errors around

the mean of the four replicate population mean values. Significant differences between

pairs of means are indicated by labeling them with different letters. For both species, as

the interaction between sex and selection regime was not significant, only the difference

between the mean pre-adult development time of males and females, averaged over

selection regimes is highlighted.

Figure 4. Mean duration in hours of different pre-adult life-stages at low larval density in

the selected and control populations of the two species. Error bars are standard errors

around the mean of the four replicate population mean values. For each species,

significant differences between pairs of means are indicated by labeling them with

different letters. For *D. ananassae*, the interaction between selection regime and life-stage

was not significant. Consequently, only the differences in mean duration between life-

stages, averaged over block and selection regime, have been highlighted.

Figure 5. Mean pre-adult survivorship at low density with three different levels of urea in

the food medium in the selected and control populations of the two species. Error bars are

standard errors around the mean of the four replicate population mean values. For each

species, significant differences between pairs of means are indicated by labeling them with

different letters. For both species, the interaction between selection regime and urea level

was not significant. Consequently, only the differences between mean survivorship at

different urea levels, averaged over block and selection regime, have been highlighted.

Figure 6. Mean pre-adult survivorship at low density with three different levels of

ammonia in the food medium in the selected and control populations of the two species.

Error bars are standard errors around the mean of the four replicate population mean

values. For each species, significant differences between pairs of means are indicated by

labeling them with different letters. For *D. n. nasuta* the interaction between selection

regime and ammonia level was not significant. Consequently, only the differences

between mean survivorship at different ammonia levels, averaged over block and selection

regime, have been highlighted.

Figure 7. (A) Mean pre-adult survivorship after larval feeding for different amounts of

time, and (B) mean dry weight (in 10^{-3} g) at eclosion after larval feeding for different

amounts of time corresponding to four different mean pre-adult survivorship levels, in the

D. ananassae selected and control populations. Error bars are standard errors around the

mean of the four replicate population mean values. For both traits, significant differences

between pairs of means are indicated by labeling them with different letters. For both

traits, the interaction between selection regime and feeding duration/survivorship was not

significant. Consequently, only the differences in mean survivorship between the different

levels of feeding duration, and differences in mean dry weight at eclosion at different

levels of survivorship, averaged over block, selection regime and sex, have been

highlighted.

Table 1. Summary of results of four-way ANOVA on mean arcsin squareroot transformed

pre-adult survivorship at low and high density in the two sets of selected populations and

controls, in single-species cultures and in two-species competitive cultures with equal

numbers of white-eye *D. melanogaster* eggs. Since the analysis was done on population

means, block and interactions involving block were not tested for significance.

____

861		ACU/AB		NCU/NB	
862	Effect	$F_{1,3}$	*P*	$F_{1,3}$	*P*
863	culture type	17.44	0.03	1.54	0.30
864	selection	14.11	0.033	43.67	<0.01
865	density	1621.07	<0.01	51.54	<0.01
866	culture type × selection	11.81	0.04	5.50	0.10
867	culture type × density	10.03	0.05	17.46	0.03
868	selection × density	525.03	<0.01	60.88	<0.01
869	culture type × selection × density	2.94	0.19	2.12	0.24

Table 2. Summary of results of two-way ANOVA on mean pre-adult development time,

assayed at low and high larval density at different generations in the two sets of selected

populations and controls. Since the analysis was done on population means, block and

interactions involving block were not tested for significance.

____

____ACU/AB________________NCU/NB________

<u>Low density</u>

Effect	$F_{1,3}$	P	$F_{1,3}$	P
selection	49.23	<0.01	115.50	<0.01
sex	82.32	<0.01	3.75	0.15
selection × sex	1.93	0.26	18.61	0.02

881   <u>High density</u>

Effect	$F_{1,3}$	P	$F_{1,3}$	P
selection	60.37	<0.01	364.65	<0.01
sex	41.18	<0.01	13.09	0.04
selection × sex	8.58	0.06	5.49	0.10

Table 3. Summary of results of three way ANOVA on mean pre-adult lifestage (egg, L1,

L2, L3, pupa) duration in the two sets of selected populations and controls. Since the

analysis was done on population means, block and interactions involving block were not

tested for significance.

____

_________________________________ACU/AB________________NCU/NB______

Effect	$F_{1,3\ sel/\ 4,12\ else}$	P	$F_{1,3\ sel/\ 4,12\ else}$	P
selection	168.38	<0.01	115.50	<0.01
life-stage	1778.55	<0.01	9485.70	<0.01
selection × life-stage	1.46	0.28	37.05	<0.01

Table 4. Summary of results of two way ANOVA on mean larval feeding rate, foraging

path length and pupation height in the two sets of selected populations and controls. Since

the analysis was done on population means, block and interactions involving block were

not tested for significance; only the F and P values for the main effect of selection regime

are shown.

____

_____________________________ ACU/AB _____________ NCU/NB _______

Trait	$F_{1,3}$	P	$F_{1,3}$	P
feeding rate	0.54	0.52	0.06	0.82
foraging path length	94.51	<0.01	0.48	0.54
pupation height	18.12	0.02	51.62	<0.01

Table 5. Summary of results of three way ANOVA on mean arcsin squareroot transformed

pre-adult survivorship at three different concentrations of urea or ammonia in the food

medium in the two sets of selected populations and controls. Since the analysis was done

on population means, block and interactions involving block were not tested for

significance.

____

________________________________ ACU/AB ________________ NCU/NB ________

<u>Urea tolerance</u>

919 Effect	$F_{1,3\ \text{sel}/\ 2,6\ \text{else}}$	P	$F_{1,3\ \text{sel}/\ 2,6\ \text{else}}$	P
920 selection	20.39	0.02	0.14	0.73
921 concentration	54.77	<0.01	24.88	<0.01
922 selection × concentration	0.51	0.62	0.32	0.74

923 <u>Ammonia tolerance</u>

924 Effect	$F_{1,3\ \text{sel}/\ 2,6\ \text{else}}$	P	$F_{1,3\ \text{sel}/\ 2,6\ \text{else}}$	P
925 selection	0.14	0.73	45.29	0.73
926 concentration	199.34	<0.01	417.76	<0.01
927 selection × concentration	14.97	0.02	0.88	0.46

Table 6. Summary of results of three way ANOVA on mean arcisn squareroot transformed

pre-adult survivorship (factors: block, selection regime and larval feeding duration) and

mean dry weight (factors: block, selection regime, sex and pre-adult survivorship) of

eclosing flies after larvae were allowed to feed for different time periods in the AB and

ACU populations. Since the analysis was done on population means, block and

interactions involving block were not tested for significance.

____

___________________________________ Survivorship _________ Dry weight ______

936 Effect	$F_{1,3 \text{ sel}/ 3,9 \text{ else}}$	P	$F_{1,3 \text{ sel}/ 3,9 \text{ else}}$	P
937 Selection	174.79	<0.01	0.01	0.94
938 feeding duration	37.07	<0.01		
939 (for survivorship)				
940 Survivorship (for dry weight)			16.39	<0.01
941 Sex (for dry weight)			223.27	<0.01
942 selection × feeding duration	0.35	0.78		
943 (for survivorship)				
944 selection × survivorship (for dry weight)			2.76	0.10
945 selection × sex (for dry weight)			3.25	0.17
946 survivorship × sex (for dry weight)			1.40	0.30
947 selection × survivorship × sex (for dry weight)			0.48	0.71

948   Figure 1.

965   Figure 2.

984   Figure 3.

1000

1001

1002

1003

1004   Figure 4.

1005

1006

(A) *D. ananassae*

(B) *D. n. nasuta*

1021   Figure 5.

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040   Figure 6.

1041

1042

1059   Figure 7.

1060

(A) Minimum critical feeding time

(B) Dry weight at different survivorship levels
corresponding to different feeding durations