

Limits to adaptation along environmental gradients

Jitka Polechová¹, Nick Barton¹,

¹ Institute for Science and Technology (IST Austria), Klosterneuburg, Austria

Running title: Limits to species' range

Keywords: genetic drift, range margin, species' border, species' range, ecological niche, evolution, geographic range, gene flow, genetic variation, heterogeneous environment

Statement of authorship: JP and NB jointly formalised and analysed the model. JP designed and processed the simulations. JP wrote the manuscript, whilst NB tirelessly commented on it.

Corresponding author: Jitka Polechová, Redlingerhütte, Am Campus 2, Klosterneuburg, Austria, Tel.: 0043 664 374 1607, E-mail: jitka@ist.ac.at

Abstract: Why do species not adapt to ever-wider ranges of conditions, gradually expanding their ecological niche? Theories of niche evolution typically omit spatial context, yet all species experience spatially variable conditions. Gene flow across environments has two conflicting effects on adaptation: while it increases genetic variation, which is a prerequisite for adaptation, gene flow may swamp adaptation to local conditions. We show that genetic drift can generate a sharp margin to a species' range, by reducing genetic variance below the level needed for adaptation to spatially variable conditions. Dimensional arguments and separation of ecological and evolutionary time scales reveal a simple threshold that predicts when adaptation at the range margin fails. Two observable parameters describe the threshold: i) the effective environmental gradient, which can be measured by the loss of fitness due to dispersal to a different environment, and ii) the efficacy of selection relative to genetic drift. The theory predicts sharp range margins even in the absence of abrupt changes in the environment. Furthermore, it implies that gradual worsening of conditions across a species' habitat may suddenly lead to range fragmentation – as adaptation to a wide span of conditions within a single species becomes impossible.

Introduction

Why a species' range sometimes ends abruptly, even when the environment changes smoothly across space, has interested ecologists and evolutionary biologists for many decades (1–8). Haldane (2) proposed that when the environment is spatially heterogeneous, a species may be unable to adapt and expand its range because gene flow from the centre swamps the populations at the range margins, preventing their adaptation. Theory showed that when genetic variance is fixed, adaptation indeed fails if the environment changes too steeply across space (9): the population remains well adapted only in the centre of the range, and gene flow swamps variants adapted to the margins, preventing range expansion. The result also explains range margins in the presence of competitors: then, interspecific competition in effect steepens the environmental gradient (10). Yet, this limit to adaptation assumes that local genetic variation is fixed.

Current theory states that when genetic variance can evolve, there is no sharp limit to a species' range (11). With environmental heterogeneity, genetic variance is generated by gene flow across the environment which supports a gradient in trait mean. Even for shallow environmental gradients, migration generates much larger variance than would be maintained by mutation alone (12, 13). When variance is maintained by gene flow across environments, deterministic theory predicts that adaptation to environmental differences is feasible until the fitness cost of maladapted variants becomes too high (i.e., when mean Malthusian fitness (growth rate) declines below zero). However, range margins are often associated with lower genetic variance (4) and experimental evidence suggests that low genetic variance coupled with high gene flow can prevent adaptation at the edge of a species' range in nature (14–16). Thus, there is clear evidence that low genetic variation may limit adaptive range expansion. This may be because genetic drift reduces local variance (17, 18) and hence the potential of the population to adapt (19). Yet, though genetic drift is necessarily strong at the range margins, studies of range expansion with genetic drift are few, and limited to simulations (20–22). By manipulating dispersal and carrying capacity, these studies demonstrated that genetic drift may be important to theory of species' range evolution. However, without establishing the key parameters, no quantitative predictions can be made that generalise beyond specific simulation models.

We explain when genetic drift limits species' range, and give the combination of parameters that describes the dynamics of evolution of a species' range when genetic variance can evolve, environment is heterogeneous, and the populations are finite. We assume that as environment changes across space, there is a corresponding change in an optimal value of some phenotypic trait. A deviation from the optimum is associated with a fitness cost, and hence range expansion requires adaptation to this environmental

gradient. Demography and evolution are considered jointly, and both genetic variance and trait mean can freely evolve *via* change in allele frequencies. Crucially, we include both genetic and demographic stochasticity. The model is first outlined at a population level, in terms of coupled stochastic differential equations (Materials and Methods: Evolutionary and ecological dynamics). Using this formalisation, we can identify the effective dimensionless parameters. Further, if we separate time-scales of ecology (fast) from evolution (slow), these reduce to just two key driving parameters. Next, individual based simulations (SI: Individual-based model) serve to determine the driving relationship between the derived key parameters, and to test its robustness.

Results

Deterministic limit

First, we show that the individual-based model matches the predictions at the deterministic limit. Fig. 1 illustrates the joint evolution of population size and trait mean and variance *via* change in allele frequencies, when genetic drift is weak. This is when $N\sigma\sqrt{s} \gg 1$ (23, 24), where population size within a dispersal distance σ is given by $N\sigma$, and s is the strength of selection per locus. N represents the local population size within each deme, which corresponds to population density in continuous space. Genetic variance V_G evolves, and it is generated primarily by gene flow due to mixing of genes from individuals with different phenotypes, well adapted to the diverse environments (Fig. 1D); the contribution due to mutational variance is negligible.

Scaling and separation of time scales

We now proceed by reducing the number of parameters (Table 1) in the model, including stochasticity (Materials and Methods: Rescaling). Rescaling space, time, trait, and population density reveals that three dimensionless parameters fully describe the system, neglecting mutation and assuming linkage equilibrium between loci. The first parameter carries over from the phenotypic model of (9): i) the effective environmental gradient $B = \frac{b\sigma}{r^*\sqrt{2V_S}}$. Two additional dimensionless parameters come from including stochasticity: ii) the efficacy of population regulation relative to demographic stochasticity $N\sigma\sqrt{r^*}$; and, iii) the efficacy of selection relative to the strength of genetic drift $N\sigma\sqrt{s}$.

The dynamics of evolution of species' range simplifies further because selection per locus s is typ-

ically much smaller than the rate of return to equilibrium population density r^* (see (25) and (26, Appendix D); Fig. S1 for values used here). This has two consequences. First, the ecological dynamics $\partial N/\partial t$ are faster than the evolutionary dynamics $\partial p/\partial t$ of the individual loci. Importantly, the genetic variance evolves yet substantially slower: whereas the trait mean changes with the product of strength of selection and the genetic variance, the directional change in the variance is slower by the inverse of the number of locally polymorphic loci (precisely, the “effective number of loci” (27). Genetic drift only slowly degrades expected heterozygosity $\langle pq \rangle$ at each locus (Fig. S2), reducing the variance. Hence, over ecological time scales, genetic variance can be treated as constant. Second, the effect of fluctuations due to genetic drift, scaling in one-dimensional habitat with $1/(N\sigma\sqrt{s})$ (see Fig. S2), is expected to dominate over the effect of demographic fluctuations, that rise with $1/(N\sigma\sqrt{r^*})$. Hence, we expect that the basic behaviour will be described by just two of the dimensionless parameters. These are the effective environmental gradient B and the efficacy of selection relative to the strength of genetic drift $N\sigma\sqrt{s}$.

Note that we define “parameter” as a quantity that describes a full set which defines the system and predicts its behaviour. As we are interested in the higher-order dynamics of species’ range expansion or collapse, for which N (or $N\sigma$) will be a measured quantity, we denote both $1/(N\sigma\sqrt{s})$ and $1/(N\sigma\sqrt{r^*})$ as parameters. Our key arguments propose that the dynamics of species’ range evolution can be understood based on bulk parameters B and $N\sigma\sqrt{s}$, rather than by focusing on asymmetric gene flow near the range margins. Later, we show that the prediction holds even when B and $N\sigma\sqrt{s}$ change steadily across space.

Threshold for collapse of adaptation

We simulated the basic model with a linear gradient, assuming equal allelic effects α (equal phenotypic change due to substitution at every locus). Parameters were drawn at random from distributions consistent with our knowledge of the range of parameters expected in nature and ensuring that without genetic drift, all ranges would expand (see Fig. S1 and (26, Discussion). This is denoted the “random set” of 1000 runs. At the start of all simulations, the population is perfectly adapted in the central part of the available habitat; over the 5000 generations of the simulation, expanding populations never reach the margin of the available habitat. We also performed additional runs to test whether the threshold obtained from the linear gradient applies when parameters are changing gradually across space. Namely, we tested whether a stable range margin forms at the predicted value when the environmental gradient varies across space or when the carrying capacity is non-uniform. Last, we tested the model assuming allelic effects

α_i (and hence the selection coefficients s_i) are exponentially distributed.

Fig. 2 shows our key result: the effective environmental gradient, B , and the efficacy of selection relative to the strength of genetic drift, $N\sigma\sqrt{s}$, determine the threshold for collapse of adaptation. This is because the genetic variance evolves primarily in response to $N\sigma\sqrt{s}$ and B , whilst the effect of demographic stochasticity $N\sigma\sqrt{r^*}$ is relatively weak. When $B \gtrsim 0.15 N\sigma\sqrt{s}$, genetic drift strongly degrades adaptation to a steeply changing environmental optimum, and the species' range contracts. The constant 0.15 is obtained as the best fitting threshold for the data in Fig. 2. Populations typically start to contract from the margins and ultimately collapse to a state with no or very little clinal variation. However, when genetic drift is very strong, populations collapse abruptly or even fragment (Fig. S3): this second threshold, based on a "critical gradient" of a phenotypic model, is explained later. This model with idealised linear gradient and uniform carrying capacity is used to identify the relationship between the driving parameters $N\sigma\sqrt{s}$ and B , which determine whether a population can adapt to environmental gradient.

In reality, the environment does not vary precisely linearly, and the carrying capacity is not uniform. When the steepness of environmental gradient varies across space, this threshold, $B \sim 0.15 N\sigma\sqrt{s}$, indicates where a stable range margin forms (Fig. 3). Without genetic drift, the genetic variance would steadily inflate, gradually reducing local population size due to an increasing number of maladapted individuals (see dashed lines in Fig. 3D). With genetic drift, the variance is pushed below the level necessary to maintain adaptation, the trait mean abruptly fails to match the optimum, so a sharp margin to the range forms. Similarly, a sharp range margin forms when $B \gtrsim 0.15 N\sigma\sqrt{s}$, if carrying capacity declines across the habitat for extrinsic reasons (Fig. S4).

A sharp range margin forms not only when all loci have equal allelic effects (i.e., the trait changes by a fixed value due to every substitution), but also when allelic effects are exponentially distributed. Then, range expansion slows down progressively around the threshold (Figs. S5 and S6), described by $B^* = 0.15 N\sigma\sqrt{s}$, where the mean selection coefficient is $\bar{s} \equiv \frac{\alpha^2}{2V_s}$. The mean selection coefficient gives an estimate for the expected range margin because clines at weakly selected loci are degraded by genetic drift (Fig. S7 and (24), reducing the genetic variance. For the population to expand further beyond the threshold, positively selected alleles with increasingly large effect need to arise (Fig. S7). In natural populations, these become ever rarer; for any finite distribution of allelic effects (as in our model), these become exhausted.

Neglecting genetic drift, low dispersal can enhance adaptation by reducing swamping by gene flow

(11, 15). However, with genetic drift, this is no longer true. To a first approximation, both the efficacy of selection relative to genetic drift $N\sigma\sqrt{s}$ and the effective environmental gradient $B = \frac{b\sigma}{\sqrt{2V_s}r^*}$ increase at the same rate with dispersal. Only a weak dependence on dispersal remains via r^* (because $\hat{N} = K\frac{r^*}{r_m}$), which favours low to intermediate dispersal: see Fig. S10. This beneficial effect of intermediate dispersal (21) can be much stronger in one-locus models where gene flow brings both new targets for mutation and positively selected mutants but does not directly swamp adaptation: the fitness decreases due to crowding in the new habitat (28) and mutation of the advantageous mutants back to the old habitat (29). Note that inbreeding depression, implied in the experimental study (30) is not necessary for dispersal to be beneficiary – but it would strengthen its beneficial effect.

In a deterministic phenotypic model with fixed genetic variance, there is a “critical gradient” $B_c \sim \sqrt{2A}$ (where $A = \frac{V_G}{r^*V_s}$), above which the trait mean fails to track the spatially changing optimum, and the population is well adapted only in the centre of its range (9). If genetic variance evolves due to gene flow across environments, in the absence of genetic drift, such a limit does not exist (11). We show that when genetic variance evolves due to selection, dispersal and genetic drift, populations can abruptly lose genetic variation and suffer demographic collapse. As genetic drift reduces the variance (Fig. S2), first a species’ range contracts from the margins (when $B \gtrsim 0.15 N\sigma\sqrt{s}$) but further decrease of variance such that $A \lesssim B^2/2$ leads to a sudden demographic collapse (Fig. 4; the critical gradient for logistic population regulation is given in (26, Fig. 3)). This demographic collapse may lead to range fragmentation, when a species cannot maintain adaptation throughout the whole range, yet multiple isolated populations persist (Fig. S3).

Our main result, that adaptation fails when the effective environmental gradient is large relative to the efficacy of selection versus genetic drift, $B \gtrsim 0.15 N\sigma\sqrt{s}$, can be rephrased to a form that is closely related to Haldane’s cost of selection (31). Haldane showed that in a single population, each substitution requires a certain number of selective deaths (i.e., reduction in mean fitness relative to the maximum possible), which is nearly independent of the strength of selection per locus. In our model with a spatially varying environment, b/α substitutions are required per unit distance: these substitution need to also arise via selective deaths. If too many selected substitutions are needed relative to births in the population, such that $b/\alpha \gtrsim 0.15 N r^*$, adaptation fails (Fig. S8). However, this failure is due to stochastic fluctuations, and so depends on the effective number of deaths per generation, Nr^* .

Discussion

We are just becoming to be able to assess the genomics of local adaptation across a species' range (32). Hence, there is a great need for a theory that integrates both genomic and demographic data. It would be exciting to see how our prediction improves experimental tests of the causes of range limits in one-dimensional habitats such as along a deep valley or a river (15), that were assuming fixed genetic variance. The predictive parameters can in principle be measured: the effective environmental gradient B measures the loss of fitness due to dispersal σ across the environmental gradient. Transplanting individuals exactly one standard deviation away (across the environmental gradient) gives B^2/r^* . The parameter r^* , which describes the rate of return of population to equilibrium can be estimated (eg., following (33)). Neighbourhood size in a one-dimensional habitat, $\sqrt{\pi N\sigma}$, can be estimated from neutral markers (34). The efficacy of selection relative to genetic drift, $N\sigma\sqrt{s_i}$, will vary across loci. The selection coefficients $s_i = \alpha_i^2/V_s$ can then be estimated either by mapping allelic effects α_i from the QTLs (quantitative trait loci) underlying an adaptive trait and measuring the strength of stabilising selection $1/(2V_s)$, or by estimating the selection per locus directly from the steepness of the clines in allele frequencies across space. Given a fixed (finite) distribution of selection coefficients s_i , we can predict when adaptation is expected to fail: for example, when, given a reduction in population size, a species' range would become prone to fragmentation.

The interface between ecology and evolution is critical in understanding the evolution of the species' range. Yet until now, no predictive theory included the fundamental stochastic process of genetic drift and demographic stochasticity – although close to the margin of a species' range, both are necessarily strong. Our theory explains that there is an inherent limit to adaptation arising in any (finite) natural population, and identifies the key parameters that determine this limit to a species' range. It explains that a sharp range margin forms when fitness cost, induced by a spatially varying environment, becomes too high relative to the efficacy of selection in the presence of genetic drift – even in the absence of fixed genetic constraints, such as insufficient genetic variance (9, 35) or rigid fitness trade-offs between traits (36, 37). Because the threshold depends only on the fitness cost of dispersal and the efficacy of selection per locus relative to genetic drift, it readily generalises to many traits. Note the prediction does not involve any asymmetry in gene flow, emphasised by (2).

The theory also implies that gradual decrease in population size can lead to a collapse of adaptation as genetic drift erodes the genetic variation, causing a sudden collapse or a fragmentation of a species' range. This is important for management of biological invasions when adaptation is required (38, 39)

and for biological conservation (40): a species' range may collapse due to genetic drift well before demographic factors (emphasised in (41)) become significant (40). Understanding species' range limits in a constant environment is also essential before extending the model to account for temporally changing environments, such as when modelling joint adaptation and range expansion in species' responding to climate change (42). Furthermore, the predicted emergence of inherent limits to species' ranges across steadily varying environments offers extensions to the theory of ecological speciation, and eventually, may help us to elucidate macroecological patterns of biodiversity.

Materials and Methods

Evolutionary and ecological dynamics

We model the joint evolution of i) population size and ii) trait mean and its variance *via* change in allele frequencies. The fitness of an individual declines quadratically with the deviation of the trait z from an optimum θ that changes linearly across space: $\theta = bx$, where b is the gradient in the environment and x is the distance in one dimension. The phenotypic trait z is determined by many additive di-allelic loci, so that genetic variance can evolve. For simplicity, we use a haploid model: for additive allelic effects, the extension to a diploid model is straightforward.

The Malthusian fitness of a phenotype z is $r(z, N) = r_e(N) + r_g(z)$, where $r_e(N)$ is the growth rate of a perfectly adapted phenotype, and includes density dependence; $r_g(z) \leq 0$ is the reduction in growth rate due to deviation from the optimum. N is the population density. The ecological component of growth rate r_e can take various forms: we assume that regulation is logistic, so that fitness declines linearly with density N : $r_e = r_m(1 - N/K)$, where r_m is the maximum per capita growth rate in the limit of the local population density $N \rightarrow 0$. The carrying capacity K is assumed uniform across space. Stabilising selection on the optimum θ has strength $1/(2V_s)$. Hence, for any individual, the drop in fitness due to maladaptation is $r_g(z) = -\frac{(z-\theta)^2}{2V_s}$. The local mean fitness is then $\bar{r}_g(\bar{z}) = -\frac{(\bar{z}-\theta)^2}{2V_s} - \frac{V_P}{2V_s}$, where $V_P = V_G + V_E$ is the phenotypic variance. The loss of fitness due to environmental variance V_E can be included in $r_m^* = r_m - V_E/(2V_s)$; hence in this model, V_E is a redundant parameter. For any given additive genetic variance V_G (assuming a Gaussian distribution of breeding values), the trait mean \bar{z} satisfies:

$$\frac{\partial \bar{z}}{\partial t} = \frac{\sigma^2}{2} \frac{\partial^2 \bar{z}}{\partial x^2} + \sigma^2 \frac{\partial \ln(N)}{\partial x} \frac{\partial \bar{z}}{\partial x} + V_G \frac{\partial \bar{r}}{\partial \bar{z}} + \zeta \quad (1)$$

The first term gives the change in the trait mean due to migration with mean displacement of σ , the second term describes the effect of the asymmetric flow from areas of higher density and the third term gives the change due to selection (43, Eq. 2). The last term ζ gives the fluctuations in the trait variance due to genetic drift (following (23), and using $V_{G,LE} = \sum_i \alpha_i^2 p_i q_i$): $\zeta = \sqrt{V_{G,LE}/N} dW_\zeta(x, t)$, where dW represents white noise in space and time.

The trait mean is $\bar{z} = \sum_i \alpha_i p_i$ for a haploid model, where p_i is the i -th allele frequency, $q_i = 1 - p_i$ and α_i is the effect of the allele on the trait – the change of the trait mean \bar{z} as frequency of locus i changes from 0 to 1. For both haploid and diploid models, the allele frequencies p_i change as:

$$\frac{\partial p_i}{\partial t} = \frac{\sigma^2}{2} \frac{\partial^2 p_i}{\partial x^2} + \sigma^2 \frac{\partial \log(N)}{\partial x} \frac{\partial p_i}{\partial x} + p_i q_i \frac{\partial \bar{r}}{\partial p_i} - \mu(p_i - q_i) + \epsilon \quad (2)$$

The expected change of allele frequency due to a gradient in fitness and local heterozygosity is $p_i q_i \frac{\partial \bar{r}}{\partial p_i} = s_i p_i q_i (p_i - q_i - 2\Delta_i)$, where selection at locus i is $s_i \equiv \alpha_i^2 / (2V_s)$ and $\Delta_i = (\bar{z} - bx) / \alpha_i$ (11, two-allele model, Appendix 3). Here, the fourth term describes the change due to (symmetric) mutation at rate μ . The last term ϵ describes genetic drift (following (23): $\epsilon = \sqrt{\frac{p_i q_i}{N}} dW_\epsilon(x, t)$).

Equation 2 is only exact at linkage equilibrium (i.e., neglecting covariance between allele frequencies): but we will see that this is a good approximation for unlinked loci. Whereas migration across the habitat generates positive linkage disequilibrium between any pair of loci, stabilising selection drives negative disequilibrium, and these cancel precisely unless selection per locus is strong. The derivation, which generalizes an ingenious but little known argument by (44), is given in Appendix.

Population dynamics reflect diffusive migration, growth due to the mean Malthusian fitness \bar{r} , and stochastic fluctuations. The number of offspring follows a Poisson distribution with mean and variance of N . Fluctuations in population numbers are given by $\xi = \sqrt{N} dW_\xi(x, t)$ (e.g. (45)):

$$\frac{\partial N}{\partial t} = \frac{\sigma^2}{2} \frac{\partial^2 N}{\partial x^2} + \bar{r}N + \xi \quad (3)$$

Rescaling

The model can be further simplified by rescaling time relative to the strength of density dependence r^* , distance relative to dispersal, trait relative to strength of stabilising selection and population density relative to equilibrium density with perfect adaptation $\hat{N} = Kr^*/r_m$; as in (9, 11): $T = r^*t$, $X =$

$$x\sqrt{\frac{2r^*}{\sigma^2}}, \quad Z = \frac{z}{\sqrt{r^*V_s}}, \quad \tilde{N} = N/\hat{N}.$$

The rescaled equations for evolution of allele frequencies and for demographic dynamics are:

$$\frac{\partial \tilde{N}}{\partial T} = \frac{\partial^2 \tilde{N}}{\partial X^2} + \bar{R}\tilde{N} + \sqrt{\frac{\tilde{N}}{\hat{N}\sigma\sqrt{r^*}}} dW_{\tilde{\xi}(X,T)} \quad (4)$$

$$\begin{aligned} \frac{\partial p}{\partial T} &= \frac{\partial^2 p}{\partial X^2} + 2\frac{\partial \log(\tilde{N})}{\partial X} \frac{\partial p}{\partial X} + \frac{s}{r^*}(pq - 2\frac{\bar{Z} - BX}{\alpha^*}) - \\ &- \frac{\mu}{r^*}(p - q) + \sqrt{\frac{pq}{\hat{N}\sigma\sqrt{s}}} dW_{\tilde{\epsilon}(X,T)} \end{aligned} \quad (5)$$

where $\bar{R} \equiv \bar{r}/r^* = 1 - \tilde{N} - (BX - Z)^2/2$.

The rescaled equations show that four parameters fully describe the system. The first two are i) the effective environmental gradient is $B \equiv b\sigma/(r^*\sqrt{2V_s})$ and ii) the strength of genetic drift relative to selection $1/(\hat{N}\sigma\sqrt{s})$. As a third parameter, one can either take iii) demographic stochasticity relative to strength of density-dependence $1/(\hat{N}\sigma\sqrt{r^*})$ or the ratio of $\hat{N}\sigma\sqrt{s}$ to $\hat{N}\sigma\sqrt{r^*}$, which is the strength of selection relative to density dependence s/r^* . Note, that the scaled effect of a single substitution, $\alpha^* \equiv \alpha/\sqrt{r^*V_s} = \sqrt{2s/r^*}$. The effect of this third parameter is small, because $s \ll r^*$. The fourth parameter, μ/r^* , will typically be very small, and hence we will neglect it. Note, that \hat{N} gives population density, where population is still well adapted: $N = \hat{N}$ throughout most of the species' range away from sink regions near the existing margins. Table 1 (bottom) summarises the full set that describes the system.

[Clines are at linkage equilibrium] Stabilising selection on a quantitative trait generates negative linkage disequilibrium, whereas dispersal generates positive linkage disequilibrium. Felsenstein's (44) analysis of variance components showed that at equilibrium, the linkage disequilibrium generated by dispersal cancels out with the negative linkage disequilibrium generated by stabilising selection.

The argument extends to a quantitative trait determined by many bi-allelic loci, here demonstrated for a haploid genome. The genetic variance is $V_G = \sum \alpha_i^2(p_i(x)q_i(x)) + 2\sum_{i \neq j} \alpha_i\alpha_j D_{ij}$. The increase of linkage disequilibrium at QLE (47) between dispersal with variance σ^2 and recombination r_{ij} is given by $D_{ij|disp} = \frac{\sigma^2}{r_{ij}} \frac{dp_i}{dx} \frac{dp_j}{dx}$ (46). With allele frequencies at equilibrium, the linkage disequilibrium generated by stabilising selection alone is $D_{ij|sel} = \frac{-r + \sqrt{r_{ij}^2 + 4(1-r_{ij})\alpha_i\alpha_j/V_s p_i q_i p_j q_j}}{2(1-r_{ij})} \sim -\alpha_i\alpha_j/(r_{ij}V_s) p_i q_i p_j q_j$ for D_{ij} small.

In this first order approximation, the terms cancel for each pair of loci when the cline shape is the same as that under linkage equilibrium (11, two-allele model) – independently of the cline spacing across

space: $D_{ij} \equiv D_{ij|disp} + D_{ij|sel} = 0$. This is because $\frac{dp_i}{dx} = p_i q_i 4/w_i$, and the cline width at linkage equilibrium is $w_i = \frac{4\sigma\alpha_i}{\sqrt{V_s}}$.

It may be that the cline width w or the linkage disequilibrium is distorted by additional forces, and/or by strong selection. However, unless selection is strong, the first order approximation gives a simple prediction for the pairwise disequilibrium: $D_{ij} = \frac{\sigma^2}{r_{ij}} p_i q_i p_j q_j 16 / (w_i w_j) - \alpha_i \alpha_j / (r_{ij} V_s) p_i q_i p_j q_j$.

Acknowledgments

We would like to thank S. Baird, A. Betancourt, J.P. Bollback, J.R. Bridle, D. Field, A. Hancock, F. Jesse, T. Paixão, T. Priklopil, R.A. Nichols, S. Novak, S. Sarikas, H. Uecker and M.G.J. de Vos for discussions and comments on the earlier drafts, and European Research Council grant (250152) to N.B. for funding.

References

1. Merriam CH (1894) *Laws of temperature control of the geographic distribution of terrestrial animals and plants*. National Geographic 6:229–238.
2. Haldane JBS (1956) *The relation between density regulation and natural selection*. Proc R Soc B 145:306–308.
3. Antonovics J (1976) *The nature of limits to natural selection*. Annals of the Missouri Botanical Garden, 63(2):224–247.
4. Gaston K (2003) *The structure and dynamics of geographic ranges*. Oxford University Press.
5. Blows MW, Hoffmann AA (2005) *A reassessment of genetic limits to evolutionary change*. Ecology 86(6):1371–1384.
6. Bridle J, Vines T (2007) *Limits to evolution at range margins: when and why does adaptation fail?*. Trends Ecol Evol 22(3):140–147.
7. Kawecki TJ (2008) *Adaptation to marginal habitats*. Annu Rev Ecol Evol Syst. 39:321–342.
8. Sexton J, McIntyre P, Angert A, Rice K (2009) *Evolution and ecology of species' range limits*. Annu Rev Ecol Evol Syst. 40:415–436.

9. Kirkpatrick M, Barton N (1997) *Evolution of a species' range*. Am Nat 150(1):1–23.
10. Case T, Taper M, (2000) *Interspecific competition, environmental gradients, gene flow, and the co-evolution of species' borders*. Am Nat 155(5):583–605.
11. Barton N (2001) *Adaptation at the edge of a species' range*. Integrating Ecology and Evolution in a Spatial Context, eds Silvertown J, Antonovics J (Blackwell, London): pp. 365–392.
12. Houle D, Morikawa B, Lynch M (1996) *Comparing mutational variabilities*. Genetics, 143(3):1467–1483.
13. Johnson T, Barton N (2005) *Theoretical models of selection and mutation on quantitative traits*. Philos Trans R Soc B 360(1459):1411–1425.
14. Magiafoglou A, Carew M, Hoffmann A (2002) *Shifting clinal patterns and microsatellite variation in Drosophila serrata populations: a comparison of populations near the southern border of the species range*. J Evol Biol 15(5):763–774.
15. Hendry AP, Taylor EB (2004) *How much of the variation in adaptive divergence can be explained by gene flow? an evaluation using lake-stream stickleback pairs*. Evolution 58(10):2319–2331.
16. Bridle JR, Gavaz S, Kennington WJ (2009) *Testing limits to adaptation along altitudinal gradients in rainforest Drosophila*. Proc R Soc B 276(1661):1507–1515.
17. Fisher R (1930) *The genetical theory of natural selection*. (Clarendon Press, Oxford)
18. Wright S (1931) *Evolution in Mendelian populations*. Genetics 16(2):97–159.
19. Lande R (1976) *Natural selection and random genetic drift in phenotypic evolution*. Evolution 30(2):314–334.
20. Butlin R, Bridle J, Kawata M (2003) *Genetics and the boundaries of species' distributions*. Concepts and consequences: the 43rd Annual symposium of the British Ecological Society, eds Blackburn TM, Gaston KJ (Blackwell, London):pp. 274–295.
21. Alleaume-Benharira M, Pen I, Ronce O (2006) *Geographical patterns of adaptation within a species? range: interactions between drift and gene flow*. J Evol Biol 19(1):203–215.
22. Bridle J, Polechová J, Kawata M, Butlin R (2010) *Why is adaptation prevented at ecological margins? New insights from individual-based simulations*. Ecol Lett 13(4):485–494.

23. Nagylaki T (1978) *Random genetic drift in a cline*. Proc Natl Acad Sci USA 75(1):423–426.
24. Polechová J, Barton N (2011) *Genetic drift widens the expected cline but narrows the expected cline width*. Genetics, 1(189):227–235.
25. Sæther B, Lande R, Engen S, Weimerskirch H, Lillegård M, et al. (2005) *Generation time and temporal scaling of bird population dynamics*. Nature 436(7047):99–102.
26. Polechová J, Barton N, Marion G (2009) *Species' range: Adaptation in space and time*. Am Nat 174(5):E186–204.
27. Barton N (1999) *Clines in polygenic traits*. Genet Res 74(3):223–236.
28. Gomulkiewicz R, Holt RD, Barfield M (1999) *The effects of density dependence and immigration on local adaptation and niche evolution in a black-hole sink environment*. Theor Pop Biol 55(3):283–296.
29. Uecker H, Otto SP, Hermisson J (2014) *Evolutionary rescue in structured populations*. Am Nat 183(1):E17–E35.
30. Sexton JP, Strauss SY, Rice KJ (2011) *Gene flow increases fitness at the warm edge of a species' range*. Proc Natl Acad Sci USA 108(28):1704–1709.
31. Haldane JBS (1957) *The cost of natural selection*. Journal of Genetics 55(3):511–524.
32. Savolainen O, Lascoux M, Merilä J (2013) *Ecological genomics of local adaptation*. Nat Rev Genet 14(11):807–820.
33. Lande R, Engen S, Saether B, Filli F, Matthysen E, et al. (2002) *Estimating density dependence from population time series using demographic theory and life-history data*. Am Nat 159(4):321–337.
34. Rousset (1997) *Genetic differentiation and estimation of gene flow from f-statistics under isolation by distance*. Genetics 145(4):1219–1228.
35. Chevin LM, Lande R, Mace GM (2010) *Adaptation, plasticity, and extinction in a changing environment: towards a predictive theory*. PLoS Biol 8(4):e1000357.
36. Etterson J, Shaw R (2001) *Constraint to adaptive evolution in response to global warming*. Science 294(5540):151–154.

37. Colautti RI, Eckert CG, Barrett SC (2010) *Evolutionary constraints on adaptive evolution during range expansion in an invasive plant*. Proc R Soc B 277(1689):1799–1806.
38. Lee C (2002) *Evolutionary genetics of invasive species*. Trends Ecol Evol 17(8):386–391.
39. Alexander JM, Edwards PJ (2010) *Limits to the niche and range margins of alien species*. Oikos, 119(9):1377–1386.
40. Spielman D, Brook BW, Frankham R (2004) *Most species are not driven to extinction before genetic factors impact them*. Proc Natl Acad Sci USA, 101(42):15261–15264.
41. Lande R (1988) *Genetics and demography in biological conservation*. Science, 241(4872):1455–1460.
42. Hoffmann AA, Sgrò CM (2011) *Climate change and evolutionary adaptation*. Nature 470(7335):479–485.
43. Pease C, Lande R, Bull J (1989) A model of population growth, dispersal and evolution in a changing environment. Ecology 70(6):1657–1664.
44. Felsenstein J (1977) *Multivariate normal genetic models with a finite number of loci*. Proceedings of the International Conference on Quantitative Genetics, eds Pollak E, Kempthorne O, Bailey Jr T (Iowa State University Press, Iowa), pp. 227–245.
45. Lande R, Engen S, Sæther BE (2003) *Stochastic population dynamics in ecology and conservation* (Oxford University Press, Oxford).
46. Barton N (1986). *The maintenance of polygenic variation through a balance between mutation and stabilizing selection*. Genetical Research 47(3):209–216.
47. Barton N & Turelli M (1991) *Natural and sexual selection on many loci*. Genetics 127(1):229–255.




Figure 1: Illustration of the individual based model at the limit of weak genetic drift, when the species' range keeps expanding as predicted by the deterministic model (11). (A) Trait mean \bar{z} matches the optimum $\theta = bx$, shown for the starting population (light blue) and after 5000 generations (dark blue). The spread of the trait values z for all individuals is shown with dots. (B) Local population size is close to the deterministic prediction $\hat{N} = Kr^*/r_m = K(1 - \frac{\sigma b}{2\sqrt{V_s r_m}})$. (C) Clines for allele frequencies are shown by thin black lines, the predicted clines (dashed) have widths $w_s = 4\sigma/\sqrt{2s}$ and are spaced α/b apart. (D) Total genetic variance is shown in blue, the linkage-equilibrium component in black; the dashed line gives the prediction $V_G = V_{LE} = b\sigma\sqrt{V_s}$: each cline contributes genetic variance $V_{G,i} = \alpha_i \sigma\sqrt{V_s}$ and per unit distance, there must be b/α clines if the trait mean matches into the optimum (11, p. 378). Parameters, defined in Table 1: $b = 0.1$, $\sigma^2 = 1/2$, $V_s = 2$, $r_m = 1.025$, $K = 300$, $\alpha = 1/\sqrt{20}$, $\mu = 10^{-6}$, 5000 generations.




Figure 2: Species' range starts to contract when the effective environmental gradient is steep relative to the efficacy of selection relative to genetic drift: $B \gtrsim 0.15 N\sigma\sqrt{s}$, where $B = \frac{b\sigma}{r^*\sqrt{2V_s}}$. This threshold is shown by a dashed line. The rate of expansion increases from light to dark blue and rate of range contraction increases from orange to red. Grey dots denote populations for which neither expansion nor collapse is significant at $\alpha = 2\%$. Open dots indicate fragmented species' ranges (illustrated at Fig. S3). The ranges of the underlying (unscaled) parameters are in the following intervals: $b = [0.01, 1.99]$, $\sigma = [0.5, 4.8]$, $V_s = [0.006, 8.4]$, $K = [4, 185]$, $r_m = [0.27, 2]$ and $\alpha = [0.01, 0.39]$, $\mu = [10^{-8}, 8 \cdot 10^{-5}]$; the number of genes is between 7 and 3971. The selection coefficient per locus is hence in the interval of $s = [3 \cdot 10^{-4}, 0.66]$, with median of 0.007. Parameters are defined in Table 1; parameter distributions are shown in Fig. S1.




Figure 3: With a steepening environmental gradient, a stable range margin forms when $B \gtrsim 0.15N\sigma\sqrt{s}$ (red dots). (A) The gradient in trait mean follows the environmental optimum (dashed line) until the gradients steepens so that $B \gtrsim 0.15N\sigma\sqrt{s}$, where expansion stops. (B) Population density drops off sharply when the predicted threshold (red dots) is reached. Dashed line gives the predicted population size assuming variation is not eroded by genetic drift. (C) Three representative clines are shown in black, other clines form the gray background. (D) Adaptation fails when genetic variance fails to increase fast enough to match the steepening environmental gradient. For all subfigures, dashed lines give deterministic predictions ((11) and Fig. 1). Parameters: central gradient $b_0 = 0.12$; $\sigma^2 = 1/2$, $V_s = 1/2$, $r_m = 1.06$, $K = 53$, $\mu = 2 \cdot 10^{-7}$. Population is depicted after 100 000 generations, whilst it stops expanding after 40 000 generations (subfigure A also shows the initial stage, in light blue).




Figure 4: Phenotypic model predicts a second sharp transition in the dynamics. As the effective environmental gradient $B \equiv b\sigma/(r^*\sqrt{2V_s})$ increases, the scaled variance $A \equiv V_G/(r^*V_s)$ increasingly deviates from the deterministic prediction with evolvable variance ((11), gray dotted line). The variance decreases due to the combined forces of genetic drift (see Fig. S2 A) and selection on small transient deviations of the trait mean from the optimum (see Fig. S9). Once A becomes smaller than $B^2/2$ (dashed curve, which gives the “critical gradient” $B_c \sim \sqrt{2A}$), the population collapses abruptly. Furthermore, no adaptation is maintained beyond the solid line $B_e = (2 + A)/\sqrt{2}$, where the phenotypic model (9) predicts extinction. Open dots denote fragmented species’ range (Fig S3). The colours are as in Fig. 2, the threshold when range starts to contract, $B \sim 0.15N\sigma\sqrt{s}$, is illustrated by the vertical dashed line. Parameters: b increases from 0.025 to 1.25, $\sigma^2 = 1/2$, $V_s = 1/2$, $K = 50$, $\alpha = 1/10$, $r_m = 1.06$, 5000 generations. Ten replicates are shown for each B .

| parameter | dimension | |
|---------------------|--------------|--|
| b | Z/D | gradient in the environmental optimum |
| V_s | Z^2T | variance of stabilising selection |
| σ | D/\sqrt{T} | dispersal per generation |
| K | T/D | carrying capacity per deme |
| r_m | $1/T$ | maximum exponential growth rate |
| α | Z | allelic effect |
| μ | $1/T$ | mutation rate |
| s | $1/T$ | selection per locus: $s \equiv \alpha^2/(2V_s)$ |
| r^* | $1/T$ | rate of return to equilibrium pop. size: $r^* = r_m - \frac{V_G}{2V_s}$ |
| B | – | effective environmental gradient, $B = b\sigma/(r^*\sqrt{2V_s})$ |
| $N\sigma\sqrt{s}$ | – | efficacy of selection relative to strength of genetic drift |
| $N\sigma\sqrt{r^*}$ | – | efficacy of population regulation relative to demographic stochasticity |

Table 1: **Three scale-free parameters: B , $N\sigma\sqrt{s}$ and $N\sigma\sqrt{r^*}$ describe the system; top section gives seven parameters of the model before rescaling.** Dimensions: T stands for time, D for distance and Z for trait. Note that with a Poisson number of offspring, the effective population size N_e (which measures rate of genetic drift / coalescence) is identical to the N that regulates population growth due to crowding: hence both carrying capacity K and population size N have units of T/D . Mutation rate μ is set to be small, with minimal contribution to the dynamics, and hence μ/r^* is neglected in the rescaled parameterization.

SI: INDIVIDUAL-BASED MODEL

We test the predictive power of the derived key parameters using individual based simulations. Individuals are distributed among demes that form a one-dimensional habitat, with the phenotypic optimum varying along the habitat, and experience a life cycle consisting of selection, mutation, recombination, and then dispersal. Distance is measured in demes, time in generations. Every generation, each individual mates with a partner drawn from the same deme, with probability proportional to its fitness, to produce a number of offspring drawn from a Poisson distribution with mean of $Exp[r]$, where r is the individual's Malthusian fitness in continuous time. Individual fitness declines due to deviation of the phenotypic trait (z) from the optimum and due to crowding (fitness is density-dependent). The trait is determined by a number of additive di-allelic loci, which permits genetic variation to evolve. All parameters are described in Table 1. The model is derived as a limit to continuous time, and so applies to a wide range of models that reduce to this limit. In this limit, the rate of spatial dispersal depends only on the variance of distance moved, and the effective population density (for both allele frequency and demographic fluctuations) depends only on the variance of offspring number.

Life cycle: Discrete-time individual based simulations are set to correspond to the model with continuous time and space. The life-cycle is selection \rightarrow mutation \rightarrow recombination \rightarrow migration.

Dispersal: The habitat is formed by a one-dimensional array of demes. With deme spacing $\delta x = 1$, the population size per deme corresponds to the population density. We assume diffusive migration with a Gaussian dispersal kernel. The tails of the dispersal kernel need to be truncated: we choose truncation at two standard deviations of the dispersal kernel throughout, and adjust the dispersal probabilities following [48, p. 1209] so that the discretised dispersal kernel sums to 1, and the variance of dispersal is adjusted correctly. For dispersal per generation at $\sigma = \sqrt{1/2}$, dispersal reduces to a nearest neighbour migration with a probability of migration left and right of $m = 1/4$.

Selection: Every generation, each individual produces a Poisson number of offspring with mean of the individual's fitness $Exp[r]$; where $r = r_m(1 - N/K) - \frac{(z-\theta)^2}{2V_s}$, as defined earlier.

Mutation: Mutation rate is set to be small so that its contribution to genetic variance is negligible, but large enough to in principle enable expansion of a species' range over the total time of 5000 generations. Specifically, it is set to one substitution per the whole population and generation. Genetic variance maintained in a population due to dispersal across environments can be substantially larger than genetic variance maintained by mutation-selection balance in uniform environments. In uniform environments, mutational variance $V_m = \sum_i^{n_l} \mu \alpha_i^2$ (where α_i are the allelic effects, and n_l is the number of loci) is robustly estimated to be between about $5 \cdot 10^{-5}V_E$ and $5 \cdot 10^{-3}V_E$ [49]. Taking a heritability $h^2 \equiv V_G/(V_G + V_E) = 1/3$ we get V_m between $10^{-4}V_G$ and $10^{-2}V_G$. In our model, genetic variance is inflated due to dispersal across environments, and so V_m/V_G must be yet smaller. Taking the higher limit of $V_m = 10^{-2}V_G$, follows that μ should be smaller than about $4 \cdot 10^{-3}$. It turns out that in general, the increase of genetic variance due to mutation cannot be fully included in the predictions, as the contribution of mutation-selection balance cannot be robustly separated from the clinal variation. A considerably higher

40 genetic variance than $V_{G,mut} = 2\mu n_l V_s$ (up to the limit of $1/4\alpha^2 n_l$) can arise due inflation of the variance by mutation along existing clines. Therefore, we concentrate on a parameter range where the contribution of mutation to genetic variance is low, which is a biologically plausible range. In our model, genetic variance is maintained by gene flow across the environment.

45 **Reproduction, recombination:** The mating partner is drawn from the same deme, with the probability proportional to its fitness. Selfing is allowed at no cost. The genome is haploid with unlinked loci (the probability of recombination between any two loci is $1/2$); the allelic effects α_i of the loci combine in an additive fashion.

Simulation runs: Evolution starts with a well adapted population at the centre of the habitat. The habitat is about 50 10 cline widths wide; the number of genes is chosen so that with all genes adapted, the population spans the whole habitat, and that there are enough genes to maintain the “optimal” variance $V_G = b\sigma\sqrt{V_S}$ at the central part of the habitat. At the start of the simulation, half of the genes are adapted: their clines take the form and spacing as assumed for the deterministic model under linkage equilibrium.

55 The population evolves for 5000 generations; in total, we recorded over a thousand runs where without genetic drift the local population density would be greater than 4 assuming uniform adaptation (such that trait mean matches the optimum). We a-priori eliminated very small local population sizes ($N < 4$) so that the population size within a generational dispersal is not excessively small. The parameters were first varied one at a time, and then we tested the threshold drawing the parameters from distributions consistent with our knowledge of the range 60 expected in nature (see [26, Discussion]). The latter will be referred to as the “random” set, with 1000 runs. The Mathematica code for the simulations, including the distributions used to draw the unscaled parameters for the “random” set is provided in FileS1. Fig. S1 shows the realised distributions for both the unscaled parameters and compound parameters.

65 REFERENCES FOR SI: INDIVIDUAL BASED MODEL

[48] J. Polechová, N. Barton, *Evolution* **59**, 1194 (2005).

[49] D. Houle, B. Morikawa, M. Lynch, *Genetics* **143**, 1467 (1996).

SUPPLEMENTARY FIGURES AND FIGURE LABELS




Figure S1: **Distribution of parameters used in the “random set”, as described in Table 1.** Grey: all unscaled parameters. Light blue: Composite parameters. Dark green: Scale-free parameters (both s/r^* and $1/(\hat{N}\sigma\sqrt{r^*})$ are given for a reference though one of them is always redundant; see Materials and Methods: Rescaling).




Figure S2: Cline width decreases as genetic drift gets stronger relative to selection. (A): Cline width, defined as a measure of total heterozygosity across space ($w \equiv 4 \int p q$), is approximated by $w = w_s(1 - 0.392/(N\sigma\sqrt{s}))$ (thick dotted line; method from [24, supplementary text]). Dots give the median cline width across loci with their positive and negative standard deviations. The dashed horizontal line gives the deterministic cline width, $w_s = 4\sigma/\sqrt{2s}$ [50, 51]. As the genetic drift gets very strong relative to selection, the predictions diverges (bottom right): for very weak selection, the neutral limit $\langle w_0 \rangle \rightarrow 4\sigma^2 N$ is approached (solid diagonal line; from [52]). Note that genetic variance decreases with the same factor as cline width. (B): The decrease in cline width can be understood from the rise in fluctuations of each cline – as fluctuations increase, allele frequencies fix locally and the cline steepens. Fluctuations in clines $\langle F \rangle$ rise approximately with $\langle F \rangle = 0.392/(N\sigma\sqrt{s})$, shown by a dashed line. $\langle F \rangle$ is the variance in allele frequencies scaled by the expected allele frequency, averaged across space, therefore ranging between 0 and 1: $\langle F \rangle \equiv \frac{\int_{-\infty}^{\infty} \text{var}(p(x,t)) dx}{\int_{-\infty}^{\infty} \langle p(x,t) \rangle \langle q(x,t) \rangle dx}$ [24, p. 228]. Note that these formulae apply only to one-dimensional habitats: as the width of the second (selectively neutral) spatial dimension of the habitat increases, the predictions will start to differ because the effect of genetic drift on a cline depends only weakly on selection in two-dimensional habitats; see [53]). Parameters: $b = 0.1$, $\sigma^2 = 1/2$, $V_s = 2$, $r_m = 1.025$, $\alpha = 1/\sqrt{20}$, $\mu = 10^{-4}$, carrying capacity K increases from 4 to 260.




Figure S3: **Species range can fragment when $B > 0.15N\sigma\sqrt{s}$ (Fig. 2), and additionally, $B > \sqrt{2A}$ (Fig. 4).** Exact conditions that always lead to range fragmentation were not determined. Fragmented populations are shown in open circles in Figs. 2 and 4. (A) Typically, there is no adaptation in trait mean left within each sub-range. (B) Populations are disjunct, and across the habitat, the population size is considerably smaller than predicted for “perfect” adaptation. (C) Typically, there is no clinal variation although transiently, a few clines are maintained. (D) Correspondingly, genetic variance is mostly near zero. Parameters: $b \doteq 1.18$; $\sigma^2 \doteq 1$, $V_s \doteq 0.44$, $r_m = 1.97$, $K = 29.2$, $\mu = 6 \cdot 10^{-8}$, $\alpha = 0.0093$, time = 5000 generations (shown at generation 4800). Note, that this fragmentation is not driven by edge effects that arise when population at the carrying capacity reaches the edge of the available habitat, where there is less maladaptive gene flow, which leads to local increase of density, followed by suppression of nearby populations towards the centre – and the effect propagates [54]. This effect is explained in [48]. Here, the simulations are set up such that the population never reaches the margins of the available habitat.




Figure S4: Non-uniform carrying capacity generates a stable range margin. (A) The optimum changes across the environment with a constant gradient $b = 0.3$ – the population starts well adapted at the more central part of the habitat (lighter blue). (B) The population density declines away from the centre - red dots give the predicted failure of the adaptation, based on $B = 0.15 N \sigma \sqrt{s}$. (C) Three representative clines are shown in black, other clines form the gray background (every tenth cline displayed). (D) Genetic variance is substantially lower than the deterministic prediction (black dashed line). The wiggly green line gives the predicted V_{LE}^* including the effect of genetic drift: $V_{LE}^* = V_{LE}(1 - 0.392/(N \sigma \sqrt{s}))$ (see Fig. S2). Parameters: $b = 0.3$, $\sigma^2 = 1/2$, $V_s = 1$, $r_m = 1.1$, $\mu = 10^{-7}$. Population is shown after a stable range margin is reached at time = 100 000 generations (with the exception of the initial distribution of trait values, shown in light blue).




Figure S5: Sharp margin to a species' range forms even when allelic effects α_i are non-uniform. With exponentially distributed allelic effects α_i , the expansion slows down after 40 000 generations (see Fig. S6), at the threshold predicted by mean selection per locus $B^* = 0.15N\sigma\sqrt{s} = 0.34$ (red dots). As in this example the allelic effects are not bounded, over very long times, as rare alleles with large effect are recruited (see Fig. S7), the species' range slowly stretches behind the threshold B^* . Parameters and subplot descriptions are the same as in Fig. 3 – with the exception that the allelic effects are exponentially distributed, with mean $\alpha = \sqrt{1/100}$. (C) Note that when allelic effects vary across loci, rarely a cline may establish in a reverse direction, correcting a substitution with a large effect on the trait mean. (D) As genetic variance V_G increases towards the margins, it evolves to match the ever steeper environmental gradient: $V_G = b\sigma\sqrt{V_s}$, prediction shown in dashed line. Gradient in the central habitat is $b = 0.12$; $\sigma^2 = 1/2$, $V_s = 1/2$, $r_m = 1.06$, $K = 50$, $\mu = 2 \cdot 10^{-7}$, time = 100 000 generations (with the exception of the initial distribution of trait values, shown in light blue).




Figure S6: **Range expansion slows down near the threshold based on mean selection coefficient even when the allelic effects α_i are non-uniform (black).** Yet, over very long times, further alleles with large effects can be recruited as they are under stronger selection, and species' range expands a little further. (Example from Fig. S5.) The extent of range expansion is only fully bounded by the substitution with the largest selection coefficient that can arise over a given time. For comparison, the rate of range expansion with equal allelic coefficients with $\alpha_i = \bar{\alpha}$ is given in gray (keeping all other parameters same, example from Fig. 3).




Figure S7: **Over time, genetic drift degrades clines with small allelic effects α .** As more alleles with larger effect α contribute to adaptation, clines become narrower and under stronger selection. Parameters as in Fig. S5. Note that $\bar{s} \equiv \frac{\bar{\alpha}^2}{2V_s}$ and in this example, $s = \alpha$.




Figure S8: **Alternatively, the threshold for range collapse can be expressed as $b/\alpha \gtrsim 0.15 N r^*$.** The threshold holds well unless spacing between the clines, α/b , is smaller than about $1/10$ - this reflects the limits of our simulation system rather than a biological boundary – the deme-spacing is fixed to $\Delta x \equiv 1$. Data and their depiction are the same as in Fig. 2.




Figure S9: Selection on minute deviations from trait mean steepens the clines. Even for negligible genetic drift, clines get steeper than predicted for “perfect” adaptation as the effective environmental gradient B increases. Predicted clines are given by dashed curves in (B, D). The blue and purple curves in (A, C) show the components of selection for a single central cline: $s p q (p - q)$ (blue) vs. $-2 s p q \Delta$ (purple), where $\Delta = (\bar{z} - bx)/\alpha$ gives a scaled deviation from the optimum. The “perfect” adaptation assumes the trait mean matches the optimum everywhere ($\Delta = 0$) - when the predicted and realised clines ought to match (for weak drift). As the effective environmental gradient B steepens, the effects of small deviations of the trait mean from the optimum increasingly affect the clines. (A, B) Example with weak environmental gradient $B = 0.035$, where the cline widths are only 4% narrower than predicted because selection acts mainly (77%) via the diversifying term $s p q (p - q)$. (C, D) Example with very steep effective environmental gradient $B = 1.2$, where the cline width is nearly twice as steep as predicted - correspondingly, the two components of selection are nearly the same: mean integrated selection components across all clines are 0.007 and 0.006, respectively. Both examples are chosen with very weak drift: $N\sigma\sqrt{s} = 23.7$ and $N\sigma\sqrt{s} = 22$ and very similar selection per locus $s = 1/80$ and $s = 1/60$.




Figure S10: **Adaptation may suddenly fail if dispersal is too large.** The threshold for collapse of adaptation (dashed line, $B \sim 0.15 N\sigma\sqrt{s}$) is weakly dependent on dispersal: to the first order, the effect cancels. Yet, for our model, the strength of density dependence r^* decreases with genetic variance ($r^* = V_G/(2V_s)$), which in turn increases with σ ; ($r^* \rightarrow b\sigma/(2\sqrt{V_s})$). When r^{*2}/r_m becomes smaller than 1, further increase in dispersal is detrimental as it brings the population closer to the predicted threshold. This is because both $B = \frac{b\sigma}{\sqrt{2V_s}r^*}$ and $N\sigma\sqrt{s} = Kr^*/r_m \sigma\sqrt{s}$ are both dependent on r^* – and the effects multiply. The scale for the colouring is adjusted (different to Fig. 2) so that the differences in the rate of range expansion (light to dark blue) and contraction (orange to red) are visible; gray dots again denote population that expanded less than one deme over 5000 generations. Parameters: $\sigma = [0.1, 4.24]$, $b = 0.45$, $V_s = 1$, $r^* \equiv 1$ hence $r_m = [1.02, 1.95]$; $K = 28$, $\mu = 4 \cdot 10^{-7}$, $\alpha = \sqrt{1/35}$, time = 5000 generations.




Figure S11: Effect of mutation rate on the rate of range expansion and local genetic variance. (A) Rate of range expansion increases about linearly with mutation rate μ per locus and generation ($5 \cdot 10^{-8} < \mu < 10^{-3}$). When mutation becomes too high, then the rate of expansion first decelerates, and for yet higher mutation rates ($\mu \gtrsim 10^{-2}$), the population starts to collapse. Note that in general, mutation rate per locus and generation is expected to be lower than about 10^{-4} (see Supplementary Text 3). (B) Local genetic variance V_G increases with mutation rate. For low to moderate mutation rates, genetic variance is maintained by mixing across the phenotypic gradient, $V_G = b\sigma\sqrt{V_s}$. The dashed curve gives the prediction $b\sigma\sqrt{V_s} + 2\mu n_l V_s$, which assumes the components of genetic variance due to gene flow (first term) and mutation-selection balance (second term) combine additively, whilst all n_l loci are at linkage equilibrium: the mismatch between the dashed curve and the realised genetic variance implies that with clinal variation, $\mu \ll s$ is not a sufficient condition for the contribution of mutation to be negligible. The top dashed line gives the maximum variance possible, $1/4 n_l \alpha^2$ (where α is a phenotypic effect of a single substitution). With increasing variance, population density drops steadily (not shown): eventually, species' range starts to contract and population collapses. Parameters: $b = 0.4$, $\sigma^2 = 1/2$, $V_s = 1/2$, $r_m = 1.06$, $K = 50$, $\alpha = s = 0.01$, 800 genes. Initial population spans over 100 demes, population evolves over 5000 generations. For both plots, error bars give the standard deviations.

REFERENCES: SUPPLEMENTARY FIGURE LEGENDS

- [50] Haldane, J. The theory of a cline. *Journal of Genetics* **48**, 277–284 (1948).
- [51] Slatkin, M. Gene flow and selection in a cline. *Genetics* **75**, 733–756 (1973).
- [52] Hallatschek, O. & Korolev, K. Fisher waves in the strong noise limit. *Physical Review Letters* **103**, 108103–1–4 (2009).
- [53] Barton, N., Depaulis, F. & Etheridge, A. Neutral evolution in spatially continuous populations. *Theoretical Population Biology* **61**, 31–48 (2002).
- [54] Doebeli, M. & Dieckmann, U. Speciation along environmental gradients. *Nature* **421**, 259–264 (2003).