The effect of carbon subsidies on marine planktonic niche partitioning and recruitment during biofilm assembly

Charles Pepe-Ranney [1]**, Edward Hall** [2,*]

[1] *Cornell University, Department of Crop and Soil Sciences, Ithaca, NY, USA*
[2] *Colorado State University, Natural Resource and Ecology Laboratory, Fort Collins, CO, USA*

Correspondence*:
Edward Hall
Colorado State University, Natural Resource and Ecology Laboratory, Fort Collins, CO 80523-1499, USA, ed.hall@colostate.edu

2 ABSTRACT

The influence of resource availability on planktonic and biofilm microbial community
membership is poorly understood. Heterotrophic bacteria derive some to all of their organic
carbon (C) from photoautotrophs while simultaneously competing with photoautotrophs for
inorganic nutrients such as phosphorus (P) or nitrogen (N). Therefore, C inputs have the
potential to shift the competitive balance of aquatic microbial communities by increasing the
resource space available to heterotrophs (more C) while decreasing the resource space
available to photoautotrophs (less mineral nutrients due to increased competition from
heterotrophs). To test how resource dynamics affect membership of planktonic communities
and assembly of biofilm communities we amended a series of flow-through mesocosms with C
to alter the availability of C among treatments. Each mesocosm was fed with unfiltered seawater
and incubated with sterilized microscope slides as surfaces for biofilm formation. The highest C
treatment had the highest planktonic heterotroph abundance, lowest planktonic photoautotroph
abundance, and highest biofilm biomass. We surveyed bacterial 16S rRNA genes and
plastid 23S rRNA genes to characterize biofilm and planktonic community membership and
structure. Regardless of resource additions, biofilm communities had higher alpha diversity
than planktonic communities in all mesocosms. Heterotrophic plankton communities were
distinct from heterotrophic biofilm communities in all but the highest C treatment where
heterotrophic plankton and biofilm communities resembled each other after 17 days. Unlike
the heterotrophs, photoautotrophic plankton communities were different than photoautotrophic
biofilm communities in composition in all treatments including the highest C treatment.
Our results suggest that although resource amendments affect community membership and
structure, microbial lifestyle (biofilm versus planktonic) has a stronger influence on community
composition.

Keywords: microbial ecology, 16S, 23S, planktonic, biofilm, carbon subsidies, resource stoichiometry

1 INTRODUCTION

Biofilms are diverse and complex microbial consortia, and, the biofilm lifestyle is the rule rather than the
exception for microbes in many environments. Large and small-scale biofilm architectural features play
an important role in their ecology and influence their role in biogeochemical cycles (**Battin et al.**, 2007).

Fluid mechanics impact biofilm structure and assembly (**Hodl et al.**, 2011; **Besemer et al.**, 2009; **Battin**
et al., 2003), but it is less clear how other abiotic factors such as resource availability affect biofilm
assembly. Aquatic biofilms initiate with seed propagules from the planktonic community (**Hodl et al.**,
2011; **McDougald et al.**, 2011). Thus, resource amendments that influence planktonic communities may
also influence the recruitment of microbial populations during biofilm community assembly.

In a crude sense, biofilm and planktonic microbial communities divide into two key groups: oxygenic
phototrophs including eukaryotes and cyanobacteria (hereafter "photoautotrophs"), and heterotrophic
bacteria and archaea. This dichotomy, admittedly an abstraction (e.g. non-phototrophs can also be
autotrophs), can be a powerful paradigm for understanding community shifts across ecosystems of varying
trophic state (**Cotner and Biddanda**, 2002). Heterotrophs meet some to all of their organic carbon (C)
requirements from photoautotroph produced C while simultaneously competing with photoautotrophs for
limiting nutrients such as phosphorous (P) (**Bratbak and Thingstad**, 1985). The presence of external C
inputs, such as terrigenous C leaching from the watershed (**Jansson et al.**, 2008; **Karlsson et al.**, 2012)
or C exudates derived from macrophytes (**Stets and Cotner**, 2008a,b), can alleviate heterotroph reliance
on photoautotroph derived C and shift the heterotroph-photoautotroph relationship from commensal and
competitive to strictly competitive (see **Stets and Cotner**, 2008a, Figure 1). Therefore, increased C supply
should increase the resource space available to heterotrophs and increase competition for mineral nutrients
decreasing nutrients available for photoautotrophs (assuming that heterotrophs are superior competitors
for limiting nutrients as has been observed (see **Cotner and Wetzel**, 1992, Figure 1)). These dynamics
should result in the increase in heterotroph biomass relative to the photoautotroph biomass along a
gradient of increasing labile C inputs. We refer to this differential allocation of limiting resources among
components of the microbial community as niche partitioning.

While these gross level dynamics have been discussed conceptually (**Cotner and Biddanda**, 2002) and
to some extent demonstrated empirically (**Stets and Cotner**, 2008a), the effects of biomass dynamics on
photoautotroph and heterotroph membership and structure has not been directly evaluated in plankton or
biofilms. In addition, how changes in planktonic communities propagate to biofilms during community
assembly is not well understood. We designed this study to test if C subsidies shift the biomass balance
between autotrophs and heterotrophs within the biofilm or its seed pool (i.e. the plankton), and, to measure
how changes in biomass pool size alter composition of the plankton and biofilm communities. Specifically,
we amended marine mesocosms with varying levels of labile C input and evaluated differences in
photoautotroph and heterotrophic bacterial biomass in plankton and biofilm samples along the C gradient.
In each treatment we characterized plankton and biofilm community composition by PCR amplifying and
DNA sequencing 16S rRNA genes and plastid 23S rRNA genes.

2   MATERIALS AND METHODS

2.0.1 Experimental Design   Test tube racks were placed in one smaller (185 L, control) and 3 larger
(370 L) flow-through mesocosms. All mesocosms were fed directly with marine water from an inflow
source in Great Bay, Woods Hole, MA, approximately 200 m from the shore. Each mesocosm had
an adjustable flow rate that resulted in a residence time of approximately 12 h. Irregular variation in
inflow rate caused the actual mesocosm flow rate to vary around the targeted flow rate throughout the day.
However, regular monitoring ensured that the entire volume of each system was flushed approximately
two times per day (i.e. maintained a residence time of $\sim$12 h). To provide a surface for biofilm formation,
we attached coverslips to glass slides using nail polish and then attached each slide to the test tube racks
using office-style binder clips. Twice daily 10 mL of 37 mM KPO_4 and 1, 5 and 50 mL of 3.7 M glucose
were added to each of 3 mesocosms to achieve target C:P resource amendments of 10, 100 and 500. The
goal of the resource amendments was to create a gradient of labile carbon. The same amount of P was
added to each treated mesocosm to ensure that response to additions of C were not inhibited by extreme
P limitation. The control mesocosm did not receive any C or P amendments.

2.0.2 DOC and Chlorophyll Measurements   To assess the efficacy of the C additions, we sampled
each mesocosm twice daily during the first week of the experiment to evaluate dissolved organic C
(DOC) content. After the initiation of the experiment we collected plankton on filters regularly to evaluate
planktonic Chl *a* and bacterial abundance. Once it was clear that pool size of each community had been
altered (day 8) we filtered plankton onto 0.2 μm filters and harvested coverslips to assess bacterial and
algal biofilm community composition (16S and plastid 23S rDNA). In addition, all mesocosms were
analyzed for community composition a second time (day 17) to assess how community composition of
both the plankton and biofilm communities had changed over time. Control samples were only analyzed
for community composition on day 17.

Samples for dissolved organic C (DOC) analysis were collected in acid washed 50 mL falcon tubes
after filtration through a 0.2 μm polycarbonate membrane filter (Millipore GTTP GTTP02500, Sigma
Aldrich P9199) attached to a 60 mL syringe. Syringes and filters were first flushed multiple times with
the control sample to prevent leaching of C from the syringe or the filter into the sample. Samples were
then frozen and analyzed for organic C content with a Shimadzu 500 TOC analyzer (**Wetzel and Likens**,
2000). Biomass of all biofilm samples were measured by difference in pre-(without biofilm) and post-
(with biofilm) weighed GF/F filters after oven drying overnight at 60 °C.

For Chl *a* analysis, we collected plankton on GF/F filters (Whatman, Sigma Aldrich Cat. # Z242489) by
filtering between 500 mL and 1L from the water column of each mesocosm for each treatment. For biofilm
samples, all biofilm was gently removed from the complete area of each coverslip (3 coverslips for each
treatment per sampling event) before being placed in a test tube for extraction with 90-95% acetone for
∼32 hours at -20 °C and analyzed immediately using a Turner 10-AU fluorometer (**Wetzel and Likens**,
2000).

Bacterial abundance of the planktonic samples was analyzed using DAPI staining and direct
visualization on a Zeis Axio epifluorescence microscope after the methods of **Porter and Feig** (1980).
Briefly, 1-3 mL of water was filtered from three separate water column samples through a 0.2 μm black
polycarbonate membrane filter and post stained with a combination of DAPI and Citifluor mountant media
(Ted Pella Redding, Ca) to a final concentration of 1 μL mL^{-1}.

2.0.3 DNA extraction   For plankton, cells were collected by filtering between 20 – 30 mL of water
onto a 0.2 μm pore-size polycarbonate filter (Whatman Nucleopore 28417598, Sigma-Aldrich cat#
WHA110656). For biofilm communities, biomass from the entire coverslip area of three separate slides
was collected and combined in an Eppendorf tube by gently scraping the slip surface with an ethanol
rinsed and flamed razor blade. DNA from both the filter and the biofilm was extracted using a Mobio
Power Soil DNA isolation kit (MoBio Cat. # 12888).

2.0.4 PCR   Samples were amplified for 454 sequencing using a forward and reverse fusion primer. The
forward primer was constructed with (5'-3') the Roche A linker, an 8-10 bp barcode, and the forward gene
specific primer sequence. The reverse fusion primer was constructed with (5'-3') a biotin molecule, the
Roche B linker and the reverse gene specific primer sequence. The gene specific primer pair for bacterial
SSU rRNA genes was 27F/519R (**Lane**, 1991). The primer pair p23SrV_f1/p23SrV_r1 was used to target
23S rRNA genes on plastid genomes (**Sherwood and Presting**, 2007). Amplifications were performed in
25 μL reactions with Qiagen HotStar Taq master mix (Qiagen Inc, Valencia, California), 1 μL of each
5 uM primer, and 1 μL of template. Reactions were performed on ABI Veriti thermocyclers (Applied
Biosytems, Carlsbad, California) under the following thermal profile: 95 °C for 5 min, then 35 cycles
of 94°C for 30 sec, 54°C for 40 sec, 72°C for 1 min, followed by one cycle of 72°C for 10 min and
4 °C hold. Amplification products were visualized with eGels (Life Technologies, Grand Island, New
York). Products were then pooled equimolar and each pool was cleaned with Diffinity RapidTip (Diffinity
Genomics, West Henrietta, New York), and size selected using Agencourt AMPure XP (BeckmanCoulter,
Indianapolis, Indiana) following Roche 454 protocols (454 Life Sciences, Branford, Connecticut). Size
selected pools were then quantified and 150 ng of DNA was hybridized to Dynabeads M-270 (Life

Technologies) to create single stranded DNA following Roche 454 protocols (454 Life Sciences). Single
stranded DNA was diluted and used in emPCR reactions, which were performed and subsequently
enriched. Sequencing followed established manufacture protocols (454 Life Sciences).

2.0.5 Sequence quality control The 16S rRNA gene and plastid 23S rRNA gene sequence collections
were demultiplexed and sequences with sample barcodes not matching expected barcodes were discarded.
We used the maximum expected error metric (**Edgar**, 2013) calculated from sequence quality scores to
cull poor quality sequences from the dataset. Specifically, we discarded any sequence with a maximum
expected error count greater than 1 after truncating the sequence to 175 nt. The forward primer and
barcode was trimmed from the remaining reads. We checked that all primer trimmed, error screened,
and truncated sequences were derived from the targeted region of the LSU or SSU rRNA gene (23S
and 16S sequences, respectively) by aligning the reads to Silva LSU or SSU rRNA gene alignment
("Ref" collection, release 115) with the Mothur (**Schloss et al.**, 2009) NAST-algorithm (**DeSantis et al.**,
2006) aligner and inspecting the alignment coordinates. Reads falling outside the expected alignment
coordinates were culled from the dataset. Remaining reads were trimmed to consistent alignment
coordinates such that all reads began and ended at the same position in the SSU rRNA gene and
screened for chimeras with UChime in "denovo" mode (**Edgar et al.**, 2011) via the Mothur UChime
wrapper. 19,978 of 56,322 16S rRNA gene sequencing reads and 44,719 or 78,695 plastid 23S rRNA
gene sequencing reads passed quality control.

2.0.6 Taxonomic annotations Sequences were taxonomically classified using the UClust (**Edgar**,
2010) based classifier in the QIIME package (**Caporaso et al.**, 2010) with the Greengenes database and
taxonomic nomenclature (version "gg_13_5" provided by QIIME developers, 97% OTU representative
sequences and corresponding taxonomic annotations, (**McDonald et al.**, 2012)) for 16S reads or the Silva
LSU database ("Ref" set, version 115, EMBL taxonomic annotations, (**Quast et al.**, 2013)) for the 23S
reads as reference. We used the default parameters for the algorithm (i.e. minimum consensus of 51% at
any rank, minimum sequence identity for hits at 90% and the maximum accepted hits value was set to 3).

2.0.7 Sequence clustering Reads were clustered into OTUs following the UParse pipeline. Specifically
USearch (version 7.0.1001) was used to establish cluster centroids at a 97% sequence identity level from
the quality controlled data and to map quality controlled reads to the centroids. The initial centroid
establishment algorithm incorporates a quality control step wherein potentially chimeric reads are not
allowed to become cluster seeds. Additionally, we discarded singleton reads. Eighty-eight and 98% of
quality controlled reads could be mapped back to our cluster seeds at a 97% identity cutoff for the 16S
and 23S sequences, respectively.

2.0.8 Alpha and Beta diversity analyses Alpha diversity calculations were made using PyCogent
Python bioinformatics modules (**Knight et al.**, 2007). Rarefaction curves show average OTU counts
from 25 re-samplings at intervals of 10 sequences for each sample. Beta diversity analyses were made
using Phyloseq (**McMurdie and Holmes**, 2014) and its dependencies (**Oksanen et al.**, 2013). A sparsity
threshold of 25% was used for ordination of both plastid 23S and bacterial 16S libraries. Additionally, we
discarded any OTUs from the plastid 23S rRNA gene data that could not be annotated as belonging in the
Eukaryota or cyanobacteria for differential abundance, ordination and Adonis analyses. Cyanobacterial
DNA sequences were removed from 16S rRNA gene sequence collections for ordination, Adonis and
differential abundance analyses. We operated under the assumption that non-cyanobacterial bacteria
are predominantly heterotrophs in our mesocosm setup and refer to non-cyanobacterial bacteria as
"heterotrophs" in the manuscript (this abstraction is useful however there are likely exceptions – i.e.
autotrophs among the non-cyanobacterial bacteria). All DNA sequence based results were visualized
using GGPlot2 (**Wickham**, 2009). Adonis tests and principal coordinate ordinations were performed using
the Bray-Curtis similarity measure for pairwise library comparisons. Adonis tests employed the default
value for number of permutations (999) ("adonis" function in Vegan R package, **Oksanen et al.** (2013)).

Principal coordinates of OTUs were found by averaging site principal coordinate values for each OTU
with OTU relative abundance values (within sites) as weights. The principal coordinate OTU weighted
averages were then expanded to match the site-wise variances (**Oksanen et al.**, 2013).

2.0.9   Identifying enriched OTUs   We used an RNA-Seq differential expression statistical framework to
find OTUs enriched in the given sample classes (R package DESeq2 developed by **Love et al.** (2014))
(for review of RNA-Seq differential expression statistics applied to microbiome OTU count data see
McMurdie and Holmes (2014)). We use the term differential abundance coined by **McMurdie and**
Holmes (2014) to denote OTUs that have different relative abundance across sample classes. We were
particularly interested in two sample classes: 1) lifestyle (biofilm or planktonic) and, 2) high C (C:P
= 500) versus not high C (C:P = 10, C:P = 100 and C:P = control). A differentially abundant OTU is
enriched on one side of a sample class (e.g. enriched in the biofilm versus the plankton). Differential
abundance could mark an enrichment of the OTU towards either side of the sample class and the direction
of the enrichment is apparent in the sign (positive or negative) of the enrichment. Differential abundance
was moderated (see **Love et al.** (2014)) such that the fold change OTU of an OTU across two categories
of a sample class can be used to rank the enrichment of OTUs that span a wide range of base abundance.
The DESeq2 RNA-Seq statistical framework has been shown to improve power and specificity when
identifying differentially abundant OTUs across sample classes in microbiome experiments **McMurdie**
and Holmes (2014).

The specific DESeq2 (**Love et al.**, 2014) parameters we used were as follows: All dispersion estimates
from DESeq2 were calculated using a local fit for mean-dispersion. Native DESeq2 independent filtering
was disabled in favor of explicit independent filtering by sparsity. The sparsity thresholds that produced
the maximum number of OTUs with adjusted p-values for differential abundance below a false discovery
rate of 10% were selected. Cook's distance filtering was also disabled when calculating p-values with
DESeq2. We used the Benjamini-Hochberg method to adjust p-values for multiple testing (**Benjamini**
and Hochberg, 1995). Identical DESeq2 methods were used to assess enriched OTUs from relative
abundances grouped into high C (C:P = 500) or low C (C:P < 500 and control) categories.

IPython Notebooks with computational methods used to create all figures and tables as well as taking
raw sequences through quality control preprocessing are provided at the following URL:

`https://github.com/chuckpr/BvP_manuscript_figures.`

Version information for all R libraries is provided at the end of each IPython Notebook.

3   RESULTS

3.1   BULK COMMUNITY CHARACTERISTICS

We first assessed the effect of the resource treatments on the dissolved chemistry and bulk community
characteristics of the plankton and the biofilms. In the control treatment the mean DOC level was 0.12
$+/-$ 0.02 μmoles L^{-1}. The lower C treatment (C:P $=$ 10) was statistically indistinguishable from the
control at 0.10 $+/-$ 0.02 μmoles C L^{-1}. The intermediate treatment (C:P $=$ 100) increased in DOC to
0.70 μmoles C L^{-1} before decreasing to 0.12 μmoles C L^{-1} at the end of the experiment, with a mean of
0.28 $+/-$ 0.16 μmoles C L^{-1} over the course of the experiment. Only the highest C treatment (C:P $=$ 500)
had DOC levels that were significantly higher (2.53 $+/-$ 1.6 μmoles C L^{-1}) than the control treatment,
over the course of the experiment. The high DOC levels in the highest C treatment were consistent with
C being supplied in excess of the metabolic requirements of the community (i.e. C saturation), but not
higher than what has been observed in coastal marine ecosystems.

This increase in DOC in the higher C treatments was associated with decreases in planktonic Chl *a* in
each treatment (Figure 2A), however there was no significant difference in biofilm Chl *a* among treatments

(Figure 2B). In combination with the decrease in planktonic Chl *a* on the 6th day of the experiment the
highest C treatment had approximately 4-fold higher planktonic heterotroph abundance than the control
and the 10 μM C treatment (Figure 2D). Similarly, biofilms had significantly higher total biomass in the
high C treatment compared to the other treatments (Figure 2D). Thus the shift in resource C:P altered the
pool size of both the photoautotroph and heterotroph communities. Clear differences in heterotroph and
photoautotroph pool size among treatments allowed us to address how shifts in pool sizes were related to
community membership and structure within and among plankton and biofilm communities.

3.2   PLANKTONIC AND BIOFILM COMMUNITY STRUCTURE

3.2.1   Alpha diversity   We used rarefaction curves to evaluate alpha diversity in all treatments for both
the plankton and the biofilm communities. Rarefaction curves showed heterotroph and photoautotroph
OTU richness was consistently higher in the biofilm compared to the planktonic communities (Figure 3).
For both the photoautotroph and heterotroph sequence datasets the biofilm and plankton communities
had the fewest OTUs in the highest C treatment (C:P = 500) (Figure 3). When planktonic rRNA gene
sequences from all planktonic samples were pooled, individual biofilm heterotroph community richness
still generally exceeded the pooled planktonic heterotroph richness. For photoautotrophs, the individual
biofilm richness was similar to the pooled photoautotroph planktonic richness (Figure 4).

3.2.2   Community membership biofilm versus plankton   Heterotroph community membership between
the plankton and biofilm communities was notably different for all treatments except for the highest C
treatment where the plankton and biofilm communities during the second sampling event (day 17) were
more similar to each other than any other community (Figure 5). Photoautotroph plankton and biofilm
communities were also different in OTU composition, however, the similarity among photoautotroph
plankton and biofilm communities in the highest C treatment was not observed as it was for the heterotroph
communities (Figure 5).

In heterotroph libraries, 19,978 sequences were distributed into 636 OTUs and 58% of quality controlled
sequences fell into the top 25 OTUs in order of decreasing sum of relative abundance across all samples.
In photoautotroph libraries 44,719 23S plastid rRNA gene sequences were distributed into 359 OTUs and
71% of sequences fell into the top 25 OTUs sorted by mean relative abundance across all samples.

To investigate differences in community structure and membership between the heterotroph biofilm
and overlying planktonic communities we identified the most enriched OTUs in biofilm compared to the
planktonic communities and vice versa. The most enriched OTUs were enriched in planktonic samples
(with respect to biofilm) as opposed to biofilm samples (Figure 6). This is consistent with the higher alpha
diversity in biofilm communities compared to planktonic communities and evidence that sequence counts
were spread across a greater diversity of taxa in the biofilm libraries compared to the planktonic libraries
(i.e. biofilm communities had higher evenness than planktonic communities). Of the top 5 enriched
heterotroph OTUs between the two lifestyles (biofilm or plankton), 1 is annotated as *Bacteroidetes*, 1
Gammaproteobacteria, 1 *Betaproteobacteria*, 1 *Alphaproteobacteria* and 1 *Actinobacteria* and all 5 were
enriched in the planktonic libraries relative to biofilm (Table 1). Of the 25 most enriched OTUs among
lifestyles only 2 heterotroph OTU centroid DNA sequences shared high sequence identity ($>= 97\%$) with
cultured isolates ("OTU.32" and "OTU.48", Table 1).

We similarly assessed membership among biofilm and plankton in the photoautotroph communities.
Photoautotroph 23S plastid rRNA gene sequence libraries also clustered strongly by lifestyle (Figure 5).
Biofilm libraries were predominantly enriched in *Stramenopile* OTUs whereas planktonic libraries were
enriched in *Haptophyceae*, *Cryptophyta* and *Viridiplantae* OTUs based on OTU positions in sample
ordination space (Figure 5, see Ordination Methods). When photoautotroph OTUs were ordered by
differential abundance between lifestyles (see Figure 6), 16 of the 25 most enriched OTUs were enriched
in the biofilm and 9 were enriched in the planktonic samples. Fourteen of these 16 biofilm enriched
OTUs were *Stramenopiles* of class *Bacillarophyta*, the remaining OTUs were classified as members of

the *Chlorophyta* and *Dinophyceae*. The 9 planktonic enriched OTUs (above) were distributed into the *Viridiplantae* (5 OTUs), *Cryptophyta* (1 OTUs), *Haptophyceae* (1 OTU), *Stramenopiles* (1 OTU) and cyanobacteria (1 OTU). The 10 most enriched photoautotroph OTUs between lifestyles were evenly split between planktonic and biofilm enriched OTUs. As with the heterotrophs, photoautotroph OTU fold change between lifestyles are qualitatively consistent with OTU positions in sample ordination space (see Figures 6 and 5).

In both the heterotroph and photoautotroph communities low abundance members of the planktonic communities became highly abundant members of the biofilm (Figure 7). The separation in community membership among biofilm and planktonic communities is supported statistically by the Adonis test (**Anderson**, 2001) for both the heterotroph and photoautotroph libraries (p-value 0.003 and 0.002, respectively). The lifestyle category represents 18% and 45% for pairwise sample distance variance in heterotroph and photoautotroph libraries, respectively. The Adonis test result is also consistent with lifestyle (biofilm versus planktonic) clustering along the first principal component for the photoautotroph libraries but not for the heterotroph libraries (Figure 5).

3.2.3 Heterotroph community membership changes with C amendments Although community membership was predominately driven by lifestyle, we also investigated how resource amendments affected community membership and structure. Because the abiotic (e.g. DOC) and all biomass indicators (e.g. heterotroph abundance, Chl *a*) were only significantly different in the highest resource C:P treatment we compared resource C:P = 500 (high C) to all other mesocosms (i.e. control, C:P = 10, and C:P = 100 – low C). The high and low carbon amended mesocosms had statistically different heterotroph communities (Adonis p-value 0.018) but not photoautotroph communities (Adonis p-value 0.59). Nine heterotroph OTUs were enriched in the high C treatment relative to low C. Four of the 9 high C enriched OTUs were annotated as *Alteromonadales*, 3 as *Campylobacterales* and 1 each into *Vibrionales* and *Pseudomonadales*. The most enriched OTU in low C mesocosms was annotated as belonging to the "HTCC2188" candidate order and shared 99% identity with a 16S sequence annotated as "marine gamma proteobacterium HTCC2089" (accession AY386332). We only observed differences at the community level between high and low C amendments in the heterotroph communities and therefore did not assess differential abundance of OTUs between high and low C treatments in photoautotroph communities.

4   DISCUSSION

4.1   BIOMASS POOL SIZE

The goal of this study was to evaluate how changes in available C affected the biomass pool size and composition of planktonic and biofilm communities. Our results suggest that C subsidies increased heterotroph biomass in both plankton and biofilm communities. C amendments also resulted in decreased photoautotroph biomass in the plankton community, but there was no significant change in biofilm photoautotroph biomass between resource treatments. Although the DOC concentration in the highest C treatment was significantly higher than the other treatments, the concentrations we measured were in the range of those reported in natural marine ecosystems (**Mopper et al.**, 1980) and it is has been noted that glucose concentrations in coastal marine ecosystems may fluctuate over several orders of magnitude (**Alonso and Pernthaler**, 2006). The changes in the biomass pool size that did occur were consistent with changing relationships (commensal to competitive) between the autotrophic and heterotrophic components of the plankton communities but not necessarily of the biofilm communities. While we recognize that other mechanisms may drive the shift in biomass pool size of these two components of the microbial community (e.g. increased grazing pressure on the photoautotrophs with C additions, or production of secondary metabolites by the heterotrophs that inhibit algal growth) previous studies **Stets and Cotner** (2008a); **Cotner and Biddanda** (2002) and the data reported here suggest that altered nutrient competition among heterotrophic and photoautotrophic members of the plankton is the most parsimonious explanation for the observed shift in biomass pool size.

4.2   BIOFILM AND PLANKTON ALPHA AND BETA DIVERSITY

Beyond changes in the biomass pool size of each community, we explored how shifts in resource C
affected the membership and structure of each community, and the recruitment of plankton during biofilm
community assembly. Intuitively, shifts in planktonic community composition should alter the available
pool of microorganisms that can be recruited into a biofilm. For example, if planktonic diversity increases,
the number of potential taxa that can be recruited to the biofilm should also increase, potentially increasing
diversity within the biofilm. Similarly, a decrease in mineral nutrients available to photoautotrophs
should decrease photoautotroph pool size, potentially decreasing photoautotroph diversity and therefore
candidate photoautotroph taxa that are available for biofilm formation. In addition, C in excess of resource
requirements may increase the production of extracellular polysaccharides (EPS) by planktonic cells thus
increasing the probability that planktonic cells are incorporated into a biofilm by adhesion. Each of these
mechanisms suggest that an increase in labile C to the system should result in increased alpha diversity
in heterotrophic plankton and heterotrophic biofilm communities while decreasing alpha diversity within
both planktonic and biofilm photoautotroph communities.

We highlight three key results that we find important for understanding aquatic biofilm assembly.
First, biofilm community richness exceeded planktonic community richness (Figure 3) in all mesocosms.
Second, for the control, C:P = 10 and C:P = 100 resource treatments the membership and structure of
the heterotroph biofilm and plankton communities were more similar within a lifestyle (plankton versus
biofilm) than within a resource treatment. However, for the bacteria in the highest C treatment (C:P =
500) both membership and structure of biofilm and planktonic communities at day 17 were more similar
to each other than to communities from other treatments (Figure 5). Third, C subsides acted differently
on the photoautotroph and heterotroph communities. Specifically, while the highest level of C subsides
(C:P = 500) resulted in a merging of membership in the heterotroph plankton and heterotroph biofilm
communities the same merging of membership was not observed for the photoautotroph biofilm and
plankton communities which had distinct membership in all treatments.

We propose two potential mechanisms for the increased richness of the biofilm communities relative
to the planktonic community richness. First, it is possible that the planktonic community composition of
our flow through incubators was dynamic in time. In this case the biofilm community would represent
a temporally integrated sample of the planktonic organisms moving through the reactor resulting in
higher apparent alpha diversity (i.e. mass effects would be the dominant assembly mechanism). Second,
the biofilm environment may disproportionately enrich the low abundance members of the planktonic
community. In this case it is probable that the biofilm would incorporate the most abundant members
from the planktonic community (i.e. mass effects) but also select and enrich (i.e. species sorting) the
least abundant members of the planktonic community resulting in a higher level of detectable alpha
diversity. The second mechanism would result if the biofilm environment represented a more diverse
microhabitat including sharply delineated oxygen, nutrient and pH gradients that are not present in the
planktonic environment. In this case the more diverse microhabitat would be able to support a more diverse
community due to an abundance of additional environmental habitats (i.e. niches).

We evaluated the first mechanism by comparing membership among the plankton samples taken 9
days apart (day 8 and day 17). While heterotrophic plankton communities were not identical between
the time points (Figure 5), communities within a treatment were more similar to each other than other
heterotroph plankton communities regardless of time. In addition, the control and two lowest C treatments
(C:P = 10 and C:P = 100) separated completely from biofilm communities in principle coordinate space
(Bray-Curtis distance metric). This suggests that the biofilm community was not integrating variable
bacterioplankton community membership, but rather was at least in part selecting for a community that
was composed of distinct populations when compared to the most abundant members of the plankton
community. As noted above, in the highest C treatment (C:P = 500) the heterotroph biofilm and plankton
community membership had significant overlap at the final timepoint (Figure 5). However, heterotrophic
plankton community composition for the highest C treatment among timepoints (8 and 17 days) were
also qualitatively as similar to each other as any other community. Thus, variable planktonic community

composition among timepoints would not explain the higher diversity observed in the biofilm compared to the planktonic community. Rather, two results point to enrichment of planktonic community members within the biofilm as the mechanism for higher diversity in the biofilm compared to the plankton. First, the increasing similarity between the plankton and the biofilm communities between each time point in the highest resource C treatment suggests that *in situ* resource conditions were sufficient to alter the relative abundance of the populations within each community. Second, an analysis of the OTU relative abundance in biofilm and planktonic libraries where OTUs are sorted by planktonic sample rank (Figure 7) shows that the least abundant members of the plankton community were routinely highly abundant within the biofilm community. This was true for both photoautotroph and heterotroph communities, at all treatment levels and both timepoints. While we did not (could not) specifically measure niche diversity within the biofilm communities our results suggest that the biofilm habitat selected for unique members of the photoautotroph and heterotrophic community that were in low abundance in the planktonic habitat but readily became major constituents of the biofilm community.

Few studies have simultaneously evaluated the relationship among membership and/or diversity of the plankton and the biofilm community from complex environmental microbial communities. One notable study looked at planktonic community composition and biofilm formation on glass beads placed for three weeks in three boreal freshwater streams (**Besemer et al.**, 2012). While that study system is markedly different than our study, the analyses and questions addressed in each study were sufficiently similar to merit comparison. **Besemer et al.** (2012) concluded that the biofilm community membership was most likely driven by species sorting over mass effects. This is consistent with what we report here. However, in the **Besemer et al.** (2012) study the authors reported that planktonic diversity was significantly higher relative to biofilm diversity (the opposite of what we found in our study). Given the differences in the source of the planktonic community among studies, this result is not surprising. While biofilm communities were established on glass beads in **Besemer et al.** (2012) and glass slides (this study) over a similar time period (21 days, **Besemer et al.** (2012) and 17 days this study) the origin of the planktonic community in each study was different. The **Besemer et al.** (2012) study was conducted in three boreal streams during snow melt when connectivity between the terrestrial and aquatic habitats was high and potentially highly variable depending on how hydrologic pathways differed among precipitation events. In this study the source community was a marine intake located approximately 200 m from the shore during July when communities are more stable over the 17 day period of the incubation. A separate study conducted in alpine and sub-alpine streams clearly showed that stream plankton communities reflected localized precipitation events and could be traced largely to soil source communities from drainages within the watershed (**Portillo et al.**, 2012). While planktonic communities in lake ecosystems can be linked to soil communities in the watershed, as residence time of the system slows the relative influence of species sorting increases. Thus, in headwater ecosystems stream plankton communities can often be composed primarily of soil organisms (**Crump et al.**, 2012). In addition to the diverse source communities the **Besemer et al.** (2012) study sampled the plankton community at multiple timepoints and integrated the samples before sequencing further increasing community richness as compared to the current study where the plankton community was sampled and analyzed only at two independent timepoints. Indeed, when we pool OTU counts from all planktonic libraries and compare the rarefaction curve of the pooled planktonic libraries (photoautotrophs and heterotrophs) against sample-wise biofilm libraries, we found more total heterotroph and photoautotroph planktonic OTUs than in any given single biofilm sample. It appears, however, that sample-wise heterotroph biofilm rarefaction curves may exceed the integrated planktonic curve upon extrapolation and most exceed the integrated planktonic curve at sampling depths where data is present for the biofilm and pooled planktonic library (Figure 4). This result is consistent with our conclusion that temporal heterogeneity in the plankton was not sufficient to explain the higher diversity in the biofilm sample but would explain the relative differences between planktonic and biofilm diversity found in **Besemer et al.** (2012) compared to this study.

In addition, for this study, it is important to note that biofilm community richness peaked at the intermediate treatment (C:P = 100) and appeared to decrease between each time point although with only two time points it was unclear how pronounced the temporal effect was nor is it possible assess the statistical significance of this effect (Figure 3). Since biomass of the plankton and the biofilm

increased with increasing C subsidies the intermediate peak in OTU richness is consistent with a classic
productivity-diversity relationship that has been shown for many ecosystems and communities both
microbial and otherwise. However, as with other experiments our experimental design did not allow
us to tell whether resources drove productivity that subsequently drove changes in diversity or whether
resources drove diversity which altered productivity. Rather, we note that as diversity decreased in the
highest C treatment, heterotrophic plankton and biofilm membership became increasingly similar. This
suggests that environments that contained high amounts of labile C selected for fewer dominant taxa,
overwhelming the lifestyle species sorting mechanisms that appeared to dominate biofilm community
assembly in all other treatments. Similarly, while we did not measure extracellular polymeric substances
(EPS), direct microscopy showed that planktonic cells in the highest C treatment (C:P = 500) were
surrounded by what appeared to be EPS. Because biofilm EPS also appeared to increase from the low
to high C treatments it is possible that abundant planktonic cells were more readily incorporated into
biofilms due both to increased "stickiness" of the planktonic cells as well as the biofilm itself. While we
did not observe flocculating DOC which has been shown to dominate high DOC environments in nature,
we did measure a substantial increase in DOC in the C:P = 500 treatment which was more than 2-fold
higher than any of the other treatments. Thus additional adhesion of the plankton and the biofilm may also
explain the merging of the planktonic and biofilm heterotroph membership in the highest C treatment.

4.3 LIFESTYLE (BIOFILM OR PLANKTONIC) ENRICHED OTUS

There are only a few studies that attempt to compare biofilm community composition and the overlying
planktonic community (**Besemer et al.**, 2007, 2012; **Jackson et al.**, 2001; **Lyautey et al.**, 2005). Those
studies illustrate community composition among the two habitats are unique, with few taxa found in
both. This is consistent with our findings in this experimental system with a natural marine planktonic
source community. In addition, our study also evaluated photoautotroph community composition which
showed a similar result suggesting that both the photoautotroph and heterotroph biofilm communities
are comprised of phylogenetically distinct organisms that exist in low abundance in the surrounding
habitat (i.e. the plankton) but are readily enriched in the biofilm lifestyle. Most of the biofilm enriched
photoautotroph OTUs were *Bacillariophyta* although there were also many *Bacillariophya* OTUs enriched
in the planktonic libraries. We also found more *Cryptophyta* and *Viridiplantae* were enriched in the
planktonic photoautotroph libraries. It appears that these broad taxonomic groups were selected against in
biofilms under our experimental conditions. Heterotroph OTUs enriched in planktonic samples displayed
more dramatic differential abundance patterns than heterotroph OTUs enriched in biofilm samples, but,
biofilm enriched heterotroph OTUs were spread across a greater phylogenetic breadth (Figure 6). This
is also consistent with the idea of greater niche diversity in the biofilm environment as opposed to the
plankton. Greater niche diversity should select for a more diverse set of taxa but individual taxa would not
be as numerically dominant in a more uniform environment like the planktonic environment. At the Order
level, enriched heterotroph OTUs tended to have members that were enriched in both the plankton and the
biofilm suggesting the phylogenetic coherence of lifestyle is not captured at the level of Order. It should be
noted, however, that taxonomic annotations in reference databases and therefore environmental sequence
collections show little equivalency in phylogenetic breadth between groups at the same taxonomic rank
(**Schloss and Westcott**, 2011). Unfortunately, at higher taxonomic resolution (e.g. Genus-level), groups
did not possess a sufficient numbers of OTUs to evaluate coherence between taxonomic annotation and
lifestyle. Carbon amendments did not affect photoautotroph library membership and structure to the same
degree as it affected heterotroph library composition. As expected, heterotroph OTUs enriched in the high
C amended mesocosm (C:P = 500) include OTUs in classic copiotroph families such as *Altermonodales*
and *Pseudomonadaceae*. Interestingly, the most depleted OTU in the high C treatments is annotated
as being in the HTCC2188 order of the *Gammaproteobacteria* and shares 99% sequence identity with
another "HTCC" strain (accession AY386332). HTCC stands for 'high throughput culture collection' and
is a prefix for strains cultured under low nutrient conditions (**Cho and Giovannoni**, 2004; **Connon and**
Giovannoni, 2002).

4.4 CONCLUSION

In summary this study shows that changes in low resolution community level dynamics are concurrent
with changes in the underlying constituent populations that compose them. We found that autotrophic
pools and heterotrophic pools responded differently to amendments of labile C as hypothesized. Notably
while C amendments altered both pool size and membership of the heterotroph communities we did not
see similar dynamics within the photoautotroph communities. Planktonic photoautotrophs decreased in
response to C amendments presumably in response to increased competition for mineral nutrients from
a larger heterotroph community, however there was not a similar decrease in biofilm photoautotroph
community. In addition membership of the photoautotroph communities between the plankton and biofilm
lifestyles did not become more similar in the photoautotrophs as it did for the bacterial heterotrophs
in the highest C treatment. Consistent with a growing body of work our results suggest that complex
environmental biofilms are a unique microbial community that form from taxa that are found in low
abundance in the neighboring communities. This membership was affected by C amendments for
heterotrophic but not photoautotrophic microbes and then only in the most extreme resource environment.
This suggests that lifestyle is a major division among environmental microorganisms and although biofilm
forming microbes must travel in planktonic form at some point, reproductive success and metabolic
contributions to biogeochemical processes comes from those taxa primarily if not exclusively while they
are part of a biofilm. Our results point to lifestyle (planktonic or biofilm) as an important trait that explains
a portion of the exceptional diversity found in snapshots used to characterize environmental microbial
communities in space and time.

5 ACKNOWLEDGEMENTS

472 This research was conducted as part of the 2010 Marine Biological Laboratory's Microbial Diversity
473 Course. Funding was supplied by the MBL, the Bernard Davis Endowed Scholarship Fund, and the
474 Selman A. Waksman Endowed Scholarship. We would like to thank Dan Buckley and Steve Zinder for
475 organizing the course, Marshall Otter, Mathew Erickson, Hugh Ducklow and Jay T. Lennon for analytical
476 support and the Austrian FWF MICDIF award to Tom Battin for salary support.

REFERENCES

Alonso, C. and Pernthaler, J. (2006), Roseobacter and SAR11 dominate microbial glucose uptake in
coastal north sea waters, *Environ. Microbiol.*, 8, 11, 2022–2030, doi:10.1111/j.1462-2920.2006.01082.
x
Anderson, M. J. (2001), A new method for non-parametric multivariate analysis of variance, *Austral
Ecology*, 26, 1, 32–46, doi:10.1111/j.1442-9993.2001.01070.pp.x
Battin, T. J., Kaplan, L. A., Denis, N. J., and Hansen, C. M. (2003), Contributions of microbial biofilms
to ecosystem processes in stream mesocosms., *Nature*, 426, 439–42
Battin, T. J., Sloan, W. T., Kjelleberg, S., Daims, H., Head, I. M., Curtis, T. P., et al. (2007), Microbial
landscapes: new paths to biofilm research., *Nat. Rev. Microbiol.*, 5, 76–81
Benjamini, Y. and Hochberg, Y. (1995), r, *J. R. Stat. Soc.*, 57, 1, 289–300, doi:10.2307/2346101
Besemer, K., Hödl, I., Singer, G., and Battin, T. J. (2009), Architectural differentiation reflects bacterial
community structure in stream biofilms., *ISME J*, 3, 1318–24
Besemer, K., Peter, H., Logue, J. B., Langenheder, S., Lindstrm, E. S., Tranvik, L. J., et al. (2012),
Unraveling assembly of stream biofilm communities., *ISME J*, 6, 1459–68
Besemer, K., Singer, G., Limberger, R., Chlup, A. K., Hochedlinger, G., Hödl, I., et al. (2007),
Biophysical controls on community succession in stream biofilms, *Appl. Environ. Microbiol.*, 73, 15,
4966–4974, doi:10.1128/aem.00588-07

Bratbak, G. and Thingstad, T. F. (1985), Phytoplankton-bacteria interactions: an apparent paradox? Analysis of a model system with both competition and commensalism, *Marine Ecology - Progress Series*, 25, 23–30

Caporaso, J. G., Kuczynski, J., Stombaugh, J., K., B., Bushman, F. D., Costello, E. K., et al. (2010), QIIME allows analysis of high-throughput community sequencing data., *Nat. Methods*, 7, 335–6

Cho, J. C. and Giovannoni, S. J. (2004), Cultivation and growth characteristics of a diverse group of oligotrophic marine gammaproteobacteria, *Appl. Environ. Microbiol.*, 70, 1, 432–440, doi:10.1128/aem.70.1.432-440.2004

Connon, S. A. and Giovannoni, S. J. (2002), High-throughput methods for culturing microorganisms in very-low-nutrient media yield diverse new marine isolates, *Appl. Environ. Microbiol.*, 68, 8, 3878–3885, doi:10.1128/aem.68.8.3878-3885.2002

Cotner, J. B. and Biddanda, B. A. (2002), Small players large role: microbial influence on biogeochemical processes in pelagic aquatic ecosystems, *Ecosystems*, 5, 2, 105–121, doi:10.1007/s10021-001-0059-3

Cotner, J. B. and Wetzel, R. G. (1992), Uptake of dissolved inorganic and organic bphosphorus compounds by phytoplankton and bacterioplankton, *Limnol. Oceangr.*, 37, 2, 232–243, doi:10.4319/lo.1992.37.2.0232

Crump, B. C., Amaral-Zettler, L. A., and Kling, G. W. (2012), Microbial diversity in arctic freshwaters is structured by inoculation of microbes from soils., *ISME J*, 6, 1629–39

DeSantis, T. Z. J., Hugenholtz, P., Keller, K., L., B. E., Larsen, N., Piceno, Y. M., et al. (2006), NAST: a multiple sequence alignment server for comparative analysis of 16S rRNA genes., *Nucleic Acids Res.*, 34, W394–9

Edgar, R. C. (2010), Search and clustering orders of magnitude faster than BLAST., *Bioinformatics*, 26, 2460–1

Edgar, R. C. (2013), UPARSE: highly accurate OTU sequences from microbial amplicon reads., *Nat. Methods*, 10, 996–8

Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., and Knight, R. (2011), UCHIME improves sensitivity and speed of chimera detection., *Bioinformatics*, 27, 2194–200

Hodl, I., Hodl, J., Worman, A., Singer, G., K., B., and Battin, T. J. (2011), Voronoi tessellation captures very early clustering of single primary cells as induced by interactions in nascent biofilms, *PLoS ONE*, 6, 10, e26368, doi:10.1371/journal.pone.0026368

Jackson, C. R., Churchill, P. F., and Roden, E. E. (2001), Successional changes in bacterial assemblage structure during epilithic biofilm development, *Ecology*, 82, 2, 555–566, doi:10.1890/0012-9658(2001)082[0555:scibas]2.0.co;2

Jansson, M., Hickler, T., Jonsson, A., and Karlsson, J. (2008), Links between terrestrial primary production and bacterial production and respiration in lakes in a climate gradient in subarctic sweden, *Ecosystems*, 11, 3, 367–376, doi:10.1007/s10021-008-9127-2

Karlsson, J., Berggren, M., Ask, J., Byström, P., A., J., Laudon, H., et al. (2012), Terrestrial organic matter support of lake food webs: Evidence from lake metabolism and stable hydrogen isotopes of consumers, *Limnol. Oceangr.*, 57, 4, 1042–1048, doi:10.4319/lo.2012.57.4.1042

Knight, R., Maxwell, P., Birmingham, A., Carnes, J., Caporaso, J. G., Easton, B. C., et al. (2007), PyCogent: a toolkit for making sense from sequence., *Genome Biol.*, 8, R171

Lane, D. J. (1991), 16S/23S rRNA sequencing, *Nucleic Acid Techniques in Bacterial Systematics*, 125–175

Love, M. I., Huber, W., and Anders, S. (2014), Moderated estimation of fold change and dispersion for RNA-Seq data with DESeq2, doi:10.1101/002832

Lyautey, E., Jackson, C. R., Cayrou, J., Rols, J., and F., G. (2005), Bacterial community succession in natural river biofilm assemblages, *Microb. Ecol.*, 50, 4, 589–601, doi:10.1007/s00248-005-5032-9

McDonald, D., Price, M. N., Goodrich, J., Nawrocki, E. P., DeSantis, T. Z., Probst, A., et al. (2012), An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea., *ISME J*, 6, 610–8

McDougald, D., Rice, S. A., Barraud, N., Steinberg, P. D., and Kjelleberg, S. (2011), Should we stay or should we go: mechanisms and ecological consequences for biofilm dispersal., *Nat. Rev. Microbiol.*, 10, 39–50

McMurdie, P. J. and Holmes, S. (2014), Waste not, want not: why rarefying microbiome data is
inadmissible., *PLoS Comput. Biol.*, 10, e1003531
Mopper, K., Dawson, R., Liebezeit, G., and Ittekkot, V. (1980), The monosaccharide spectra of natural
waters, *Marine Chemistry*, 10, 1, 55–66, doi:10.1016/0304-4203(80)90058-4
Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., et al. (2013), vegan:
Community Ecology Package, R package version 2.0-10
Porter, K. G. and Feig, Y. S. (1980), The use of DAPI for identifying and counting aquatic microflora,
Limnol. Oceangr., 25, 5, 943–948, doi:10.4319/lo.1980.25.5.0943
Portillo, M. C., Anderson, S. P., and Fierer, N. (2012), Temporal variability in the diversity and
composition of stream bacterioplankton communities., *Environ. Microbiol.*, 14, 2417–28
Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., T., S., Yarza, P., et al. (2013), The SILVA ribosomal RNA
gene database project: improved data processing and web-based tools., *Nucleic Acids Res.*, 41, D590–6
Schloss, P. D. and Westcott, S. L. (2011), Assessing and improving methods used in operational taxonomic
unit-based approaches for 16s rrna gene sequence analysis, *Appl. Environ. Microbiol.*, 77, 10, 3219–
3226, doi:10.1128/aem.02810-10
Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., et al. (2009),
Introducing mothur: open-source, platform-independent, community-supported software for describing
and comparing microbial communities., *Appl. Environ. Microbiol.*, 75, 7537–41
Sherwood, A. R. and Presting, G. G. (2007), Universal primers amplify a 23s rdna plastid marker in
eukaryotic algae and cyanobacteria, *Journal of Phycology*, 43, 3, 605–608, doi:10.1111/j.1529-8817.
2007.00341.x
Stets, E. G. and Cotner, J. B. (2008a), Littoral zones as sources of biodegradable dissolved organic carbon
in lakes, *Can. J. Fish. Aquat. Sci.*, 65, 11, 2454–2460, doi:10.1139/f08-142
Stets, E. G. and Cotner, J. B. (2008b), The influence of dissolved organic carbon on bacterial phosphorus
uptake and bacteria-phytoplankton dynamics in two Minnesota lakes, *Limnol. Oceangr.*, 53, 1, 137–
147, doi:10.4319/lo.2008.53.1.0137
Wetzel, R. G. and Likens, G. E. (2000), Limnological Analyses (Springer New York), doi:10.1007/
978-1-4757-3250-4
Wickham, H. (2009), ggplot2 (Springer New York), doi:10.1007/978-0-387-98141-3

FIGURES

Table 1. Results for BLAST search against Living Tree Project (top 25 lifestyle enriched bacterial OTUs (Operational Taxonomic Unit)

OTU	Phylum	$log_2(plankton : biofilm)$	Species Name	%Identity	Accession
OTU.103	Bacteroidetes	7.78	*Zunongwangia profunda*	89.66	DQ855467
OTU.105	Proteobacteria	8.09	*Microbulbifer yueqingensis*	90.14	GQ262813
OTU.11	Proteobacteria	9.59	*Methylobacillus glycogenes*	93.96	FR733701
OTU.123	Proteobacteria	8.96	*Flexibacter roseolus*	83.46	AB078062
			Flexibacter elegans	83.46	AB078048
OTU.165	Proteobacteria	-7.05	*Kangiella spongicola*	92.05	GU339304
			Kangiella marina	92.05	JN559388
OTU.166	Proteobacteria	-7.52	*Halomonas halocynthiae*	92.62	AJ417388
OTU.19	Proteobacteria	9.31	*Neptuniibacter caesariensis*	90.07	AY136116
OTU.195	Proteobacteria	7.17	*Methylobacillus glycogenes*	94.63	FR733701
OTU.20	Proteobacteria	9.07	*Ruegeria halocynthiae*	96.15	HQ852038
			Phaeobacter daeponensis	95.49	DQ981486
OTU.207	Proteobacteria	9.30	*Methylobacillus glycogenes*	91.28	FR733701
OTU.223	Proteobacteria	7.94	*Methyloferula stellata*	87.02	FR686343
			Methylocapsa aurea	87.02	FN433469
			Beijerinckia indica subsp. lacticogenes	87.02	AJ563931
			Beijerinckia indica subsp. indica	87.02	CP001016
			Beijerinckia derxii subsp. venezuelae	87.02	AJ563934
OTU.26	Actinobacteria	8.58	*Corallomonas stylophorae*	88.17	GU569894
OTU.31	Bacteroidetes	9.63	*Sediminitomix flava*	91.33	AB255370
			Kordia algicida	91.33	AY195836
OTU.32	Bacteroidetes	8.90	*Bizionia echini*	97.32	FJ716799
OTU.36	Actinobacteria	9.55	*Pseudoclavibacter soli*	95.95	AB329630
OTU.369	Actinobacteria	7.93	*Agrococcus terreus*	96.0	FJ423764
OTU.40	Bacteroidetes	7.68	*Aureitalea marina*	91.33	AB602429
OTU.44	Proteobacteria	8.92	*Glaciecola mesophila*	92.62	AJ488501
			Aestuariibacter salexigens	92.67	AY207502
			Aestuariibacter halophilus	92.67	AY207503
OTU.48	Actinobacteria	7.28	*Microterricola viridarii*	97.33	AB282862
			Leifsonia pindariensis	97.33	AM900767
OTU.62	Proteobacteria	8.75	*Haliea rubra*	91.55	EU161717
			Congregibacter litoralis	91.55	AAOA01000004
			Chromatocurvus halotolerans	91.55	AM691086
OTU.69	Proteobacteria	9.34	*Sneathiella glossodoripedis*	87.94	AB289439
OTU.71	Bacteroidetes	7.40	*Aequorivita sublithincola*	95.77	AF170749
OTU.83	Actinobacteria	7.27	*Microbacterium invictum*	92.47	AM949677
OTU.84	Proteobacteria	-7.44	*Alcanivorax dieselolei*	93.29	AY683537
			Alcanivorax balearicus	93.29	AY686709
OTU.89	Actinobacteria	7.17	*Corallomonas stylophorae*	87.91	GU569894

Figure 1. Carbon subsidies in the form of glucose alleviate the dependence of heterotrophic bacteria on photoautotroph derived C exudates. This should result in an increase in resource space and biomass for heterotrophs and a decrease in resource space and biomass for photoautotrophs due to increased competition for mineral nutrients (for simplicity we illustrate competition for P but this is equally applicable other elements that may limit primary production). We hypothesized that this predicted change in biomass pool size of these two groups will result in changes in the plankton community composition of both groups that will propagate to to the composition of biofilm communities for both groups. We refer to shifts in the demand and availability of resources among components of the microbial community as 'partitioning. Blue rods represent heterotrophs, green stars represent photoautotrophs, brown diamonds represent EPS or other cohesive components of a biofilm.

Figure 2. Increased C amendments diminished A) planktonic photoautotroph biomass (estimated as Chl *a*) but B) not biofilm photoautotroph biomass. In contrast, both C) biofilm total biomass and D) number of planktonic bacterial cells increased with increasing C subsidies. Only the highest C treatment produced biomass that was significantly greater than ($p < 0.05$) the other treatments (significant differences among treatments are denoted by different letters). The bacterial abundance sample for the C:P = 100 treatment was lost before analysis and is therefore not reported in panel D.

Figure 3. Rarefaction curves for all biofilm versus plankton libraries. Each panel represents a single C:P treatment and time point. Richness is greater for all biofilm communities when compared to corresponding planktonic communities. Gray ribbons are 99% confidence intervals around each rarefaction point based on variance from 25 re-samplings.

Figure 4. Rarefaction plots for all samples. Planktonic libraries have been integrated such that the count for each OTU is the sum of counts across all samples. Gray ribbons are 99% confidence intervals around each rarefaction point based on variance from 25 re-samplings.

Figure 5. Principal coordinates ordination of bray-curtis distances for 23S rRNA plastid libraries and 16S rRNA gene libraries. OTU points are weighted principal coordinate averages (weights are relative abundance values in each sample) and the variance along each principal axis is expanded to match the site variance. Point annotations denote the amended C:P ratio for the mesocosm from which each sample was derived.

Figure 6. log_2 of lifestyle OTU abundance fold change between biofilm and plankton communities. Each point represents one OTU and points are grouped along the x-axis by Order. Outlined points have adjusted p-values below a false discovery rate of 0.10. Positive fold change values represents enrichment in planktonic samples.

Figure 7. Rank abundance plots. Each panel represents a single time point and C:P. The "rank" of each OTU is based on planktonic sample relative abundance. Each position along the x-axis represents a single OTU. Both the x and y axes are scaled logarithmically.