Speciation in *Heliconius* Butterflies: Minimal Contact Followed by Millions of Generations of Hybridisation

Simon H. Martin*[1], Anders Eriksson*[1,2], Krzysztof M. Kozak[1], Andrea Manica[1] and Chris D. Jiggins[1]

[1] Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, United Kingdom

[2] Integrative Systems Biology Laboratory, King Abdullah University of Science and Technology, Thuwal 23955-6900, Kingdom of Saudi Arabia

* These authors contributed equally

Corresponding Author: S.H. Martin, shm45@cam.ac.uk

Running Head: Change in the rate of gene flow during butterfly speciation

Abstract

Documenting the full extent of gene flow during speciation poses a challenge, as species ranges change over time and current rates of hybridisation might not reflect historical trends. Theoretical work has emphasized the potential for speciation in the face of ongoing hybridisation, and the genetic mechanisms that might facilitate this process. However, elucidating how the rate of gene flow between species may have changed over time has proved difficult. Here we use Approximate Bayesian Computation (ABC) to fit a model of speciation between the Neotropical butterflies *Heliconius melpomene* and *Heliconius cydno*. These species are ecologically divergent, rarely hybridize and display female hybrid sterility. Nevertheless, previous genomic studies suggests pervasive gene flow between them, extending deep into their past, and potentially throughout the speciation process. By modelling the rates of gene flow during early and later stages of speciation, we find that these species have been hybridising for hundreds of thousands of years, but have not done so continuously since their initial divergence. Instead, it appears that gene flow was rare or absent for as long as a million years in the early stages of speciation. Therefore, by dissecting the timing of gene flow between these species, we are able to reject a scenario of purely sympatric speciation in the face of continuous gene flow. We suggest that the period of minimal contact early in speciation may have allowed for the accumulation of genomic changes that later enabled these species to remain distinct despite a dramatic increase in the rate of hybridisation.

Introduction

Speciation is widely viewed as the development of reproductive isolation between

lineages. However, there is now considerable evidence that reproductive isolation is not

necessarily a genome-wide phenomenon, but rather that species integrity can be

maintained despite gene flow affecting a considerable proportion of the genome [1–6].

What remains less clear is the importance of gene flow (or lack thereof) for the

establishment of new species. Theory has shown that it may be possible, under certain

genetic and selective conditions, for species to become established in the face of

ongoing gene flow [7–14]. To test this theory, it is necessary either to observe speciation

in real time, or to reconstruct the historical extent and timing of gene flow between

existing species.

In geographic terms, speciation can be described as sympatric, parapatric or

allopatric. We follow Mallet et al. [15] in defining these terms: sympatric populations

share the same geographic area (but not necessarily the same niche), such that

individuals from the two populations are liable to encounter one-another frequently over

much of their range. Parapatric populations "occupy separate but adjoining geographic

regions," such that only a small fraction of individuals at the edge of each range are

liable to encounter the other. Allopatric populations are geographically separated, such

that encounters between them are very rare or impossible. Despite the abundance of

closely related sympatric species, there are very few cases in which it can be stated with

any certainty that speciation occurred in sympatry [e.g. crater-lake cichlids [16]]. In

terms of gene flow, we can predict that sympatric speciation might involve a gradual

decline over time, with higher rates of historic than contemporary gene flow [17]. In

allopatric speciation, gene flow would be absent until the populations came into

secondary contact. Parapatric speciation might fall somewhere in between these

extremes, with a low level of gene flow throughout, potentially decreasing over time,

but possibly also increasing if the populations later return to a sympatric distribution.

Genomic data now offers the exciting possibility of reconstructing the history of gene

flow between existing species, illuminating the roles of gene flow and geography in the

origin of new species.

A number of methods exist to fit a model of "isolation with migration" (IM) using

patterns of DNA sequence variation, thereby testing for post-speciation gene flow. This

can be achieved by maximising the likelihood of observed genetic data in a coalescent

framework, either directly [18–22] or using Markov Chain Monte Carlo (MCMC)

approximation [23–26]. However, these approaches lack power and accuracy to

examine change in the rate of gene flow over time [27,28], owing to characteristics of

the standard IM model itself [29]. This limitation could be overcome through the

implementation of more complex models [30], but this is currently not feasible in a

likelihood framework. Approximate Bayesian Computation (ABC) offers a tractable

means to fit such complex genetic models by avoiding the need to derive likelihoods

[31]. ABC is therefore suited to the problem of reconstructing changes in the rate of

gene flow during speciation [32,33], offering the potential to resolve long-standing

debates in the speciation literature.

Here we investigate the history of gene flow during speciation in *Heliconius*

butterflies. This Neotropical genus is well known for its broad diversity of aposematic

wing patterns, and multiple instances of Müllerian mimicry – where unrelated species

converge in wing pattern, providing a unified signal of toxicity to predators. Closely

related species usually differ in wing pattern, and it is thought that pattern divergence

between populations adapting to mimic different locally-abundant patterns could lead to

parapatric speciation [34,35]. We examine *Heliconius melpomene* and *Heliconius*

cydno, closely related species that have diverged in wing pattern and other ecological

traits during the past million years, but continue to hybridise at low frequency

[36] where their ranges overlap in the western parts of South America and Central

America. The ability to compare sympatric and allopatric populations of *H. cydno* and

H. melpomene provides an ideal opportunity to detect the genetic signatures of recent

gene flow. Indeed, whole-genome studies have found evidence of abundant gene flow

between these species, affecting a large proportion of the genome and extending deep

into the past [3,4]. However, it has remained uncertain whether this pair diverged in the

face of ongoing gene flow, or experienced a period of isolation early during speciation.

We used ABC to reconstruct the genealogical history of three populations: *H. cydno*

from Panama, *H. melpomene rosina* from Panama (sympatric with *cydno*) and *H.*

melpomene melpomene from French Guiana (allopatric) (Fig. 1A). Our model allowed

for hybridisation between *H. cydno* and *H. melpomene* throughout speciation, and

accounted for the possibility of a change in the rate of hybridisation during this time by

considering two separate periods with distinct migration rates, the duration or which

could vary. This enabled us to test various hypotheses under a single model, including a

clean split without gene flow, continuous gene flow throughout speciation or gene flow

restricted to ancient or recent time periods.

Figure 1. Species distributions, sample locations, summary statistic and model

design. A. Distributions (shaded) of *H. melpomene* (light red) and *H. cydno* (grey), based

on Rosser et al. [37]. Sampling locations in Panama and French Guiana are indicated. **B.**

The composite summary statistic used consisted of the proportions of the 13 possible

biallelic genotype combinations, where one individual is sampled from each population

(given to the right). '0' and '2' indicate alternative homozygous states and '1' indicates the

heterozygous state. Given that four individuals were sampled from each population, the

proportion of biallelic SNPs carrying each pattern was averaged over all 64 possible sets

of three samples. Although 25 SNP states are theoretically possible, twelve of these can

be folded if we ignore major and minor alleles (e.g. 2-0-1 is equivalent to 0-2-1) and so

these were counted together, to give 13 unique states. **C.** The model had ten free

parameters: four population sizes (N_a, N_c, N_{mr} and N_{mm}) (the ancestral *H. melpomene*

population size was assumed to be the average of N_{mr} and N_{mm}); migration rates between

H. cydno and *H. melpomene* in Periods 1 and 2 (M_{cm1} and M_{cm2}) the time dividing Periods

1 and 2 (t_d); migration rate between the two *H. melpomene* populations (M_m) , and the

split times for the two species (t_1) and the two *H. melpomene* populations (t_2).

Results

Genotype data and summary statistics

Whole genome resequence data from twelve wild-caught butterflies, with four

representatives from each of the three sampled populations, *H. cydno*, *H. m. rosina* and

H. m. melpomene, were used for model fitting (S1 Table, data from Martin et al. 2013).

Only intergenic regions, as defined by the *Heliconius melpomene* reference genome

annotation v1.1 [38], with high-quality genotype calls (see Materials and Methods) for

all twelve samples, were considered. We also excluded all scaffolds on the Z

chromosome, which is known to experience strongly reduced gene flow [4], as well as

putative CpG clusters, which can have unusual mutation rates. These criteria gave ~60

million sites (22% of the genome), of which approximately 10% were polymorphic

(Single Nucleotide Polymorphisms, SNPs). The composite summary statistic used for

model fitting consisted of the proportion of bi-allelic sites carrying each possible

combination of genotypes among three diploid individuals. These proportions were

averaged over all possible triplets, where each population is represented by one

individual, and folded such that major and minor alleles were not distinguished (Fig.

1B, see Materials and Methods for details). This composite summary statistic is similar

to a three-dimensional site frequency spectrum. It provides a nearly exhaustive

summary of the available SNP data among the ingroup taxa, is independent of linkage

effects and scalable to any number of sites.

As expected, the most common SNP patterns were singletons, where one individual

was heterozygous and the other two were homozygous for the same allele (0-0-1, 0-1-0,

and 1-0-0; Fig. 1B). The most common pattern overall was 1-0-0, where *H. cydno* is

heterozygous and both *H. melpomene* individuals are homozygous. This is unsurprising,

given the longer branch leading to *H. cydno* (Fig. 1B). The pattern 0-0-1, where *H. m.*

melpomene from French Guiana is heterozygous, was also considerably more common

than 0-1-0, where *H. m. rosina* is heterozygous. This is consistent with increased shared

variation between *H. cydno* and the sympatric *H. m. rosina*. Similarly, 1-1-0 was more

common than 1-0-1, and 0-0-2 was more common than 0-2-0.

Estimating the timing and extent of gene flow

We consider a model with three populations (Fig. 1C), corresponding to *H. cydno*

(which splits from the ancestral *melpomene* population at time t_1), and *H. m. rosina* and

H. m. melpomene (which split at time t_2). Each lineage has a separate population size,

except for the ancestral *melpomene* population, which is assumed to have a size equal to

the mean of the two *melpomene* populations. Because the two *melpomene* populations

represent extremes of a somewhat continuous range, migration between them is allowed

at a continuous rate M_m. Migration is also allowed between *H. cydno* and *H.*

melpomene, although after the split between the *melpomene* populations (t_2) only *H. m.*

rosina is able to exchange migrants with *H. cydno*. Two distinct periods of between-

species migration are modelled, with rates M_{cm1} and M_{cm2}. These periods are divided at a

time t_d such that Period 1 begins at t_1 and ends at t_d, and Period 2 runs from t_d to the

present. This model had ten free parameters: four population sizes, two split times, three

migration rates and one time dividing the migration periods. Model parameters were

estimated using Approximate Bayesian Computation (ABC) based on the summary

statistics described above (see Materials and Methods for details). Uniform priors were

used for all parameters except for t_1 and t_2, for which prior distributions were estimated

by analysis of mitochondrial sequence data (see Materials and Methods for details).

Our model pointed toward a dramatic change in the rate of inter-specific migration

(i.e. hybridisation resulting in gene flow) from early to later stages of speciation (Fig.

2). Migration was minimal in Period 1 (M_{cm1}~0.08 migrants per year [posterior mean]),

and around tenfold greater in Period 2 (M_{cm2}~0.81 migrants per year) (Fig. 2A, Table 1).

The date of transition between these two periods (t_d) had a fairly wide posterior

distribution, with a mean of 0.5 million years ago (Ma), but a 90% posterior density

interval extending from 0.1 Ma to 1.2 Ma (Fig. 2A, Table 1). This was nevertheless

considerably more recent than the inferred time of speciation (t_1), which had a fairly

narrow posterior distribution centred around 1.5 Ma (Fig. 2A, Table 1). Therefore, our

results support a case of stronger isolation during the early stages of speciation, with a

large increase in the rate of hybridisation later. The posterior mean of 0.5 Ma for the

date dividing the two periods would imply that hybridisation was rare or absent during

the first two thirds of the time since initial divergence (~1 million years). However, we

note that this transition cannot be dated with great accuracy, and indeed our model does

not allow us to infer whether the increase in hybridisation was sharp or gradual.

Nevertheless, it is notable that the posterior mean for the onset of more frequent

hybridisation coincides roughly with the split between the two *H. melpomene*

populations (t_2~0.53 Ma, Fig. 2A). Interestingly, the inferred rate of gene flow between

H. cydno and *H. m. rosina* during this second period is several times greater than that

between the two *H. melpomene* populations (Fig. 2A, Table 1).

 Figure 2. Posteriors for time and migration rate parameters, and a schematic
representation of the inferred model. A. Posterior distributions for the three time
perameters (left) and three migration rates (right). Densities are scaled for ease of
comparison. See S2 Figure for all posterior distributions. Posterior means are indicated by
vertical dashed lines. **B.** A schematic phylogeny, where the posterior mean for each
parameter is indicated. Colours correspond to those in A. Times are indicated by
horizontal dashed lines. Posteriors for the times are given on the right-hand-side for
reference. Migration rates are indicated by arrows, with the width of the arrow scaled
according to the migration rate. Relative population sizes are indicated by branch widths.
See Table 1 for all values.

Table 1. Properties of posterior distributions for the ten model parameters

Parameter	Mean	Mode	Median	HPD90 Lower[1]	HPD90 Upper[1]
t_1 (Ma)	1.5	1.4	1.5	1.2	1.8
t_2 (Ma)	0.53	0.53	0.53	0.42	0.65
t_d (Ma)	0.5	0.3	0.4	0.1	1.2
M_{cm1} (ind. /year)	0.08	0.01	0.06	0.01	0.86
M_{cm2} (ind. /year)	0.81	0.95	0.84	0.36	1.6
M_m (ind. /Year)	0.26	0.34	0.28	0.06	0.77
N_c (M ind.)	5.3	5.3	5.3	4.6	5.9
N_{mr} (M ind.)	1.8	1.8	1.8	1.2	2.4
N_{mm} (M ind.)	3.9	3.9	3.9	3.4	4.4
N_A (M ind.)	1.1	1.3	1.1	0.3	2.0

[1] Upper and lower bounds of the 90% Highest Posterior Density (HPD) interval. The

HPD is defined as the set of values making up 90% of the density distribution and within

which all values have higher density than outside.

Estimates of N_e were generally larger than previous estimates for these species [3],

with relatively narrow posterior distributions (Table 1, S2 Figure). This difference may

be driven by a lower mutation rate used in the present study. Consistent with this

previous study, the estimated N_e for *H. cydno* (~5.3 M individuals) was larger than both

H. m. rosina (~1.8 M) and *H. m. melpomene* (~3.9 M).

Overall, simulated summary statistics from the retained parameter combinations

matched the observed summary statistic well (S3 Figure). To investigate the robustness

of our conclusions, we repeated the ABC analysis using the split time priors from model

F (S3 Table and see above). The general findings were largely the same (data not

shown), indicating that our conclusions are not strongly influenced by the split time

priors.

Discussion

The existence of distinct species that share genetic material through hybridisation, or

have done so in the recent past, is no longer disputable. Genome-scale data have

provided overwhelming evidence of pervasive gene flow between species, in taxa as

diverse as fruit flies [39], flycatchers [2], and hominids [19,40]. However,

understanding the timing of gene flow during speciation, and the importance of

geographical isolation for species establishment remain difficult. Here, we combine a

large-scale genomic dataset with Approximate Bayesian Computation (ABC) to

reconstruct speciation in *Heliconius* butterflies. Our findings support recent studies

showing abundant gene flow between *H. melpomene* and *H. cydno* [3,4], and suggest

that gene flow has been ongoing for approximately half a million years (two million

generations). However, we also find that hybridisation was rare or absent during the

roughly the first million years of divergence between these species, a factor that may

have played an important role in their establishment.

In order to reduce the parameter space explored, we used a fairly narrow joint prior

distribution for the two split times, t_1 and t_2. This joint distribution was inferred using

mitochondrial sequence data, under the assumption that mitochondrial introgression

between *H. cydno* and *H. melpomene* should be unlikely. This is reasonable given the

fact that female hybrids are sterile [41]. Although one *H. melpomene* population from

Colombia is known to carry *H. cydno*-like mitochondrial haplotypes [42], an analysis of

all 449 unique haplotypes available on Genbank has not indicated any other

mitochondrial introgression events between these species (see Materials and Methods).

Our *cydno-melpomene* split time of 1.5 Ma is within the range of values inferred in

previous analyses using IM-based methods [3,42,43]. In addition, posterior distributions

for both split times were not strongly skewed toward the edge of the priors. It is

nevertheless possible that the mitochondrial split times provide inadequate estimates of

the nuclear split times. Even if this were the case, it is likely that our general conclusion

of reduced migration earlier in speciation will still hold. Indeed, when we repeated the

ABC analysis with a different, but overlapping, set of split time priors, the results were

largely unchanged.

Allopatric speciation is often the null hypothesis in speciation studies [44]. One

scenario that would be consistent with our results is that speciation began in allopatry,

possibly with the emerging species separated by the Andes mountains. Subsequent

range expansion of *H. melpomene* into the western Andes and Central America would

have lead leading to secondary contact. The lower N_e of *H. m. rosina* compared to *H. m.*

melpomene is consistent with range expansion from east to west, which has been

proposed previously [45]. However, our results may also be consistent with parapatric

speciation, which is probably more common in *Heliconius* [35,46]. In fact, most pairs of

sister species in the genus are sympatric or parapatric (Rosser et al. In Review).

Allopatric populations of extant species, for example those on Caribbean islands, tend

not to display phenotypic and ecological divergence from their mainland progenitors. In

contrast, many species, including *H. melpomene* and *H. cydno*, are divided into

numerous parapatric wing-pattern races across their mainland ranges. *H. melpomene*

and *H. cydno* are also partially segregated by altitude, so it is plausible that parapatric

adaptation to altitude in the Andes played a role in their speciation. The evolution of

strong assortative mating associated with wing pattern might then have led to nearly

complete reproductive isolation between the parapatric populations. Indeed, loci

affecting both mate preference and hybrid sterility are known to be physically linked to

wing patterning loci in these species, which might have enhanced reproductive isolation

following divergence in wing pattern [47,48]. Ecological divergence, most notably in

host plant use but perhaps also microhabitat preference [49], would then have followed

later, permitting sympatric coexistence without competition [34]. The inferred increase

in gene flow later in speciation might therefore reflect increased contact associated with

the transition from parapatric to sympatric ranges. One final piece of evidence for

parapatric speciation is the existence of several other species pairs that may represent an

intermediate step in this process. The best studied are *Heliconius himera* and *H. erato,*

which are largely parapatric with only narrow zones of overlap. They are strongly

differentiated genetically [50] and display assortative mating based on colour pattern

[51]. However, they have not diverged in host plant usage [52], which perhaps prevents

their sympatric coexistance through competitive exclusion.

Regardless of its cause, we can speculate that an initial period of reduced gene flow

contributed to the formation of these species. Reduced gene flow can facilitate the

accumulation of Dobzhanzky-Muller incompatibilities [53,54], which would help to

maintain species integrity even after the rate of hybridisation increased. For example,

gene flow is minimal across the entire Z chromosome [4], consistent with a high density

of incompatibility loci in this part of the genome. Interestingly, in these species there are

also genetic associations between wing pattern, host preference and mate preference loci

which likely facilitate coexistence in sympatry [47,48]. However it is unclear whether

such associations have arisen since hybridisation became widespread, or whether they

fortuitously pre-dated the period of extensive contact. Finer-scale analysis of the

patterns of introgression across the genome, combined with mapping incompatibility

loci and structural differences in the genome will help to dissect the various factors

contributing to species persistence.

Despite our finding that hybridisation was rare or absent for approximately two

thirds of the time since speciation, this nevertheless implies that the hybridisation has

now been ongoing for around two million generations. Our model assumes a single

change in the rate of gene flow, while it is highly probably that this rate has changed

more gradually through time. The wide posterior distribution for the time of this

transition is consistent with a gradual increase over perhaps hundreds of thousands of

generations. There is also reason to believe that the rate of gene flow has recently begun

to decrease. The existence of character displacement (sympatric males display stronger

mate discrimination than allopatric males [55], suggests that selection may have acted to

reinforce reproductive isolation in sympatry. Nevertheless, there remains strong evidence that gene flow continues today, both in the occurance of natural F1 hybrids [36], and in geographic patterns of shared variation [4]. Specifically, *H. cydno* samples from Panama share an excess of variation with *H. melpomene* samples from the same location compared to those from about 100 km away [4]. A recent study of the hybridising mouse subspecies *Mus musculus musculus* and *M. m. domestica* supported a similar scenario to that described here, with gene flow occurring over the last 25% of time since initial divergence, although at a lower rate (<0.2 ind/gen) [32]. Our estimate of ~0.84 migrants per generation represents the effective number of hybrids, but it is certain that the number of actual F1 hybrid butterflies produced exceeds this value considerably, for two reasons. Firstly, in accordance with Haldane's Rule, female (ZW) F1 hybrids are sterile [41], and therefore do not contribute to observed gene flow. Secondly, F1 hybrids are subject to increased predation owing to their non-mimetic wing patterns [56].

Finally, it is worth considering the consequences of continued hybridisation between these species. Although a whole-genome phylogeny groups the *H. melpomene* populations as monophyletic, currently 40% of 100 kb windows group *H. m. rosina* with *H. cydno*, to the exclusion of the French Guianan *H. m. melpomene* [4]. Nevertheless, these sympatric populations retain the phenotypic, behavioural and ecological traits specific to their respective species, implying that species integrity is surprisingly resilient to gene exchange. It is certain that gene flow is inhibited by selection in some genomic regions, most obviously the wing pattern loci. However, natural selection has also favoured the occasional exchange of wing pattern alleles between certain populations of these clades, producing the paired mimetic races of *H. melpomene* and *H. timareta* found on the eastern slopes of the Andes [38,57]. It seems likely that much of the genome is neutral with respect to gene flow, and that most of the signal seen here is due to neutral exchange of alleles in sympatry, although we have not attempted to test for evidence of adaptive introgression. It is therefore possible that

319 ongoing hybridisation, even at a low rate, might eventually lead to a situation where the

320 majority of the genome clusters populations by geography rather than by species,

321 making one or both species paraphyletic. It seems inevitable that genomic studies will

322 reveal such species pairs in the near future, posing a challenge to species definitions

323 based on aggregate genetic ancestry.

Materials and Methods

Samples and genotyping

We used published whole-genome resequence data for twelve wild-caught butterflies

(S1 Table, data from Martin et al. [4], www.datadryad.com doi:10.5061/dryad.dk712).

Details of the sequencing, mapping and genotyping procedures are described by Martin

et al. [4]. Briefly, 100bp paired-end Illumina reads were mapped to the *H. melpomene*

reference genome [38], version 1.1, using Stampy [58]. Local realignment around

indels and genotyping were both performed using The Genome Analysis Toolkit

(GATK) [59]. For the purpose of this study, we considered only intergenic SNPs,

identified based on the *H. melpomene* genome annotation, version 1.1. CpG islands

were identified using the program CpGcluster [60], and these sites were excluded. Only

high quality genotype calls were considered. High quality genotypes met the following

conditions: quality (QUAL) $\geq$ 30, 10 $\leq$ read depth per individual $\leq$ 200, and GQ $\geq$ 30

for SNPs. Processed genotype calls data are available from www.datadryad.com

doi:XXX.

Summary statistic

The summary statistic used for model fitting was a composite of the proportion of

sites representing each of the possible combinations of bi-allelic genotypes among three

diploid individuals, with one individual representing each population (Fig. 1B). For

example, a SNP would be assigned the pattern 0-1-2, if the *H. cydno* individual was

homozygous, carrying zero copies of the minor allele, the *H. m. rosina* individual was

heterozygous, carrying one copy, and the *H. m. melpomene* individual was homozygous

with two copies of the minor allele. The counts of all patterns were then folded, such

that major and minor alleles were not taken into account. For example the pattern 0-1-2

was taken as equivalent to 2-1-0. This gave 13 unique SNP patterns (Fig. 1B). Because

four individuals were sampled from each population, the counts of each pattern were

averaged over all 64 possible triplets with one individual from each population. Custom

scripts used to calculate and plot summary statistics are avauialable from

www.datadryad.com doi:XXX. Since having too many summary statistics is a known

problem with ABC, we used Partial Least Squares [implemented in the findPLS.r script

in the ABC Toolbox [61]] to find the eight most informative linear combinations of the

original summary statistics.

Model

A three-population model of isolation with migration was used (Fig. 1C). An

ancestral population divides at time t_1 into two populations (corresponding to *H. cydno*

and *H melpomene,* respectively). At time t_2, the *melpomene* population further divides

into the *H. m. rosina* and *H. m. melpomene* races, which remain connected by limited

gene flow at a constant rate M_m. The two *H. melpomene* populations, the *H. cydno*

population, and the ancestral population, all have unique population sizes, but the size

of the ancestral *melpomene* population is assumed to be the average of the two

melpomene populations. Migration is allowed between *H. cydno* and the ancestral *H.*

melpomene population, and between *H. cydno* and *H. m. rosina* after the two *H.*

melpomene populations diverge. Two distinct periods of hybridisation are modelled,

with rates M_{cm1} and M_{cm2}. These two periods occupy the entire speciation time from t_1 to

the present, and are divided at time t_d. Hence, Period 1 runs from t_1 to t_d and Period 2

from t_d to the present. The division between the periods, t_d, may fall anywhere between

t_1 and the present. A constant mutation rate of 1.9×10^{-9} per site per generation was used.

This corresponds to the estimated per-generation mutation rate for *H. melpomene* [62],

corrected for weak purifying selection on intergenic regions by multiplying by the

relative level of interspecific divergence at intergenic and putatively neutral four-fold

degenerate sites (data not shown). A generation time of 0.25 years was assumed [63].

Priors for split times

To reduce the dimensionality of the model, we used fairly narrow priors for the two

split times t_1 and t_2 (S2 Table). These were inferred using analysis of mitochondrial

sequence data, which should be resistant to gene flow between these taxa. This

assumption was first tested with a Maximum Likelihood analysis of all 847 publicly

available sequences (449 unique haplotypes) in RAxML v.8 [64]. This identified a

single, previously known [42] case of mitochondrial introgression between these

species, which does not involve the populations considered here. Sequence data for

1606 bp of *CoI/II* for 125 samples from several populations of *H. melpomene*, *H. cydno*

and the outgroup silvaniform clade were obtained from Genbank and the data of from

Martin et al. [4], and aligned with MUSCLE [65]. Strict and relaxed molecular clock

models and codon-partitioning schemes were fitted to the data in BEAST v. 1.8.

[66] and compared with a posterior analog of AIC (AICM) and Bayes Factors calculated

by the Stepping Stone Analysis [67] (S2 and S3 Table). For root-calibrated analyses, the

split time between the *H. melpomene* and Silvaniform caldes inferred by Kozak et al.

[68] was used. For fixed rate analyses a mutation rate of 0.0024 per million years was

used, as inferred under a relaxed-clock model applied to the complete *Heliconiini*

alignment of Kozak et al. [68]. While the exact split dates varied, all approaches

converged on the same topology and a similar ratio of split times t_1/t_2 (S3 Table). Bayes

Factors and AICM [67] favoured a strict clock model (model E, S2 Table) with a

separate partition for third codon positions. We present results obtained using the split

times from model E, although we also tested the split times from model F (S3 Table).

The resulting joint posterior distributions for the two split times formed the priors for

the ABC simulations, with pairs of times for t_1 and t_2 being drawn together from this

joint distribution.

Model fitting using ABC

398
Approximate Bayesian Computation (ABC) was used to estimate parameters of the
model. Briefly, ABC fits a model by evaluating the distance between observed and
simulated summary statistics, allowing the estimation of posterior probability
distributions without calculating likelihood functions.

Uniform priors were used for all parameters except for the split times t_1 and t_2 (see
above). Two million parameter combinations were generated over the parameter space
by sampling from the prior distributions randomly and independently (except for
parameters t_1 and t_2, which had a joint prior distribution). A custom program (written in
C/C++) was then used to simulate 100,000 unlinked SNPs from our model under the
standard coalescent framework [69] for each sampled parameter set, and then to
calculate the summary statistic. SNPs were simulated independently (i.e. unlinked) as
the composite summary statistic used here, which is based on the genome-wide joint
frequency spectrum, should not be strongly influenced by linkage disequilibrium,
especially given the fairly rapid decline in LD in *Heliconius melpomene* [4]. Using the
standard ABC method, we used the Euclidian distance between simulated and observed
values to identify parameter combinations that fit the data well. We used a cut-off of
0.01 for accepting parameter combinations, yielding 27377 good parameter
combinations for ABC. To account for variation in goodness of fit obtained among
retained parameters, the distribution of retained parameters was adjusted using the
General Linear Model method of [70], as implemented by ABCestimator of the ABC
Toolbox [61].

Acknowledgements

421   We thank Nick Barton, Aylwyn Scally and Konrad Lohse, for helpful comments on
422 earlier drafts of the manuscript. Computational support was provided by Jenny Barna,
423 School of Life Sciences, University of Cambridge.

References

1. Kulathinal RJ, Stevison LS, Noor M a F (2009) The genomics of speciation in Drosophila: diversity, divergence, and introgression estimated using low-coverage genome sequencing. PLoS Genet 5: e1000550. doi:10.1371/journal.pgen.1000550.

2. Ellegren H, Smeds L, Burri R, Olason PPI, Backström N, et al. (2012) The genomic landscape of species divergence in Ficedula flycatchers. Nature 491: 756–760. doi:10.1038/nature11584.

3. Kronforst MRR, Hansen MEB, Crawford NGG, Gallant JRR, Zhang W, et al. (2013) Hybridization reveals the evolving genomic architecture of speciation. Cell Rep 5: 666–677. doi:10.1016/j.celrep.2013.09.042.

4. Martin SH, Dasmahapatra KK, Nadeau NJ, Salazar C, Walters JR, et al. (2013) Genome-wide evidence for speciation with gene flow in Heliconius butterflies. Genome Res 23: 1817–1828. doi:10.1101/gr.159426.113.

5. Fontaine MC, Pease JB, Steele A, Waterhouse RM, Neafsey DE, et al. (2014) Extensive introgression in a malaria vector species complex revealed by phylogenomics. Science 347: 1258524 – . doi:10.1126/science.1258524.

6. Lamichhaney S, Berglund J, Almén MS, Maqbool K, Grabherr M, et al. (2015) Evolution of Darwin's finches and their beaks revealed by genome sequencing. Nature advance on. doi:10.1038/nature14181.

7. Maynard Smith J (1966) Sympatric speciation. Am Nat 100: 637–650.

8. Lande R (1982) Rapid Origin of Sexual Isolation and Character Divergence in a Cline. Evolution (N Y) 36: 213–223.

9. Slatkin M (1982) Pleiotropy and Parapatric Speciation. 36: 263–270.

10. Kirkpatrick M, Ravigné V (2002) Speciation by natural and sexual selection: models and experiments. Am Nat 159 Suppl : S22–S35. doi:10.1086/338370.

11. Navarro A, Barton NH (2003) Accumulating postzygotic isolation genes in parapatry: a new twist on chromosomal speciation. Evolution 57: 447–459.

12. Gavrilets S (2004) Fitness landscapes and the origin of species. Princeton University Press.

13. Kirkpatrick M, Barton N (2006) Chromosome inversions, local adaptation and speciation. Genetics 173: 419–434. doi:10.1534/genetics.105.047985.

14. Van Doorn GS, Dieckmann U, Weissing FJ (2004) Sympatric speciation by sexual selection: a critical reevaluation. Am Nat 163: 709–725. doi:10.1086/383619.

15. Mallet J, Meyer a., Nosil P, Feder JL (2009) Space, sympatry and speciation. J Evol Biol 22: 2332–2341. doi:10.1111/j.1420-9101.2009.01816.x.

16. Barluenga M, Stölting KN, Salzburger W, Muschick M, Meyer A (2006) Sympatric speciation in Nicaraguan crater lake cichlid fish. Nature 439: 719–723. doi:10.1038/nature04325.

17. Wu C (2001) The genic view of the process of speciation. J Evol Biol 14: 851–865.

18. Lohse K, Harrison RJ, Barton NH (2011) A general method for calculating likelihoods under the coalescent process. Genetics 189: 977–987. doi:10.1534/genetics.111.129569.

19. Lohse K, Frantz L a F (2014) Neandertal admixture in eurasia confirmed by maximum-likelihood analysis of three genomes. Genetics 196: 1241–1251. doi:10.1534/genetics.114.162396.

20. Wilkinson-Herbots HM (2008) The distribution of the coalescence time and the number of

pairwise nucleotide differences in the "isolation with migration" model. Theor Popul
Biol 73: 277–288. doi:10.1016/j.tpb.2007.11.001.

21. Wang Y, Hey J (2010) Estimating divergence parameters with small samples from a large
number of loci. Genetics 184: 363–379. doi:10.1534/genetics.109.110528.

22. Zhu T, Yang Z (2012) Maximum likelihood implementation of an isolation-with-migration
model with three species for testing speciation with gene flow. Mol Biol Evol 29: 3131–
3142. doi:10.1093/molbev/mss118.

23. Nielsen R, Wakeley J (2001) Distinguishing migration from isolation: a Markov chain
Monte Carlo approach. Genetics 158: 885–896.

24. Hey J, Nielsen R (2004) Multilocus methods for estimating population sizes, migration rates
and divergence time, with applications to the divergence of Drosophila pseudoobscura
and D. persimilis. Genetics 167: 747–760.

25. Hey J, Nielsen R (2007) Integration within the Felsenstein equation for improved Markov
chain Monte Carlo methods in population genetics. Proc Natl Acad Sci U S A 104:
2785–2790. doi:10.1073/pnas.0611164104.

26. Hey J (2010) Isolation with Migration Models for More Than Two Populations. Mol Biol
Evol 27: 905–920.

27. Becquet C, Przeworski M (2009) Learning about modes of speciation by computational
approaches. Evolution (N Y) 63: 2547–2562.

28. Strasburg JL, Rieseberg LH (2011) Interpreting the estimated timing of migration events
between hybridizing species. Mol Ecol 20: 2353–2366. doi:10.1111/j.1365-
294X.2011.05048.x.

29. Sousa VC, Grelaud A, Hey J (2011) On the nonidentifiability of migration time estimates in
isolation with migration models. Mol Ecol 20: 3956–3962.

30. Strasburg JL, Rieseberg LH (2013) Methodological challenges to realizing the potential of
hybridization research. J Evol Biol 26: 259–260. doi:10.1111/jeb.12006.

31. Beaumont M a. (2010) Approximate Bayesian Computation in Evolution and Ecology. Annu
Rev Ecol Evol Syst 41: 379–406. doi:10.1146/annurev-ecolsys-102209-144621.

32. Duvaux L, Belkhir K, Boulesteix M, Boursot P (2011) Isolation and gene flow: inferring the
speciation history of European house mice. Mol Ecol 20: 5248–5264.
doi:10.1111/j.1365-294X.2011.05343.x.

33. Li J-W, Yeung CKL, Tsai P-W, Lin R-C, Yeh C-F, et al. (2010) Rejecting strictly allopatric
speciation on a continental island: prolonged postdivergence gene flow between Taiwan
(Leucodioptron taewanus, Passeriformes Timaliidae) and Chinese (L. canorum canorum)
hwameis. Mol Ecol 19: 494–507. doi:10.1111/j.1365-294X.2009.04494.x.

34. Mallet J, McMillan WO, Jiggins CD (1998) Mimicry and warning color at the boundary
between races and species. Endless forms species Speciat: 390–403.

35. Jiggins C (2008) Ecological speciation in mimetic butterflies. Bioscience 58: 541–548.

36. Mallet J, Beltrán M, Neukirchen W, Linares M (2007) Natural hybridization in heliconiine
butterflies: the species boundary as a continuum. BMC Evol Biol 7: 28.
doi:10.1186/1471-2148-7-28.

37. Rosser N, Phillimore AB, Huertas B, Willmott KR, Mallet J (2012) Testing historical
explanations for gradients in species richness in heliconiine butterflies of tropical
America. Biol J Linn Soc 105: 479–497. doi:10.1111/j.1095-8312.2011.01814.x.

38. The Heliconius Genome Consortium 1 (2012) Butterfly genome reveals promiscuous
exchange of mimicry adaptations among species. Nature 487: 94–98.
doi:10.1038/nature11041.

39. Garrigan D, Kingan SSB, Geneva AJ, Andolfatto P, Clark AG, et al. (2012) Genome
sequencing reveals complex speciation in the Drosophila simulans clade. Genome … 22:
1499–1511. doi:10.1101/gr.130922.111.

40. Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, et al. (2010) A draft sequence of the
Neandertal genome. Science 328: 710–722. doi:10.1126/science.1188021.

41. Naisbit RE, Jiggins CD, Linares M, Salazar C, Mallet J (2002) Hybrid Sterility, Haldane's
Rule and Speciation in. Race 1526: 1517–1526.

42. Salazar C, Jiggins CD, Taylor JE, Kronforst MR, Linares M (2008) Gene flow and the
genealogical history of Heliconius heurippa. BMC Evol Biol 8: 132. doi:10.1186/1471-
2148-8-132.

43. Bull V, Beltrán M, Jiggins CD, McMillan WO, Bermingham E, et al. (2006) Polyphyly and
gene flow between non-sibling Heliconius species. BMC Biol 4: 11. doi:10.1186/1741-
7007-4-11.

44. Coyne JA, Orr HA, others (2004) Speciation. Sinauer Associates Sunderland, MA.

45. Quek S-P, Counterman B a, Albuquerque de Moura P, Cardoso MZ, Marshall CR, et al.
(2010) Dissecting comimetic radiations in Heliconius reveals divergent histories of
convergent butterflies. Proc Natl Acad Sci U S A 107: 7365–7370.
doi:10.1073/pnas.0911572107.

46. Jiggins CD, Estrada C, Rodrigues a (2004) Mimicry and the evolution of premating isolation
in Heliconius melpomene Linnaeus. J Evol Biol 17: 680–691. doi:10.1111/j.1420-
9101.2004.00675.x.

47. Merrill RM, Van Schooten B, Scott J a, Jiggins CD (2011) Pervasive genetic associations
between traits causing reproductive isolation in Heliconius butterflies. Proc Biol Sci 278:
511–518. doi:10.1098/rspb.2010.1493.

48. Merrill RM, Naisbit RE, Mallet J, Jiggins CD (2013) Ecological and genetic factors
influencing the transition between host-use strategies in sympatric Heliconius butterflies.
J Evol Biol 26: 1959–1967. doi:10.1111/jeb.12194.

49. Estrada C, Jiggins CD (2002) Patterns of pollen feeding and habitat preference among
Heliconius species. Ecol Entomol 27: 448–456. doi:10.1046/j.1365-2311.2002.00434.x.

50. Jiggins CD, McMillan WO, King P, Mallet J (1997) The maintenance of species differences
across a Heliconius hybrid zone. Heredity (Edinb) 79: 495–505.
doi:10.1038/sj.hdy.6882230.

51. Merrill RM, Chia A, Nadeau NJ (2014) Divergent warning patterns contribute to assortative
mating between incipient Heliconius species. Ecol Evol 4: 911–917.
doi:10.1002/ece3.996.

52. Jiggins CD, McMillan WO, Mallet J (1997) Host plant adaptation has not played a role in
the recent speciation of Heliconius himera and Heliconius erato. Ecol Entomol 22: 361–
365. doi:10.1046/j.1365-2311.1997.00067.x.

53. Kondrashov AS (2003) Accumulation of Dobzhansky-Muller incompatibilities within a
spatially structured population. Evolution 57: 151–153. doi:10.1111/j.0014-
3820.2003.tb00223.x.

54. Bank C, Bürger R, Hermisson J (2012) The limits to parapatric speciation: Dobzhansky-
Muller incompatibilities in a continent-island model. Genetics 191: 845–863.
doi:10.1534/genetics.111.137513.

55. Jiggins CD, Naisbit RE, Coe RL, Mallet J (2001) Reproductive isolation caused by colour
pattern mimicry. Nature 411: 302–305. doi:10.1038/35077075.

56. Merrill RM, Wallbank RWR, Bull V, Salazar PC a, Mallet J, et al. (2012) Disruptive

ecological selection on a mating cue. Proc Biol Sci. doi:10.1098/rspb.2012.1968.

57. Pardo-Diaz C, Salazar C, Baxter SW, Merot C, Figueiredo-Ready W, et al. (2012) Adaptive
introgression across species boundaries in Heliconius butterflies. PLoS Genet 8:
e1002752. doi:10.1371/journal.pgen.1002752.

58. Lunter G, Goodson M (2011) Stampy: a statistical algorithm for sensitive and fast mapping
of Illumina sequence reads. Genome Res 21: 936–939. doi:10.1101/gr.111120.110.

59. DePristo M a, Banks E, Poplin R, Garimella K V, Maguire JR, et al. (2011) A framework for
variation discovery and genotyping using next-generation DNA sequencing data. Nat
Genet 43: 491–498. doi:10.1038/ng.806.

60. Hackenberg M, Previti C, Luque-Escamilla PL, Carpena P, Martínez-Aroza J, et al. (2006)
CpGcluster: a distance-based algorithm for CpG-island detection. BMC Bioinformatics
7: 446. doi:10.1186/1471-2105-7-446.

61. Wegmann D, Leuenberger C, Neuenschwander S, Excoffier L (2010) ABCtoolbox: a
versatile toolkit for approximate Bayesian computations. BMC Bioinformatics 11: 116.

62. Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, et al. (2014)
Estimation of the Spontaneous Mutation Rate in Heliconius melpomene. Mol Biol Evol:
1–5. doi:10.1093/molbev/msu302.

63. Mallet J (1986) Hybrid zones of Heliconius butterflies in Panama and the stability and
movement of warning colour clines. Heredity (Edinb) 56: 191–202.

64. Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis
of large phylogenies. Bioinformatics 30: 1312–1313. doi:10.1093/bioinformatics/btu033.

65. Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res 32: 1792–1797. doi:10.1093/nar/gkh340.

66. Drummond AJ, Suchard MA, Xie D, Rambaut A (2012) Bayesian phylogenetics with
BEAUti and the BEAST 1.7. Mol Biol Evol 29: 1969–1973.

67. Baele G, Lemey P, Bedford T, Rambaut A, Suchard MA, et al. (2012) Improving the
accuracy of demographic and molecular clock model comparison while accommodating
phylogenetic uncertainty. Mol Biol Evol 29: 2157–2167. doi:10.1093/molbev/mss084.

68. Kozak KM, Wahlberg N, Neild A, Dasmahapatra KK, Mallet J, et al. (2015) Multilocus
Species Trees Show the Recent Adaptive Radiation of the Mimetic. Syst Biol.

69. Kingman J (1982) The coalescent. Stoch Proc Appl 13: 235–248.

70. Leuenberger C, Wegmann D (2010) Bayesian computation and model selection without
likelihoods. Genetics 184: 243–252.

Financial Disclosure

This work was funded by the Biotechnology and Biological Sciences Research Council (www.bbsrc.ac.uk/) grants BB/H01439X/1 to CDJ and H005854/1 to AM; and the European Research Council (erc.europa.eu/) SpeciationGenomics grant to CDJ. SHM is funded by St John's College, Cambridge (www.joh.cam.ac.uk/). KMK was supported by the Herchel Smith and Balfour Studentships to the Department of

List of supplemental files

S1 Table. Sample information and sequencing read depth

S2 Table. Model comparison of various strategies for estimating the divergence

times in BEAST. Models E and F are equally good based on Bayes Factors. Log Bayes

Factors (BF) are calculated based on the log Marginal Likelihood estimates (MLE) from

Path Sampling (PS) and Stepping Stone Analysis (SSA) (Baele et al. 2012). The

molecular clock rates were calibrated by either modelling the age of the root or setting a

constant rate of substitution. UCLD= Uncorrelated Lognormal clock.

S3 Table. Parameter values for the three best Bayesian models of divergence

between the *CoI/II* sequences.

S1 Figure. Bayesian phylogeny of the *Cytochrome Oxidase I/II* (*CoI/II*) haplotypes

from the *Heliconius melpomene* / *H. cydno* clade. This tree was estimated under the

relaxed molecular clock model (F) with calibrated age of the root. The sampling is based

on a balanced design with 25 samples per group, including *H. timareta* as the sister

species of *H. cydno*. Blue: *H. melpomene* from Central America; green: *H. melpomene*

from French Guiana (clade including allopatric samples used in the analysis of the

nuclear genomes); pink: *H. cydno*; grey: Silvaniform outgroups. Time scale in millions of

years, bars represent the 95% HPD intervals around split ages. Samples with whole

mitogenome data from Martin et al. (2013) indicated in capital letters. NCBI GI numbers

are provided for all sequences obtained from GenBank.

S2 Figure. Posterior density plots inferred by ABC for the 10 model parameters.

Parameters labels match Fig. 1 in the main text, with the three time parameters given in

the left-hand column (A-C), migration rates in the middle column (D-F), and population

sizes in the right-hand column (G-J). Grey shading indicates the 90% highest posterior

density (HPD) interval, and posterior means are indicated by vertical dashed lines.

S3 Figure. Summary statistics from retained parameter combinations compared to

the observed values. Patterns 1-13 correspond respectively to 0-0-1, 0-0-2, 0-1-0, 0-1-1,

0-1-2, 0-2-0, 0-2-1, 2-0-0, 1-0-0, 1-0-1, 1-0-2, 1-1-0 and 1-1-1. Box plots indicate the

distribution of simulated values from the 27377 retained parameter combinations. Red

points indicate outliers (outside of the 10th and 90th percentiles). Yellow diamonds

indicate observed frequencies, as in Fig. 1B. Here the y-axis indicates the the scaled

frequency of each pattern, in units of million generations. This can be obtained by

dividing the observed absolute frequency of each pattern (as in Fig. 1B) by the mutation

rate per site per million years.