1

3   Running head: Hidden state speciation and extinction

5   **Detecting hidden diversification shifts in models of trait-dependent speciation and**

6   **extinction**

8   Jeremy M. Beaulieu[1,2] and Brian C. O'Meara[2]

10   *[1]National Institute for Biological and Mathematical Synthesis, University of Tennessee,*

11   *Knoxville, TN 37996, USA*

12   *[2]Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville,*

13   *TN, 37996-1610, USA*

Email for correspondence: jbeaulieu@nimbios.org

Abstract – The distribution of diversity can vary considerably from clade to clade.

Attempts to understand these patterns often employ state-dependent speciation and

extinction models to determine whether the evolution of a particular novel trait has

increased speciation rates and/or decreased their extinction rates. It is still unclear,

however, whether these models are uncovering important drivers of diversification, or

whether they are simply pointing to more complex patterns involving many unmeasured

and co-distributed factors. Here we describe an extension to the popular state-dependent

speciation and extinction models that specifically accounts for the presence of

unmeasured factors that could impact diversification rates estimated for the states of any

observed trait, addressing at least one major criticism of BiSSE methods. Specifically,

our model, which we refer to as HiSSE (Hidden-State Speciation and Extinction),

assumes that related to each observed state in the model are "hidden" states that exhibit

potentially distinct diversification dynamics and transition rates than the observed states

in isolation. We also demonstrate how our model can be used as character-independent

diversification (CID) models that allow for a complex diversification process that is

independent of the evolution of a character. Under rigorous simulation tests and when

applied to empirical data, we find that HiSSE performs reasonably well, and can at least

detect net diversification rate differences between observed and hidden states and detect

when diversification rate differences do not correlate with the observed states. We

discuss the remaining issues with state-dependent speciation and extinction models in

general, and the important ways in which HiSSE provides a more nuanced understanding

of trait-dependent diversification.

Key words – hidden states, diversification, comparative methods, speciation, extinction

A key question in biology is, why are some groups are much more diverse than

others? Discussions of such questions are often focused on whether there is something

unique about exceptionally diverse lineages, such as the presence of some novel trait,

which has increased their speciation rate and/or decreased their extinction rates. The

BiSSE model (Binary-State Speciation and Extinction; Maddison et al. 2007) was derived

specifically as a means of examining the effect that the presence or absence of a single

character could have on diversification rates, while also accounting for possible

transitions between states. In theory, this model could be used not only for identifying

differences in diversification, but also detecting differences in transitions between

character states, or even some interplay of the two. In practice, however, it has mainly

been used to focus on diversification rate differences (e.g., Goldberg et al. 2010; Wilson

et al. 2011; Price et al. 2012; Beaulieu and Donoghue 2013; Weber and Agrawal, 2014).

It is somewhat surprising, perhaps, that studies that employ BiSSE often find that

the prediction of a trait leading to higher diversification rates is supported by the data. In

fact, all sorts of traits have been implicated as potential drivers of diversity patterns,

ranging from the evolution of herbivory in mammals (Price et al. 2012), to the evolution

of extra-floral nectaries in flowering plants (Weber and Agrawal 2014), to even the

evolution of particular body plans in fungi (i.e., gasteroid vs. nongasteroid forms; Wilson

et al. 2011). Some crucial caveats have recently been identified, however. First,

Maddison and FitzJohn (2014) raise a statistical concern regarding the inability of these

methods to properly account for independence. Consider, for instance, that the carpel,

which encloses the seeds of angiosperms, has evolved only once. They argue that the

inheritance and pseudoreplication of a single event becomes problematic – even if BiSSE

uncovers a significant correlation between the carpel and diversification, it is unclear

whether the carpel is an important driver of the immense diversity of flowering plants, or

whether this diversity is simply coincidental. It was also pointed out by Beaulieu and

Donoghue (2013) that even with many origins of a trait, it could be that only one clade

actually has a higher diversification rate associated with the focal trait, which is strong

enough to return higher diversification rates for that trait as a whole. In their case, it

appeared that plants with an achene (a fruit resembling a bare seed, as in "sunflower

seeds") had a higher diversification rate, but upon subdividing the tree it appeared that

this was from the inclusion of one clade in particular, Asterales, which contain the highly

diverse Asteraceae, the sunflower family. They argue that it is far more likely that some

combination of the achene and another unmeasured, co-distributed trait within Asterales

led to a higher diversification rate for achenes as a whole (on this point, also see

Maddison et al. 2007). Finally, Rabosky and Goldberg (2015) recently showed that for

realistically complex data sets, BiSSE methods almost always find that a neutral trait is

correlated with higher diversification. This is, however, largely a consequence of using

fairly "trivial" null models that assume equal rates of diversification between character

states, which cannot discern the possibility that a complex diversification process is

entirely independent from the focal trait.

All these caveats relate to the broader issue of whether the proximate drivers of

diversification are really ever just the focal traits themselves. At greater phylogenetic

scales this issue seems the most relevant, where the context of a shared trait is unlikely to

be consistent across many taxonomically distinct clades. In other words, a character's

effect on diversification will often be contingent on other factors, such as the assembly of

particular combinations of characters (e.g., a "synnovation" as defined by Donoghue and

Sanderson 2015) and/or movements into new geographic regions (e.g., de Querioz 2002;

Moore and Donoghue 2007). Recent generalizations of the BiSSE model (i.e., MuSSE:

Multistate Speciation and Extinction; FitzJohn 2012) do allow for additional binary

characters to be accounted for when examining the correlation between a binary trait and

net diversification. However, it may not always be clear what the exact characters might

be, and in the absence of such information, it should be difficult to ever confidently view

any one character state as the *true* underlying cause of increased diversification rates.

Here we describe a potential way forward for trait-dependent models of

diversification by extending the BiSSE framework to account for the presence of

unmeasured factors that could impact diversification rates estimated for the states of any

observed trait. Our model, which we refer to as HiSSE (Hidden State Speciation and

Extinction), assumes that related to each observed state in the model are "hidden" states

that exhibit potentially distinct diversification dynamics and transition rates than the

observed states in isolation. In this way, HiSSE is a state-dependent speciation and

extinction form of a Hidden Markov model (HMM). As we will show, HiSSE can

distinguish higher net diversification rates nested within clades exhibiting a particular

character state, can provide more meaningful tests of character-dependent diversification

(CID) through its use as a different kind of null model, and thus can provide a much more

refined understanding of how particular observed character states may influence the

diversification process.

117      Despite the important methodological advancement and power afforded by the

118   BiSSE model, it can provide a rather coarse-grained view of trait-based patterns of

119   diversification. Specifically, what may appear like a connection when examining

120   particular characters in isolation may actually be due to other unmeasured factors, or

121   because the analysis included a nested clade that exhibits both the focal character plus

122   "something" else (Maddison et al. 2007; Beaulieu and Donoghue 2013; Beaulieu and

123   O'Meara 2014). This particular point is illustrated in Figure 1. Here the true underlying

124   model is one in which state 0 and 1 have identical diversification rates. However, there is

125   some other trait with states A and B, and state B has twice the diversification rate of A.

126   We call this trait a "hidden" trait, because it is not observed in the tip data (though it

127   could be observable if we knew what it was). If state 1 happens to be a prerequisite for

128   evolving state B (or even by chance), all the state 0 branches will have state A, but some

129   branches in state 1 will have state A and some will have state B. Thus state 1 actually

130   takes on two states, $1A$ when the hidden state with higher diversification rate is absent,

131   and $1B$ when the hidden state with higher diversification rate is present. As indicated by

132   the model, transitions to this unmeasured variable produce nested shifts towards higher

133   rates of diversification *within* clades comprised of species in state 1. Necessarily, BiSSE

134   can only infer parameters for characters that we can observe, and since all origins of state

135   1 are lumped together, the model infers state 1 as being associated with significantly

136   higher diversification rates. It is associated with higher diversification, but only due to its

137   association with trait B.

We attempt to solve this problem by deriving an expansion of the BiSSE model

that allows for the inference of these hidden states. For the example in Figure 1, we can

assume that all observations of state 1 are actually ambiguous for being in each of the

possible hidden states, 1*A* (i.e., hidden state absent) or state 1*B* (i.e., hidden state present).

We then include transition rates and parameters for the diversification process associated

with this hidden state. Our model, which we refer to hereafter as the HiSSE model, is

actually a modified form of the MuSSE model (Multi-State Speciation and Extinction;

FitzJohn 2012) that extends BiSSE type analyses to allow for multiple binary characters

or characters with more than two states. Thus, the HiSSE model can, in theory, have a

number of observed states and a number of hidden states (i.e., observed states 0, 1, 2, and

hidden states *A, B, C*, resulting in nine possible state combinations).

Formally, the state space in our model is defined as *o* being the index of the

observed state, $o \in 0,1,...,\alpha$, and *h* as the index of the hidden state, $h \in A,B,...,\beta$. A

lineage at time *t* evolves under diversification rates $\lambda_i = \lambda_{oh}$ and $\mu_i = \mu_{oh}$. Thus, the model

has, in general, $\alpha\beta$ different diversification rate categories. The likelihood $D_{N,i}(t)$ is

proportional to the probability that a lineage in state *i* at time *t* before the present (*t*=0)

evolved the exact branching structure and character states as observed. Changes in $D_{N,i}$

over time along a branch are thus described by the following ordinary differential

equation:

$$\frac{dD_{N,i}(t)}{dt} = -\left(\lambda_i + \mu_i + \sum_{j \neq i} q_{ij}\right)D_{N,i}(t) + 2\lambda_i E_i(t)D_{N,i}(t) + \sum_{j \neq i} q_{ij}D_{N,j}(t) \qquad \text{Eq. (1a)}$$

where $E_i(t)$ is the probability that a lineage in state i at time, t, goes extinct by the present,

and is described by:

$$\frac{dE_i(t)}{dt} = \mu_i - \left(\lambda_i + \mu_i + \sum_{j \neq i} q_{ij}\right) E_i(t) + \lambda_i E_i(t)^2 + \sum_{j \neq i} q_{ij} E_j(t) \qquad \text{Eq. (1b)}$$

These series of equations are modified from Maddison et al. (2007); they are solved

numerically along each edge starting at the tips and moving rootward. The initial

conditions for $D_{N,i}(0)$ are set to 1 when the trait is consistent with the observed data, and

0 otherwise. For example, we would set the probability to 1 for both state 1A and 1B for

all species exhibiting state 1 – in other words, the probability of observing a tip

demonstrating state 1 is 1 if the true underlying state is 1A or 1B (note this could be done

with states 1C, 1D, and so forth in this model, although the current implementation in

software currently only allows a binary observed character with four or fewer total

hidden state combinations). The initial conditions for $E_i(0)$ are all set to zero (i.e., we

observe the tip at the present). Incomplete sampling can be allowed by incorporating a

state-specific or even clade-specific sampling frequency, f, and setting the initial

conditions for $D_{N,i}(0)$ as f_i if the corresponding tip, n, is in state i, and 0 otherwise, and

for 1- f_i for $E_i(0)$ (FitzJohn et al., 2009). We assume the sampling frequency of the hidden

state to be identical to the state with which it is associated (e.g., $f_{1A} = f_{1B}, f_{0A} = f_{0B}$).

179   At each internal node, A, that joins descendant lineages N and M, we multiply the

probabilities of both daughter lineages together with the rate of speciation:

$D_{A,i}(t) = D_{N,i}(t) \, D_{M,i}(t) \, \lambda_i$ Eq. (2)

and the resulting values become the initial conditions for the subtending branch. The

overall likelihood is the product of $D_{R,i}(t)$ calculated at the root. We condition this

likelihood by $(1-E_i(t))^2$, which is the probability that the two descendant lineages of the

root, evolving under the same estimates of speciation and extinction rates, survived to the

present and were sampled (Nee et al. 1994). Finally, we follow the procedure described

by FitzJohn et al. (2009) and weight the overall likelihood by the probability that each

possible state gave rise to the observed data, although other weights, such as assuming

equal probabilities or even "known" fixed probabilities, can also be used. Note that in the

absence of a hidden state, our likelihood calculation reduces exactly to the BiSSE model.

We also note that it is rather straightforward to use this framework to implement a

SSE version of the "precursor" model described by Marazzi et al. (2012) or the "hidden

rates" model (HRM) described by Beaulieu et al. (2013). In the latter case, consider a

HiSSE model with "two rate classes", A and B. We can define speciation and extinction

parameters for states $0A$, $1A$, $0B$, and $1B$, and then define a set of transitions to account

for transitioning between all character state combinations:

$$\mathbf{Q} = \begin{array}{c} 0A \\ 1A \\ 0B \\ 1B \end{array} \left[\begin{array}{cccc} - & q_{0A \to 1A} & q_{0A \to 0B} & 0 \\ q_{1A \to 0A} & - & 0 & q_{1A \to 1B} \\ q_{0B \to 0A} & 0 & - & q_{0B \to 1B} \\ 0 & q_{1B \to 1A} & q_{1B \to 0B} & - \end{array} \right]$$ Eq. (3)

Thus, HiSSE can be used to account for differences in the diversification process while

simultaneously identifying different classes of transition rates within a binary character.

Our implementation allows for this and more complex models, including those that allow

for dual transitions between both the observed trait and the hidden trait (e.g., $q_{0A} \leftrightarrow q_{1B}$).

206   MULTIMODEL INFERENCE AND THE ISSUE OF APPROPRIATE MODELS

207   An important issue was recently raised by Rabosky and Goldberg (2015), who

demonstrated that on empirical trees even traits evolving under a neutral, diversification-

independent model will still tend to be best fit by a BiSSE model. FitzJohn (2012)

discussed a similar issue of MuSSE falsely assigning diversification rate effects to a

character with no such effect. While this behavior is seemingly troubling, it is important

to bear in mind that BiSSE, MuSSE, HiSSE, and any other model of state-dependent

speciation and extinction are *not* models of trait evolution, but rather joint models for the

underlying tree *and* the traits. A trait evolution model like those in Pagel (1994) or

Hansen (1997) maximizes the probability of the observed states at the tips, given the tree

and model – the tree certainly affects the likelihood of the tip data, but that is the only

way it enters the calculation. A trait-based diversification model, on the other hand,

maximizes the probability of the observed states at the tips and the observed tree, given

the model. If a tree violates a single regime birth-death model due to any number of

causes (e.g., mass extinction events, trait-dependent speciation or extinction, maximum

carrying capacity, climate change affecting speciation rates, etc.), then even if the tip data

are perfectly consistent with a simple transition model, the tip data *plus* the tree are not.

In such a case, it should not be surprising that a more complex model will tend to be

chosen over a nested simpler model, particularly if the underlying tree is large enough.

225   Furthermore, as is well known in statistics, rejecting the null model does not

imply that the alternative model is true. It simply means that the alternative model fits

better. This will often be the case when looking at models in any complex system where

the true model may not be one of the included models. For example, Rabosky and

Goldberg (2015) showed, among several other empirical examples, that binary characters

simulated under a model with each character having no effect on diversification rate on

an empirical tree of cetaceans (i.e., whales, dolphins, and relatives) almost always

rejected the supposed null model. Though presented as a Type I error (i.e., incorrectly

rejecting a true null), it is not. While the chosen character model is wrong, the cetacean

tree is almost certainly not evolving with a single speciation and extinction rate for the

entire clade (see Slater et al. 2010; Morlon et al. 2011). In other words, BiSSE is correct

in saying that the simple model is not correct, but it is very wrong in assigning rate

differences to the simulated traits. Nonetheless, even when presented with two bad

models, where one assumes constant rates and the other assumes rates are changing rate

exactly with character states, BiSSE must still select which model fits the data better than

the other.

A simple example may further illustrate this point. Imagine simulating data under

a hypothetical model, called "A", and then comparing the fit of this model against

another hypothetical model, called "B". If model A is a "null model", but we end up

choosing model B, then this would an instance of a "Type I" error – we are incorrectly

rejecting the null. Now, imagine simulating data under another distinct model, "C", but,

like before, we only evaluate the fit of model A and model B. Would choosing model B

be considered a Type I error? Of course not, because neither model is correct. So clearly

the term Type I error (or even Type II error) is outside the region of having any valid

meaning in this context.

The reality is that it will be always be a problem comparing any two models and

taking the accepted one as the "truth" given a complex underlying reality. Regardless of

whether the behavior they observe is properly called Type I error or not, this is the key

point of Rabosky and Goldberg (2015). When comparing a simpler model to a more

complex model, merely rejecting the simple one does not mean we should accept all the

assumptions and interpretations of the alternate one. It is a simple exercise, for example,

to simulate a tree using the model of Rabosky (2010) and then fit alternate models such

as age-dependent diversification (Alexander et al. 2015) or logistic growth (Rabosky and

Lovette 2008) and compare them to a constant rate model; in most simulated trees, the

alternate model is chosen (results not shown), even though the processes each assumes

were not used to generate the tree.

One potential way forward, at least in regards to the issues Rabosky and Goldberg

(2015) pointed out for SSE models, is to provide equally complex models that fit

different biological explanations. Take, for instance, the likely complex diversification

history of cetaceans. As Rabsosky and Goldberg (2015) demonstrate, when evaluating

neutral data simulated on the cetacean tree using a simple, equal diversification rates

model, and a complex one with trait-dependent diversification, the data choose the

complex one. However, what if there were a complex model that also allowed different

diversification rates on the tree, but was completely independent of the diversification

process? Then, if the true model were trait-correlated diversification, a model of trait-

dependent diversification would be chosen, but if the tree contained some additional

complexity, this new trait-independent model might also be chosen some proportion of

the time. Again, it bears repeating: *SSE models are not models of trait evolution, but*

rather, a joint modeling framework for the underlying tree and the trait. In the case of

cetaceans, or really any empirical tree, we do not know the true underlying model that

generated the tree, so picking between models of equal rates, trait-dependent, or trait-

independent diversification some percentage of the time in no way represents a kind of

Type I error. Furthermore, if statistics are done with a focus on parameter estimates rather

than model rejection, even if there were some weight for trait-dependent rates, there

would also be substantial weight for trait-independent rates as well, and so the average

rates across these models should tend to be similar across the possible character states.

281   Here we propose two character independent diversification (CID) models that are

devised to equal the complexity with respect to the number of parameters for the

diversification process (i.e., same number of free speciation and extinction rates under the

Maddison et al. (2007) parameterization) as a general BiSSE or HiSSE model, but

without actually linking them to the observed traits. Thus, these models explicitly assume

that the evolution of a binary character is independent of the diversification process

without forcing the diversification process to be constant across the entire tree, which is

the normal CID used in these types of analyses. The first kind of model, which we refer

to as "CID-2", contains four diversification process parameters that account for trait-

dependent diversification solely on the two states of an unobserved, hidden trait (e.g., for

speciation rates, $\lambda_{0A} = \lambda_{1A}$, $\lambda_{0B} = \lambda_{1B}$). In this way, CID-2 contains the same amount of

complexity in terms of diversification as a BiSSE model. As with BiSSE or HiSSE, these

models include the possibility of a variety of constraints for transition rates. These

transition rates can be general by allowing them all to be freely estimated, or simplified in

various ways such as assuming they are all equal. The second kind of model, which we

refer to as "CID-4" contains the same number of diversification parameters as in the

general HiSSE model that are linked across four hidden states (e.g., for speciation rates,

$\lambda_{0A} = \lambda_{1A}$, $\lambda_{0B} = \lambda_{1B}$, $\lambda_{0C} = \lambda_{1C}$, $\lambda_{0D} = \lambda_{1D}$). The transition rates under this model are set up to

account for transitions between the four different rate categories, as well as between the

states of the binary character and, in matrix form, they are set up as follows:

302
$$
\mathbf{Q} = \begin{array}{c} 0A \\ 0B \\ 0C \\ 0D \\ 1A \\ 1B \\ 1C \\ 1D \end{array}
\begin{bmatrix}
- & q_{0A\to0B} & q_{0A\to0C} & q_{0A\to0D} & q_{0A\to1A} & 0 & 0 & 0 \\
q_{0B\to0A} & - & q_{0A\to0C} & q_{0B\to0D} & 0 & q_{0B\to1B} & 0 & 0 \\
q_{0C\to0A} & q_{0C\to0B} & - & q_{0C\to0D} & 0 & 0 & q_{0C\to1C} & 0 \\
q_{0D\to0A} & q_{0D\to0B} & q_{0D\to0C} & - & 0 & 0 & 0 & q_{0D\to1D} \\
q_{1A\to0A} & 0 & 0 & 0 & - & q_{1A\to1B} & q_{1A\to1C} & q_{1A\to1D} \\
0 & q_{1B\to0B} & 0 & 0 & q_{1B\to1A} & - & q_{1B\to1C} & q_{1B\to1D} \\
0 & 0 & q_{1C\to0C} & 0 & q_{1C\to1A} & q_{1C\to1B} & - & q_{1C\to1D} \\
0 & 0 & 0 & q_{1D\to0D} & q_{1D\to1A} & q_{1D\to1B} & q_{1D\to1C} & -
\end{bmatrix}
$$

303   Eq. (4)

To simplify the number of transitions in the model there are two natural assumptions:

assume either all transition rates are equal, or assume there are three distinct transition

rates: one rate describing transitions among the different hidden states (i.e., the rates in

columns and rows 1-4, and columns and rows 5-8), and two rates for transitions between

the observed character states (i.e., one rate for columns 5-8, rows 1-4, and one rate for

columns 1-4, and rows 5-8). These or other constraints can be used as part of the CID

family of models.

312   IMPLEMENTATION

We implemented the above models in the R package "*hisse*" available through

CRAN. As input all that *hisse* requires is a phylogeny with branch lengths and a data file

that contains the observed states of a binary character. Note that this is an entirely new

implementation, not a fork of the existing *diversitree* package, as we employ modified

optimization procedures and model configurations. For example, rather than optimizing λ_i

and μ_i separately, *hisse* optimizes transformations of these variables: we let $\tau_i = \lambda_i + \mu_i$

define "net turnover", and we let $\varepsilon_i = \mu_i / \lambda_i$ define the extinction fraction. This

reparameterization alleviates problems associated with over-fitting when λ_i and μ_i are

highly correlated, but both matter in explaining the diversity pattern (e.g., Goldberg et al.

2010; Beaulieu and Donoghue 2013). With empirical data we often see good estimates

for diversification rate but correlations for birth and death rate estimates. For example,

Beaulieu and Donoghue (2013) (their Figure S2) showed that the confidence region for

birth and death rates tightly follows a diagonal line, with different characters having lines

of the same slope but different intercepts. This leads to a behavior where looking at the

confidence or credibility intervals for birth or death show overlap between the characters

but intervals for diversification rates do not overlap between characters. However, the

only way to fit this in the common parameterization is with two birth rates and two death

rates. Reparameterizing, as we do, allows us to have the same turnover rate for both states

but estimate different diversification rates, resulting in a less complex, but better fitting,

model. Thus, users specify configurations of models that variously fix τ_i and ε_i. Note that

estimates of τ_i and ε_i can be easily backtransformed to reflect estimates of λ_i and μ_i by

$\lambda_i = \tau_i / (1 + \varepsilon_i),$ Eq. (5a)

$\mu_i = (\tau_i \, \varepsilon_i) / (1 + \varepsilon_i),$ Eq. (5b)

which can be used to derive other measures of diversification process parameters, such as

net diversification (λ_i-μ_i). One of the *diversitree* authors (Sally Otto, personal

communication) noted that Hardy and Otto (2014) tested a hidden model that only

affected transition rates within the MuSSE framework. Note that we have not directly

compared the two implementations, though we have confirmed that MuSSE and HiSSE

return the same likelihood when all states are observed.

The *hisse* package assigns the probability of each ancestral state to an internal

node using marginal ancestral state reconstruction (Yang et al.1995; Schluter et al., 1997)

under the chosen diversification with trait evolution model. These probabilities can be

used to "paint" areas of a phylogeny where the diversification rate has increased or

decreased due to some unmeasured character. In our implementation the marginal

probability of state *i* for a focal node is the overall likelihood of the tree and data when

the state of the focal node is fixed in state *i*. Note that the likeliest tip states can also be

estimated: we observe state 1, but the underlying state could either be state 1*A* or 1*B*.

Thus, for any given node or tip we traverse the entire tree as many times as there are

states in the model. As the size of the tree grows, however, these repeated tree traversals

can slow the calculation down considerably. For this reason, we also include a function

that allows the marginal calculation to be conducted in parallel across any number of

independent computer processors.

357   SIMULATIONS

We evaluated the performance of the HiSSE model by simulating trees and

character states under various scenarios and then estimating the fit and bias of the

inferred rates from these trees. Our initial simulations first tested scenarios of the single

hidden state situation described in Figure 1; the known parameters for each scenario are

described in detail in Table 1. We included BiSSE scenarios to test whether we could

correctly conclude that there was no support for a HiSSE model in the absence of a

hidden state in the generating model. For each of these scenarios, trees and trait data were

simulated using *diversitree* (FitzJohn 2012) to contain 50, 100, 200, or 400 species, with

100 replicates for each taxon set. When the generating model included a hidden state, we

simulated trees that could transition between three possible states: 0, 1, or 2. After each

simulation replicate was completed, we created the hidden state by simply switching the

state of all tips observed in state 2 to be in state 1.

Each simulated data set was evaluated under the generating model, as well as 13

additional models that variously added, removed, or constrained certain parameters. The

entire model set is described in Table 1. For all models under a given scenario, model fit

was assessed by calculating the average Akaike weight (w_i), which represents the relative

likelihood that model i is the best model given a set of models (Burnham and Anderson

2002). We also calculated a null expectation of the Akaike weight across our model set,

as the average Akaike weight assuming an equal likelihood across all models. Thus, our

null expectation is based solely on the penalty term in the AIC calculation – in the

absence of information from the model, we would expect to see these weights returned,

rather than equal weights for all models. Moreover, since BiSSE is nested within HiSSE,

they could return the same likelihood, so even with infinite amounts of data the weight of

the HiSSE model should drop to this null expectation, but, counterintuitively, not to zero.

We also conducted a second set of simulations that specifically evaluated the

performance of the general HiSSE model and our two CID models of CID diversification.

Our goal was to test how much model weight the trait-independent models exhibited

under scenarios of trait-dependent diversification. We were concerned at the outset that

our CID models would always fit at least as well, if not better, than a trait-dependent

BiSSE or HiSSE model. That is to say, these models were constructed such that they are

not constrained by character states and can assign rates wherever they want in order to

maximize the likelihood. Table 2 describes the known parameters for the three scenarios

evaluated. Again we relied on *diversitree* to simulate trees and trait data that contained

400 species, repeated 100 times, with transitions allowed between four possible states, 0,

1, 2, or 3. After each simulation replicate was completed, we created the hidden state by

simply switching the state of all tips observed in state 2 to be in state 0, and all tips in

state 3 to be in state 1.

We also included a scenario that was designed to test whether the general HiSSE

model is immune to empirical issues of spurious assignment of importance to state

combinations that have no actual effect on diversification. In other words, is HiSSE still

favored in situations where the trees and traits evolved under a very different model than

the one used for inference? Here we generated trees containing 400 taxa using code from

Rabosky (2010). A symmetric Markov model for trait evolution alone (no influence by

diversification) was used to simulate binary traits on this tree (using the R package

geiger; Harmon et al. 2008; Pennell et al. 2014). The Rabosky (2010) model was used, as

it is very different from the model assumed by BiSSE/HiSSE; speciation rates evolve

gradually on branches, rather than moving discretely between distinct levels based on a

trait (hidden or not). Though the rate change is gradual under the Rabosky (2010) model,

the speciation rate does not evolve under a Brownian, Ornstein-Uhlenbeck, or similar

processes, but in a heterogeneous way that depends on the timing of speciation events

(Beaulieu and O'Meara 2015). Thus, as with many empirical data sets, this

diversification model is quite different from HiSSE's model, providing a difficult

challenge. The Rabosky (2010) model has also been very influential in affecting

biologists' attitudes towards estimating extinction rates, and so we include it as a semi-

realistic "worst-case" scenario. Each simulated data set was evaluated under multiple

models that variously added, removed, or constrained certain parameters, with model fit

again being assessed by calculating the average Akaike weight (w_i). The entire model set

is described in Table 2. For simplicity, all models in this set assumed equal transition

rates.

In regards to the parameter estimates in all our simulations, comparisons between

the model-average of the parameters against the known parameters provided an

assessment of the bias and precision of the inferred parameters. However, rather than

averaging parameters across the entire model set, we only averaged across models that

included similar parameters. For example, when estimating the bias in the HiSSE

scenarios we only model-averaged parameters for models that included the hidden state.

This required reformulating the Akaike weights to reflect the truncated model set.

Finally, we also assessed the reliability of the ancestral state reconstructions by

comparing the true node states from each simulated tree to the marginal probabilities

calculated from the model-averaged parameter estimates.

Our main focus is estimating parameters well, but this is aided by picking the true

model with high weight when it is in the set (though, of course, for real data the true

model is always more complex than any examined one). From a model comparison

perspective, our first set of simulations indicated that data sets that lack hidden states

could generally be distinguished from those that do, especially with larger data sets

(Table 1). When the generating model is a BiSSE model with only two observable states,

there were low levels of support for all seven HiSSE models. In fact, as sample size

increased, the average Akaike weight of the HiSSE models converged towards the null

Akaike weight based only on the penalty term (Table 1). When evaluating data sets that

included a hidden state, the ability to correctly favor a HiSSE model over any of the

BiSSE models depended not just on the size of the data set, but also the underlying

generating model. For example, when the generating model assumed a hidden state with

higher speciation rates, data sets that contained 200 or more taxa were required to provide

strong evidence for a HiSSE model that varied the turnover rate (Table 1). However,

when the main effect of the generating model assumed lower extinction fractions for the

hidden state there remained strong support for a BiSSE model that assumed both equal

turnover rates and extinction fractions. Interestingly, when we simulated under a HiSSE

scenario that combines the processes of higher speciation rates and asymmetrical

transition rates, the issue of incorrectly favoring a BiSSE model disappears (Table 1).

We also found that HiSSE was able to distinguish between generating models that

assumed diversification rates differences are trait-dependent (e.g., when speciation is

two-fold greater for state $1B$), or when the diversification rate difference is trait-

independent and due to simply to the presence of a hidden trait only (e.g., when

speciation is two-fold greater for hidden state *B*) (Table 2; Table S1). And, contrary to

our concerns, when the generating model assumed some form of trait-dependent

diversification neither the CID-2 nor the CID-4 model had much support.

For the "worst-case" scenario, which simulated a neutral binary character along

trees generated from a complex heterogeneous rate branching process, it clearly falls into

the zone where BiSSE has unwelcome behavior (94% of data sets favored a BiSSE model

over a single-rate diversification model, and the average Akaike weight for state-

dependent models across data sets was 92.0%; see Table S1). Thus, these simulation

conditions create difficult data sets of the sort used by Rabosky and Goldberg (2015).

The addition of merely the CID-2 model dramatically improved performance, where the

average weight for the BiSSE models went from 92.0% to 1.3%. When the full HiSSE

model set was examined, we found that the CID-4 trait independent model had the

highest support on average across the entire set of models (Table 2; Table S1). However,

if we combine support for both this CID model and the CID-2 model, they cumulatively

account for 70% of the total model weight. Both of these models are independent of any

particular character state, but assume shifts among distinct levels, and therefore more

closely resemble the conditions by which these data sets were generated.

We note, however, that there was some support for a trait-dependent HiSSE

model with the same number of diversification parameters (Table 2; Table S1). And, in

spite of the Akaike weight across all data sets showing greater support for the CID-4

model on average, 29% of the individual data sets would favor some form of the general

HiSSE model (Table S1). However, there was only a 1.02% difference in the mean

diversification rate between states 0 and 1 (though, focusing on just the set of trees for

which a HiSSE model was best, the percent difference is 16% – still relatively small for

diversification studies, but clearly not as good). Taken together, these results indicate that

in these "worst-case" trait-independent scenarios, likely encountered in many empirical

data sets, the inclusion of the CID models can attenuate the issue of spuriously accepting

trait-dependent diversification. The addition of models that allow some dependence with

the observed traits and some with hidden traits, the rates of falsely assessing some state-

dependence somewhat increases, but with the benefit of being able to better detect hidden

effects when they are true (see earlier simulation results).

485   ESTIMATING MODEL PARAMETERS

It is important to point out that in all simulation scenarios we never recovered

strong support for a model consistent with the model that actually generated the data.

That is, whether we vary the speciation rate or the extinction rate, which affects both net

turnover and extinction fraction, we always found support for a simplified model that

either allowed τ to vary or ε to vary, but not both. This is likely a consequence of the

uncertainty and upward biases in estimates of extinction fraction (Fig. 2-4), making it

difficult for the model to infer multiple extinction fractions, at least with smaller data

sets.

It is well known that there can be difficulties, generally, in obtaining precise

estimates of the extinction rates (i.e., μ_i) under the BiSSE framework (Maddison et al.

2007). Backtransforming estimates of net turnover and extinction fraction shows that

HiSSE suffers the same precision issues in regards to the rate of extinction (Fig. 3, Fig.

S2). Recently, it was reported that biases in the tip state ratios could also impact all

parameter estimates (see Davis et al. 2013). Our simulations did not specifically test

issues related to tip state biases. However, we did find that with HiSSE, the precision in

extinction rate estimates remains fairly low regardless of the ratio of the states at the tips,

especially when estimating extinction for the hidden state (Fig. 3). Interestingly, the lack

of precision for extinction rates seems to have a relatively minor impact on estimates of

net turnover or net diversification (Fig. 2-4; Fig. S1), though there is a general bias for

net diversification to be underestimated as a consequence of inflated extinction rates.

Nevertheless, it appears HiSSE correctly and qualitatively distinguishes differences in

diversification among the various states in the model. And, as the number of species

increases, the trajectory of the downward bias in net diversification suggests it will

eventually disappear, and rate differences can be distinguished even if the differences are

trivial (e.g., Fig. 2c).

For parameter estimates from the simulations testing both the CID models and the

general HiSSE model we found very similar results as those described above, in that the

model-averaged net turnover and net diversification parameters largely resembled the

generating model (Fig. 4, S4). However, similar to the first set of simulations, there is a

large amount of uncertainty and upward biases in estimates of extinction fraction. Thus,

HiSSE generally has very low power in distinguishing multiple extinction fractions

regardless of the number of states included in the model.

One issue of concern from our first set of simulations is that transition rates are

almost always overestimated. This behavior appears unique to the HiSSE model in our

simulations (Fig. 2), given that when evaluating data sets under BiSSE scenarios,

transition rates are estimated reasonably well (Fig. S3). One suggestion from Rich

FitzJohn (personal communication) is that this can occur when some states are present in

low frequency, and since HiSSE has more states than BiSSE, it is likely that many state

combinations are in very low frequencies. There are also relatively large confidence

intervals surrounding each of the transition rate estimates that naturally favor models that

assume equal transition rates, which should be reflected in the model-averaged rates.

Indeed, as in the case of the HiSSE scenario that assumed pronounced differences in the

transition rates, even 800 taxa was still not enough to unequivocally reject models that

assumed equal transition rates (Table 1). We also examined the impact each parameter

had on a fixed set of equilibrium frequencies, by randomly sampling sets of values and

retaining those that estimated the same frequency within a small measure of error (see

Supplemental Materials). The proportional range of values for the transition rates were

more than double those found for the speciation rate, indicating that state frequencies are

fairly resilient to changes in the transition rates. Taken together, estimating unique

transition rates (and to some extent the rate of extinction) appears to be a difficult

problem, which is not made any easier by HiSSE's increase in state space. That is to say,

HiSSE requires more from the data by including additional parameters without providing

any more observable information. It is likely that in many cases far larger data sets with

many more state origins than the ones we have generated here may be required to

adequately estimate these particular parameters.

Finally, in regards to inference of ancestral states from the model-averaged

parameter estimates, the simulations indicate that HiSSE correctly identifies and locates

regions of the tree where supposed diversification rate differences have taken place (Fig.

5). The degree of reliability does, of course, depend on the size of the data set. For the

HiSSE scenario that assumed a doubled speciation rate for state 1*B*, for example, data

sets comprised of 50 taxa, 84.1% of the nodes, on average, have the state correctly

inferred, and data sets comprised of 400 taxa 92.4% are correct. However, we note that

there is also a general tendency for HiSSE to infer high marginal probabilities for the

incorrect state (e.g., Fig. 5), which could provide misleadingly confident state

reconstructions.

551   MODEL REJECTION PROPERTIES

552   We have strongly advocated (e.g., Beaulieu et al. 2012), as have many others

(e.g., Anderson et al. 2000; Nickerson 2000; Nakagawa and Cuthill 2007), that parameter

estimation is more important for understanding biology than is rejecting models. Model

selection is part of this, of course, but it is more appropriate to examine a set of models

and average or integrate their results, either through the use of information theory (e.g.,

Burnham and Anderson 2002), as we do here, or Bayesian approaches (e.g., Huelsenbeck

et al. 2004). However, there are clearly still many studies whose main focus is rejecting

trivial null models. The use of AIC or Akaike weights are inappropriate for these

purposes (Burnham and Anderson 2002), despite being used frequently. Here, we

examine how HiSSE and its various models perform in this way, especially in relation to

the situation highlighted by Rabosky & Goldberg (2015), where a species tree evolves

under a complex, unknown diversification process but traits evolve under a simple model

independent of diversification parameters (e.g., our "worst-case" scenario).

565   Even though examining ΔAIC values are not recommended for model rejection,

the rule of thumb from Burnham and Anderson (2002) remains popular for this use in

phylogenetics: 0-2 means a model has substantial support, 4-7 means considerably less

support, and >10 means essentially no support. Using these guidelines, we examined two

of our simulations, the CID-2 and our "worst-case" scenario, both of which assume trait-

independent diversification model (see Table S1). If we were to compare the fit of a

constant birth-death BiSSE with a BiSSE model that assumes trait-dependent

diversification, as Rabosky and Goldberg (2015) did, we found that when the true

generating model was our CID-2 model, 38% of the time the BiSSE model had

substantial support; in our "worst-case" scenario, 80% of the time BiSSE had substantial

support. However, if we were to add just our CID-2 model into the set, in both cases the

number of simulated data sets that show substantial support for the BiSSE case drop

dramatically: just 7% in the CID-2 simulations, and just 1% in the "worst-case" scenario.

If we examine the full set, which also includes CID-4 and the HiSSE model, 10% of

simulated data sets that favor a trait-dependent scenario for CID-2, and 16% of the time

under the "worst-case" scenario (Table S1).

We also conducted similar tests by simulating neutral characters on two empirical

trees – the cetacean tree used by Rabosky and Goldberg (2015), and our own empirical

tree (see below) – and found very similar behavior (see Table S2). More importantly, for

parameter estimation, using a set of models and doing parameter estimation using a

weighted average results in largely accurate inferences. For the 87-taxon cetacean tree,

across simulated trees the model-averaged diversification rate in state 0 was within 10%

of the rate for state 1 in 87% of the simulations (and we expect substantial uncertainty in

these estimates); for Dipsidae, in 95% of the simulations the estimated diversification rate

for state 0 was within 0.5% of the rate for state 1. In other words, even if a trait-

dependent model were chosen, a careful biologist who looks at parameter estimates

would be unlikely to find a rate difference of half a percent biologically significant.

592   In summary, we have gone from 94% of the time choosing the "wrong" model, to

anywhere from 7-16% when the model set includes any number of our HiSSE models. Of

course, if the goal is to simply find the best model, the rather marked improvement

afforded by our HiSSE framework may remain somewhat unsatisfying. Future attention

could, therefore, be paid to determining the appropriate ΔAIC that would constitute

"substantial support" when using HiSSE, as opposed to simply using a ΔAIC<2 cutoff, as

we have done here. Nevertheless, if the goal is to understand the potential biological

implications provided by the parameters estimated from these models, even though the

"best" model may at times be incorrect, taking into account model uncertainty when

summarizing the parameter estimates will ameliorate spurious interpretations of trait-

dependent diversification when it does not exist (e.g., Fig. 4, Fig. S5).

603   THE EVOLUTION OF ACHENE FRUITS

604   The development of this model was inspired by results from recent empirical

work that applied BiSSE to understand the macroevolutionary consequences of evolving

particular fruit types within a large flowering plant clade (i.e., campanulids; Beaulieu and

Donoghue 2013). This study investigated whether diversification rates differences could

explain why more than 80% of campanulid species exhibit fruits that are indehiscent (i.e.,

do not open mechanically), dry, and contain only a single seed. From a terminological

standpoint, these fruits were broadly referred to as "achene" or "achene-like" to unify the

various terms used to identify the same basic fruit character configuration (e.g.,

"cypselas" of Asteraceae – sunflowers and their relatives – or the single-seeded

"mericarps" of Apiaceae – carrots and their relatives). According to the BiSSE model, the

preponderance of achene fruits within campanulids can be explained by strong

differences in diversification rates, with achene lineages having a rate that was roughly

three times higher than non-achene lineages.

While these results are seemingly straightforward, they are complicated by the

fact that the correlation between net diversification rates and the achene character state

differed among the major campanulid lineages and was driven entirely by the inclusion of

Asterales clade (Beaulieu and Donoghue 2013). Within Apiales and Dipsacales, the two

remaining major achene-bearing clades, the diversification rate differences were not

significant. However, in both these clades there were qualitative differences in the

predicted direction, likely as a consequence of one or more shifts in diversification nested

within one of the major achene-bearing clades. Together, these point to a more complex

scenario for the interaction between achene fruits and diversification patterns that is not

being adequately explained by BiSSE.

We illustrate an empirical application of HiSSE by examining the Dipsidae

(Paracryphiales+Dipsacales; Tank and Donoghue 2010) portion of the achene data set of

Beaulieu and Donoghue (2013). Specifically, we used HiSSE to locate and "paint"

potential areas within Dipsidae that may be inflating the estimates of net diversification

rates for achene lineages as a whole. We modified the original data set of Beaulieu and

Donoghue (2013) in three important ways. First, we re-estimated the original molecular

branch lengths using PAUP* (Swofford 2000), as opposed to relying on the branch

lengths from the original RAxML (Stamatakis 2006) analysis, because PAUP* provides

better optimization precision. Second, the molecular branch lengths were re-scaled in

units of time using *treePL* (Smith and O'Meara 2012), an implementation of the

penalized likelihood dating method of Sanderson (2002) specifically designed for large

trees. We applied the same temporal constraints for Dipsidae as in the original Beaulieu

and Donoghue (2013) study, and used cross-validation to determine the smoothing value

that best predicted the rates of terminal branches pruned from the tree. Third, we

conservatively removed various taxa of dubious taxonomic distinction, taxa considered

varietals or subspecies of a species already contained within the tree, and tips that *treePL*

assigned very short branch lengths (i.e., <1.0 Myr) – all of which have the tendency to

negatively impact accuracy of estimating diversification rates (though in this case,

rerunning the analyses including such tips did not have a qualitative effect on the results).

The exclusion of these taxa resulted in a data set comprised of 417 species from the

original 457.

648   We fit 24 different models to the achene data set for Dipsidae. Four of these

models corresponded to BiSSE models that either removed or constrained particular

parameters, 16 corresponded to various HiSSE models that assumed a hidden state

associated with both the observed states (i.e., non-achene+, achene+), and four

corresponded to various forms of our trait-independent models (i.e., CID-2, CID-4). In all

cases, we incorporated a unique sampling frequency scheme to the model. Rather than

assuming random sampling for the entire tree (see above), we included the sampling

frequency for each major clade included in the tree (see Table S3; note, that the results

are qualitatively similar regardless of sampling scheme used). In order to generate a

measure of confidence for parameters estimated under a given model, we implemented an

"adaptive search" procedure that provides an estimate of the parameter space that is some

pre-defined likelihood distance (e.g., 2 lnL units) from the maximum likelihood estimate

(MLE), which follows from Edwards (1992). We also took into account both state

uncertainty and uncertainty in the model when "painting" diversification rates across the

tree. Our procedure first calculated a weighted average of the likeliest state and rate

combination for every node and tip for each model in the set, using the marginal

probability as the weights, which were then averaged across all models using the Akaike

weights. All analyses were carried out in *hisse*.

The best model, based on Akaike weights, was a relatively simple HiSSE model

with regards to the number of free parameters it contained (Table 3). The model suggests

character-dependent diversification with fruit type, where only rates for non-achene+ and

achene+ are allowed to be free, and where transitions between state $0B$ and $1B$ were

disallowed. This model had a pronounced improvement over the set of BiSSE models,

where none had an Akaike weight that was greater than 0.001. However, before

describing the parameter estimates of the best model, we note that, with a modified tree

and character set, the parameter estimates under the BiSSE models were different from

those reported by Beaulieu and Donoghue (2013). Specifically, the higher diversification

rates estimated for achene lineages (r_{achene}=0.148, support range: [0.121,0.161])

compared to non-achene lineages ($r_{non-achene}$=0.065, support range: [0.0498,0.083]) were

indeed significant based on a sampling of points falling within 2lnL units away from the

MLE. The parameters estimated under the HiSSE model, on the other hand, suggest a

more nuanced interpretation of this result. The higher diversification rates of clades

bearing achenes as a whole is likely due to a hidden state nested within some of these

lineages that is associated with exceptionally high diversification rates ($r_{achene+}$=0.199,

support region: [0.179,0.221]). In fact, the model suggests that achene lineages not

associated with the high diversification hidden state have a diversification rate that is

indistinguishable from the non-achene diversification rate ($r_{\text{non-achene-}}= r_{\text{achene-}}$=0.059,

support range: [0.049,0.068]). Non-achene lineages associated with the high

diversification hidden state also show elevated diversification rates ($r_{\text{non-achene+}}$=0.158,

support region: [0.138,0.185]), relative to the rate of the non-achene state with the other

hidden state, suggesting strong rate heterogeneity even in Dipsidae lineages that bear

other fruit types.

Character reconstructions identified two transitions to the hidden fast state in

achene lineages, and thus a higher diversification rate: one shift occurred along the stem

leading to crown Dipsacaceae and the other occurred along the stem leading to "core

Valerianaceae" (the most inclusive clade that excludes *Patrinia*, *Nardostachys*, and

Valeriana celtica) (Fig. 6). It is important to note that in the case of non-achene lineages,

the model identified four shifts to the hidden fast state – one at the base of Caprifolieae

(*Lonicera*, *Symphoricarpus*, *Leycesteria*, and *Triosteum*), and three within *Viburnum*

(Fig. 6). The several shifts detected in *Viburnum* are noteworthy in that they correspond

almost exactly with the inferred shifts detected in a more focused study of the genus that

applied various models of diversification (Spriggs et al. 2015).

In regards to achene lineages, the general location of the inferred shifts is

intriguing, as they appear to coincide with specific clades that exhibit specialized

structures related to achene dispersal. In Dipsacaceae, for example, there is tremendous

diversity in the shape of the "epicalyx", a tubular structure that subtends the calyx and

encloses the ovary (Donoghue et al. 2003; Carlson et al. 2009), which is often associated

with elaborated structures (e.g., "wings", "pappus-like" bristles) that accompany their

achene fruits. Interestingly, many of these same general forms are observed throughout

core Valerianaceae, although they arise from modifications to the calyx (Donoghue et al.

2003; Jacobs et al. 2010). While the significance of these structures is thought to improve

protection, germination, and dispersal of the seed, we emphasize that, at this stage, it is

difficult to confidently rule out other important factors, such important biogeographic

movements due to increased dispersal abilities (i.e., "key opportunities" *sensu* Moore and

Donoghue 2007), or even purely genetic changes, such as gene and genome duplications

(Hildago et al. 2010; Carlson et al. 2011). But, we can at least confidently conclude that

the achene by itself is likely not a strong correlate of diversity patterns within Dipsidae.

715   DISCUSSION

Progress in biology comes from confronting reality with our hypotheses and

either confirming that our view of the world is correct, or, more excitingly, finding out

that we still have a lot left to discover. With studies of diversification, we have mostly

been limited to doing the former – we have an idea about a trait that may affect

diversification rates based on intuition about its potential effect (e.g., an achene might

allow greater dispersal, and thus easier colonization of new habitats to form new species)

as well as some knowledge of its distribution (e.g., some very large plant clades have

achenes) and then run an analysis. Typical outcomes are yes, there is a difference in

diversification in the way we expected, or there is no difference but maybe we just lack

the necessary power. In any case, chances are we are at least vaguely correct that the trait

we think credibly has a mechanism for increasing diversification rate actually has an

effect (at least once there is enough power), or, less compellingly, clades we identify for

such a test from eyeballing the data return a significant result despite no underlying

reality.

Surprise is a necessary part of discovery that, to put it bluntly, has been relatively

lacking in trait-dependent diversification studies until now. With HiSSE we can still test

our intuitions about a particular character, but we can also discover that rates seem to be

driven by some unknown and unmeasured character state, allowing the data to help us

generate new hypotheses – Is the diversification rate correlated with the achene, or is the

achene simply a necessary precursor to some other trait that is more likely to be driving

diversification? This lets us go from a scenario where we simply reject trivial nulls, such

as whether diversification rates of clades with and without some focal trait are precisely

equal, to being potentially surprised by the results – No, it is not the achene *per se*, but it

is something else nested within these particular clades in addition to the achene fruit type.

Currently, diversification models are divided between those that look at one or

more focal traits only, integrating over any other factors (e.g., BiSSE, Maddison et al.

2007; BiSSE-ness, Magnuson-Ford and Otto, 2012; ClaSSE, Goldberg and Igic, 2012;

sister group comparisons, Mitter et al. 1988), and those that fit rates to trees but ignore

trait information altogether (e.g., MEDUSA, Alfaro et al. 2009; BAMM, Rabosky 2014).

Our HiSSE framework spans this range. If the true model is strictly trait-dependent

diversification, it can detect this (Fig. 1-5), as a BiSSE analysis would. If the true model

is rates varying due to some other unexamined factor (a physical trait, a property of the

environment, etc.) HiSSE recovers this too, as in the case of achene evolution in the

Dipsidae clade (Fig. 6). Uniquely, HiSSE can also give an intermediate answer where a

focal trait explains some but not all of the diversification difference.

The HiSSE framework also addresses some of the recent important criticisms

levied against state speciation and extinction models (Rabosky and Goldberg 2015).

Specifically, our method no longer requires the assumption that focal character states are

associated with diversification rate differences. Instead, it allows this assumption to be

explored as part of a more flexible overall model, as opposed to relying on separate tests

for uncovering character-dependent and CID rates of diversification (e.g., Beaulieu and

Donoghue 2013; Weber and Agrawal 2014; Spriggs et al. 2015). Including our models of

trait-independent diversification in a set of alternative trait-dependent models should also

alleviate concerns of spurious assignments of diversification rate differences between

observed character states in cases where trees are evolving separately from the focal trait

(c.f., Rabosky and Goldberg 2015). These CID diversification models are designed

specifically to be as complex as competing BiSSE or HiSSE models, which provides a

fairer comparison over more trivial equal-rate "nulls."

We do, however, highlight one important statistical concern in which HiSSE

requires significant caution in its use. This involves its indifference to number of changes

(c.f., Maddison and FitzJohn 2014). As with BiSSE, we should find it more credible that

a particular character state enhances diversification rates if we see a rate increase in each

of the 10 times it evolves than if it evolved just once but had the same magnitude of rate

increase. A good solution to this problem has not yet been proposed, and we urge a

healthy skepticism of any result based on a trait that has evolved only a few times.

There are other practical concerns in empirical applications of the HiSSE model.

While it could be used over a set of trees (i.e., bootstrap or Bayesian post-burnin tree

samples), the model assumes that the branch lengths, topology, and states are known

without error. Certain kinds of phylogenetic errors, such as terminal branch lengths that

are too long (as may occur with sequencing errors in the data used to make the tree) can

result in particular biases in estimates of speciation and/or extinction (also see Beaulieu

and O'Meara 2015). Similarly, if one clade were reconstructed to be younger than it

actually is, due to a substitution rate slowdown caused by some other trait (e.g., life-

history; Smith and Donoghue 2008), it could be interpreted as having a faster

diversification rate, perhaps even inferred to have its own hidden state. For trees that

come from Bayesian dating analyses, whether a birth-death or Yule prior was used may

affect results unless the data is strong enough to overwhelm the prior (Condamine et al.

2015). Future simulation studies will focus on better understanding the impact that these

and other branch length error scenarios can have on the interpretation and estimation of

various model parameters. Also, we point out that the HiSSE model assumes discrete

characters, whether they be hidden or observed, but it could be that a continuous

parameter is the cause of a diversification rate difference (e.g., perhaps extinction risk

varies inversely with mean of the dispersal kernel). However, this may only enter the

model as an unseen discrete character, perhaps corresponding to low and high values of

the continuous character.

Our HiSSE model is part of a long tradition in comparative methods of

identifying, and then addressing, perceived shortcomings. For instance, Felsenstein

(1985) pointed out the pitfalls of treating species values as independent data points and

provided two alternatives (i.e., independent contrasts and dividing a tree into pairs of tips)

to address those issues. Maddison (2006) realized the issue of not accounting for

diversification in transition-based methods (i.e., Pagel 1994) and not accounting for

differential transitions in diversification models, which led to the development of state-

dependent speciation and extinction models (Maddison et al. 2007; FitzJohn et al. 2009).

Recently, Rabosky and Goldberg (2015) pointed out a serious problem with

interpretations in state speciation and extinction analyses in general, and our work

largely, but not entirely, addresses these concerns. We caution users that it is important to

keep a perspective: all methods have flaws, and all will fail given a strong enough

violation of their underlying assumptions. For example, it appears difficult for HiSSE to

adequately estimate different transition rates when the model assumes any number of

hidden states, and so many estimates will be biased. Even with the increasing efforts to

test new methods (in our case, over 17,000 computer-days were devoted to conducting

simulations and analyses) there will be flaws that may have gone undetected. We urge

skepticism towards all models, but also skepticism towards statements of fatal flaws in

some models while leaving newer, competing models relatively untested.

There is no question that state-dependent speciation and extinction models are an

important advancement for understanding characters' impact on diversification patterns.

They have greatly improved statistical power over older, simpler sister-clade

comparisons, and the explicit inference of differences in speciation and extinction has the

potential for a much more fine-grained analysis of diversification. But, in a way, these

models have also allowed us to retreat to the old comforts of reducing complex organisms

into units of single, independently evolving characters, and offering adaptive

interpretations to each (c.f., Gould and Lewontin 1979). To be fair, of course, it is

unlikely that any trait of great interest to biologists has exactly zero effect on speciation

and/or extinction rates, but it is certainly unlikely that this trait acts in isolation. Thus, we

hope HiSSE is viewed as a step away from this line of thinking, as we no longer have to

necessarily focus analyses, or even interpret the results, by reference to the focal trait by

itself, but can instead estimate how important it is as a component of diversification

overall. It is in this way that analyses focused on "hidden" factors promoting

diversification will afford us a more refined understanding of why certain clades become

extraordinarily diverse, while still allowing us to examine our hypotheses about effects of

observed traits.

828   SUPPLEMENTAL MATERIAL

829 Data available from the Dryad Digital Repository: http://dx.doi.org/10.5061/dryad.52hp1

831   FUNDING

832 This work was supported by the National Institute for Mathematical and Biological

833 Synthesis [to J.M.B.], an Institute sponsored by the National Science Foundation, the

834 U.S. Department of Homeland Security, and the U.S. Department of Agriculture through

835 NSF Award #EF-0832858, with additional support from The University of Tennessee,

836 Knoxville.

838   ACKNOWLEDGEMENTS

We thank Nathan Jackson, Nick Matzke, Kathryn Massana, Chuck Bell, Andrew Leslie,

Jim Fordyce, and Chris Hamm for helpful discussions at various stages of this

manuscript. Rich FitzJohn, Sally Otto, and an anonymous reviewer provided very useful

comments on the manuscript, and we are deeply indebted to Tanja Stadler for her

suggestion to consider more complex null models, which led to the CID models here. We

also thank Jeffrey Oliver and Elizabeth Spriggs for feedback on the method

implementation. JMB would like to thank Michael Donoghue for wonderful discussions

over the years regarding some of the concepts described here.

REFERENCES

Alexander H.K., Lambert A., Stadler T. 2015. Quantifying age-dependent extinction

from species phylogenies. Systematic Biology In press.

doi:10.1093/sysbio/syv065.

Alfaro M.E., Santini F., Brock C., Alamillo H., Dornbug A., Rabosky D.L., Carnevale

G., Harmon L.J. 2009. Nine exceptional radiations plus high turnover explain

species diversity in jawed vertebrates. Proceedings of the National Academy of

Sciences of the U.S.A. 106:13410-13414.

Anderson D.R., Burnham K.P., Thompson W.L. 2000. Null hypothesis testing: problems,

prevalence, and an alternative. Journal of Wildlife Management 64:912-923.

Beaulieu J.M., Donoghue M.J. 2013. Fruit evolution and diversification in campanulid

angiosperms. Evolution 67: 3132-3144.

Beaulieu J.M., Jhueng D.-C., Boettiger C., O'Meara B.C. 2012. Modeling stabilizing

selection: expanding the Orstein-Uhlenbeck model of adaptive evolution.

Evolution 66:2369-2383.

Beaulieu J.M., O'Meara B.C., Donoghue M.J. 2013. Identifying hidden rate changes in

the evolution of a binary morphological character: the evolution of plant habit in

campanulid angiosperms. Systematic Biology 62:725-737.

Beaulieu J.M., O'Meara B.C. 2014. Hidden Markov models for studying the evolution of

binary morphological characters. In *Modern Phylogenetic Comparative Methods*

and their Application in Evolutionary Biology, Garamszegi L.Z. editor.

London:Springer.

Beaulieu J.M., O'Meara B.C. 2015. Extinction can be estimated from moderately sized

molecular phylogenies. Evolution 69:1036-1043.

Burnham K.P., Anderson D.R. 2002. Model selection and multimodel inference: a

practical information-theoretic approach. New York:Springer.

Carlson S.E., Mayer V., Donoghue M.J. 2009. Phylogenetic relationships, taxonomy, and

morphological evolution in Dipsacaceae (Dipsacales) inferred by DNA sequence

data. Taxon 58:1075-1091.

Carlson S.E., Howarth D.G., Donoghue M.J. 2011. Diversification of CYCLOIDEA-like

genes in Dipscaceae (Dipsacales): implications for the evolution of capitulum

inflorescences. BMC Evolutionary Biology 11:325.

Condamine F.L., Nagalingum N.S., Marshall C.R., Morlon H. 2015. Origin and

diversification of living cycads: a cautionary tale on the impact of the branching

process prior in Bayesian molecular dating. BMC Evolutionary Biology 15:65.

Davis M.P., Midford P.E., Maddison W. 2013. Exploring power and parameter

estimation of the BiSSE method for analyzing species diversification. BMC

Evolutionary Biology 13:38.

de Querioz A. 2002. Contingent predictability in evolution: Key traits and diversification.

Systematic Biology 51:917-929.

Donoghue M.J., Bell C.D., Winkworth R.C. 2003. The evolution of reproductive

characters in Dipsacales. International Journal of Plant Sciences. 164:S453-S464.

Donoghue M.J., Sanderson M.J. 2015. Confluence, synnovation, and depauperons in

plant diversification. New Phytologist doi:10.1111/nph.13367.

Edwards A.W.F. 1992. Likelihood. Cambridge:Cambridge University Press.

Felsenstein J. 1985. Phylogenies and the comparative method. The American Naturalist

125:1-15.

FitzJohn R.G. 2012. Diversitree: comparative phylogenetic analyses of diversification in

R. Methods in Ecology and Evolution 3:1084-1092.

FitzJohn R.G., Maddison W.P., Otto S.P. 2009. Estimating trait-dependent speciation and

extinction rates from incompletely resolved phylogenies. Systematic Biology

58:595-611.

Goldberg E.E., Kohn J.R., Lande R., Robertson K.A., Smith S.A., Igíc B. 2010. Species

selection maintains self-incompatibility. Science 330:493-495.

Goldberg E. E., Igic B. 2012. Tempo and mode in plant breeding system evolution.

Evolution 66:3701-3709.

Gould S.J., Lewontin R.C. 1979. The spandrels of San Marco and the Panglossian

paradigm: a critique of the adaptationist programme. Proceedings of the Royal

Society, B. 205:581-598.

Hansen T.F. 1997. Stabilizing selection and the comparative analysis of adaptation.

Evolution 51:1341-1351.

Hardy N.B. and Otto S.P. 2014. Specialization and generalization in the diversification of

phytophagous insects: tests of the musical chairs and oscillation hypotheses.

Proceedings of the Royal Society B. 281: 20132960.

Harmon L.J., Weir J.T., Brock C.D., Glor R.E., Challenger W. 2008. GEIGER:

investigating evolutionary radiations. Bioinformatics 24:129-131.

Hildago O., Mathez J., Garcia S., Garnatje T., Pellicer J., Valles J. 2010. Genome size

study in the Valerianaceae: First results and new hypotheses. Journal of Botany.

doi:10.1155/2010/797246.

Huelsenbeck J.P., Larget B., Alfaro M.E. 2004. Bayesian phylogenetic model selection

using reversible jump Markov chain Monte Carlo. Molecular Biology and

Evolution 21:1123-1133.

Jacobs B., Bell C., Smets E. 2010. Fruits and seeds of the Valeriana clade (Dipsacales):

Diversity and evolution. International Journal of Plant Sciences 171:421-434.

Maddison, W. P. 2006. Confounding asymmetries in evolutionary diversification and

character change. Evolution 60:1743–1746.

Maddison W.P., Midford P.E., Otto S.P. 2007. Estimating a binary character's effect on

speciation and extinction. Systematic Biology 56:701-710.

Maddison W.P., FitzJohn R.G. 2014. The unsolved challenge to phylogenetic correlation

tests for categorical characters. Systematic Biology 64:127-136.

Magnuson-Ford K., Otto S.P. Linking the investigations of character evolution and

species diversification. American Naturalist 180:225-245.

Marazzi B., Ane C., Simon M.F., Delgado-Salinas A., Luckow M., Sanderson M.J. 2012.

Locating evolutionary precursors on a phylogenetic tree. Evolution 66:3918-3930.

Mitter C., Farrell B., Wiegmann B. 1988. The phylogenetic study of adaptive zones: Has

phytophagy promoted insect diversification? American Naturalist 132:107-128.

Moore B.R., Donoghue M.J. 2007. Correlates of diversification in the plant clade

Dipsacales: geographic movement and evolutionary innovations. American

Naturalist 170:S28-S55.

Morlon H., Parsons T.L., Plotkin J.B. 2011. Reconciling molecular phylogenies with the

fossil record. Proceedings of the National Academy of Sciences of the U.S.A.

108:16327-16332.

Nakagawa S., Cuthill I.C. 2007. Effect size, confidence interval and statistical

significance: a practical guide for biologists. Biological Reviews 82:591-605.

Nee S., Holmes E.C., May R.M., Harvey P.H. 1994. Extinction rates can be estimated

from molecular phylogenies. Philosophical Transactions of the Royal Society, B.

344:77-82.

Nickerson R.S. 2000. Null hypothesis significance testing: a review of an old and

continuing controversy. Psychological Methods 5:241-301.

Pagel M. 1994. Detecting correlated evolution on phylogenies: a general method for the

comparative analysis of discrete characters. Proceedings of the Royal Society, B.

255:37-45.

Pennell M.W., Eastman J.M., Slater G.J., Brown J.W., Uyeda J.C., FitzJohn R.G., Alfaro

M.E., Harmon L.J. 2014. Geiger v2.0: an expanded suite of methods for fitting

macroevolutionary models to phylogenetic trees. Bioinformatics doi:

10.1093/bioinformatics/btu181

Price S.A., Hopkins S.S.B, Smith K.K., Roth V.L. 2012. Tempo of trophic evolution and

its impact on mammalian diversification. Proceedings of the National Academy of

Sciences of the U.S.A. 109:7008-7012.

Rabosky D.L. 2010. Extinction rates should not be estimated from molecular

phylogenies. Evolution 64:1816-1824.

Rabosky D.L. 2014. Automatic detection of key innovations, rate shifts, and diversity-

dependence on phylogenetic trees. PLoS One doi: 10.1371/journal.pone.0089543

Rabosky D.L., Lovette I.J. 2008. Explosive evolutionary radiations: decreasing speciation

or increasing extinction through time? Evolution 62:1866-1875.

Rabosky D.L., Goldberg E.E. 2015. Model inadequacy and mistaken inferences of trait-

dependent speciation. Systematic Biology 64:340-355.

Sanderson M.J. 2002. Estimating absolute rates of molecular evolution and divergence

times: a penalized likelihood approach. Molecular Biology and Evolution 19:101-

109.

Schluter D., Price T., Mooers A.O., Ludwig D. 1997. Likelihood of ancestor states in

adaptive radiation. Evolution 51:1699-1711.

Slater G.J., Price S.A., Santini F., Alfaro M.E. 2010. Diversity versus disparity and the

radiation of modern cetaceans. Proceedings of the Royal Society, B. 277:3097-

3104.

Smith S. A., Donoghue M.J. 2008. Rates of molecular evolution are linked to life history

in flowering plants. Science 322:86-89.

Smith S.A. O'Meara B.C. 2012. treePL: Divergence time estimation using penalized

likelihood for large phylogenies. Bioinformatics 28:2689-2690.

Spriggs E.L., Clement W.L., Sweeney P.W., Madriñán, Edwards E.J., Donoghue M.J.

2015. Temperate radiations and dying embersof a tropical past: the diversification

of Viburnum. New Phytologist. In press, doi:10.1111/nph.13305.

StamatakisA. 2006. Maximum likelihood-based phylogenetic analyses with thousands of

taxa and mixed models. Bioinformatics 22:2688–2690.

Swofford D.L. 2000. PAUP*: Phylogenetic analysis using parsimony (and other

methods) version 4. Sinauer Associates, Sunderland, Massachusetts.

Tank D.C., Donoghue M.J. 2010. Phylogeny and phylogenetic nomenclature of the

Campanulidae based on an expanded sample of genes and taxa. Systematic

Botany 35:425-441.

Weber M.G., Agrawal A.A. 2014. Defense mutualisms enhance plant diversification.

Proceedings of the National Academy of Sciences of the U.S.A. 111:16442-

16447.

Wilson A.W., Binder M., Hibbett D.S. 2011. Effects of gasteroid fruiting body

morphology on diversification rates in three independent clades of fungi estimated

using binary state speciation and extinction analysis. Evolution 65:1305-1322.

Yang Z., Kumar S., Nei M. 1995. A new method of inference of ancestral nucleotide and

amino acid sequences. Genetics 141:1641–1650.

1013

1014

TABLE 1. Summary of model support for simulated scenarios with and without a hidden trait, where calculating the average Akaike weight (w_i) for all models assessed the fit. We also calculated a null expectation of the Akaike weight as the average Akaike weight if we assumed an equal likelihood across all models. Thus, the null expectation is based solely on the penalty term in the AIC calculation.

Generating Model	Ntaxa	BiSSE Free	q's equal	ε0=ε1 / q's equal	τ0=τ1	τ0=τ1 / q's equal	ε0=ε1	τ0=τ1 / ε0=ε1	HiSSE Free	q's equal	ε0=ε1A=ε1B / q's equal	τ0=τ1A=τ1B	τ0=τ1A=τ1B / q's equal	ε0=ε1A=ε1B	τ0=τ1A=τ1B / ε0=ε1A=ε1B
Null w_i expectation		0.0260	0.0708	0.1923	0.0708	0.1923	0.0707	0.1923	0.0004	0.0096	0.0708	0.0035	0.0708	0.0035	0.0260
BiSSE, transition rates vary	50	0.0416	0.0821	***0.1504***	0.0864	***0.1639***	0.0925	***0.1737***	0.0022	0.0179	0.0739	0.0076	0.0725	0.0010	0.0257
λ0=λ1=0.1	100	0.0464	0.0837	0.1322	0.0791	0.1319	0.0930	***0.1641***	0.0046	0.0289	0.0944	0.0123	0.0811	0.0185	0.0298
μ0=μ1=0.03	200	0.0499	0.0623	0.0927	0.0969	0.0982	0.0989	***0.1812***	0.0069	0.0374	0.0919	0.0198	0.0874	0.0313	0.0450
q01=0.015, q10=0.005	400	0.0662	0.0635	0.0578	0.1147	0.0668	0.1119	***0.2116***	0.0070	0.0320	0.0771	0.0227	0.0801	0.0284	0.0599
BiSSE, state 1 2x speciation	50	0.0370	0.0848	***0.1896***	0.0803	***0.1681***	0.0930	0.1339	0.0117	0.0160	0.0781	0.0075	0.0685	0.0113	0.0203
λ0=0.1, λ1=0.2	100	0.0450	0.0918	***0.1920***	0.0783	0.1545	0.1066	0.0997	0.0030	0.0218	0.0933	0.0089	0.0706	0.0182	0.0163
μ0=μ1=0.03	200	0.0508	0.1006	***0.2129***	0.0820	0.1544	0.1194	0.0508	0.0069	0.0226	0.0837	0.0113	0.0643	0.0314	0.0086
q01=q10=0.01	400	0.0690	0.1172	***0.2028***	0.0911	0.1503	0.1324	0.0186	0.0091	0.0268	0.0637	0.0161	0.0615	0.0356	0.0057
BiSSE, state 1 .5x extinction	50	0.0480	0.0866	***0.1465***	0.0866	***0.1451***	0.0958	***0.1714***	0.0040	0.0219	0.0769	0.0092	0.0684	0.0132	0.0264
λ0=λ1=0.1	100	0.0502	0.0934	***0.1332***	0.0879	***0.1545***	0.0874	***0.1576***	0.0039	0.0214	0.0784	0.0128	0.0779	0.0147	0.0267
μ0=0.06, μ1=0.03	200	0.0596	0.1015	0.1219	0.0904	***0.1652***	0.0826	0.1324	0.0048	0.0286	0.0717	0.0101	0.0824	0.0144	0.0344
q01=q10=0.01	400	0.0710	0.1145	0.1008	0.0996	***0.1605***	0.0825	0.1113	0.0082	0.0325	0.0618	0.0156	0.0983	0.0163	0.0270
HiSSE, state 1B 2x speciation	50	0.0486	0.0808	***0.1573***	0.0936	***0.1488***	0.1069	***0.1520***	0.0030	0.0200	0.0768	0.0089	0.0621	0.0128	0.0285
λ0A=λ1A=0.01, λ1B=0.2	100	0.0533	0.0786	***0.1264***	0.0777	***0.1147***	0.0931	0.1074	0.0095	0.0386	***0.1438***	0.0146	0.0957	0.0248	0.0216
μ0A=μ1A=μ1B=0.03	200	0.0666	0.0553	0.0719	0.0726	0.0742	0.0791	0.0743	0.0134	0.0611	***0.2158***	0.0225	0.1263	0.0469	0.0202
q0A1A=q1A0A=q1A1B=q1B1A=0.01	400	0.0321	0.0275	0.0359	0.0254	0.0283	0.0370	0.0236	0.0315	0.1094	***0.3119***	0.0420	0.1876	0.1024	0.0053
HiSSE, state 1B 3x speciation	50	0.0524	0.0796	***0.1594***	0.0821	0.1218	0.1124	0.1003	0.0091	0.0364	***0.1295***	0.0156	0.0544	0.0317	0.0153
λ0A=λ1A=0.01, λ1B=0.3	100	0.0504	0.0604	0.0998	0.0629	0.0817	0.0893	0.0686	0.0112	0.0660	***0.2250***	0.0156	0.1048	0.0464	0.0177
μ0A=μ1A=μ1B=0.03	200	0.0414	0.0316	0.0405	0.0304	0.0397	0.0556	0.0146	0.0241	0.1101	***0.4102***	0.0215	0.0906	0.0871	0.0026
q0A1A=q1A0A=q1A1B=q1B1A=0.01	400	0.0220	0.0140	0.0112	0.0140	0.0090	0.0437	0.0078	0.0347	0.1654	***0.4646***	0.0175	0.0703	0.1244	0.0015
HiSSE, state 1B .5x extinction	50	0.0487	0.0911	***0.1453***	0.0918	***0.1472***	0.1019	***0.1711***	0.0023	0.0230	0.0694	0.0074	0.0667	0.0096	0.0246
λ0A=λ1A=λ1B=0.1	100	0.0518	0.0745	0.1251	0.0939	0.1236	0.0968	***0.1743***	0.0068	0.0286	0.0697	0.0147	0.0728	0.0223	0.0450
μ0A=μ1A=0.06, μ1B=0.03	200	0.0538	0.0648	0.1002	0.0951	***0.1128***	0.0918	***0.1402***	0.0083	0.0435	0.0877	0.0207	0.1032	0.0347	0.0433
q0A1A=q1A0A=q1A1B=q1B1A=0.01	400	0.0591	0.0461	0.0464	0.0986	0.0660	0.1018	***0.1200***	0.0235	0.0604	0.0920	0.0441	***0.1523***	0.0352	0.0546
HiSSE, state 1B 2x speciation, transition rates vary	50	0.0432	0.0743	***0.1452***	0.0834	***0.1338***	0.0981	***0.1403***	0.0070	0.0391	0.1095	0.0156	0.0682	0.0212	0.0209
λ0A=λ1A=0.01, λ1B=0.3	100	0.0379	0.0518	0.0737	0.0594	0.0672	0.0778	0.0835	0.0181	0.0701	***0.2457***	0.0222	0.0823	0.0886	0.0216
μ0A=μ1A=μ1B=0.03	200	0.0399	0.0184	0.0190	0.0399	0.0202	0.0500	0.0535	0.0310	0.1267	***0.3259***	0.0229	0.1090	0.1323	0.0111
q0A1A=0.015,q1A0A=q1A1B=0.005,	400	0.0098	0.0027	0.0048	0.0108	0.0067	0.0233	0.0118	0.0883	0.1706	***0.3261***	0.0291	0.0596	***0.2545***	0.0019
q1B1A=0.01	800	0.0020	0.0002	0.0004	0.0022	0.0004	0.0021	0.0038	0.1040	0.1647	***0.3156***	0.0230	0.0638	***0.3172***	0.0005

Bold italics indicate significance based on 95% confidence intervals

TABLE 2. Summary of the model support for simulated scenarios that tested the performance of the general HiSSE model and two CID models (CID-2, CID-2; see text). All data sets tested contained 400 species, and calculating the average Akaike weight (w_i) for all models assessed the fit. As with Table 1, we calculated a null expectation of the Akaike weight as the average Akaike weight if we assumed an equal likelihood across all models.

	Equal rates	BiSSE		CID-2		CID-4		HiSSE			
	$\tau_0=\tau_1$			τ_A, τ_B	τ_A, τ_B	$\tau_A, \tau_B, \tau_C, \tau_D$	$\tau_A, \tau_B, \tau_C, \tau_D$			$\tau_{0A}=\tau_{1A}=\tau_{0B}$	$\tau_{0A}=\tau_{1A}=\tau_{0B}$
Generating Model	$\varepsilon_0=\varepsilon_1$	Free	$\varepsilon_0=\varepsilon_1$	$\varepsilon_A, \varepsilon_B$	$\varepsilon_A=\varepsilon_B$	$\varepsilon_A, \varepsilon_B, \varepsilon_C, \varepsilon_D$	$\varepsilon_A=\varepsilon_B=\varepsilon_C=\varepsilon_D$	Free	$\varepsilon_{0A}=\varepsilon_{1A}=\varepsilon_{0B}=\varepsilon_{1B}$	$\varepsilon_{0A}=\varepsilon_{1A}=\varepsilon_{0B}$	$\varepsilon_{0A}=\varepsilon_{1A}=\varepsilon_{0B}=\varepsilon_{1B}$
Null w_i expectation	0.1426	0.0865	0.1426	0.0865	0.1426	0.0193	0.0865	0.0117	0.0525	0.0865	0.1426
BiSSE, state 1 2x speciation $\lambda_0=0.1, \lambda_1=0.2$ $\mu_0=\mu_1=0.03$ All q's = 0.01	0.0085	0.1995	***0.3313***	0.0046	0.0089	0.0001	0.0012	0.0255	0.0917	0.1298	0.1989
CID-2, state B 2x speciation $\lambda_{0A}=\lambda_{1A}=0.1, \lambda_{0B}=\lambda_{1B}=0.2$ $\mu_{0A}=\mu_{1A}=\mu_{0B}=\mu_{1B}=0.03$ All q's = 0.01	0.0505	0.0445	0.0568	0.1836	***0.3107***	0.0063	0.0656	0.0322	0.1461	0.0407	0.0631
HiSSE, state 1B 2x speciation $\lambda_{0A}=\lambda_{1A}=\lambda_{0B}=0.1, \lambda_{1B}=0.2$ $\mu_{0A}=\mu_{1A}=\mu_{0B}=\mu_{1B}=0.03$ All q's = 0.01	0.0049	0.0957	0.1327	0.0073	0.0135	0.0002	0.0032	0.0417	0.1199	0.2265	***0.3544***
"Worst-case" trait-independence Initial $\lambda=0.1$, drawn from lognormal $\varepsilon=0.3$, q01=q10=0.01	0.0005	0.0003	0.0007	0.0897	0.1562	0.0544	***0.3996***	0.0563	0.2213	0.0067	0.0142

Bold italics indicate significance based on 95% confidence intervals that bracket the mean Akaike weight.

TABLE 3. The fit of alternative models of achene fruit evolution in the flowering plant clade Dipsidae. The best model, based on ΔAIC and Akaike weights (w_i) is denoted in bold.

Model	np	lnLik	AIC	ΔAIC	wi
BiSSE: All free	6	-1486.5	2984.9	90.3	<0.001
BiSSE: ε0=ε1	5	-1488.6	2987.2	92.6	<0.001
BiSSE: q's equal	5	-1487.3	2984.5	89.9	<0.001
BiSSE: ε0=ε1, q's equal	4	-1490.2	2988.4	93.8	<0.001
Null-two: q's equal	5	-1449.1	2908.3	13.7	0.001
Null-two: ε's, q's equal	4	-1449.7	2907.4	12.8	0.001
Null-four: q's equal	9	-1445.1	2908.1	13.5	0.001
Null-four: ε's equal, q's equal	6	-1445.3	2902.6	7.98	0.015
HiSSE: q's equal	9	-1446.6	2911.2	16.6	<0.001
HiSSE: ε's equal, q's equal	6	-1447.2	2906.5	11.9	0.002
HiSSE: τ0A=τ1A=τ0B, ε0A=ε1A=ε0B, q's equal	5	-1463.3	2936.6	42.0	<0.001
HiSSE: τ0A=τ1A=τ0B, ε's equal, q's equal	4	-1471.0	2950.0	55.4	<0.001
HiSSE: q0B1B=0, q1B0B=0, all other q's equal6	9	-1451.7	2921.5	26.9	<0.001
HiSSE: ε's equal, q0B1B=0, q1B0B=0, all other q's equal	6	-1451.7	2915.5	20.9	<0.001
HiSSE: τ0A=τ1A=τ0B, ε0A=ε1A=ε0B, q0B1B=0, q1B0B=0, all other q's equal	5	-1462.2	2934.3	39.7	<0.001
HiSSE: τ0A=τ1A=τ0B, ε's equal, q0B1B=0, q1B0B=0, all other q's equal	4	-1469.5	2947.1	52.5	<0.001
HiSSE: τ0A=τ0B, ε0A=ε0B, q's equal	7	-1459.6	2933.2	38.6	<0.001
HiSSE: τ0A=τ0B, ε's equal, q's equal	5	-1466.4	2942.8	48.2	<0.001
HiSSE: τ0A=τ0B, ε0A=ε0B, q0B1B=0, q1B0B=0, all other q's equal	7	-1458.4	2930.8	36.2	<0.001
HiSSE: τ0A=τ0B, ε's equal, q0B1B=0, q1B0B=0, all other q's equal	5	-1464.7	2939.4	44.8	<0.001
HiSSE: τ0A=τ1A, ε0A=ε1A, q's equal	7	-1446.7	2907.3	12.7	<0.001
HiSSE: τ0A=τ1A, ε's equal, q's equal	5	-1469.1	2948.3	53.7	<0.001
HiSSE: τ0A=τ1A, ε0A=ε1A, q0B1B=0, q1B0B=0, all other q's equal	7	-1442.2	2898.3	3.71	0.130
HiSSE: τ0A=τ1A, ε's equal, q0B1B=0, q1B0B=0, all other q's equal	5	*-1442.3*	*2894.6*	*0.00*	*0.848*

a)

c)

e)

b)

d)

f)

Frequency of state 0A

Frequency of state 1A

Frequency of state 1B

a)

Triplo.+
Dipsac.

Valerian.

Morin.

Zab.

Linn.

Para.

Capr.+
Hepta.

Adox.

Dier.

BiSSE

Net diversification

0.059

0.200

Tip frequency

achene

non-achene

States

b)

Dipsac.

core Valerian.

Viburnum

Capr.

HiSSE

Net diversification

0.059

0.200

Tip frequency

achene

non-achene

States