Eight thousand years of natural selection in Europe

Iain Mathieson* [1], Iosif Lazaridis [1,2], Nadin Rohland [1,2], Swapan Mallick [1,2], Bastien Llamas [3], Joseph Pickrell [4], Harald Meller [5], Manuel A. Rojo Guerra [6], Johannes Krause [7,8,9], David Anthony [10], Dorcas Brown [10], Carles Lalueza Fox [11], Alan Cooper [3], Kurt W. Alt [5,12,13,14], Wolfgang Haak [3], Nick Patterson [1,2], David Reich* [1,2,15]

1. Harvard Medical School, Department of Genetics, Boston MA, USA

2. Broad Institute of MIT and Harvard, Cambridge MA, USA

3. Australian Centre for Ancient DNA, School of Biological Sciences & Environment Institute, The University of Adelaide, Adelaide South Australia, Australia

4. New York Genome Center, New York NY, USA

5. State Office for Heritage Management and Archaeology Saxony-Anhalt and State Museum of Prehistory, Halle, Germany

6. Departament of Prehistory and Archaeology. University of Valladolid, Spain

7. Institute for Archaeological Sciences, University of Tübingen, Tübingen, 72074, Germany

8. Department of Paleoanthropology, Senckenberg Center for Human Evolution and Paleoenvironment, University of Tübingen, Tübingen, Germany

9. Max Planck Institute for the Science of Human History, D-07745 Jena, Germany

10. Hartwick College, Department of Anthropology, Oneonta NY, USA

11. Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), Barcelona, Spain

12. Danube Private University, Krems, Austria

13. Institute for Prehistory and Archaeological Science, University of Basel, Basel, Switzerland

14. Institute of Anthropology, Johannes Gutenberg University of Mainz, Mainz, Germany

15. Howard Hughes Medical Institute, Harvard Medical School, Boston MA, USA

*To whom correspondence should be addressed:

Iain Mathieson (iain_mathieson@hms.harvard.edu) or David Reich (reich@genetics.med.harvard.edu)

The arrival of farming in Europe beginning around 8,500 years ago required adaptation to
new environments, pathogens, diets, and social organizations. While evidence of natural selection
can be revealed by studying patterns of genetic variation in present-day people[1-6], these pattern are
only indirect echoes of past events, and provide little information about where and when selection
occurred. Ancient DNA makes it possible to examine populations as they were before, during and
after adaptation events, and thus to reveal the tempo and mode of selection[7,8]. Here we report the
first genome-wide scan for selection using ancient DNA, based on 83 human samples from Holocene
Europe analyzed at over 300,000 positions. We find five genome-wide signals of selection, at loci
associated with diet and pigmentation. Surprisingly in light of suggestions of selection on immune
traits associated with the advent of agriculture and denser living conditions, we find no strong
sweeps associated with immunological phenotypes. We also report a scan for selection for complex
traits, and find two signals of selection on height: for short stature in Iberia after the arrival of
agriculture, and for tall stature on the Pontic-Caspian steppe earlier than 5,000 years ago. A
surprise is that in Scandinavian hunter-gatherers living around 8,000 years ago, there is a high
frequency of the derived allele at the *EDAR* gene that is the strongest known signal of selection in
East Asians and that is thought to have arisen in East Asia. These results document the power of
ancient DNA to reveal features of past adaptation that could not be understood from analyses of
present-day people.
19
We analyzed genome-wide data from 83 ancient individuals who lived in Europe over the last
eight thousand years. The data for 64 samples was obtained by using in-solution hybridization with
synthesized oligonucleotide probes to enrich ancient DNA libraries for 394,577 single nucleotide
polymorphisms (SNPs), including 29,075 chosen based on evidence of functional importance[9]. We merged
this data with shotgun sequence data from 19 published samples[7,10-13] and co-analyzed these ancient
samples with 503 present-day European samples from five populations from phase 3 of the 1000 Genomes
Project[14]. After quality control, we were left with 364,618 SNPs genome-wide, and a mean effective
sample size of 81 chromosomes at each SNP (Table 1, Methods).
28
Present-day European populations can be modeled as a mixture of three ancient populations:
western hunter-gatherers, early European farmers, and Yamnaya pastoralists from the Pontic-Caspian
steppe[9,11], all of which are represented by samples in our study. To scan for selection, we tested every SNP
to evaluate whether its observed frequency distribution across present-day Europeans from the 1000
Genomes Project was consistent with this model (Figure 1, Extended data Figure 1, Methods). SNPs with
frequencies in present-day populations that are inconsistent with a neutral mixture of the ancestry of the
ancient populations are likely to be influenced by selection. A strength of this test is that it does not make
the simplifying assumption of population continuity over time, which is problematic in light of the
population replacements that are now known to have occurred in many places in the world including in
Europe[9,11,15]. Of the 307,494 non-monomorphic SNPs that we tested, the 25,017 that were in the set of
potentially functional SNPs were, as a group, significantly more inconsistent with the model than neutral
SNPs (Figure 1A). This result held across all SNP categories, including genome-wide association study
(GWAS) hits and human leukocyte antigen (HLA) haplotype tag SNPs (Figure 1B), suggesting pervasive
selection on polymorphisms of biological importance.
43
Using a conservative significance threshold of $p = 1.6 \times 10^{-7}$ (0.05/307,494), and a genomic control
correction of 1.21 (Ref. 16), we identified five loci that contained SNPs achieving genome-wide
significance. Three of these have previously been identified as targets of selection[8,17-19]: the *LCT* lactase
persistence locus (rs4988235, $p = 4.8 \times 10^{-33}$), the *SLC45A2* pigmentation locus (rs16891982, $p = 6.9 \times 10^{-25}$)
and the *HERC2/OCA2* eye color locus (rs12913832, $p = 3.2 \times 10^{-11}$). We also identified two novel signals at
NADSYN1 (rs7940244, $p = 7.1 \times 10^{-8}$) and *FADS1* (rs174546, $p = 1.6 \times 10^{-7}$) (Figure 2C, Extended Data
Figure 1). All of these signals except *FADS1* remained significant in a less powerful but more robust

analysis in which we called a single majority allele for each sample at each site instead of computing
genotype likelihoods (Extended Data Figure 2).
53
The SNP (rs4988235) responsible for lactase persistence in Europe[20,17] gives the strongest signal in
our analysis. We estimated the selection coefficient on the derived allele to be 0.015 (95% confidence
interval; CI=0.010-0.034) using a method that fits a hidden Markov model to the population allele
frequencies as they change over time[21]. Our data strengthens previous reports of the late appearance of
lactase persistence in Europe[7,22], with the earliest appearance of the allele in a central European Bell
Beaker sample (individual I0112) who lived approximately 4,300 years ago. We detect no evidence of
lactase persistence in Early Neolithic farming populations like the *Linearbandkeramik* (LBK), or in the
steppe pastoralist Yamnaya, despite their use of domesticated cattle (Figure 2).
62
We also found evidence of selection at two loci that affect skin pigmentation. The derived alleles
of rs1426654 at *SLC24A5* and rs16891982 at *SLC45A2* are, respectively, fixed and almost fixed in present-
day Europeans[23,24]. As previously reported[7,11,12], both derived alleles are absent or very rare in western
hunter-gatherers. suggesting that mainland European hunter-gatherers may have had dark skin
pigmentation. *SLC45A2* first appears in our data at low frequency in the Early Neolithic, and increases
steadily in frequency until the present, when it has frequency 1 in all populations except Spanish (IBS,
Figure 2; estimated selection coefficient $\hat{s}$ =0.020, CI=0.011-0.031). In contrast, the derived allele of
SLC24A5 increases rapidly in frequency to around 0.9 in the Early Neolithic, suggesting that most of the
increase in frequency of this allele is due to its high frequency in the early farmers who migrated to Europe
from the southeast at this time, although there is still strong evidence of ongoing selection after the arrival
of farming (p=5.5× 10^{-7}, $\hat{s}$ =0.025, CI=0.003-0.21).
74
The derived allele of rs12913832 at the *HERC2/OCA2* locus is the primary determinant of blue
eyes in Europeans, and may also contribute to light skin and hair pigmentation[25-28]. Our analysis detects a
genome-wide signal of selection at this locus, but instead of the signal being one of positive selection with
a coefficient of 0.036 as in a previous study of ancient DNA in the eastern Europe steppe[8], our signal is of
weakly negative selection ($\hat{s}$ =-0.007, CI = -0.011 to -0.001). One possible explanation is local adaptation:
that the allele is advantageous in the north and disadvantageous in the south of Europe. This hypothesis is
supported by the fact that our data shows that an extreme north-south gradient in allele frequencies has
been maintained in Europe for the last 8,000 years (Figure 2C, Extended data Figure 3).
83
The remaining two genome-wide signals are both located on chromosome 11 in the genes *FADS1*
and *NADSYN1*. *FADS1* (and its linked family member *FADS2*), are involved in fatty acid metabolism, and
variation at this locus is associated with plasma lipid and fatty acid concentration[29-31]. The derived allele of
the most significant SNP in our analysis, rs174546, is associated with decreased triglyceride levels[31]. This
locus is therefore a plausible target of selection related to changes in diet. Variants at *NADSYN1* (and the
nearby *DHCR1*), have been associated with circulating vitamin D levels[32,33] and the most associated SNP
rs7940244 in our dataset is highly differentiated across closely related Northern European populations[34,35],
suggesting the possibility of selection related to variation in environmental sources of vitamin D.
92
We find a surprise in seven Scandinavian hunter-gatherers from the Motala site in southern
Sweden who lived around 7,700 years before present. While the western hunter-gatherers of central and
southern Europe largely have the ancestral allele at the two major European skin pigmentation loci, the
closely related Scandinavian hunter-gatherers have both the derived alleles contributing to light skin
pigmentation at high frequency (Figure 2B). Thus, the derived allele of *SLC24A5* was common in both the
Scandinavian hunter-gatherers and Early European farmers, but not in the geographically intermediate
western hunter-gatherers. Further, in four out of seven Motala samples, we observe the derived allele of
rs3827760 in the *EDAR* gene, which has effects on tooth morphology and hair thickness[36,37] This allele has
been the subject of a selective sweep in East Asia[38], and today it is at high frequency in East Asians and

Native Americans. The *EDAR* derived allele is largely absent in present-day Europe except in Scandinavia,
which is plausibly due to Siberian movements into the region associated with the arrival of the Finno-
Ugric languages many millennia after the date of the Motala samples - we estimate a date of admixture of
East Asian and Sardinian like ancestry in present-day Finns at 56 ± 7 generations using ALDER[39]. The
derived allele in the Motala samples lies on the same haplotype as in modern East Asians (Extended Data
Figure 4) implying a shared origin. The statistic f_4(Yoruba, Scandinavian hunter-gatherers, Han, Onge
Andaman Islanders) is significantly negative (Z=-3.9) implying gene flow between the ancestors of
Scandinavian hunter-gatherers and Han so this shared haplotype is likely the result of ancient gene flow
between groups ancestral to these two populations.
111

We also tested for selection on complex traits, which are controlled by many genetic variants, each
with a weak effect. Under the pressure of natural selection, these variants are expected to experience small
but correlated directional shifts, rather than any single variant changing dramatically in frequency, and
recent studies have argued that this may be a predominant mode of natural selection in humans[40]. The best
documented example of this process in humans is height, which has been shown to have been under recent
selection in Europe[41]. At alleles known from GWAS to affect height, northern Europeans have, on average,
a significantly higher probability of carrying the height-increasing allele than southern Europeans, which
could either reflect selection for increased height in the ancestry of northern Europeans or decreased height
in the ancestry of southern Europeans. To test for this signal in our data, we used a statistic that tests
whether trait-affecting alleles are more differentiated than randomly sampled alleles, in a way that is
coordinated across all alleles consistent with directional selection[42]. We applied the test to all populations
together, as well as to pairs of populations in order to localize the signal (Figure 3, Extended Data Figure
5, Methods).
125

We detect a significant signal of directional selection on height in Europe (p=0.002), and our
ancient DNA data allows us to determine when this occurred and also to determine the direction of
selection. Both the Iberian Early Neolithic and Middle Neolithic samples show evidence of selection for
decreased height relative to present-day European Americans (Figure 3A; p=0.002 and p < 0.0001,
respectively). Comparing populations that existed at the same time (Figure 3B), there is a significant signal
of selection between central European and Iberian populations in each of the Early Neolithic, Middle
Neolithic and present-day periods (p=0.011, 0.012 and 0.004, respectively). Therefore, the selective
gradient in height in Europe has existed for the past 8,000 years. This gradient was established in the Early
Neolithic, increased into the Middle Neolithic and decreased at some point thereafter. Since we detect no
significant evidence of selection or change in genetic height among Northern European populations, our
results further suggest that selection operated mainly on Southern rather than Northern European
populations. There is another possible signal in the Yamnaya, related to people who migrated into central
Europe beginning at least 4,800 years ago and who contributed about half the ancestry of northern
Europeans today[9]. The Yamnaya have the greatest predicted genetic height of any population, and the
difference between Yamnaya and the Iberian Middle Neolithic is the greatest observed in our data (Z=6.2,
p<0.0001, Extended data Figure 6). This observation is consistent with archaeological evidence that the
Yamnaya were taller than populations contemporary to them[43].
143

We also detect a significant signal of selection on body mass index (BMI) around the central
European Late Neolithic/Bronze Age (LNBA; selection for increased BMI in CEU, Central_LNBA-CEU,
p=0.022), although this does not persist when analyzed with the more weakly powered majority-called data
(Extended Data Figure 5) and when we test all populations together the result is not significant (p=0.14).
This could be due to reduced power for BMI relative to height, since the variance explained by the tested
BMI SNPs is much lower than that is explained by the height SNPs. We observe no other significant
signals of polygenetic selection among the six complex traits we tested, which included not just height[44]
and BMI[45], but also waist-hip ratio[46], type 2 diabetes[47], inflammatory bowel disease[48] and lipid levels[31].
152

Our results show that ancient DNA can be used to perform genome-wide scans for selection, and
to reveal the histories of phenotypically important alleles. This study has power to detect strong selection,
with selective advantages greater than one percent, over the last eight thousand years of European history
(Extended Data Figure 7). The negative results – the fact that we detect only five alleles with genome-wide
evidence of selection – are equally interesting, as they suggest that there were only a small number of
strong selective sweeps on single alleles since the advent of agriculture in Europe. None of these strong
sweeps involved immune-related loci despite the fact that it has been argued that very strong selection on
disease resistance would be common following the introduction of agriculture. However there are signs in
our analysis of weaker selection on immune phenotypes, for example at the HLA or in the region around
TLR1 and *TLR6* (rs7661887, p=9.8×10^{-7}). From the genome-wide analysis we also see evidence for a long
tail of more weakly selected alleles implying that, with additional ancient samples, similar studies would
identify many loci with smaller or more complex effects. These results demonstrate the power of ancient
DNA to reveal the nature of natural selection, and highlight the importance of applying approaches such as
the one here beyond Europe, and to non-human species.
167

Acknowledgments

169
We thank Paul de Bakker, Joachim Burger, Christos Economou, Elin Fornander, Fredrik Hallgren, Karola
Kirsanow, Alissa Mittnik, Adam Powell, Pontus Skoglund and Shervin Tabrizi for helpful discussions,
suggestions about SNPs to include in the capture reagent, or contribution to sample preparation. We also
thank Julio Manuel Vidal Encinas and María Encina Prada for allowing us access to the La Braña sample
and the 1000 Genomes Project for allowing use of the phase 3 data. IM is supported by a long-term
fellowship from the Human Frontier Science Program. DR is supported by U.S. National Science
Foundation HOMINID grant BCS-1032255, U.S. National Institutes of Health grant GM100233, and the
Howard Hughes Medical Institute. This research was supported by an Australian Research Council grant
to WH and BL. (DP130102158).

Methods

Selection of potentially functional sites

The in-solution capture reagent targets 394,577 SNPs[9]. Of these, 34,747 are "potentially functional" sites chosen as follows: 1,290 SNPs identified as targets of selection in Europeans by the Composite of Multiple Signals (CMS) test[2]; 14,814 SNPS identified as significant hits by genome-wide association studies, or with known phenotypic effect (GWAS); 1,289 SNPs with extremely differentiated frequencies between HapMap populations[49] (HiDiff); 9,116 immunochip SNPs chosen for study of immune phenotypes and largely centered at the HLA locus (Immune); 347 SNPs phenotypically relevant to South America (mostly altitude adaptation SNPs in EGLN1 and EPAS1) and 5,387 SNPs which tag HLA haplotypes (HLA). The reagent also targets 2,524 Y chromosome SNPs, which we do not analyze in this study.

Sample selection and merging

From the set of 69 ancient samples first published by Ref. 9, we excluded five from our main analyses. Two of these were Eastern hunter-gatherers (I0061 and I0124), two were relatives of others in the set (I0411 and I0114) and one had low coverage and uncertain assignment (I0550). The Loschbour and Stuttgart samples had been genotyped at every callable site in the genome so we took the intersection of these sites with the 394,577 targeted sites[9]. For other published samples we obtained shotgun sequence data from the authors and counted reads at the 394,577 sites.

For the western hunter-gatherer sample from La Braña in present-day Spain[12], we generated additional data beyond the published data, using the extraction, library preparation and capture protocols described in Ref. 9. We generated a total of 94,246,843 sequences passing standard Illumia filters, of which 42,952,907 mapped to the SNP targets and 22,389,493 remained after removing duplicate molecules. The fraction of terminal sites that were C→T substitutions from the reference sequence (indicating authentic ancient DNA) was 12.3%, and the average coverage on the SNP targets was 58.8-fold. The BAM file associated with these data is available at <mark>xx</mark>.

We downloaded integrated phase 3 variant calls from the 1,000 Genomes ftp site (ftp://ftp-trace.ncbi.nih.gov/1000genomes/ftp/release/20130502/; downloaded on 17th September 2014) and merged these with the ancient data. This resulted in a loss of 29,995 sites that were not present in the 1,000 Genomes VCF. For each sample we took the midpoint of its date range (usually calibrated [14]C dates) as a point estimate of its age. For each population, we took the mean of the sample age estimates and used that as the age of the population.

Genotype calls and likelihood

For most ancient samples we did not have sufficient coverage to make reliable diploid calls. We therefore used the read counts at each SNP to compute the likelihood of the allele frequency in each population. Suppose that at a particular site, for population i, we have N samples with read level data, and M samples for which we made hard genotype calls (Loschbour, Stuttgart and the 1,000 Genomes samples). For samples $i = 1 \dots N$, with genotype data, we observe X copies of the reference allele out of $2N$ total chromosomes. For each of samples $i = (N + 1) \dots (N + M)$, with read data, we observe R_i reads with the reference allele out of T_i total reads. Then, the likelihood of the population reference allele frequency, p given data $D = \{X, N, \bar{R}, \bar{T}\}$ is given by

$$L(p; D) = B(X, 2N, p) \prod_{i=N+1}^{N+M} \{p^2 B(R_i, T_i, \varepsilon) + 2p(1-p)B(R_i, T_i, 0.5) + (1-p)^2 B(R_i, T_i, 1-\varepsilon)\},$$

where $B(k, n, p) = \binom{n}{k} p^{pk}(1-p)^{n-k}$ is the binomial probability distribution and ε is a small probability
of error, which we set to 0.001. We write $\ell(p; D)$ for the log-likelihood. To estimate allele frequencies, for
example in Figure 2 or for the polygenic selection test, we maximized this likelihood numerically for each
population.

We also constructed a dataset where we called a single allele at each site for each sample by choosing the
allele most commonly seen in the reads mapped to that site. This results in a reduction in effective sample
size of around 40% but should be more robust to small levels of contamination. We reran our main
analyses on this "majority-called" dataset to check that they were not confounded by systematic
contamination.

Genome-wide scan for selection

To scan for selection across the genome, we used the following test. Consider a single SNP. Assume that
we can model the allele frequencies p_{mod} in A modern populations as a linear combination of allele
frequencies in B ancient populations p_{anc}. That is, $p_{mod} = C p_{anc}$, where C is an A by B matrix with
columns summing to 1. We have data D_j from population j which is some combination of read counts and
genotypes as described above. Then, writing $\bar{p} = [p_{anc}, p_{mod}] = [p_1 \ldots p_{A+B}]$ the log-likelihood of the
allele frequencies equals the sum of the log-likelihoods for each population.

$$\ell(\bar{p}, \overline{D}) = \sum_{j=1}^{A+B} \ell(p_j; D_j)$$

To test for deviations in allele frequency, we test the null hypothesis $H_0: p_{mod} = C \, p_{anc}$ against the
alternative $H_1: p_{mod}$ unconstrained. We numerically maximized this likelihood in both the constrained
and unconstrained model and used the fact that twice the difference in log-likelihood is approximately χ^2_A
distributed to compute a test statistic and a p-value.

As in Ref. 9, we used western hunter-gatherers, Early Neolithic, and Yamnaya as the ancient source
populations and the 1000 Genomes CEU, GBR, IBS and TSI as the present-day populations. We did not
use FIN, because they do not fit this three-population model well. We estimated the proportions of
(western hunter-gatherers, Early Neolithic, Yamnaya) to be CEU=(0.164, 0.366, 0.470),
GBR=(0.213,0.337,0.450), IBS= (0, 0.773, 0.227) and TSI=(0, 0.712, 0.288). In practice we found that
there was substantial inflation in the test statistic, most likely due to unmodeled ancestry or additional drift.
To correct for this we applied a genomic control correction, dividing all the test statistics by a constant, λ
chosen so that the median p-value matched the median of the null χ^2_4 distribution[16]. Excluding sites in the
potentially functional set, we estimated $\lambda = 1.21$ and used this value as a correction throughout.

To estimate the power of this test, we randomly sampled allele counts from the full dataset, restricting to
polymorphic sites with a mean frequency across modern populations of less than 0.1. We then simulated
what would happen if the allele had been under selection in one of the modern populations by simulating a
Wright-Fisher trajectory for 50, 100 or 200 generations. We took the final frequency from this simulation,
sampled observations to replace the actual observations in that population, and counted the proportion of
simulations that gave a genome-wide significant result after GC correction. We also performed a similar
simulation, assuming that the site was selected in every population (Extended data Figure 7A).

We investigated how robust the test was to misspecification of the mixture matrix C. We reran the power
simulations using a matrix $C' = pC + (1 - p)R$ for $p \in [0,1]$ where R was a random matrix chosen so
that for each modern population, the mixture proportions of the three ancient populations were jointly
normally distributed on [0,1]. Increasing p increases the genomic inflation factor and reduces power,
demonstrating the advantage of explicitly modeling the ancestries of the modern populations (Extended
data Figure 7B). Finally, we investigated how the genomic control correction responded when we
simulated small amounts of admixture from a highly diverged population (Yoruba; 1000 Genomes YRI)
into to a randomly chosen modern population. The genomic inflation factor increased from around 1.2 to
around 1.6 with 10% admixture (Extended data Figure 7C).

Estimating selection coefficients
281

Using the time-serial component of this dataset, we investigated in detail the trajectories of loci which
were genome-wide significant, or for which strong evidence of selection in Europe had previously been
reported. We estimated selection coefficients by fitting a hidden Markov model to the observed allele
counts using an EM algorithm[21]. Since this model relies on a binomial sampling likelihood, we used the
majority call alleles for this analysis. While this model assumes a continuous, constant size population,
which is not the case for Europe, we reasoned that the effect of turnover and admixing populations would
be to increase the variance of the allele frequency trajectory, and that we could compensate for this to some
extent by excluding western hunter-gatherers and estimating the effective population size from neutral
sites. Using 1,000 randomly sampled sites, the maximum likelihood estimate of the diploid effective
population size $2N_e$ was 5,998 (95% confidence interval, CI=5,199-6,979). For each sample we assumed
its age was given by the midpoint of the estimated age range (archaeological or ^{14}C), and converted to
generations by assuming a mean generation time of 29 years.
294

295   ## Test for polygenic selection
296

We implemented the test for polygenic selection described by Ref. 42. This test evaluates whether trait-
associated alleles, weighted by their effect size, are overdispersed compared to randomly sampled alleles,
in the directions associated with the effects measured by genome-wide association studies (GWAS). For
each trait, we obtained a list of significant SNP associations and effect estimates from GWAS data, and
then applied the test both to all populations combined and to selected pairs of populations. We estimated
frequencies in each population by computing the MLE, using the likelihood described above. For each test,
we sampled SNPs frequency matched in 20 bins, computed the test statistic Q_X and for ease of comparison,
converted these to Z scores, signed according the direction of the genetic effects. Theoretically Q_X has a
χ^2 distribution but in practice it is overdispersed so we report bootstrap p-values computed by sampling
10,000 sets of frequency matched SNPs. For this analysis we grouped populations together by period and
location so, for example, "Central_MN" means all Middle Neolithic populations from Central Europe
(present-day Germany, see Table 1). We repeated the analysis using the majority-called dataset.

	Population	Location	Sample size	Majority size	Diploid size	Date (kybp)	Reference (samples)
Western Hunter-gatherers			3	3.60	4.85	7.7-8.1	
WHG	Loschbour	Luxembourg	1	1.98	1.98	8.1	11
	LaBrana1	Spain	1	0.94	1.60	7.8	12
	HungaryGamba_HG	Hungary	1	0.69	0.88	7.7	7 (K01)
Early Neolithic			26	15.27	22.77	6.9-7.2	
EN	Starcevo_EN	Hungary	1	0.28	0.32	7.6	9
	Stuttgart	Germany	1	1.97	1.97	7.2	11
	Spain_EN	Spain	4	3.08	5.21	7.2	9
	LBK_EN	Germany	12	6.56	10.41	7.1	9
	LBKT_EN	Hungary	1	0.09	0.09	7.1	9
	HungaryGamba_EN	Hungary	7	3.29	4.77	6.9	7 (K02,NE1-NE7)
Middle Neolithic			9	5.27	9.62	5.2-5.8	
MN	Spain_MN	Spain	4	2.84	4.99	5.8	9
	Baalberge_MN	Germany	3	0.64	0.71	5.4	9
	Iceman	Italy	1	0.99	1.93	5.3	10
	Esperstedt_MN	Germany	1	0.79	1.99	5.2	9
Late Neolithic			15	8.90	14.01	4.2-4.8	
LN	HungaryGamba_CA	Hungary	1	0.62	0.77	4.8	7 (C01)
	Alberstedt_LN	Germany	1	0.99	1.93	4.4	9
	Corded_Ware_LN	Germany	4	2.56	4.47	4.4	9
	Bell_Beaker_LN	Germany	6	3.19	4.63	4.3	9
	BenzigerodeHeimburg_LN	Germany	3	1.55	2.21	4.2	9
Bronze Age			10	6.66	10.86	3.6-4.8	
BA	Unetice_EBA	Germany	7	4.20	6.48	4.0	9
	HungaryGamba_BA	Hungary	2	1.51	2.59	3.6	7 (BR1,2)
	Halberstadt_LBA	Germany	1	0.95	1.79	2.1	9
Scandinavian Hunter-gatherers			11	6.79	10.88	4.8-7.7	
SHG	SwedenSkoglund_MHG	Sweden	1	0.09	0.09	7.4	13 (StoraForvar11)
	Motala_HG	Sweden	7	5.55	9.14	7.7	9
	SwedenSkoglund_NHG	Sweden	3	1.14	1.64	4.8	13 (Ajvide52,58,70)
Yamnaya			9	5.11	7.60	5.0	
	Yamnaya	Russia	9	5.11	7.60	5.0	9
Modern			503	1006	1006	0.0	
	CEU	NW Europe	99	198	198	0.0	14
	FIN	Finland	99	198	198	0.0	14
	GBR	Great Britain	91	182	182	0.0	14
	IBS	Spain	107	214	214	0.0	14
	TSI	Italy	107	214	214	0.0	14

Table 1: Samples analyzed in this study. **Population**: samples grouped by a combination of date, archaeology and genetics. **Population:** Labels used in Ref. [9]. **Location**: Present-day country where samples originated. In the main text, we refer to samples from present-day Spain, Germany, Hungary and Sweden as Iberian, Central European, Eastern European and Scandinavian, respectively. **Sample size**: Number of individuals sampled. **Majority size**: Average over sites of the number of chromosomes observed at each SNP targeted, if we make a single majority call at each site for each individual. Equivalently, the average number of samples hit at least once. **Diploid size**: Average over sites of the effective number of chromosomes when we use genotype likelihoods. Computed as 2 per sample for samples with genotype calls, or $2 - 0.5^{(c-1)}$ for samples with read depth c. **Date**: Mean of the best date available for the samples in each population, thousands of years before present. **Reference**: Original reference for each sample (and specific sample names, where appropriate).

Figure 1: Genome-wide scan for selection. **A**: Quantile-quantile (QQ) plot of -$\log_{10}$ p-value for potentially functional SNPs (red) and probably neutral SNPs (blue), after genomic control (GC) correction. **B**: QQ plots for different categories of potentially functional SNPs (Methods). All curves are significantly different from neutral expectation. **C**: Plot showing the GC-corrected –$\log_{10}$ p-value for each marker. The red dashed line represents a genome-wide significance level of $10^{-6.79}$. Insets show chromosomes 11 and 15 on a larger scale.

Figure 2: Time series of derived allele frequencies for alleles with genome-wide significant signals of selection, or otherwise mentioned in the text. **A**: Estimated mainland European frequencies. Boxes show the estimated frequency and approximate time range of the observations. Small numbers give the effective sample size as described in Table 1. **B**: Frequencies in other ancient populations. **C**: Frequencies in modern European populations from the 1000 Genomes Project.

Figure 3: Evidence for polygenic selection in Europe. **A-C**: Each row represents a different trait (Height, body mass index, waist-hip ratio, type 2 diabetes, irritable bowel disease, and lipid levels), and each point represents a test statistic. Red, labeled points have bootstrap p-values < 0.01. **A**: Z scores for each population tested against CEU **B**: Z scores for the difference between populations existing at approximately the same time. Populations ordered so the difference is positive. **C**: Z scores for the difference between populations existing at the same location. Populations ordered so the difference is positive. **D**: A hypothesis, based on these results, for when selection on height may have occurred (see Table 1 for abbreviations).

References

Bustamante, C. D. *et al.* Natural selection on protein-coding genes in the human genome. *Nature* **437**, 1153-1157 (2005).
Grossman, S. R. *et al.* Identifying recent adaptations in large-scale genomic data. *Cell* **152**, 703-713 (2013).
Nielsen, R. *et al.* A scan for positively selected genes in the genomes of humans and chimpanzees. *PLoS biology* **3**, e170 (2005).
Voight, B. F., Kudaravalli, S., Wen, X. & Pritchard, J. K. A map of recent positive selection in the human genome. *PLoS biology* **4**, e72 (2006).
Pickrell, J. K. *et al.* Signals of recent positive selection in a worldwide sample of human populations. *Genome research* **19**, 826-837 (2009).
Grossman, S. R. *et al.* A composite of multiple signals distinguishes causal variants in regions of positive selection. *Science* **327**, 883-886 (2010).
Gamba, C. *et al.* Genome flux and stasis in a five millennium transect of European prehistory. *Nature communications* **5**, 5257 (2014).
Wilde, S. *et al.* Direct evidence for positive selection of skin, hair, and eye pigmentation in Europeans during the last 5,000 y. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 4832-4837 (2014).
Haak, W. *et al.* Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature* **Advance Online Publication** (2015).
Keller, A. *et al.* New insights into the Tyrolean Iceman's origin and phenotype as inferred by whole-genome sequencing. *Nature communications* **3**, 698 (2012).
Lazaridis, I. *et al.* Ancient human genomes suggest three ancestral populations for present-day Europeans. *Nature* **513**, 409-413 (2014).
Olalde, I. *et al.* Derived immune and ancestral pigmentation alleles in a 7,000-year-old Mesolithic European. *Nature* **507**, 225-228 (2014).
Skoglund, P. *et al.* Genomic diversity and admixture differs for Stone-Age Scandinavian foragers and farmers. *Science* **344**, 747-750 (2014).
The 1000 Genomes Project Consortium. An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**, 56-65 (2012).
Skoglund, P. *et al.* Origins and genetic legacy of Neolithic farmers and hunter-gatherers in Europe. *Science* **336**, 466-469 (2012).
Devlin, B. & Roeder, K. Genomic control for association studies. *Biometrics* **55**, 997-1004 (1999).
Bersaglieri, T. *et al.* Genetic signatures of strong recent positive selection at the lactase gene. *American journal of human genetics* **74**, 1111-1120 (2004).
Donnelly, M. P. *et al.* A global view of the OCA2-HERC2 region and pigmentation. *Human genetics* **131**, 683-696 (2012).
Sabeti, P. C. *et al.* Positive natural selection in the human lineage. *Science* **312**, 1614-1620 (2006).
Enattah, N. S. *et al.* Identification of a variant associated with adult-type hypolactasia. *Nat Genet* **30**, 233-237 (2002).
Mathieson, I. & McVean, G. Estimating selection coefficients in spatially structured populations from time series data of allele frequencies. *Genetics* **193**, 973-984 (2013).
Burger, J., Kirchner, M., Bramanti, B., Haak, W. & Thomas, M. G. Absence of the lactase-persistence-associated allele in early Neolithic Europeans. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 3736-3741 (2007).
Lamason, R. L. *et al.* SLC24A5, a putative cation exchanger, affects pigmentation in zebrafish and humans. *Science* **310**, 1782-1786 (2005).
Quillen, E. E. *et al.* OPRM1 and EGFR contribute to skin pigmentation differences between Indigenous Americans and Europeans. *Human genetics* **131**, 1073-1080 (2012).

Sturm, R. A. *et al.* A single SNP in an evolutionary conserved region within intron 86 of the HERC2 gene determines human blue-brown eye color. *American journal of human genetics* **82**, 424-431 (2008).

Eiberg, H. *et al.* Blue eye color in humans may be caused by a perfectly associated founder mutation in a regulatory element located within the HERC2 gene inhibiting OCA2 expression. *Human genetics* **123**, 177-187 (2008).

Han, J. *et al.* A genome-wide association study identifies novel alleles associated with hair color and skin pigmentation. *PLoS genetics* **4**, e1000074 (2008).

Branicki, W., Brudnik, U. & Wojas-Pelc, A. Interactions between HERC2, OCA2 and MC1R may influence human pigmentation phenotype. *Annals of human genetics* **73**, 160-170 (2009).

Bokor, S. *et al.* Single nucleotide polymorphisms in the FADS gene cluster are associated with delta-5 and delta-6 desaturase activities estimated by serum fatty acid ratios. *Journal of lipid research* **51**, 2325-2333 (2010).

Tanaka, T. *et al.* Genome-wide association study of plasma polyunsaturated fatty acids in the InCHIANTI Study. *PLoS genetics* **5**, e1000338 (2009).

Teslovich, T. M. *et al.* Biological, clinical and population relevance of 95 loci for blood lipids. *Nature* **466**, 707-713 (2010).

Ahn, J. *et al.* Genome-wide association study of circulating vitamin D levels. *Human molecular genetics* **19**, 2739-2745 (2010).

Wang, T. J. *et al.* Common genetic determinants of vitamin D insufficiency: a genome-wide association study. *Lancet* **376**, 180-188 (2010).

Price, A. L. *et al.* The impact of divergence time on the nature of population structure: an example from Iceland. *PLoS genetics* **5**, e1000505 (2009).

Wellcome Trust Case Control, C. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* **447**, 661-678 (2007).

Fujimoto, A. *et al.* A scan for genetic determinants of human hair morphology: EDAR is associated with Asian hair thickness. *Human molecular genetics* **17**, 835-843 (2008).

Kimura, R. *et al.* A common variation in EDAR is a genetic determinant of shovel-shaped incisors. *American journal of human genetics* **85**, 528-535 (2009).

Kamberov, Y. G. *et al.* Modeling recent human evolution in mice by expression of a selected EDAR variant. *Cell* **152**, 691-702 (2013).

Loh, P. R. *et al.* Inferring admixture histories of human populations using linkage disequilibrium. *Genetics* **193**, 1233-1254 (2013).

Pritchard, J. K., Pickrell, J. K. & Coop, G. The genetics of human adaptation: hard sweeps, soft sweeps, and polygenic adaptation. *Current biology* **20**, R208-215 (2010).

Turchin, M. C. *et al.* Evidence of widespread selection on standing variation in Europe at height-associated SNPs. *Nature genetics* **44**, 1015-1019 (2012).

Berg, J. J. & Coop, G. A population genetic signal of polygenic adaptation. *PLoS genetics* **10**, e1004412 (2014).

Kruts, S. *Палеоантропологические исследования степного Приднепровья: эпокха бронзы (Paleoanthropological investigation of the Pridnepr steppe)*. (Наук. думка, 1984).

Lango Allen, H. *et al.* Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature* **467**, 832-838 (2010).

Speliotes, E. K. *et al.* Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nature genetics* **42**, 937-948 (2010).

Heid, I. M. *et al.* Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. *Nature genetics* **42**, 949-960 (2010).

Morris, A. P. *et al.* Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nature genetics* **44**, 981-990 (2012).

Jostins, L. *et al.* Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* **491**, 119-124 (2012).

International HapMap Consortium. A second generation human haplotype map of over 3.1 million SNPs. *Nature* **449**, 851-861 (2007).

Pruim, R. J. *et al.* LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics* **26**, 2336-2337 (2010).

Extended data

Extended data Figure 1: Fine scale maps of genome-wide significant signals of selection. In each case, the most significant SNP is labeled, and other SNPs are colored according to their LD with the most significant SNP. The blue line shows the recombination rate and the lower panel in each plot shows gene locations.

Generated using locuszoom[50] (http://csg.sph.umich.edu/locuszoom/).

Extended data Figure 2: Genome wide scan for selection using majority called alleles. As Figure 1 in the main text but calling a single majority allele for each site for the ancient samples. **A**: QQ plot of neutral (blue) and possibly functional (red) sites. **B**: QQ plot of possibly functional sites broken down by category. **C**: Manhattan plot of scan results.

Extended data Figure 3: Estimated frequencies of the derived allele of rs12913832 in HERC2. We divided our samples by date and location. Boxes show the maximum likelihood estimate of the frequency as in main text Figure 2 and vertical lines show approximate 95% confidence intervals.

rs3827760

Extended data Figure 4: Motala haplotypes carrying the derived, selected EDAR allele. This figure compares the genotypes at all sites within 100kb of rs3827760 (in blue) for the 7 Motala samples and 20 randomly chosen CHB (Chinese from Beijing) and CEU (Central European) samples. Each row is a sample, and each column is a SNP. Grey means homozygous for the minor (in CEU) allele. Pink denotes heterozygotes and red homozygotes for the other allele. For the Motala samples, an open circle means that there is only a single read, otherwise the circle is colored according to the number of reads observed. Note that the original haplotype on which rs3827760 arose appears to be common in CEU (marked with arrows on the right). Four of the Motala samples appear to be heterozygous for the rs3827760 haplotype, and one of the samples carries the original haplotype without the derived rs3827760 allele.

A

B

C

Extended data Figure 5: Test for polygenic selection using frequencies estimated from majority-called alleles rather than genotype likelihoods. As panels A-C in main text Figure 3. Comparisons with bootstrap p<0.01 are in red and labeled. **A**: Z scores for tests relative to CEU. **B**: Z scores for tests between populations that exist at the same times. **C**: Z scores for tests between populations that exist in the same place.

Extended data Figure 6: Predicted genetic heights for populations split up by region and date. We computed the maximum likelihood allele frequencies for each SNP that was significant in the analysis in Ref. 44 and multiplied by the effect size from that study. The genetic height here is expressed in terms of regression effect sizes (roughly, number of standard deviations after correction for age and sex), translated so that the mean across populations is zero.

Extended data Figure 7: Power and robustness of the selection scan. **A**: Estimated power to detect selection at genome-wide significance under different scenarios, as a function of selection coefficent. Dashed lines indicate slelection in all populations and dotted lines indicate selection in only a single population. Different colors indicate selection over different periods of time: 50, 100 and 200 generations. **B**: Effect of misspecifying the mixture matrix C, over 10,000 simulations. In blue (left hand axis), the genomic inflation factor as a function of the proportion of the matrix that is random. In orange (right hand axis), the corresponding power, for a selection coefficient of 0.02 over a period of 100 generations. **C**: Genomic inflation constant as a function of admixture from Yoruba into a random modern population.