Sex chromosome dosage compensation in *Heliconius* butterflies: global yet still incomplete?

James R. Walters[1]*, Thomas J. Hardcastle[2], and Chris D. Jiggins[3]

1) Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS, USA
2) Department of Plant Biology, University of Cambridge, Cambridge, UK
3) Department of Zoology, University of Cambridge, Cambridge, UK

*Author for correspondence: James R Walters, Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS, USA
phone: 785-864-6341
email: jrwalters@ku.edu

Abstract

The evolution of heterogametic sex chromosome is often – but not always – accompanied
by the evolution of dosage compensating mechanisms that mitigate the impact of sex-
specific gene dosage on levels of gene expression. One emerging view of this process is that
such mechanisms may only evolve in male-heterogametic (XY) species but not in female-
heterogametic (ZW) species, which will consequently exhibit "incomplete" sex
chromosome dosage compensation. However, some recent results from moths suggest that
Lepidoptera (moths and butterflies) may prove to be an exception to this prediction. Here
we report an analysis of sex chromosome dosage compensation in *Heliconius* butterflies,
sampling multiple individuals for several different adult tissues (head, abdomen, leg,
mouth, and antennae). Methodologically, we introduce a novel application of linear mixed-
effects models to assess dosage compensation, offering a unified statistical framework that
can estimate effects specific to chromosome, to sex, and their interactions (*i.e.,* a dosage
effect). Our results show substantially reduced Z-linked expression relative to autosomes
in both sexes, as previously observed in bombycoid moths. This observation is consistent
with an increasing body of evidence that at least some species of moths and butterflies
possess an epigenetic sex chromosome dosage compensating mechanism that operates by
reducing Z chromosome expression in males. However, this mechanism appears to be
imperfect in *Heliconius*, resulting in a modest dosage effect that produces an average 5-
20% male-bias on the Z chromosome, depending on the tissue. Strong sex chromosome
dosage effects have been previously in a pyralid moth. Thus our results reflect a mixture of
previous patterns reported for Lepidoptera and bisect the emerging view that female-
heterogametic ZW taxa have incomplete dosage compensation because they lack a
chromosome-wide epigenetic mechanism mediating sex chromosome dosage
compensation. In the case of *Heliconius*, sex chromosome dosage effects persist apparently
despite such a mechanism. We also analyze chromosomal distributions of sex-biased genes
and show an excess of male-biased and a dearth of female-biased genes on the Z
chromosome relative to autosomes, consistent with predictions of sexually antagonistic
evolution.

Introduction:

Dosage compensation is a gene-regulatory mechanism that equalizes levels of gene expression in response to differences in gene dose (i.e. copy number). Without dosage compensation, changes in gene dose can substantially affect gene expression, potentially resulting in detrimental effects on finely-tuned gene networks (Birchler et al. 2001). For this reason, it was long assumed that the evolution of a chromosome-wide dosage compensating mechanism was an essential step in the evolution of heteromorphic sex chromosomes (Vicoso & Bachtrog 2009; Disteche 2012; Mank 2013).

Heteromorphic sex chromosomes typically evolve from homologous autosomes that acquire a sex-determining locus, accumulate sexually-antagonistic alleles, and suppress recombination (B Charlesworth 1996; B Charlesworth & D Charlesworth 2000; Bachtrog 2006) . Eventually, substantial gene loss and degeneration occurs on the nascent Y chromosome (or W in female-heterogametic ZW taxa) (Rice 1984; B Charlesworth & D Charlesworth 2000; Bachtrog 2013). The erosion of genes from the Y/W presents the problem of balancing gene expression with the autosomes. Monosomy of the X/Z in one sex means the dose (*i.e.,* copy number) of most sex-linked genes differs by half between the sexes. If the resulting sex-linked gene expression becomes similarly unbalanced, degradation of the Y/W could impose a substantial fitness cost for the heterogametic sex due to impaired dosage-sensitive interactions with autosomal loci (Ohno 1967; Mank 2009; Pessia et al. 2013). It is thus predicted that a global dosage-compensating mechanism should evolve to balance the expression of X/Z loci relative to autosomes as the Y/W degrades (Ohno 1967; Brian Charlesworth 1978; Mank 2013; Veitia et al. 2015). This process should also result, indirectly, in balanced gene expression between the sexes for the X/Z. One important working assumption in this framework is that average expression is approximately equal across autosomes, therefore *complete* dosage compensation should yield X:A (or Z:A) expression ratios of ~1 in both sexes (Mank 2009; 2013; Nguyen & Disteche 2006; Vicoso & Bachtrog 2011b; Smith et al. 2014; Walters & Hardcastle 2011).

Efforts to evaluate this hypothesis have expanded greatly in the last decade with the application of genome-wide expression analyses via microarray or RNA-seq, and have yielded several unexpected results. The history and contemporary findings of research on sex chromosome dosage compensation are extensively reviewed in several recent publications (Disteche 2012; Mank 2013; Pessia et al. 2013; Ferrari et al. 2014). Here we briefly highlight details particularly relevant to our current results. Where investigated during the pre-genomic decades, the limited results obtained tended to support theoretical predictions. Initial genome-wide investigations using microarrays in established model organisms such as mouse, humans, fruit flies, and nematodes – all male heterogametic species – yielded patterns consistent with global sex chromosome dosage compensation. As predicted, average expression on the X was comparable to autosomes (X:A ~ 1) in both

sexes and the canonical view of dosage compensation and sex chromosome evolution
appeared robust (Hamada 2005; Gupta et al. 2006; Nguyen & Disteche 2006). Moreover,
the recognition that distinct molecular mechanism underlay dosage compensation in flies,
worms, and humans added further support to the universality of dosage compensation
evolving concomitantly with differentiated sex chromosomes (Deng et al. 2011; Straub &
Becker 2011). However, more recent research has added substantial complexity and
controversy to the issue. In particular, evaluating dosage compensation in the context of
ancestral expression levels indicates that eutherian mammals should be considered to have
X:A expression ratio of ~0.5 (Julien et al. 2012; F Lin et al. 2012). Importantly, in all of these
cases, sex-linked expression appears to be balanced between males and females and there
is no evidence of a gene dosage-effect on X chromosome expression.
Other striking exceptions to the canonical theory of dosage compensation were
observed when investigations of sex chromosome dosage compensation expanded into
novel taxa. Notably, several female-heterogametic taxa exhibit incomplete dosage
compensation: Male birds, snakes, and schistosomes are homogametic (ZZ) with Z:A $\sim$ 1,
but in females the Z:A ratio is significantly less than 1 (Itoh et al. 2007; Vicoso & Bachtrog
2011a; Vicoso, Emerson, et al. 2013a; Mank & Ellegren 2009; Naurin et al. 2011). The
apparently dichotomous pattern of completely dosage compensated XY taxa versus
incompletely compensated ZW species catalyzed strong suggestions and some nascent
theory claiming that global sex chromosome dosage compensation should be universally
absent from ZW taxa and occur only XY species (Bachtrog et al. 2011; Naurin et al. 2012;
Mank 2013; Vicoso, Emerson, et al. 2013a). We call this the *heterogametic dichotomy*
hypothesis.
Despite the emerging evidence supporting the heterogametic dichotomy hypothesis,
there is at least one conspicuous female-heterogametic taxon that may prove to be an
exception: Lepidoptera (moths and butterflies). The status of sex chromosome dosage
compensation in Lepidoptera is currently ambiguous, primarily due to inconsistent results
reported by the few studies currently available. The first genome-wide assessment of sex
chromosome dosage compensation in Lepidoptera was based on a microarray dataset in
the silkmoth, *Bombyx mori* (Xia et al. 2007). An initial analysis of these data reported
incomplete dosage compensation similar to other ZW taxa (Zha et al. 2009). However,
analytical flaws were later identified in this initial effort and a subsequent reanalysis
indicated the Z:A ratio was equal in males and females, offering evidence for global,
complete sex chromosome dosage compensation (Walters & Hardcastle 2011).
Intriguingly, this reanalysis further revealed a Z:A ratio of ~0.7, significantly less than one.
This result is not anticipated by current theory concerning sex chromosome evolution.
Additional evidence for complete sex chromosome dosage compensation in Lepidoptera
was more recently reported in another bombycoid moth, *Manduca sexta* (tobacco
hormworm), using RNA-seq (Smith et al. 2014). Again the Z:A ratio was equal between
sexes, but in this case the Z:A ratio was only marginally less then 1, with significance

depending on filtering thresholds. In stark contrast to results from Bombycoid moths,
RNA-seq analysis of the Pyralid moth *Plodia interpunctella* (Indian meal moth) showed no
evidence for dosage compensation, with female Z:A~0.5 while male Z:A was ~ 1 (Harrison
et al. 2012). This is the largest magnitude of sex chromosome dosage effect yet reported,
exceeding patterns in observed in birds and snakes (Itoh et al. 2007; Vicoso & Bachtrog
2011a; Vicoso, Emerson, et al. 2013a; Mank & Ellegren 2009; Naurin et al. 2011).

These studies differed considerably in what tissues or body parts were assayed. The *B.*
mori microarray data included samples from 10 different larval body parts (Xia et al. 2007;
Walters & Hardcastle 2011). The *M. sexta* study sampled only adult heads while pools of
whole adult *P. interpunctella* were used (Harrison et al. 2012; Smith et al. 2014). These
latter two studies both constructed *de novo* transcriptome assemblies and assigned
chromosomal linkage based on contig homology to *B. mori.* This approach of assigning Z-
linkage is seemingly robust because synteny is highly conserved in Lepidoptera (Pringle et
al. 2007; Heliconius Genome Consortium 2012; Yue et al. 2013; Ahola et al. 2014).

However, only the *B. mori* dataset, which included isolated gonads, provided any
opportunity to assess patterns of dosage compensation separately in somatic and
reproductive tissues. It is well-established that, at least in *B. mori*, the Z chromosome is
enriched for highly-expressed, testes-specific genes but depleted of ovary-specific
transcripts (Arunkumar et al. 2009; Suetsugu et al. 2013). Thus, inclusion of gonadal tissue
may substantially skew patterns of Z:A ratios to appear "uncompensated" even when
somatic Z:A ratios are otherwise comparable between the sexes (Walters & Hardcastle
2011).

In this manuscript we report the first genomic analysis of sex chromosome dosage
compensation in a butterfly. Using RNA-seq, we assay male and female gene expression in
several body parts of *Heliconius melpomene* and its closely related sibling species, *H. cydno*
(Martin et al. 2013; Quek et al. 2010). The existence of a complete reference genome and
linkage map for *H. melpomene* facilitates a nuanced inference of chromosome and sex
specific effects on gene expression, which we achieve through the novel application of
mixed-effects linear models to analyze dosage compensation (Heliconius Genome
Consortium 2012). Our results show substantially reduced Z-linked expression relative to
autosomes in both sexes, but also modest dosage effect on the Z chromosome, and thus
reflect a mixture of previous patterns reported for Lepidoptera.
161

Methods:

Samples and Sequencing

Two groups of RNA-seq data were used to estimate gene expression. First, we used the
paired-end data sequenced from *H. melpomene* generated by Briscoe et al. (Briscoe et al.

2013) ArrayExpress ID: E-TAB-1500. This data set includes three male and three female
samples from mouth, leg, and antennae. Additionally, we generated new RNA-seq data from
adult head (excluding antennae) and abdomen. For these samples, *Heliconius* butterflies
were reared in large insectaries in Gamboa, Panama. Insectary populations were recently
established from local natural populations. Males and females were kept separate after
eclosion and aged 6 days before collection to allow reproductive tissues to develop
(Dunlap-Pianka et al. 1977); all samples were virgins. Head and abdomen tissues were
collected into RNAlater and stored frozen before RNA purification. Total RNA was
extracted with TRIzol reagent (Invitrogen, Carlsbad, CA), purified using RNeasy columns
(Qiagen, Valencia, CA), and treated with TURBO DNase (Life Technologies, Grand Island,
NY) following manufacturer's instructions. Messenger RNA was isolated from total RNA via
poly-A pulldown and subsequently transformed into a cDNA library using the Illumina
TruSeq sample preparation kits. Paired-end 100 basepair sequencing was performed on an
Illumina HiSeq.

Read mapping and normalization
Read mapping and estimation of fragment counts per gene were performed using RSEM
(v1.2.11) running Bowtie2 (v2.1.0) (Li & Dewey 2011; Langmead & Salzberg 2012). Reads
were mapped as paired-end data, with the first 9 bp and the final 40 bp trimmed before
mapping to remove low and variable-quality bases. All subsequent statistical analyses
were performed using R and BioConductor, especially the baySeq package for assessing
differential expression (Hardcastle & Kelly 2010; R Development Core Team 2014). The
library scaling factors were calculated as the sum of non-zero gene expression levels below
the 75[th] percentile of gene expression, following the example of Hardcastle et al. (2012) as
an amendment to Bullard *et al.* (2010) . Because of the potential for substantial Z
chromosome effects on gene expression, only known autosomal loci were considered when
calculating library scaling factors. For non-parametric statistical analyses, we further
normalized expression levels as fragments per kilobase per million mapped reads (FPKM).
193
Failing to remove transcriptionally inactive genes from RNA-seq data sets assayed for sex
chromosome dosage compensation can result in problematic biases, yet the most
appropriate filtering method to apply is not well-established (Xiong et al. 2010;
Kharchenko et al. 2011; Jue et al. 2013; Smith et al. 2014). Here we employ a probabilistic
approach to assessing whether a given locus is expressed. The baySeq framework
calculates the posterior likelihood (ranging from 0 to 1) that a given locus has no true
expression. (i.e., any observed reads should be considered "noise", not signal) (Hardcastle
2014).   We primarily report results filtered at a value of 0.5, but results are comparable
across a range thresholds tested, from 0.25 to 0.9 (see Supplementary Information)
203

Assessing Z chromosome and dosage effects: non-parametric statistics

To test for Z chromosome dosage effects, we compared the expression of Z-linked and autosomal loci in males and females. Greater than 80% of predicted coding loci have been mapped to the 21 chromosomes of *H. melpomene*, with Z-linkage validated or corrected based on sex-specific genome sequencing coverage as reported in Martin *et al.* 2013 (Martin et al. 2013; Heliconius Genome Consortium 2012). For each body part sampled, replicates were averaged by sex to give mean male and female FPKM values for each locus. Within each sex, Z-linked versus autosomal (Z:A) expression was compared, with differences in average expression evaluated via Mann-Whitey U test (MWU). The average male:female (M:F) expression ratio of Z-linked versus autosomal loci was also compared via MWU. Only loci actively expressed in both sexes were included when analyzing M:F ratios.

We further explored the effects of Z chromosome dosage by comparing the average expression of Z-linked genes in males and females, split by quartiles of expression magnitude. Within quartile, differences between sex in average Z-linked expression was tested via MWU. Only loci actively expressed in both sexes were included in this analysis. This analysis closely resembles an analysis performed by Harrison *et al.* (2012) aimed at assessing how Z chromosome dosage effects depend on expression magnitude. However, we have slightly modified the analysis to avoid a bias we believe is inherent in the analysis as originally performed by Harrison *et al.* (2012). Rather than basing expression quartiles solely on male expression as was previously done, we calculated quartiles based on the maximum of male or female expression for each locus. The reasoning for this modification and the potential biases arising from ranking genes using data from only one sex is provided in the Appendix.

Assessing sex-chromosome and dosage effects: Linear modeling of expression levels

In addition to the application of non-parametric MWU tests, as is typically employed for investigations of sex-chromosome dosage compensation, we implemented a linear modeling framework to test for dosage and Z-specific effects on gene expression.

The count data (after filtering) were fitted via maximum likelihood methods (Bates et al. 2014) to a generalized linear mixed-effects model assuming a Poisson distribution and a log linkage function, using library scaling factor and gene length as offsets. We fit a series of models using various effects of chromosome, Z-linkage and sex, together with relevant interactions between these effects. In all models, a per gene random effect was applied over sex, simultaneously allowing for variation in individual gene expression and differential expression of genes between the sexes.

Using this modeling framework, we first tested for a global effect of Z-linkage on expression
by comparing a model with both **sex** and **Zlink** as fixed effects (but no interaction)

$X_{\{gene,sample,sex,Z\text{-}linkage\}} \sim$ Z-linkage + sex + (sex|gene)

versus a model with **sex** as the sole fixed effect.

$X_{\{gene,sample,sex,Z\text{-}linkage\}} \sim$ sex + (sex|gene)

We then additionally tested for sex-specific effect on Z-expression (*i.e.,* a dosage effect), by
fitting the full model

$X_{\{gene,sample,sex,Z\text{-}linkage\}} \sim$ Z-linkage + sex + Z-linkage $\times$ sex + (sex|gene)

to one without the interaction term.

Chromosomal distribution of sex-biased genes

Genes differentially expressed between males and females were identified using baySeq
(Hardcastle & Kelly 2010). We applied a false-discovery rate of 0.05 and required at least a
1.5-fold change in expression between sexes. To examine the chromosomal distribution of
sex-biased genes, we counted the number of male, female, and unbiased genes among all
actively expressed genes on each chromosome and also the unmapped scaffolds not yet
assigned to chromosome. Gene activity was based on the probabilistic criteria, and
assessed independently in males and females, so a gene expressed in males but not females
was counted as male biased (and vice versa). Differences between the Z and autosomes in
proportion of sex-biased genes were tested using a Fisher's exact test.

Results

Sequencing and read mapping

Data sets generated new for this project were considerably larger that those generated by
Briscoe et al. (2013). For the 40 samples sequenced new here, the mean number of total
reads sequenced was ~130 M, ranging from ~30 M to ~ 284 M reads. The 18 samples from
Briscoe et al. (2013) gave a mean of ~23 M, ranging from ~8 M to ~46 M reads. The
proportion of reads aligned with RSEM was also generally higher for the abdomen and
head samples compared to the Briscoe et al samples from antennae, mouth, and leg. The 40
abdomen and head samples yielded a mean of 49% aligned, ranging from 21 to 65%. Mean

percent mapping in the 18 Briscoe et al. (2013) samples was 37%, ranging from 9 to 55%.
A complete summary of read counts and alignment statistics calculated with Picard's
CollectAlignmentSummaryMetrics utility (http://picard.sourcefource.net) is provided in
supplementary Table S1. Newly generated data sets for *H. melpomene* and *H. cydno* head
and abdomen are available from the Sequencing Read Archive (SRA) preojct: XXXXX.
283

Z-chromosome to autosome comparisons
For both sexes, all body parts in both species yielded average Z expression significantly
lower than autosomal expression, as is evident from both linear modeling and non-
parametric analyses (Fig 1 and Tables 1 & 2). Comparing median expression, the Z:A ratio
ranged from about 0.5 to 0.75; ratios based on mean values are even lower. Linear
modeling of these data showed Z-linkage had a significant negative effect on gene
expression relative to autosomal average (Table 2). Average expression of individual
autosomes varied somewhat in both sexes, but in nearly all cases the median Z expression
was lower than any autosome (supplementary Fig 1)
293

Notably, in several body parts sampled, the Z:A ratio was slightly greater in males than in
females (Table 1). This apparent sex-effect on Z-chromosome expression was statistically
confirmed with linear modeling. All tissues except antennae showed a significant
interaction between sex and Z-linked expression, with greater male Z-linked expression
(Table 2). This interaction can be interpreted as dosage effect of the Z chromosome on
average gene expression. Note, however, that the magnitude of this interaction (dosage)
effect is distinctly less than the effect of the Z-chromosome on gene expression levels.
301

Male:Female expression ratios
Consistent with the dosage effect observed in the linear models, the Z chromosome showed
a modest but consistent male bias in expression relative to autosomes in all samples
(Figure 2 and Table 3). The distribution of M:F expression ratios was significantly greater
on the Z (MWU $p < 0.05$ after bonferroni correction) in all samples except antennae. The
average magnitude of this Z-linked male expression bias ranged from 5-20% relative to
autosomes based on the median M:F expression ratios (Table 2).

Quartile analysis of Z expression
Comparing Z-linked expression between sexes split by quartiles of expression showed no
obvious pattern of discrepancy in male versus female Z-linked expression for tissues other
than abdomen (Figure 3; supplemental Figure S2). In both abdomen samples, male
expression was significantly greater than female in the fourth (highest expression) quartile,
assuming a bonferroni correction for four tests.

Chromosomal distribution of sex-biased genes

The amount of sex-biased gene expression differed substantially between body parts. Tissues other than in the abdomen showed almost no differential expression between sexes using the criteria we applied (Table 4). In contrast, roughly 30% of active genes in abdomen were differentially expressed. We therefore analyzed the chromosomal distribution of sex-biased genes only in the abdomen. We observed a significant difference in the proportions of sex-biased genes on the Z versus the autosomes in both *H. melpomene* and *H. cydno* (Fisher's exact test, p-value << 0.001). Figure 4 shows that the Z chromosome is distinctly enriched for male-biased and has a paucity of female-biased.

Discussion

Dosage compensation

Patterns of sex chromosome dosage compensation in *Heliconius* butterflies reflect an interesting amalgam of previous results from Lepidoptera. Similarly to the bombycoid moths *B. mori* and *M. sexta*, *Heliconius* males reduce expression of Z chromosomes distinctly below autosomal expression levels (Walters & Hardcastle 2011; Smith et al. 2014). In the two bombycoid species, male Z expression was low enough to be comparable to the female Z; there was no detectable dosage effect on Z-linked expression. Yet despite having Z:A < 1 in both sexes, *Heliconius* shows a dosage effect such that the Z chromosome retains a modest but consistent male-bias in expression. This result in *Heliconius* echoes the very substantial dosage effects reported for *P. interpunctella*. Unfortunately, the magnitude of these *Heliconius* dosage effects cannot easily be contrasted with *P. interpunctella* because direct male:female expression ratios were not reported for that species (Harrison et al. 2012). However, the discrepancy in Z:A ratios between sexes of *P. interpunctella* is so large that we presume the dosage effects reflected in M:F expression ratios are far greater than we observe in *Heliconius*. Notably, no reduction in male Z-linked expression relative to autosomes was reported for *P. interpuctella*.

Thus in *Heliconius* we have further evidence that some Lepidoptera appear to have a molecular mechanism for globally reducing Z-linked expression in males that compensates for the difference in Z-chromosome dosage between sexes. Curiously, in *Heliconius* this mechanism appears to be operating imperfectly, resulting in a measurable dosage effect that could arguably be called "incomplete" dosage compensation. However, the 5-20% Z-linked male bias in *Heliconius* is distinctly less than the ~50% bias typically reported for vertebrate ZW species (*e.g.,* snakes, birds) identified as having "incomplete" sex chromosome dosage compensation (Ellegren et al. 2007; Itoh et al. 2010; Vicoso, Emerson, et al. 2013a). In the case of the female-heterogametic vertebrates, it is assumed that dosage compensation is "incomplete" because no global epigenetic mechanism exists to offset dosage effects, in contrast to the epigenetic mechanisms known in fruit flies,

nematodes, and mammals. In the case of *Heliconius*, we would argue that "incomplete"
dosage compensation occurs despite a global mechanism operating to balance Z:A ratios
between the sexes, in this case reducing male Z-linked expression similar to that in females.
Further evidence in *Heliconius* that a global mechanism mitigates dosage effects
comes from the lack of detectable transcriptional saturation on the Z chromosome. None of
the purely somatic tissues sampled showed an effect of expression magnitude on
differences between male and female Z-linked expression. This result contrasts with *P.*
interpunctella, where the male expression bias increased with expression level, suggesting
substantial transcriptional saturation due to uncompensated Z chromosome dosage.
However, in abdomens we did observe significantly greater expression in males for the
highest expression quartile, reminiscent of the *P. interpunctella* result. While this could be
regarded as the effect of transcriptional saturation due to gene imbalance (*i.e.,* a dosage
effect), we tend to think it primarily reflects the unusually high expression levels of male-
biased reproductive genes in the testes. If male-biased genes tend to have very high
expression, this could produce the pattern observed. Robustly testing this hypothesis
would require sequencing transcriptomes of gonads separately from somatic tissue; such
data are currently not available in *Heliconius*. However, we note that in male abdomens,
autosomal male-biased genes are expressed at significantly greater levels on average than
unbiased or female biased genes (Kruskal-Wallis test, p << 0.001; Supplemental Figure S3).
In female abdomen, average expression of autosomal sex-biased genes does not differ from
unbiased genes (Kruskal-Wallis test, N.S.). These patterns are consistent with our
hypothesis of highly-expressed testes genes generating a pattern of male over-expression
in the top quartile. Furthermore, this pattern would be exacerbated on the Z chromosome
due to the over-representation of male-biased genes on the Z, as observed here (see below)
and also reported in *B. mori* (Arunkumar et al. 2009; Walters & Hardcastle 2011).
Finally, we note there is additional experimental evidence for a global sex
chromosome dosage mechanism that down-regulates male Z expression, at least in *B. mori*.
Kiuchi *et al.* (2014) recently reported the characterization of a Z-linked *masculinizing*
protein that plays a fundamental role in *B. mori* sex determination. RNAi disruption of this
masc protein does not much alter autosomal expression, but causes chromosome-wide up-
regulation of Z-linked expression. Thus *masc* apparently controls a switch that initiates a
global epigenetic reduction in Z-linked male expression.
What does this current set of results from Lepidoptera mean for the *heterogametic*
dichotomy hypothesis? These different lines of evidence pointing to a global sex
chromosome dosage compensation mechanism in several lepidopteran species undermine
the simple notion that female heterogametic taxa do not evolve such mechanisms.
However, there is substantial room for nuance here. First, despite emerging support for
Lepidoptera as an exception, there is still substantial evidence that global dosage
compensation is much less common in ZW than XY taxa (Mank 2013).   Thus, while

seemingly not a universal rule, the *heterogametic dichotomy hypothesis* still reflects a
compelling and unexplained trend in sex chromosome evolution.
Second, results presented here from *Heliconius* suggest there is a need to separately
consider observed patterns, proximate mechanism, and evolutionary process. In most
cases, reporting that a species shows a pattern of "incomplete" dosage compensation has
been assumed to indicate the organism lacks a global mechanism to mitigate sex
chromosome dosage differences between sexes. We now observe that *Heliconius* butterflies
appear to break this assumption. *Heliconius* seems to share a mechanism with bombycoid
moths that reduces male Z-linked expression in order to balance Z:A expression between
sexes (Walters & Hardcastle 2011; Kiuchi et al. 2014; Smith et al. 2014). In bombycoids
this balancing is "perfect", resulting in "complete" sex chromosome dosage compensation
with no male bias in Z-linked M:F expression ratios. Yet *Heliconius* shows a pattern of
"incomplete" dosage compensation despite apparently deploying this mechanism of down-
regulating the male Z. This result defies a simple characterization as being consistent, or
not, with the *heterogametic dichotomy hypothesis*. On the one hand, opposing the
hypothesis, *Heliconius* butterflies apparently have a global mechanism of dosage
compensation. On the other hand, supporting the hypothesis, dosage effects persist and
compensation is "incomplete". A similar scenario was recently reported for the neo-X of *D.*
pseudoobscura, where the well-characterized dosage-compensation complex appears in
some tissues and developmental stages to be "incompletely" compensating male neo-X
expression to an X:A ratio ~0.85, much greater than 0.5 expected without compensation
but still short of X:A ~1 "complete" compensation observed in other *Drosophila* species and
the ancestral portion of the *D. pseudoobscura* X (Nozawa et al. 2014). The functional and
evolutionary significance of a lingering dosage effect in the presence of a global
compensating mechanism is certainly a promising area for future research addressing the
evolution of sex chromosome dosage compensation, especially the *heterogametic*
dichotomy hypothesis.
The presence of a mechanism that globally reduces Z-linked expression in males is
another point requiring nuanced consideration in light of evolutionary processes. A
reasonable argument can be made that reducing Z-linked expression in males to balance
Z:A expression in females cannot legitimately be considered sex chromosome dosage
compensation, *sensu stricto*. The canonical evolutionary model of dosage compensation
invokes stabilizing selection to maintain ancestral expression levels of sex-linked genes in
the heterogametic sex as gametologs erode from the W (or Y) (Ohno 1967; Charlesworth
1978; Mank 2009a). This model assumes that substantial reduction in the Z:A ratio is
deleterious. Following this theory, sex chromosome dosage compensation should be
defined as dosage *conservation* retaining the ancestral expression patterns where Z:A~1.
(This assumes all autosomes have roughly comparable average expression levels, a pattern
that appears to be true in most species examined (Gupta et al. 2006; Deng et al. 2011; Hong

Lin et al. 2011; Walters & Hardcastle 2011; Vicoso, Emerson, et al. 2013a; Vicoso, Kaiser, et
al. 2013b).)
Thus, following this narrow and canonical view, sex chromosome dosage
compensation should be limited only to situations where the heterogametic sex has Z:A $\sim$ 1
(and, comparably, X:A $\sim$1). Notably, this would include cases where sexual antagonism of
sex-linked expression presumably remains unresolved and results in hyper-expression of
sex-linked genes (i.e., X:A > 1) in the homogametic sex (Prince et al. 2010; Mank et al. 2011;
Allen et al. 2013). But a mechanism that globally reduces Z-linked expression in males to
balance Z:A expression in females, as appears to exist in Lepidoptera, is not consistent with
canonical evolutionary theory regarding dosage compensation and sex chromosomes.
Specifically, if Z:A < 1 in heterogametic females is problematic, why would similarly
reducing Z:A expression ratios in homogametic males resolve this problem? Doing so
would compensate sex chromosome expression relative to autosomes between sexes, but
presumably without dosage *conservation*. These results from Lepidoptera present a
conundrum in light of current evolutionary theory (Walters & Hardcastle 2011).
Some resolution to this conundrum may come from better understanding the
evolutionary history of the lepidopteran Z chromosome. Substantial ambiguity currently
exists around the evolutionary relationship between the Z and W chromosomes. Basal
Lepidoptera and Trichoptera (caddisflies, the sister group to Lepidoptera) are female-
heterogametic but completely lack a W chromosome; females are ZO (Lukhtanov 2000;
Sahara et al. 2011). The W chromosome now shared by most "advanced" (ditrysian)
Lepidoptera arose long after the split between Lepidotera and Trichoptera. One
hypothetical origin of the W invokes a fusion between an autosome and the basal Z
chromosome, with the remaining free autosome following the canonical evolutionary
degradation into the W allosome shared by ditrysian species (Traut & Marec 1996). In this
case, standard expectations of dosage compensation apply and the current observations of
Z:A < 1 in males remain theoretically inconsistent.
However, karyotypes of basal pre-ditrysian and ditrysian species do not obviously
indicate a lost autosome, a fact consistent with an alternative scenario that posits the W
originating as a supernumerary B chromosome that acquired a sex-determining locus and
meiotic pseudo-bivalence with the Z (Lukhtanov 2000). In this latter scenario, origins of
the Z chromosome are quite ancient, likely involve the transition to female-heterogamety,
and existing theory offers few expectations about how sex chromosomes and dosage effects
might evolve.
It should not be overlooked that available data indicate substantial variation among
lepidopteran lineages in patterns of sex chromosome dosage compensation. Considered in
light of the most recent lepidopteran phylogenies (Regier et al. 2013; Kawahara &
Breinholt 2014), the patterns appear to reflect a couple of evolutionary transitions in
dosage compensation (Table 5). Two possible histories could explain these patterns. Either
a similar dosage compensating mechanism evolved independently in butterflies and the

ancestor of bombycoids, or an "incomplete" dosage compensating mechanism present in
butterflies was refined to "completeness" in bombycoids while also lost in pyralids.
Regardless of how dosage compensation has evolved among moths and butterflies,
there are some intriguing similarities between the patterns emerging among Lepidoptera
and the emerging understanding of sex chromosome dosage compensation – or the lack
thereof – in mammals. Although still controversial, there is a growing consensus the
mammalian X evolved without dosage compensation and, in light of inferred ancestral
expression levels on the X, the mammalian X:A ratio is ~0.5 in both sexes (Julien et al.
2012; F Lin et al. 2012; Pessia et al. 2013). This view means that mammalian X-
chromosome inactivation (XCI) effectively balances X-linked expression between sexes by
halving the potential dose in females. A similar pattern was also recently reported in
twisted-wing insects (Strepsiptera) (Mahajan & Bachtrog 2015). This scenario appears
analogous to Lepidoptera, where Z-linked expression is similarly balanced between sexes
by an apparent global reduction in expression in the homogametic sex. There is clearly
need for a much further investigation into the functional mechanisms of lepidopteran
dosage compensation. At this point it seems possible that Lepidoptera, Strepsiptera, and
mammals have converged on similar mechanisms of mitigating sex chromosome dosage
differences between the sexes.
488

Chromosomal distribution of sex-biased genes
The general lack of sex-biased genes in purely somatic, non-reproductive tissue in
Heliconius is unusual. There is good precedent from many other animals, particularly fruit
flies and mice, that such tissues should contain at least dozens if not hundreds of genes
with sex-biased expression (Yang et al. 2006; Catalán et al. 2012; Meisel et al. 2012). Even
among other Lepidoptera such tissues yield numerous sex-biased genes (Walters &
Hardcastle 2011; Smith et al. 2014). Thus it is tempting to explain away this result through
a lack of statistical power or other methodological artifacts. Yet this seems unlikely
because abdomens yielded thousands of differentially expressed loci. Heads and abdomens
in this study were from the same individuals, with RNA extracted and analyzed
simultaneously in parallel. Furthermore, the mouth, leg, and antennae data were generated
independently of the head and abdomen samples but yielded a similar paucity of sex-
biased genes. So it may simply be that *Heliconius* butterflies have unusually monomorphic
patterns of gene expression between sexes for tissues unrelated to reproduction. Certainly
phenotypic sexual dimorphism is quite limited in *Heliconius* compared to most moths and
many butterflies (personal observations, JRW & CJD). Resolving this issue will have to wait
for further similar genome-wide expression experiments to be conducted.
In contrast to non-reproductive tissues, the abdomens yielded a very large number
of genes with significantly dimorphic expression. The distribution of such sex-biased genes
is clearly different between the Z and autosomes, with male-biased genes more abundant

and female-biased genes less abundant on the Z relative to autosomes, which have
approximately equal amounts of male- and female-biased genes (Figure 4). A positive
association between sex-linkage and expression biased towards the homogametic sex has
been widely observed in several species of both male and female heterogametic taxa,
including Lepidoptera (Reinke et al. 2000; Khil et al. 2004; Kaiser & Ellegren 2006;
Arunkumar et al. 2009; Walters & Hardcastle 2011; Meisel et al. 2012). This pattern is
generally predicted by models of sexually antagonistic evolution, specifically when sexually
antagonistic mutations are fully or partially dominant (Rice 1984; Ellegren & Parsch 2007;
Connallon & Clark 2010). Thus our results further support for the theory that the
asymmetrical transmission of sex chromosomes favors sex-linked accumulation of
mutations benefitting the homogametic sex.
A deficit of genes biased towards the heterogametic sex is also predicted by the
same theory, but observations are much less consistent across taxa and seem to depend on
a range biological idiosyncrasies including meiotic sex chromosome inactivation, tissue
specificity of expression, and mechanisms of dosage compensation (Connallon & Clark
2010; Meisel et al. 2012; Parsch & Ellegren 2013). Nonetheless, there is at least one
promising coincidence of theory and data that has been recently noted: species with equal
rates of recombination between sexes tend to show an excess of sex-linked genes with
heterogametic bias (e.g. birds, mammals), while in *Drosophila*, where males do not
recombine, a paucity of male-biased genes on the X is often reported (Connallon & Clark
2010). Since lepidopteran females lack recombination, the paucity of female-biased genes
on the Z in *Heliconius* is consistent with this pattern. However, our data represent only an
initial and broadly scoped assessment of this phenomenon. Robust inferences about both
patterns and causes of the chromosomal distribution of sex-biased genes benefit
substantially from a much greater diversity of tissue and ontogenetic sampling than
presented here (Mank et al. 2010; Meisel et al. 2012; Perry et al. 2014).
535

Conclusion

Our results are consistent with an increasing body of evidence that many species of
moths and butterflies possess a sex chromosome dosage compensating mechanism that
operates by reducing Z chromosome expression in males. However, this mechanism
appears to be imperfect in *Heliconius*, resulting in a moderate Z chromosome dosage effect.
These results bisect the emerging view that female-heterogametic ZW taxa have
incomplete dosage compensation because they lack a chromosome-wide epigenetic
mechanism mediating sex chromosome dosage compensation (the *heterogametic
dichotomy hypothesis*). In the case of *Heliconius*, sex chromosome dosage effects persist
apparently despite such a mechanism. This result adds additional complexity to patterns of
dosage compensation observed in Lepidoptera to date and highlights the need to increase

both the breadth of taxa sampled along with the consistency of analysis. Additionally,
patterns of sex-biased expression in our data highlight the impact of sexually antagonistic
selection to shape the evolution of the genome.
550

Acknowledgements

551 **Acknowledgements**
552 This research was supported in part by a NSF post-doctoral fellowship to JRW (DBI-
553 0905698). RNA sequencing was funded by the "Capacity and Capability Challenge
554 Programme" from The Genome Analysis Centre, Norwich, UK. Luiqi (Aloy) Gu provided
555 valuable comments on the manuscript.
556
557

Appendix: Analysis of Z-linked expression quartiles between male and female

Harrison et al. (2011) analyzed patterns of sex-chromosome dosage compensation in the Indian meal moth, *Plodia interpunctella.* One aspect of their analysis contrasted male and female expression of Z-linked loci split by quartiles of expression magnitude. This analysis is intended to illustrate that dosage effects increase with gene expression level. Importantly, Harrison et al. defined quartiles of expression based solely on male expression; female expression was ignored when binning loci into subsets to assess dosage effects. Here we demonstrate that, in the context of RNA-seq data, their approach potentially leads to an artifactual bias that would result in expression appearing greater in males than females for highly expressed loci, as reported by Harrison et al. (2011).

We judge it unlikely that this bias can completely explain the patterns of reported for *P. interpunctella.* Therefore we have not re-analyzed the data of Harrison et al. and we do not question their qualitative conclusions regarding the relationship between relative dosage effects and the magnitude of gene expression. Nonetheless, for the sake of future analyses of sex chromosome dosage compensation, we feel it worthwhile to describe the flaw in their analysis and offer a correction. We suggest that quartile binning based on the *maximum of male or female expression at each locus* provides a simple adjustment to the analysis that removes bias imposed by binning on data from only one sex.

This analysis tests whether dosage effects (i.e. greater Z-linked expression in males) increases with magnitude of gene expression. It is based on the idea that, for any given locus, the maximum level of gene expression is limited by the rate of translation because the translational machinery becomes saturated. In the absence of sex chromosome dosage compensation, highly expressed genes at or near the point of maximum translation will exhibit a strong dosage effect; the single female gene copy cannot reach the transcription rate of two copies in male. In contrast, genes expressed at low to intermediate levels will exhibit little or no dosage effect because sufficient transcripts can be generated from a single female copy to balance transcript levels in males. The resulting pattern is a discrepancy between male and female Z-linked expression that increases with the magnitude of gene expression.

In principle, it should be possible capture this effect by comparing the distributions of male and female expression of Z-linked genes that are expressed at relatively high or low absolute expression levels. Following the approach of Harrison et al., binning loci by quartiles of expression should reveal increasingly greater relative expression in males with each additional quartile if translational saturation effects are occurring in the absence of dosage compensation.

However, *it is in the grouping of loci into bins of relative expression levels where a bias may arise.* Binning based on expression in only one sex can cause expression in that sex to appear relatively reduced among weakly expressed genes, but relatively high for strongly

40 expressed genes, even if the distribution of gene expression is identical for both sexes.
41 Thus grouping loci by expression quartiles using values in only one sex can cause biases in
42 which sex appears to have relatively greater expression in each quartile. This phenomenon
43 is best demonstrated with a simple simulation in which the average expression of males
44 and females is identical.

45   Described briefly, the simulation begins by generating "true" expression values for
46 10,000 loci as random draws from a log-normal distribution. Variation in expression levels
47 for male and female "observations" are created by independently adding values drawn
48 from a normal distribution with a mean of 0 to each "true" expression value. Additionally,
49 1000 loci are further modified to be differentially expressed between males and females,
50 500 in each direction. For the sake of brevity and clarity, we refer the reader seeking
51 further details to the attached R script encoding the simulation. The "MA-plot" of simulated
52 gene expression data (Figure A1) should suffice to convince most readers that the
53 simulated data reasonably represents gene expression data.

54   Using these simulated values, we can split the dataset into quartiles of expression
55 and compare the distributions between sexes (Figure A2). Intuitively, since all differences
56 between male and female expression are random and unbiased, we would expect no
57 systematic differences between sexes at any expression level. However, this outcome is
58 only observed if quartiles are based *on the maximum of male or female expression* for each
59 locus. Ranking using only one sex clearly results in a strongly biased outcome that depends
60 on which sex is used for ranking (Figure A2).

61   We emphasize that this simulation has been created primarily to illustrate our point
62 about biases that potentially arise from quartile analyses based on ranking on one sex
63 alone. We do not claim that the simulation is highly realistic and we have not extensively
64 explored the effect of changing simulation parameters. Our narrow aim is to communicate
65 an intuitive and straightforward demonstration of why we have deviated from the
66 precedent set by Harrison et al. (2012) for this type of analysis.

70   Figure A1. MA-plot of simulated male and female expression data.

Figure A2. Quartile comparisons of simulated gene expression distributions of simulated
male and female expression levels. Comparison of male (blue) and female (red) expression
levels for loci, divided into quartiles based on male expression (left panel), female
expression (center panel), and the maximum male or female expression (right panel).
Boxes indicate the interquartile range (IQR) around the median (black bar) and whiskers
extend to 1.5 times the IQR. P-values of a Mann-Whitney U test of inequality between
males and females are printed for each quartile along the x-axis, with "0" indicating P <
0.0001.

References

Ahola V et al. 2014. The Glanville fritillary genome retains an ancientkaryotype and reveals selective chromosomalfusions in Lepidoptera. Nature Communications. 5:1–9. doi: 10.1038/ncomms5737.

Allen SL, Bonduriansky R, Chenoweth SF. 2013. The Genomic Distribution of Sex-Biased Genes in Drosophila serrata: X Chromosome Demasculinization, Feminization, and Hyperexpression in Both Sexes. Genome Biology and Evolution. 5:1986–1994. doi: 10.1093/gbe/evt145.

Arunkumar KP, Mita K, Nagaraju J. 2009. The silkworm Z chromosome is enriched in testis-specific genes. Genetics. 182:493–501. doi: 10.1534/genetics.108.099994.

Bachtrog D. 2006. A dynamic view of sex chromosome evolution. Curr Opin Genet Dev. 16:578–585. doi: 10.1016/j.gde.2006.10.007.

Bachtrog D. 2013. Y-chromosome evolution:emerging insights into processesof Y-chromosome degeneration. Nat Rev Genet. 14:113–124. doi: 10.1038/nrg3366.

Bachtrog D et al. 2011. Are all sex chromosomes created equal? Trends Genet. 27:350–357. doi: 10.1016/j.tig.2011.05.005.

Bates D, Machler M, Bolker B, Walker S. 2014. Fitting Linear Mixed-Effects Models using lme4. arXive. http://arxiv.org/abs/1406.5823

Birchler JA, Bhadra U, Bhadra MP, Auger DL. 2001. Dosage-dependent gene regulation in multicellular eukaryotes: implications for dosage compensation, aneuploid syndromes, and quantitative traits. Dev Biol. 234:275–288. doi: 10.1006/dbio.2001.0262.

Briscoe AD et al. 2013. Female behaviour drives expression and evolution of gustatory receptors in butterflies. PLoS Genet. 9:e1003620. doi: 10.1371/journal.pgen.1003620.

Bullard JH, Purdom E, Hansen KD, Dudoit S. 2010. Evaluation of statistical methods for normalization and differential expression in mRNA-Seq experiments. BMC Bioinformatics. 11:94. doi: 10.1186/1471-2105-11-94.

Catalán A, Hutter S, Parsch J. 2012. Population and sex differences in Drosophila melanogaster brain gene expression. BMC Genomics. 13:654. doi: 10.1186/1471-2164-13-654.

Charlesworth B. 1978. Model for evolution of Y chromosomes and dosage compensation. P Natl Acad Sci Usa. 75:5618–5622.

Charlesworth B. 1996. The evolution of chromosomal sex determination and dosage
compensation. Curr Biol. 6:149–162.

Charlesworth B, Charlesworth D. 2000. The degeneration of Y chromosomes. Philos Trans
R Soc Lond, B, Biol Sci. 355:1563–1572. doi: 10.1098/rstb.2000.0717.

Connallon T, Clark AG. 2010. Sex linkage, sex-specific selection, and the role of
recombination in the evolution of sexually dimorphic gene expression. Evolution. 64:3417–
3442. doi: 10.1111/j.1558-5646.2010.01136.x.

Deng X et al. 2011. Evidence for compensatory upregulation of expressed X-linked genes in
mammals, Caenorhabditis elegans and Drosophila melanogaster. Nat Genet. 43:1179–1185.
doi: 10.1038/ng.948.

Disteche CM. 2012. Dosage Compensation of the Sex Chromosomes. Annu Rev Genet.
46:537–560. doi: 10.1146/annurev-genet-110711-155454.

Dunlap-Pianka H, Boggs CL, Gilbert LE. 1977. Ovarian Dynamics in Heliconiine Butterflies:
Programmed Senescence versus Eternal Youth. Science. 197:487–490. doi:
10.1126/science.197.4302.487.

Ellegren H et al. 2007. Faced with inequality: chicken do not have a general dosage
compensation of sex-linked genes. BMC Biol. 5:40. doi: 10.1186/1741-7007-5-40.

Ellegren H, Parsch J. 2007. The evolution of sex-biased genes and sex-biased gene
expression. Nat Rev Genet. 8:689–698. doi: 10.1038/nrg2167.

Ferrari F, Alekseyenko AA, Park PJ, Kuroda MI. 2014. Transcriptional control of a whole
chromosome: emerging models for dosage compensation. Nature Publishing Group.
21:118–125. doi: 10.1038/nsmb.2763.

Gupta V et al. 2006. Global analysis of X-chromosome dosage compensation. J Biol. 5:3. doi:
10.1186/jbiol30.

Hamada FN. 2005. Global regulation of X chromosomal genes by the MSL complex in
Drosophila melanogaster. Genes & Development. 19:2289–2294. doi:
10.1101/gad.1343705.

Hardcastle TJ. 2014. Generalised empirical Bayesian methods for discovery of differential
data in high-throughput biology. bioRxiv. doi: 10.1101/011890.

Hardcastle TJ, Kelly KA. 2010. baySeq: empirical Bayesian methods for identifying
differential expression in sequence count data. BMC Bioinformatics. 11:422. doi:
10.1186/1471-2105-11-422.

Hardcastle TJ, Kelly KA, Baulcombe DC. 2012. Identifying small interfering RNA loci from
high-throughput sequencing data. Bioinformatics. 28:457–463. doi:

10.1093/bioinformatics/btr687.

Harrison PW, Mank JE, Wedell N. 2012. Incomplete Sex Chromosome Dosage Compensation
in the Indian Meal Moth, Plodia interpunctella, Based on De Novo Transcriptome Assembly.
Genome Biology and Evolution. 4:1118–1126. doi: 10.1093/gbe/evs086.

Heliconius Genome Consortium. 2012. Butterfly genome reveals promiscuous exchange of
mimicry adaptations among species. Nature. 487:94–98. doi: 10.1038/nature11041.

Itoh Y et al. 2007. Dosage compensation is less effective in birds than in mammals. J Biol.
6:2. doi: 10.1186/jbiol53.

Itoh Y et al. 2010. Sex bias and dosage compensation in the zebra finch versus chicken
genomes: general and specialized patterns among birds. Genome Res. 20:512–518. doi:
10.1101/gr.102343.109.

Jue NK et al. 2013. Determination of dosage compensation of themammalian X
chromosome by RNA-seq isdependent on analytical approach. BMC Genomics. 14:1–1. doi:
10.1186/1471-2164-14-150.

Julien P et al. 2012. Mechanisms and Evolutionary Patterns of Mammalian and Avian
Dosage Compensation Barton, NH, editor. PLoS Biol. 10:e1001328. doi:
10.1371/journal.pbio.1001328.t002.

Kaiser VB, Ellegren H. 2006. Nonrandom distribution of genes with sex-biased expression
in the chicken genome. Evolution. 60:1945–1951.

Kawahara AY, Breinholt JW. 2014. Phylogenomics provides strong evidence for
relationships of butterflies and moths. Proceedings of the Royal Society B: Biological
Sciences. 281:20140970–20140970. doi: 10.1126/science.1174096.

Kharchenko PV, Xi R, Park PJ. 2011. correspondence. Nat Genet. 43:1167–1169. doi:
10.1038/ng.991.

Khil PP, Smirnova NA, Romanienko PJ, Camerini-Otero RD. 2004. The mouse X chromosome
is enriched for sex-biased genes not subject to selection by meiotic sex chromosome
inactivation. Nat Genet. 36:642–646. doi: 10.1038/ng1368.

Kiuchi T et al. 2014. A single female-specific piRNA is the primary determiner of sex in the
silkworm. Nature. 509:633–636. doi: 10.1038/nature13315.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods.
9:357–359. doi: 10.1038/nmeth.1923.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC Bioinformatics. 12:323. doi: 10.1186/1471-2105-12-
323.

Lin F, Xing K, Zhang J, He X. 2012. Expression reduction in mammalian X chromosome
evolution refutes Ohno's hypothesis of dosage compensation. doi:
10.1073/pnas.1201816109/-/DCSupplemental/pnas.201201816SI.pdf.

Lin H et al. 2011. ng.992. Nat Genet. 43:1169–1170. doi: 10.1038/ng.992.

Lukhtanov VA. 2000. Sex chromatin and sex chromosome systems in nonditrysian
Lepidoptera (Insecta). Journal of Zoological Systematics and Evolutionary Research. 38:73–
79.

Mahajan S, Bachtrog D. 2015. Partial Dosage Compensation in Strepsiptera, a Sister Group
of Beetles. Genome Biology and Evolution. 7:591–600. doi: 10.1093/gbe/evv008.

Mank JE. 2013. Sex chromosome dosage compensation: definitely not for everyone. Trends
Genet. 1–7. doi: 10.1016/j.tig.2013.07.005.

Mank JE. 2009. The W, X, Y and Z of sex-chromosome dosage compensation. Trends Genet.
25:226–233. doi: 10.1016/j.tig.2009.03.005.

Mank JE, Ellegren H. 2009. All dosage compensation is local: gene-by-gene regulation of
sex-biased expression on the chicken Z chromosome. Heredity. 102:312–320. doi:
10.1038/hdy.2008.116.

Mank JE, Hosken DJ, Wedell N. 2011. Some inconvenient truths about sex chromosome
dosage compensation and the potential role of sexual conflict. Evolution. 65:2133–2144.
doi: 10.1111/j.1558-5646.2011.01316.x.

Mank JE, Nam K, Brunstrom B, Ellegren H. 2010. Ontogenetic Complexity of Sexual
Dimorphism and Sex-Specific Selection. Molecular Biology and Evolution. 27:1570–1578.
doi: 10.1093/molbev/msq042.

Martin SH et al. 2013. Genome-wide evidence for speciation with gene flow in Heliconius
butterflies. Genome Res. 23:1817–1828. doi: 10.1101/gr.159426.113.

Meisel RP, Malone JH, Clark AG. 2012. Disentangling the relationship between sex-biased
gene expression and X-linkage. Genome Res. doi: 10.1101/gr.132100.111.

Naurin S, Hansson B, Hasselquist D, Kim Y-H, Bensch S. 2011. The sex-biased brain: sexual
dimorphism in gene expression in two species of songbirds. BMC Genomics. 12:37. doi:
10.1186/1471-2164-12-37.

Naurin S, Hasselquist D, Bensch S, Hansson B. 2012. Sex-Biased Gene Expression on the
Avian Z Chromosome: Highly Expressed Genes Show Higher Male-Biased Expression
Nurminsky, DI, editor. PLoS ONE. 7:e46854. doi: 10.1371/journal.pone.0046854.t004.

Nguyen DK, Disteche CM. 2006. Dosage compensation of the active X chromosome in
mammals. Nat Genet. 38:47–53. doi: 10.1038/ng1705.

Nozawa M, Fukuda N, Ikeo K, Gojobori T. 2014. Tissue- and Stage-Dependent Dosage
Compensation on the Neo-X Chromosome in Drosophila pseudoobscura. Molecular Biology
and Evolution. 31:614–624. doi: 10.1093/molbev/mst239.

Ohno S. 1967. *Sex chromosomes and sex-linked genes*. Springer.

Parsch J, Ellegren H. 2013. The evolutionary causes and consequences of sex-biased gene
expression. Nat Rev Genet. 14:83–87. doi: 10.1038/nrg3376.

Perry JC, Harrison PW, Mank JE. 2014. The Ontogeny and Evolution of Sex-Biased Gene
Expression in Drosophila melanogaster. Molecular Biology and Evolution. 31:1206–1219.
doi: 10.1093/molbev/msu072.

Pessia E, Engelstädter J, Marais GAB. 2013. The evolution of X chromosome inactivation in
mammals: the demise of Ohno's hypothesis? Cell. Mol. Life Sci. 71:1383–1394. doi:
10.1007/s00018-013-1499-6.

Prince EG, Kirkland D, Demuth JP. 2010. Hyperexpression of the X chromosome in both
sexes results in extensive female bias of X-linked genes in the flour beetle. Genome Biology
and Evolution. 2:336–346. doi: 10.1093/gbe/evq024.

Pringle EG et al. 2007. Synteny and chromosome evolution in the lepidoptera: evidence
from mapping in Heliconius melpomene. Genetics. 177:417–426. doi:
10.1534/genetics.107.073122.

Quek SP et al. 2010. Dissecting comimetic radiations in Heliconius reveals divergent
histories of convergent butterflies. P Natl Acad Sci Usa. 107:7365–7370. doi:
10.1073/pnas.0911572107.

R Development Core Team. 2014. R: A Language and Environment for Statistical
Computing. http://www.R-project.org/.

Regier JC et al. 2013. A Large-Scale, Higher-Level, Molecular Phylogenetic Study of the
Insect Order Lepidoptera (Moths and Butterflies) Moreau, CS, editor. PLoS ONE. 8:e58568.
doi: 10.1371/journal.pone.0058568.s012.

Reinke V et al. 2000. A global profile of germline gene expression in C. elegans. Mol Cell.
6:605–616.

Rice W. 1984. Sex chromosomes and the evolution of sexual dimorphism. Evolution.
38:735–742.

Sahara K, Yoshido A, Traut W. 2011. Sex chromosome evolution in moths and butterflies.
Chromosome Res. 20:83–94. doi: 10.1007/s10577-011-9262-z.

Smith G, Chen YR, Blissard GW, Briscoe AD. 2014. Complete Dosage Compensation and Sex-
Biased Gene Expression in the Moth Manduca sexta. Genome Biology and Evolution. 6:526–

537. doi: 10.1093/gbe/evu035.

Straub T, Becker PB. 2011. Transcription modulation chromosome-wide: universal features
and principles of dosage compensation in worms and flies. Curr Opin Genet Dev. 21:147–
153. doi: 10.1016/j.gde.2011.01.012.

Suetsugu Y et al. 2013. Large scale full-length cDNA sequencing reveals a unique genomic
landscape in a lepidopteran model insect, Bombyx mori. G3 (Bethesda). 3:1481–1492. doi:
10.1534/g3.113.006239/-/DC1.

Traut W, Marec F. 1996. Sex chromatin in lepidoptera. Q Rev Biol. 71:239–256.

Veitia RA, Veyrunes F, Bottani S, Birchler JA. 2015. X chromosome inactivation and active X
upregulation in therian mammals: facts, questions, and hypotheses. J Mol Cell Biol. 7:2–11.
doi: 10.1093/jmcb/mjv001.

Vicoso B, Bachtrog D. 2011a. Lack of Global Dosage Compensation in Schistosoma mansoni,
a Female-Heterogametic Parasite. Genome Biology and Evolution. 3:230–235. doi:
10.1093/gbe/evr010.

Vicoso B, Bachtrog D. 2011b. Lack of global dosage compensation in Schistosoma mansoni,
a female-heterogametic parasite. Genome Biology and Evolution. doi: 10.1093/gbe/evr010.

Vicoso B, Bachtrog D. 2009. Progress and prospects toward our understanding of the
evolution of dosage compensation. Chromosome Res. 17:585–602. doi: 10.1007/s10577-
009-9053-y.

Vicoso B, Emerson JJ, Zektser Y, Mahajan S, Bachtrog D. 2013a. Comparative Sex
Chromosome Genomics in Snakes: Differentiation, Evolutionary Strata, and Lack of Global
Dosage Compensation Barton, NH, editor. PLoS Biol. 11:e1001643. doi:
10.1371/journal.pbio.1001643.s024.

Vicoso B, Kaiser VB, Bachtrog D. 2013b. Sex-biased gene expression at homomorphic sex
chromosomes in emus and its implication for sex chromosome evolution. Proceedings of
the National Academy of Sciences. 110:6453–6458. doi: 10.1073/pnas.1217027110.

Walters JR, Hardcastle TJ. 2011. Getting a full dose? Reconsidering sex chromosome dosage
compensation in the silkworm, Bombyx mori. Genome Biology and Evolution. 3:491–504.
doi: 10.1093/gbe/evr036.

Xia Q et al. 2007. Microarray-based gene expression profiles in multiple tissues of the
domesticated silkworm, Bombyx mori. Genome Biol. 8:R162. doi: 10.1186/gb-2007-8-8-
r162.

Xiong Y et al. 2010. RNA sequencing shows no dosage compensation of the active X-
chromosome. Nat Genet. 42:1043–1047. doi: 10.1038/ng.711.

Yang X et al. 2006. Tissue-specific expression and regulation of sexually dimorphic genes in
mice. Genome Res. 16:995–1004. doi: 10.1101/gr.5217506.

Yue Z et al. 2013. A heterozygous moth genome provides insights into herbivory and
detoxification. Nat Genet. 1–8. doi: 10.1038/ng.2524.

Zha X et al. 2009. Dosage analysis of Z chromosome genes using microarray in silkworm,
Bombyx mori. Insect Biochem Mol Biol. 39:315–321. doi: 10.1016/j.ibmb.2008.12.003.

Table 1. Summary of average expression of Z-linked and autosomal loci across several tissues in *Heliconius* butterflies. Results reflect data filtering with a null expression likelihood of 0.5 and are representative of results using a range of likelihood thresholds (supplementary Table S2)

Statistic	Abdomen (H. cydno)		Head (H. cydno)		Abdomen (H. melpomene)		Head (H. melpomene)		Antenna		Leg		Mouth	
	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female
Mean Z-linked FPKM	40.66	26.15	29.27	23.78	36.63	29.6	28.14	29.18	30.32	27.3	47.99	41.65	40.74	35.07
Mean autosomal FPKM	85.45	54.81	70.64	53.6	63.42	53.33	54.2	68.43	75.66	67.84	63.3	59.02	71.77	68.06
Z:A ratio of mean expression	0.4758	0.4771	0.4144	0.4437	0.5776	0.555	0.5192	0.4264	0.4007	0.4024	0.7581	0.7057	0.5676	0.5153
Median Z-linked FPKM	7.15	6.15	5.48	5.23	8.53	7.38	6.8	6.59	10.38	10.26	9.12	7.59	9.03	7.81
Median autosomal FPKM	11.67	12.12	7.78	7.82	11.6	12.71	8.93	9.03	14.65	14.93	12.43	11.69	12.13	11.96
Z:A ratio of median expression	0.6127	0.5074	0.7044	0.6688	0.7353	0.5806	0.7615	0.7298	0.7085	0.6872	0.7337	0.6493	0.7444	0.653
MWU p-value: Autosomal ≠ Z-linked	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	.0007	.0001	<.0001	<.0001	.0003	<.0001	.0002	<.0001
No. expressed Z-linked loci	481	403	443	429	506	417	449	447	392	394	357	385	361	390
No. expressed autosomal loci	10638	9758	10272	10050	11020	10014	10004	10383	9296	9318	8548	9239	8792	9243

Table 2. Linear modeling analysis of Z-chromosome and dosage effects on gene expression levels across several tissues in *Heliconius* butterflies. Results reflect data filtering with a null expression likelihood of 0.5 and are representative of results using a range of likelihood thresholds (supplementary Table S3)

Sample	Z-linkage only			Z-linkage x sex interaction (Dosage effect)		
	Z-linkage Effect Size	ANOVA p-value		Z-linkage Effect Size	Interaction (Dosage) Effect Size	ANOVA p-value
Abdomen (H. cydno)	-0.56	2.16E-14		-0.65878	0.212	2.57E-05
Head (H. cydno)	-0.431	3.02E-08		-0.40678	0.0642	7.66E-08
Abdomen (H. melpomene)	-0.385	1.53E-07		-0.47591	0.189	0.0001
Head (H. melpomene)	-0.263	0.00030		-0.4581	0.289	6.90E-27
Antenna	-0.371	2.35E-07		-0.37173	0.0307	0.224
Leg	-0.275	0.00047		-0.36309	0.169	1.47E-14
Mouth	-0.340	6.72E-06		-0.36206	0.0615	0.00227

Table 3. Summary of average Male:Female gene expression ratios for Z-linked and autosomal loci in *Heliconius* butterflies for several tissues. Results reflect data filtering with a null expression likelihood of 0.5 and are representative of results using a range of likelihood thresholds (supplementary Table S4).

Tissue	No. Z-linked loci	No. Autosomal loci	Mean Z-linked log2(M:F)	Mean autosomal log2(M:F)	Median Z – linked log2(M:F)	Mean autosomal log2(M:F)	Z:A ratio of means (Not Log2)	Z:A ratio of medians (Not Log2)	MWU p-value: Autosomal ≠ Z-linked
Abdomen (H. cydno)	387	9539	0.3553	0.1101	0.1683	-0.0009	1.1853	1.1244	<.0001
Head (H. cydno)	425	9972	0.1173	0.0584	0.116	0.0543	1.0417	1.0437	0.0034
Abdomen (H. melpomene)	408	9859	0.2418	0.0106	0.1904	-0.0788	1.1738	1.2051	<.0001
Head (H. melpomene)	433	9909	0.0847	-0.1372	0.1002	-0.0904	1.1662	1.1412	<.0001
Antenna	378	8957	0.0115	-0.039	0.0731	-0.009	1.0356	1.0586	0.021
Leg	354	8457	0.1272	-0.0912	0.1731	-0.097	1.1634	1.2059	<.0001
Mouth	356	8636	0.003	-0.0741	0.0811	-0.0561	1.0549	1.0997	<.0001

Table 4. Counts of sex-biased and unbiased genes (FDR < 0.05 and a minimum fold-change of 1.5) on the Z chromosome and autosomes (A) in several tissues of *Heliconius* butterflies.

Expression Category	Abdomen (H. cydno)		Head (H. cydno)		Abdomen (H. melpomene)		Head (H. melpomene)		Antenna		Leg		Mouth	
	Z	A	Z	A	Z	A	Z	A	Z	A	Z	A	Z	A
Male bias	114	1470	3	7	130	1590	0	0	0	0	0	0	0	0
Unbiased	275	7008	385	9389	290	6888	394	9354	319	8167	301	7717	295	7715
Female bias	48	1540	0	8	58	2025	0	3	0	3	0	0	0	0

Table 5. Summary of observed patterns of sex chromosome dosage compensation in the Lepidoptera.

Phylogeny	Taxon	Male Z:A ratio	Female Z:A ratio	Z Dosage effect (average M:F expression on Z)
	Bombyx [1]	< 1	< 1	None (M = F)
	Manduca [2]	< 1	< 1	None (M = F)
	Plodia [3]	= 1	<< 1	Strong male bias (M >> F)*
	Heliconius	< 1	< 1	Weak male bias (M $\geq$ F)
	Basal ZO Lepidoptera	?	?	?

1) Walters and Hardcastle (2011)
2) Smith et al. (2014)
3) Harrison et al. (2012)
*not directly reported, but assumed based on sex-specific Z:A ratios

Fig. 1. Distributions of log2-transformed gene expression levels (FPKM) for the Z-chromosome (Z; red) and autosomes (A; grey) in male and female *Heliconius* butterflies in several tissues. Boxes indicate the interquartile range (IQR) around the median (black bar) and whiskers extend to 1.5 times the IQR; outlier points beyond 1.5 IQR are not shown. Non-parametric tests as well as linear modeling indicate a significant reduction in Z-chromosome expression relative to autosomes in both sexes for all tissues (see Tables 1 & 2). Data were filtered with a null expression likelihood of 0.5 and are representative of results using a range of likelihood thresholds (supplementary Table 2).

Fig. 2. Distribution density plots of log2(Male:Female) expression ratios for the Z-chromosome (Z; red) and autosomes (A; black) in male and female *Heliconius* butterflies in several tissues. Dashed lines indicate median values. Non-parametric tests and linear modeling indicate all tissues but antenna show a modest but significant sex-chromosome dosage effect, visualized here as a shift towards male-biased expression among Z-linked loci (see Tables 2 & 3). Antenna also shows this pattern but the effect is not statistically significant.

Fig. 3. Comparison of male (blue) and female (red) expression levels for Z-linked loci, divided into quartiles based on the maximum of male or female expression for *H. melpomene* head and abdomen. Boxes indicate the interquartile range (IQR) around the median (black bar) and whiskers extend to 1.5 times the IQR. An asterisk (*) indicates significant difference in average expression (bonferroni corrected MWU p-value < 0.05).   Plots for all seven tissue samples are given in supplemental Figure S2.

Fig. 4. Proportions of sex-biased genes across the chromosomes in abdomens for *H. cydno* and *H. melpomene*. A gene was considered sex-biased if significantly differentially expressed between males and females with FDR < 0.05 and a minimum fold-change of 1.5. The two larger bars at the bottom represent the Z chromosome individually and the combined results across autosomes (A). Smaller bars (1-20) represent data for individual chromosomes and "UM" indicates genes not mapped to chromosome.

Figure S1. Distributions of log2-transformed gene expression levels (FPKM) for each chromosome in male and female *Heliconius* butterflies. The right-hand most box reflects genes not mapped to chromosome. The horizontal redline indicates the median log2(FPKM) of the Z chromosome. Boxes indicate the interquartile range (IQR) around the median (black bar) and whiskers extend to 1.5 times the IQR; outlier points beyond 1.5 IQR are not plotted. Data were filtered with a null expression likelihood of 0.5

Figure S2. Comparison of male (blue) and female (red) expression levels for Z-linked loci, divided into quartiles based on the maximum of male or female expression in various tissues from *Heliconius* butterflies. Boxes indicate the interquartile range (IQR) around the median (black bar) and whiskers extend to 1.5 times the IQR. P-values of a Mann-Whitney U test of inequality between males and females are printed for each quartile along the x-axis

Figure S3. Distributions of log2-transformed gene expression levels (FPKM) for expressed autosomal loci, partitioned by category of sex-biased expression, in abdomens from two species of *Heliconius* butterflies. In both species males, but not females, showed significantly different levels of expression across categories (Kruskal-Wallis test, p << 0.001). A gene was considered sex-biased if significantly differentially expressed between males and females with FDR < 0.05 and a minimum fold-change of 1.5. Boxes indicate the interquartile range (IQR) around the median (black bar) and whiskers extend to 1.5 times the IQR. Data were filtered with a null expression likelihood of 0.5.