Non-dichotomous inference using bootstrapped evidence

D. Samuel Schwarzkopf

Experimental Psychology, University College London, 26 Bedford Way, London WC1H 0AP, United Kingdom &

Institute of Cognitive Neuroscience, University College London, 17 Queen Square, London, WC1N 3AR, United Kingdom

Email: mailto:s.schwarzkopf@ucl.ac.uk

Abstract

The problems with classical frequentist statistics are well established, yet the enthusiasm of researchers to adopt alternatives like Bayesian inference remains modest. Here I present the *bootstrapped evidence test*, an objective resampling procedure that takes the precision with which both the experimental and null hypothesis can be estimated into account. Simulations and reanalysis of actual experimental data demonstrate that this test minimizes false positives while maintaining sensitivity. It is equally applicable to a wide range of situations and thus minimizes problems arising from analytical flexibility. Critically, it does not dichotomize the results based on an arbitrary significance level but instead quantifies how well the data support either the alternative or the null hypothesis. It is thus particularly useful in situations with considerable uncertainty about the expected effect size. Because it is non-parametric, it is also robust to severe violations of assumptions made by classical statistics.

Introduction

To this day classical null hypothesis significance testing (NHST) remains the dominant approach for inferring the validity of an observed result in the life sciences. It rests on the probability ('p-value') that the observed effect, or a more extreme one, could have occurred under the assumption that there is no population effect. If the p-value is sufficiently low, this null hypothesis is rejected. However, p-values are frequently misinterpreted by researchers, uninformative about the evidence *for* an experimental hypothesis, highly susceptible to biased data sampling strategies, and generally prone to false positives (Cohen, 1994; Wagenmakers, 2007; Wagenmakers et al., 2011b; Masicampo and Lalande, 2012; Cumming, 2014; Colquhoun, 2014; Halsey et al., 2015; Trafimow and Marks, 2015; Gigerenzer and Marewski, 2015; Gigerenzer, 2004; Nuzzo, 2014). The most devastating effect of NHST may be that it encourages an artificial dichotomy between significant and non-significant results (Cumming, 2014).

The scatter plots in Figure 1 illustrate the problems with p-values. Figure 1A shows an almost perfect correlation between two measures (r=0.995, p<0.0004). However, there are only five observations. In contrast, the data in Figure 1B are clearly correlated even though the correlation is weaker. Yet the p-value is similar (r=0.535, p<0.0004) because the sample size is much larger. Surely the evidence for a correlation in the second example is more compelling and more likely to replicate?

Figure 1. Scatter plots showing examples of correlation analysis. A-B. Correlated Gaussian data with n=5 (A) and n=40 (B). C-D. Uncorrelated Gaussian data with n=5 (C) and n=40 (D). E-F. Severely heteroscedastic data with n=40 (E) and n=200 (F). Each black dot is one observation. The grey shading denotes the Mahalanobis distance from the bivariate mean. Above each panel the Pearson's correlation coefficient, the default Bayes factor (Wetzels and Wagenmakers, 2012) BF_{10} comparing the alternative and the null hypothesis, and the bootstrapped evidence ε are given.

One journal went so far as to ban the use of classical inference completely while proposing no viable alternative (Trafimow and Marks, 2015). Others proposed guidelines to focus on effect size estimation and confidence intervals instead (Psychological Science, 2014; Cumming, 2014). However, the use of confidence intervals is also fraught with problems (Morey et al., 2014) and may simply become a new significance testing procedure in disguise (Cumming, 2014; Hoekstra et al., 2014). Moreover, like p-values, confidence intervals are frequently misinterpreted (Hoekstra et al., 2014) and may perform inadequately (Wilcox and Muska, 2001). Evidence for a hypothesis should *compare* an experimental (alternative) hypothesis to a baseline (null) hypothesis. Bayesian hypothesis tests can achieve that but are often difficult to apply and rely on the choice of a prior, which can result in considerable debate (see e.g. (Bem, 2011; Bem et al., 2011; Wagenmakers et al., 2011a, 2011b; Savalei and Dunn, 2015)).

Here I present the *bootstrapped evidence* (BSE) test. It makes minimal assumptions and is applicable to a wide range of situations. Crucially, it quantifies the evidence for either the alternative or the null hypothesis non-dichotomously. Yet unlike Bayesian methods it is based

67 only on the existing data without any question about prior distributions. Simulations demonstrate
68 the test's efficacy and robustness and compare it to classical frequentist and Bayesian inferential
69 methods. Further, I apply this test to several concrete examples to show its advantages in
70 practice. The MATLAB source code and example data are available:
71 http://dx.doi.org/10.6084/m9.figshare.1342798

72
73 **Results**

74
75 The principle underlying the BSE test is that under both the alternative (H_1) and the null
76 hypothesis (H_0) the results follow a probability distribution (Figure 2, left panels). In classical
77 statistics, the p-value reflects the distance of the observed effect, θ (e.g. a correlation coefficient),
78 from the center of the null distribution. The one-tailed p-value is the area under the blue curve
79 (null distribution) to the right of the red diamond, which denotes the observed effect.

80
While the null distribution depends on the sample variance, it nonetheless fails to take the
variability of the effect under H_1 into account. The BSE estimates how distinct these two
distributions are from one another by bootstrapping both H_0 and H_1 and quantifying Δ, the
distribution of differences (Figure 2, right panels), between them (see Materials and Methods). If
the distribution is narrow and shifted away from zero this constitutes evidence for H_1 (Figure 2A).
However, when the distribution is narrow but centered on zero this is instead evidence for H_0
(Figure 2B). A wide Δ distribution provides only inconclusive evidence (Figure 2C).

88
The BSE is expressed by ε, which is effectively a *signal-to-noise ratio on a logarithmic scale*. It is
the ratio of the observed effect, μ, and the uncertainty, σ, with which it can be estimated (Figure
2 and Materials and Methods). When μ is smaller than σ and thus the Δ distribution overlaps zero
(as quantified by ω), ε decreases as sample size, n, increases. Since ε is the logarithm of this ratio,
it is positive when the data support H_1 and negative when they favor H_0. If ε is near zero (-
$0.5 < \varepsilon < 0.5$) the evidence is inconclusive.

95
96 For the near perfect correlation with n=5 (Figure 1A) the evidence is only $\varepsilon=0.7$. In contrast, for
97 the modest correlation with n=40 the evidence $\varepsilon=1$ (Figure 1B). The data in Figure 1C,D are
98 uncorrelated and neither correlation would reach classical significance. However, for a small
99 sample size of n=5 (Figure 1C) the evidence is inconclusive ($\varepsilon=-0.4$) while for a large sample
100 (Figure 1D) the evidence compellingly favors the null hypothesis ($\varepsilon=-1.6$).

101
The assumption made by parametric tests of normally distributed errors is often violated as in
Figure 1E. Even though the classical p-value is highly significant (r=-0.45, p=0.0034), the evidence
for H_1 is fairly weak ($\varepsilon=0.6$). The reason is that the data are heteroscedastic and thus skew
classical Pearson's correlation: the residuals of a linear fit depend on x. While y is chosen from a
random normal distribution each point is also multiplied by the absolute magnitude of its paired
value in x (Wilcox and Muska, 2001). Such situations can readily occur in real experimental data:
for instance, the proliferation of cell growth or the mean firing rate of neurons may also be
accompanied by greater variability in those measures. This in turn could skew any correlations
between these measures and an independent variable.

Figure 2. Examples of the bootstrapped evidence procedure. *Left panels* show the effect size distribution under the null (blue) and alternative (red) hypothesis estimated by bootstrapping. The red diamond denotes the observed effect size. The red shaded region denotes though the 95% confidence interval of the estimated effect. *Right panels* show the distribution of differences, Δ, between the

alternative and null distributions (see Materials and Methods). The green diamond denotes the mean,
μ. The ratio of the red and green areas under the curve is ω and quantifies the overlap between the
null and alternative distributions. The standard deviation of Δ is denoted by σ. The bootstrapped
evidence, ε, incorporates these three parameters and the sample size n and is expressed on a
logarithmic scale (see equation). A. Strongly correlated data with n=30. Evidence compellingly
supports H_1. B. Uncorrelated data with n=700. Evidence compellingly supports H_0. C. Uncorrelated
data with n=7. Evidence is inconclusive.

Notably, even increasing the sample size does not alleviate this problem. The data in Figure 1F
were drawn from the same heteroscedastic population but the sample size is five times larger
(n=200) and the correlation is again highly significant (r=0.22, p=0.0014). Even robust significance
tests, including those specifically developed to control for heteroscedasticity, do not fare any
better (skipped correlation (Wilcox, 2005; Pernet et al., 2012; Rousselet and Pernet, 2012): t=4.03,
$t_{critical}$=2.35; permutation test: r=0.22, p=0.0013; Spearman's rho: ρ=0.18, p=0.0105; Kendall's tau
τ=0.14, p=0.0032; percentage bend correlation (Wilcox, 2005): r=0.2, p=0.0042; Shepherd's pi
(Schwarzkopf et al., 2012): π=-0.23, p=0.0034). In contrast, the BSE test suggests only inconclusive
evidence for H_1 (ε=0.4) because bootstrapped distributions for H_1 and H_0 overlap substantially. In
comparison, a homoscedastic data set with the same effect and sample size would produce more
convincing evidence for H_1 (ε=0.7).

Performance on simulated data

For a more objective evaluation of the method I ran a series of simulations. For several sample
sizes (n=15, 30, 60, 120, 240, and 480) I generated 5,000 data sets each drawn from two different
distributions in which H_0 is true: an uncorrelated bivariate Gaussian and the same heteroscedastic
distribution underlying Figures 1E,F. For each simulated data set I calculated the bootstrapped
evidence, ε, the classical parametric p-value and a default Bayes factor for H_1 over H_0 (BF_{10})
(Wetzels and Wagenmakers, 2012).

Figure 3A shows the distribution of these inferential statistics for uncorrelated Gaussian data with
the various sample sizes denoted by different colors. As sample size increases the distributions for
the default Bayes factor and bootstrapped evidence become increasingly shifted towards negative
numbers, indicating increasing support for H_0. In contrast, the distributions for classical p-values
remain the same irrespective of sample size because under H_0 the distribution of p-values is
uniform (because the x-axis is logarithmic this manifests as a long leftwards tail). This ensures
that, provided the assumptions of the test are met, the false positive rate in classical statistics is
constant across sample sizes when H_0 is true. This illustrates why classical *p-values can never*
provide evidence for the null hypothesis (Rouder et al., 2009). When H_0 is true, a proportion of
tests given by the α level will be false positives. Because the estimated effect size with large
sample sizes is typically very small (i.e. close to the truth of zero effect), trivially tiny effects may
thus become statistically significant.

The grey shaded regions in each panel indicate the boundaries of commonly used criterion levels.
For classical statistics the dark grey region corresponds to p-values between 0.05-0.1, sometimes
called "marginally significant." The light grey region denotes the range between 0.01-0.05. Any p-

value to the left of the light grey region would constitute a significant result. For Bayes factors and
the bootstrapped evidence the regions are symmetric around 0. The dark grey region corresponds
to inconclusive evidence that supports neither H_1 nor H_0 (i.e. ⅓-3 for BF_{10}, -0.5-0.5 for ε). The light
grey regions refer to evidence that passed the criterion but which is still relatively weak (i.e. BF_{10}
between 10^{-1} and ⅓ or 3 and 10; ε between -1 to -0.5 or 0.5 to 1). The proportion of these
statistics to the right of the criterion becomes smaller as sample size increases.

Figure 3B shows results from simulations using the uncorrelated but severely heteroscedastic
distribution. For all sample sizes the distributions for classical p-values are biased so that false
positives are drastically inflated. The same is also somewhat true for the default Bayes factor and
bootstrapped evidence. However, for the BSE the skew is at worst modest, while the proportion
of Bayes factors exceeding "strong" evidence for H_1 is larger. This is because the default Bayes
factor is a function of the Pearson's correlation coefficient and the sample size. It is therefore
skewed by heteroscedasticity in the same way as classical statistics. However, since the BSE is
based on non-parametric resampling it is less affected by violations of parametric assumptions.

Next I performed a sensitivity analysis determining how well the BSE test detects true effects. I
repeated the same kind of simulation but now data were chosen from a Gaussian bivariate
distribution with population correlations of ρ=0.3 or ρ=0.7. Unsurprisingly, for all of the three
procedures the evidence for H_1 becomes stronger as the sample size increases. For ρ=0.3 the
evidence passes criterion only for larger samples sizes (Figure 3C) while for most data both
Bayesian and bootstrapped evidence remains inconclusive. Classical statistics are less
conservative as the peak of the distribution with n=120 (green curve) is already below the p<0.01
threshold. For ρ=0.7 the evidence with most sample sizes passes criterion (Figure 3D).

Figure 3. Distributions of statistical evidence in 5,000 simulations of uncorrelated Gaussian data (A), severely heteroscedastic data (B), weakly correlated data with ρ=0.3 (C) or strongly correlated data with ρ=0.7 (D). Six sample sizes were tested (see color code). Each panel shows distributions for the classical p-value (left), default Bayes factor (Wetzels and Wagenmakers, 2012) BF_{10} (middle), and

bootstrapped evidence ε (right). The dark shaded regions denote "inconclusive" results (see text).
The light shaded regions denote results that pass a basic criterion but provide no strong evidence for
a given hypothesis.

I summarized the false positive and correct detection rates as a function of sample size. As
expected, for classical statistics the false positive rate remains constant near the nominal level of
5% across all sample sizes, if data are Gaussian and homoscedastic (Figure 4A). For BF_{10} and ε,
false positive rates for standard criteria ($BF_{10}>3$ and ε>0.5, respectively) decrease as sample size
increases. For either method the false positive rate is already below 5% even at the smallest
sample size (n=15). When heteroscedasticity is present, false positives are dramatically inflated:
approximately one in four tests are positive at p<0.05 (Figure 4B). Both evidence-based methods
also show some inflation; however, false positives for the BSE are only about half that for the
default Bayes factor. For the smallest sample size (n=15) the worst false positive rate for ε>0.5 is
~8.5% compared to ~14.7% for $BF_{10}>3$. When there is a real effect (ρ=0.3) the detection rate rises
steeply and then saturates for all three methods but Bayes factors and BSE are more conservative
than classical p-values (Figure 4C).

2(
210

Figure 4. Detection rates from the simulations in Figure 3 plotted against sample size for classical
p<0.05 (blue), default Bayes factor (Wetzels and Wagenmakers, 2012) BF_{10} (green), and the
bootstrapped evidence ε (red). A. Uncorrelated Gaussian data. B. Severely heteroscedastic data. C.
Correlated data with ρ=0.3.

While the conclusions one would draw from all three approaches are usually similar, one notable
difference is evident between classical and Bayesian inference and the bootstrapped evidence:
distributions for ε tend to become *narrower as sample/effect size increase*. In contrast, the
distributions for p-values and BF_{10} become wider. Note that all of these plots are on logarithmic
scales (log-transformation is inherent to the calculation of ε; see Figure 2 and Materials and
Methods). Despite this, the distributions for p-values and Bayes factors display extraordinary
variability, e.g. the distribution for ρ=0.3 at the largest sample size of n=480 (Figure 3C, red
curves).

Here 95% of simulated p-values are between $8.5×10^{-18}$ and $1.8×10^{-06}$. All are highly significant at
p<0.001 but this range spans many orders of magnitude. The default Bayes factor behaves
similarly. The equivalent range spans BF_{10} between 3,212 and 378 quadrillion. Any of these would
constitute "decisive" evidence of $BF_{10}>100$ (Jeffreys, 1961; Wetzels and Wagenmakers, 2012). But
pragmatically, how much more confident should we be of the highest Bayes factor in this range

compared to the lowest? In comparison, the range for the bootstrapped evidence is between 1.2
and 1.9. Again, these are well above even a strict criterion of $\epsilon>1$ but there is no stark discrepancy
between the weakest and strongest evidence. Replicate experiments would produce very
consistent evidence for H_1.

Naturally, Bayesian analysis depends on the choice of a prior but typically with a range of default
priors the outcome does not vary qualitatively (Wagenmakers et al., 2011b). Nonetheless,
choosing a prior could theoretically also lead to substantial analytic flexibility, thus inflating the
"researcher degrees of freedom"(Simmons et al., 2011). The BSE test on the other hand is
objective. It makes no assumptions beyond the resampling strategy needed for either hypothesis.

Evidence as a function of sample size

The default Bayes factor also places undue confidence on strong effects when sample sizes are
small as in Figure 1A. The Bayes factor is rather large $BF_{10}=90.2$ while the BSE is modest ($\epsilon=0.7$).
The Bayes Factor reflects how much more probable the data are under H_1 than H_0 (Rouder, 2014).
However, from a pragmatic perspective this must nonetheless lead to an inflation of spurious
results.

Figure 5 plots the evidence for a range of effect sizes against sample size. The conclusions we
would draw from bootstrapped evidence (Figure 5A) and the default Bayes factor (Figure 5B,C)
are largely the same. For strong effects, the evidence rises continuously beyond the inconclusive
region, while for weaker effects the evidence starts off as indistinguishable from the situation
when the null hypothesis is true (black curves) until it departs and also rises. This behavior is
natural because if the true effect is weaker than what could be meaningfully detected given the
data at hand this constitutes support for the null hypothesis.

The slopes of the curves for the bootstrapped evidence are far less steep. Thus it is possible to see
the behavior for the full range of conditions within the same plot. There is however one
considerable difference: for a perfect correlation ($\rho=1$) the default Bayes factor immediately rises
even at tiny sample sizes (Figure 5C). At n=3 the BF_{10} is already 48.8. In contrast, the BSE for this
point is low ($\epsilon=-0.4$) and inconclusive. As sample size increases, so does the bootstrapped
evidence. At n=4 it is still inconclusive but favoring H_1 ($\epsilon=0.4$). At n=7 it clearly supports H_1 ($\epsilon=0.9$)
and it continues to rise as sample size increases. This behavior is far more intuitive than that of
the default Bayes factor and also the classical p-value, which would be extremely significant in all
these situations. Compare this to the earlier example of a strong correlation (r=0.9953) with a
small sample size of n=5 (Figure 1A). Classical inference would be extremely significant (p<0.001)
and the default Bayes factor would yield "very strong" evidence for H_1 ($BF_{10}=90.2$). The BSE is
however only fairly modest, especially given the strong effect ($\epsilon=0.7$). It is above the criterion for
conclusive evidence but it does not instill undue confidence in H_1.

The data in this example were in fact drawn from an uncorrelated Gaussian distribution so the
null hypothesis was true. From a pragmatic perspective at least this is problematic especially given
widespread problems with false positives and reproducibility in the scientific literature (Ioannidis,

2005). The bootstrapped evidence provides an intuitive measure of the strength of evidence and

should thus be a safeguard against weak or inconclusive results.

Figure 5. Statistical evidence for H_1 plotted against sample size for a range of effect sizes (see color

code). Individual panels show the bootstrapped evidence ε (A) or the default Bayes factor (Wetzels

and Wagenmakers, 2012) BF_{10} (B,C). Panels C replots the Bayes factor with y-axis zoomed in on zero.

The shaded grey region denotes "inconclusive" evidence (i.e. $-0.5 < \varepsilon < 0.5$ or $\frac{1}{3} < BF_{10} < 3$, respectively).

For the bootstrapped evidence (A) these data represent the mean across 100 simulations.

Simulations of optional stopping

The BSE test has further advantages over classical inference based on significance thresholds. In

classical statistics, even when there is no true effect, it is theoretically possible to reach an

arbitrarily significant p-value, if data collection continues until the p-value passes the significance

threshold. This is known as "optional stopping", which is an incorrect but possibly widespread use

of classical statistics (Masicampo and Lalande, 2012; Lakens, 2015). Under the classical framework

one should first define the expected effect size *a priori*, perform a power analysis to see how large

a sample is needed to detect this effect with sufficiently high probability, and then collect those

data without stopping until the sample is complete. However, typically this is not realistic as one

can often only make a vague guess about the expected effect size.

The bootstrapped evidence does not suffer from this conundrum. First, even if a dubious optional

stopping strategy is used, the false positive rate is not inflated substantially. I simulated this 1,000

times by drawing data repeatedly from an uncorrelated bivariate Gaussian distribution thus

successively increasing the sample size by 1, starting with a minimal sample of n=5. At each step I

applied classical statistics, the default Bayes hypothesis test (Wetzels and Wagenmakers, 2012),

and the BSE test. The first instance one of these tests passed the criterion level, that is p<0.05 for

classical statistics, $BF_{10} > 3$ or $BF_{10} < \frac{1}{3}$ for Bayes factors, and $\varepsilon > 0.5$ or $\varepsilon < -0.5$ for the bootstrapped

evidence, I recorded the measure of evidence. In addition, I also performed this procedure on the

bootstrapped uncertainty and recorded the first instance that $\sigma < 0.2$. If none of the measures

reached criterion simulated data collection would cease at n=150.

Under the assumptions of classical statistics this false positive rate should be near 5%, however,

the actual probability was much greater, 36.4%, illustrating the considerable problems optional

stopping can cause in classical inference. In contrast, the false positive rates of the evidence-

based methods was much lower and well below the classical α level (Bayes factor: 2.9%; bootstrapped evidence: 2.8%).

I repeated the same simulation but this time drawing from the heteroscedastic distribution used in previous examples (Figures 1E,F). Now classical statistics massively inflated support for the alternative hypothesis with a false positive rate of 84.6%. Default Bayes factors fared a lot better but are nonetheless strongly skewed (20.6%). For the bootstrapped evidence on the other hand the false positive rates were only half that (10.3%), again reflecting the fact that it is based on a non-parametric procedure that takes into account the anomalous distribution of the data.

This demonstrates that optional stopping based on symmetric evidence is far less problematic than for classical statistics. In particular, sequential analysis until the bootstrapped evidence reaches conclusive support for either H_1 or H_0 results in only minimal false positive rates even in extreme situations. However, there is an even better optional stopping strategy that could be employed in the bootstrapped evidence framework. When data collection continued until the bootstrapped uncertainty, σ, was 0.2, the false positive rate for using ε>0.5 in the first scenario (homoscedastic Gaussian data) was only 1.3%, while for the heteroscedastic data it was 7.9%. This suggests that using a criterion uncertainty level is the most optimal strategy for minimizing spurious findings in sequential analysis.

Example 1: Anscombe's quartet

Simulations are crucial for testing a method's performance because the ground truth is known. However, for illustration I also apply the method to Anscombe's quartet (Anscombe, 1973) a famous demonstration of the pitfalls of correlation analysis. It consists of four data sets, each comprising 11 pairs of variables, in which Pearson's correlation produces (almost) identical results (r=0.82, p=0.002). Applying the BSE test reveals that while the data afford low but sufficient confidence for the correlation in the first three data sets (Figure 6A-C), in the final example (Figure 6D) the evidence clearly supports H_0 (ε=-1.4) because one influential outlier drives the correlation but the remaining data are uncorrelated.

Interestingly, the confidence in H_1 is actually subtly greater (ε=1) for the third example (Figure 6C) than the first (ε=0.9, Figure 6A). This is because a single outlier contaminates the perfect correlation in this example, whereas the first example contains noisy but normally distributed data.

The BSE does not distinguish strongly between the first and second examples (Figure 6A,B). The second example contains a perfect relationship between x and y; however, it does not conform to the linear relationship assumed by Pearson's correlation. Curve fitting can also be implemented in the BSE framework (see Materials and Methods). Here we could compare a simple linear fit to polynomial curves. The evidence for H_1 with a second-order polynomial is considerably greater (ε=1.6) than for a standard linear model (ε=1). Interestingly, the BSE is also robust to overfitting more complex models: the evidence for higher-order polynomials is weaker than for the second-order (third-order: ε=1.3; fourth-order: ε=1).

Example 2: Links between visual cortex surface area and perceptual function

I further applied the BSE test to published experimental data that showed correlations between
the size of early visual areas and perceptual function. These studies hypothesized that the
transmission speed/strength of lateral connections running tangential to the cortical surface is
reduced for individuals with larger cortical surface areas. In the first two studies, this should
manifest as an anti-correlation between the strength of the Ebbinghaus illusion and V1 surface
area (Schwarzkopf et al., 2011; Schwarzkopf and Rees, 2013). Classical statistics confirmed this
hypothesis in both studies (Figure 6E,F). However, according to the BSE the findings of the initial
study were inconclusive (r=-0.4, p=0.028, ε=0.3). The second study used a more sophisticated
design producing more compelling evidence for this link (r=-0.38, p=0.006, ε=0.7; note, however,
that this study also normalized V1 area by the whole cortical surface area to control for non-
linearity issues and other confounds. For the sake of consistency with the other findings I chose
not to apply this correction here).

The third study (Genç et al., 2014) reported a linear relationship between the speed of travelling
waves in binocular rivalry and the surface areas of V1 and V2. Classical statistics were very similar
for both regions (r=0.67, p=0.0001). However, the bootstrapped evidence was in fact greater for
V2 (Figure 6G; ε=1.1) than V1 (Figure 6H; ε=1.1), possibly because an influential outlier affected
the latter.

Other statistical tests

The BSE test can also address many other questions. One simply needs to change how the effect
size is calculated and how data are resampled during bootstrapping (see Materials and Methods).
For example I also ran simulations for comparing the means of two samples (Supplementary
Figures 1 and 2).

Figure 6. Example data sets. A-D. Anscombe's quartet: Each data sets has approximately the same Pearson's correlation (r=0.82, p=0.0022) and thus all have a default Bayes factor (Wetzels and Wagenmakers, 2012) $BF_{10} \approx 23$. Panels show Typical Gaussian data (A), data showing a perfect non-linear relationship (B), a perfect correlation contaminated by one influential outlier (C) and uncorrelated data contaminated by one influential outlier (D). E-H. Experimental data showing correlations between visual cortical surface area and perceptual function. Correlations between V1 area and Ebbinghaus illusion strength from (Schwarzkopf et al., 2011) (E) and (Schwarzkopf and

Rees, 2013) (F). Correlations from (Genç et al., 2014) between travelling wave speed in binocular

rivalry and the surface areas of V1 (G) and V2 (H). All other conventions as in Figure 1.

Example 3: reassessing evidence for precognition

A few years ago a psychology study reported experimental evidence for the proposition that

participants had precognitive abilities (Bem, 2011). This study was criticized as an illustration of

the shortcomings of classical inference: Bayesian reanalysis found little evidence in favor of

precognition (Wagenmakers et al., 2011b). However, a Bayesian analysis by the original author

supported his claims (Bem et al., 2011). The conclusions are evidently dependent on the exact

prior chosen (Wagenmakers et al., 2011a, 2011b). A cautious prior seems advisable when

evaluating such unusual claims but the debate illustrates how Bayesian inference can appear to

lack objectivity.

Here I reevaluate claims from a more recent study on precognitive abilities (Maier et al., 2014)

with the BSE test (see Materials and Methods). It is a perfect test of the method because these

results seem biologically and physically implausible but both classical and default Bayesian

inference (Wagenmakers et al., 2011b) nonetheless support the alternative hypothesis for several

of the studies (Table 1). In contrast, the BSE does not provide convincing support for precognition.

It is noteworthy that despite a large sample size of 1222 participants, even the web-based study 3

only produced inconclusive evidence ($\varepsilon=-0.1$). This illustrates that in comparison to classical

inference, the BSE is far less susceptible to the inflation of "significant" findings with large sample

sizes. Taken together this suggests that the available data do not provide conclusive evidence for

either H_1 or H_0.

Experiment	Classical statistics	Bayes Factor, BF_{10}	Bootstrapped evidence, ε
Maier Study 1	**t(110)=-2.65, p=0.0092**	2.9 (anecdotal H_1)	0.2 (inconclusive)
Maier Study 2	**t(200)=-2.99, p=0.0032**	**5.9 (moderate H_1)**	0.3 (inconclusive)
Maier Study 3	**t(1221)=-2.36, p=0.0186**	0.5 (anecdotal H_0)	-0.1 (inconclusive)
Maier Unsuccessful 1	t(62)=0.16, p=0.8700	0.14 (moderate H_0)	-1.8 (compelling H_0)
Maier Unsuccessful 2	t(405)=0.44, p=0.6587	0.06 (strong H_0)	-1.8 (compelling H_0)
Maier Unsuccessful 3	t(639)=-1.3, p=0.1943	0.1 (strong H_0)	-0.3 (inconclusive)
Maier Study 4	t(326)=-1.81, p=0.0717	0.3 (strong H_0)	-0.2 (inconclusive)
Aspirin study[*]	**t(21998)=5.19, p<10^{-6}**	10561.9 (decisive H_1)	1.3 (compelling H_1)

Table 1. Reanalysis of data purportedly showing precognitive abilities (Maier et al., 2014) and a

simulated example of a clinical trial. For each experiment this shows the result using classical statistics, a

default Bayes factor (Rouder et al., 2009) and the bootstrapped evidence ε. For Bayes factors and

bootstrapped evidence verbal descriptions of the strength of evidence for H_1 and H_0 are also given. Cells

with bold font denote tests formally supporting H_1, that is, if evidence for the alternative hypothesis is

above criterion (i.e. $p<0.05$, $BF_{10}>3$ or $\varepsilon>0.5$). All these statistics are based on my own analysis of the raw

data. Small discrepancies between the present analysis and previously reported results may be because

statistics were misreported previously although in most cases they are most likely rounding errors. Note

that the pattern of results is qualitatively the same as in previous reports so these discrepancies do not

influence the general conclusions in any way. The asterisks indicate that the data for these studies were

simulated based on the reported effect and sample sizes because the raw data were not available.

Is the BSE test simply too conservative to reveal these rather subtle effects? To test whether the
low evidence in my reanalysis could be due to a lack in sensitivity, I also used the BSE test to
evaluate a small effect size in the clinical literature. In this study, the effect of aspirin on cardiac
health was tested in a large sample (n≈22,000) (Young et al., 1988). The effect was minute
(Cohen's d≈0.07) but highly significant, and it was thus deemed to be of practical value in
promoting health. In the absence of raw data I simulated this data set using the reported effect
and sample sizes. Here the BSE test agrees with classical statistics and Bayesian inference: the
observed data strongly support (ε=1.3) the efficacy of the drug (Table 1). The sample size in this
case is more than sufficient to conclude that this effect is *small but real*. In contrast, to provide
conclusive evidence for the tiny precognition effects, the sample sizes would have to be orders of
magnitude larger (Supplementary Figure 2).
440
441 **Discussion**
442
Here I outlined the bootstrapped evidence test, which makes minimal assumptions and is easily
applicable to numerous situations. It quantifies non-dichotomously the evidence *for the*
alternative or the null hypothesis rather than whether a particular statistic passes an arbitrary
significance threshold. A result does not stand or fall based on its exact value. Rather it allows us
to express how convincing a result is. This also allows for the use of sequential sampling
strategies, which can greatly benefit research practice especially when there is uncertainty about
the expected effect size.
450
A possible criticism of non-dichotomous alternatives to null hypothesis significance testing is that
any measures could be subject to the same thresholding dilemma as classical p-values. If the
consensus emerges that $\varepsilon > 0.5$ is sufficient evidence for H_1 then are we not merely shifting the
problem from p-values to a different measure? However, even in the classical framework many
researchers regularly make non-dichotomous judgments based on p-values. "Marginally
significant" findings are often reported that do not quite pass reach p<0.05. Many researchers are
probably more convinced by $p=10^{-17}$ than p=0.049. The bootstrapped evidence test directly
quantifies the reliability of the data in drawing conclusions about the two competing hypotheses.
Being a new measure it will make it easier for researchers to adopt non-dichotomous thinking
than with p-values.
461
It remains unclear in how far researchers need dichotomous thresholds for statistical inference
and whether labels for the strength of evidence, such as those employed for Bayes factors
(Jeffreys, 1961; Wetzels and Wagenmakers, 2012), are necessary. I would argue that they are not
and deliberately refrained from proposing a categorical stratification of ε. For replication attempts
or incremental experiments for which clear predictions can be made, ε between 0.7-1 can already
be very convincing. In contrast, for more extraordinary claims even ε=1 is still low.
468
Another problem with most commonly used statistics is that they are based on parametric
assumptions that may not hold. Anomalous data can skew the default Bayes factor to a similar
extent as inferences based on classical p-values, because it is based on the same effect size
calculations. Robustness to outliers and heteroscedasticity could be incorporated into Bayesian
hypothesis testing, e.g. by outlier removal procedures. Within the framework of classical

inference robust hypothesis testing suggest following a complicated tree of tests for the presence
of outliers and heteroscedasticity, and then applying the appropriate robust test depending on
the situation (Wilcox, 2005; Pernet et al., 2012). This is often accompanied by remonstrations that
there is a "statistical toolbox" that should be employed and that one method is not best for every
situation (Gigerenzer and Marewski, 2015).

While it is doubtless true that inference should never be made without thought (Gigerenzer,
2004), it is nevertheless advisable to seek a method that serves a universal purpose because in
practice many researchers are not statisticians. The idea of a "statistical toolbox" is fraught with
danger. It must inevitably result in increasing the underreported flexibility in the range of
methods employed by published studies (Simmons et al., 2011; Wagenmakers et al., 2012). Of
course the BSE test does not preclude the use of other statistical methods. It simply provides a
common starting point for inference. It is an objective method with minimal assumptions that in
principle works exactly the same for almost any situation.

Materials and Methods

I will refer to the distributions for the effect size under null and alternative hypothesis as θ_0 and
θ_1, respectively (Figure 2, left panels, blue and red curves). To quantify the *similarity* of these two
distributions at each bootstrapping step the difference between the effects under the two
hypotheses is calculated, that is, θ_{1i} - θ_{0i} where *i* denotes the *i*-th bootstrap step. This produces a
distribution of differences, Δ, between the effects expected under the two hypotheses (Figure 2,
right panels). Theoretically, the evidence for H_1 is thus a kind of standardized score based on this
Δ distribution:

$$z = \frac{|\mu|}{\sigma}$$

499   (1)

where μ and σ, respectively, denote the arithmetic mean and standard deviation of Δ. This
effectively normalizes the shift of this distribution relative to zero by its dispersion. When z is
greater than 1 this implies that the sample effect size, and thus our estimate of the true
population effect, is larger than the *uncertainty* of the estimate (Figure 2A). This provides
evidence supporting the alternative hypothesis. The more data we collect and the sample size, n,
becomes larger, the more accurate is the estimate of the population effect, μ, and the smaller is
the uncertainty, σ.

Unfortunately, this only holds when the alternative hypothesis is in fact true. When the null
hypothesis is true instead, that is, when the population effect is zero, the parameters of Δ do not
reflect the evidence for H_0. While increasing n reduces the uncertainty, σ, on average it also
reduces estimates of the absolute population effect size, μ. It follows that the ratio of these
parameters, z, remains more or less constant.

Thus, in order to quantify how strongly the data support *either* H_1 or H_0 we must weight the ratio
of μ and σ according to how much evidence is available. This is achieved by calculating a third
parameter describing the overlap of θ_0 and θ_1. This is given by:
518

$$\omega = \frac{P(\Delta \; sgn \; \mu \leq 0)}{P(\Delta \; sgn \; \mu > 0)}$$

519   (2)
520

521   That is, dividing the proportion of bootstraps in Δ that have the opposite sign as μ or zero (red
522   area under the curves in the right panels of Figure 2) by the proportion of iterations with the
523   same sign (green area under the curves in the right panels of Figure 2). The strength of evidence,
524   ε, for or against H_1 is then calculated as:
525

$$\varepsilon = \ln \frac{(1 - \omega)|\mu| + \; \omega\sigma}{(1 + \omega \ln n)\sigma}$$

526   (3)
527

While this equation may seem complex, it is essentially the same as z, the standardized score in
equation 1, but it is moderated by the sample size and the overlap between θ_1 and θ_0. When the
data clearly support the alternative hypothesis, the Δ distribution is shifted far away from zero
and thus ω (the ratio of the red and green areas under the curve) is very small (Figure 2A). In fact,
because it is based on the proportion of bootstraps it may even be 0. Under these circumstances
the denominator is close to σ, the numerator is close to $|\mu|$, and thus the evidence is
approximately z. However, when the null hypothesis is true, or the effect size is too small to
clearly support the alternative, the Δ distribution is centered near zero and thus ω is
approximately 1. In this situation the denominator is a multiple of σ, growing ever larger as the
sample size increases. This in turn ensures that the ratio in equation 3 becomes ever smaller and
evidence for the null hypothesis grows (Figure 2B).
539

The numerator is also moderated by the strength of the evidence. When H_0 is true and ω is near
1, the numerator is close to σ. This reflects the fact that when the data provide only weak
evidence, there is substantial uncertainty as to whether the estimate of the effect size is accurate.
When the sample size is large, this means that the denominator dominates the equation and ε
becomes very small. However, when the sample size is *small*, the ratio is close to 1 and thus the
data support neither H_1 nor H_0 very clearly – this means the evidence is inconclusive (Figure 2C).
546

547   To recap, the bootstrapped evidence, ε, measures how confident one can be of the effect size
548   estimate given the uncertainty in the data. If the observed effect is relatively large, the
549   uncertainty will decrease as sample size grows and thus support for the alternative hypothesis
550   also grows. However, when the effect remains considerably smaller than the uncertainty, a larger
551   sample instead provides greater evidence for the null hypothesis.
552

553   Finally, as equation 3 shows, ε is the natural logarithm of this ratio. Therefore, when the ratio is
554   near 1 and the evidence is inconclusive, ε is approximately 0. Positive ε indicates evidence for H_1,
555   while negative ε indicates evidence for H_0. The bootstrapped evidence thus provides a non-

dichotomous measure of the evidence for either hypothesis, similar to a Bayes factor (Wagenmakers, 2007; Rouder et al., 2009; Dienes, 2014). However, while a Bayes factor is a measure of how much one should update the prior odds due to the observed evidence, the bootstrapped evidence is in essence a signal-to-noise ratio. A strong "signal" implies strong evidence for H_1, while a negligible signal with a lot of data provides strong evidence for H_0. The only prior assumptions this procedure makes pertain to how the data are resampled under the two hypotheses.

Bootstrapped correlations

To bootstrap the evidence for a linear correlation data are resampled *with replacement* and on each step the correlation coefficient is computed. To derive the null distribution (θ_0) data are resampled without restriction as would be expected if the effect occurred by chance, that is, observations for the two variables are no longer paired but intermixed randomly. This is essentially standard procedure for non-parametric resampling methods in the classical frequentist framework (although for this purpose permutation analysis where resampling is performed *without* replacement is more common). A classical one-tailed p-value could be calculated by determining the proportion of bootstraps θ_{0i} that are at least as large as the observed effect size – that is, the area under the blue curve to the right of the red diamond.

However, to derive the alternative distribution (θ_1), quantifying the reliability of the observed effect, we restrict the resampling strategy on the alternative hypothesis that there is a correlation. In this case the pairing of data points in each variable is preserved so many resamples will show a positive linear relationship.

As described, we next calculate Δ, the distribution of differences between θ_1 and θ_0 (Figure 2B). Its standard deviation is the uncertainty, σ. For all of the examples given in this article, I used 10,000 bootstrap iterations, except in the interest of time for lengthy simulations and curve fitting examples I only used 1,000 iterations. Reducing the number of iterations only changes the precision of the estimate of ε but does not alter the general conclusions substantially.

The further apart the two distributions for θ_0 and θ_1 are, the farther Δ is from zero and the greater is the evidence for H_1. This is quantified by ε. When ε is very negative, the evidence favors H_0 because it means the two distributions for θ_0 and θ_1 overlap considerably which means that Δ is centered near zero. Intuitively this indicates that the effect size estimate under H_1 could very likely have been smaller than that under H_0. When the evidence is $-0.5 < \varepsilon < 0.5$ this provides inconclusive support for the either hypothesis. This region is somewhat arbitrary but it reflects the fact that while there is overlap between the distributions for θ_0 and θ_1, there is not sufficient data to be confident that there is no subtle effect. This corresponds to the range of ε one typically obtains with small sample sizes when the null hypothesis is true.

Bootstrapping differences

Naturally, the same procedure can be applied to other statistical comparisons in addition to tests of correlation. For instance, when *comparing the means of two independent samples* the effect

size is the difference between the sample means. To estimate the null distribution, θ_0,
observations are resampled with replacement and divided into new samples of the same size as
the original samples. To estimate the distribution for the alternative hypothesis, θ_1, the
segregation of the two samples is maintained and resampling is done *within* each sample. In all
other respects, the procedure is identical to the correlation test already described.

When testing the *difference between two repeated measures* the same underlying principle
applies. Here the effect size is the mean over the *differences* in each pair of observations. We
estimate θ_1 by maintaining the pairing but resampling with replacement. For estimating θ_0 the
pairing is also kept intact because what matters is only the variance across repeated measures.
However, under the null hypothesis the order of measures is irrelevant so the resampling
randomizes the sign for each observation. This corresponds to scrambling the order of
observations in a repeated measures design.

Similarly, the BSE can also be used to test the *difference between two correlations*. Again the data
need to be resampled based on the assumptions of the null as well as the alternative hypothesis.
In this case the null hypothesis resamples data ignoring how variables are paired while the
alternative hypothesis preserves the pairing. The estimated effect size is the difference between
the two correlation coefficients.

The situation becomes more complicated for testing the *difference of one sample from a fixed
value* (e.g., when comparing a normative sample to a patient case-study). Conceptually, it is
possible to use the same resampling strategy as for a repeated measures design: the observations
are the differences from the fixed value and for resampling the null distribution θ_0 we randomize
the sign of each observation. However, this approach is probably not sufficiently conservative.
While it is conceptually correct for repeated measures designs to assume a mean difference of 0
under the null hypothesis, in many other situations fixed values are themselves subject to
variability and/or measurement error. For instance, a measurement in a case study is subject to
within-subject variability and chance performance in a behavioral task follows a probability
distribution.

Such variance should be incorporated in the BSE framework to estimate the null distribution more
accurately. Therefore, whenever possible it is advisable to design experiments with a control or
baseline condition against which to test the experimental manipulation. For example, when
testing whether a group of individuals has above average IQ it would make more sense to
compare this sample to a well-matched control group rather than to assume an average IQ of 100.
This is not only useful for the BSE test but is generally good experimental design.

Bootstrapping tests against chance

Alternatively, the null distribution can also be simulated. This is suitable for testing a binomial
process, such as whether a coin is fair or whether a participant performed better than chance at a
behavioral task. As usual, in each bootstrap step the observed data (e.g. a series of 1s and 0s for
heads or tails) are resampled to obtain the alternative distribution θ_1. However, for estimating θ_0
we instead generate a *new* set of 1s and 0s of the same number as the observed trials using the

chance probability (i.e. 0.5 or whatever the chance level is). Alternatively, one can also permute
the raw trial data in each resampling step and recalculate the accuracy. The latter approach is
advised when the design is unbalanced or if there is any suspicion that chance may not have a
binomial distribution. In all other ways the procedure works as described.

A very similar approach for simulating a chance distribution based on the assumptions underlying
the null hypothesis can also be used for other problems, for example comparing the performance
of a group of participants against chance. In this case, we can simulate θ_0 by generating a new set
of 'chance' participants at each resampling step under the same conditions as the actual
experiment (same chance probability, number of trials, and number of participants).

Bootstrapping curve fits

The bootstrapped evidence procedure also affords itself easily for curve fitting or regression
analyses. The estimated effect size in this case is the coefficient of determination, R^2 (or
goodness-of-fit). Otherwise the procedure works in much the same way as for calculating
correlations. Under the null hypothesis the observations for the dependent and independent
variables are scrambled randomly with replacement. Under the alternative hypothesis, the pairing
is kept intact but observations are resampled with replacement.

Acknowledgements

I thank Ged Ridgway, Benjamin de Haas, and Micah Allen for comments on previous versions of
this manuscript. DSS is supported by an ERC Starting Grant.

References

Anscombe, F. J. (1973). Graphs in Statistical Analysis. *Am. Stat.* 27, 17–21. doi:10.2307/2682899.

Bem, D. J. (2011). Feeling the future: experimental evidence for anomalous retroactive influences
on cognition and affect. *J. Pers. Soc. Psychol.* 100, 407–425. doi:10.1037/a0021524.

Bem, D. J., Utts, J., and Johnson, W. O. (2011). Must psychologists change the way they analyze
their data? *J. Pers. Soc. Psychol.* 101, 716–719. doi:10.1037/a0024777.

Cohen, J. (1994). The Earth is round (p < .05). *Am. Psychol.* 49, 997–1003.

Colquhoun, D. (2014). An investigation of the false discovery rate and the misinterpretation of p-
values. *R. Soc. Open Sci.* 1, 140216. doi:10.1098/rsos.140216.

Cumming, G. (2014). The new statistics: why and how. *Psychol. Sci.* 25, 7–29.
doi:10.1177/0956797613504966.

Dienes, Z. (2014). Using Bayes to get the most out of non-significant results. *Quant. Psychol.*
Meas. 5, 781. doi:10.3389/fpsyg.2014.00781.

Genç, E., Bergmann, J., Singer, W., and Kohler, A. (2014). Surface Area of Early Visual Cortex
Predicts Individual Speed of Traveling Waves During Binocular Rivalry. *Cereb. Cortex N. Y.*
N 1991. doi:10.1093/cercor/bht342.

Gigerenzer, G. (2004). Mindless statistics. *J. Socio-Econ.* 33, 587–606.
doi:10.1016/j.socec.2004.09.033.

Gigerenzer, G., and Marewski, J. N. (2015). Surrogate Science The Idol of a Universal Method for
Scientific Inference. *J. Manag.* 41, 421–440. doi:10.1177/0149206314547522.

Halsey, L. G., Curran-Everett, D., Vowler, S. L., and Drummond, G. B. (2015). The fickle P value
generates irreproducible results. *Nat. Methods* 12, 179–185. doi:10.1038/nmeth.3288.

Hoekstra, R., Morey, R. D., Rouder, J. N., and Wagenmakers, E.-J. (2014). Robust misinterpretation
of confidence intervals. *Psychon. Bull. Rev.* doi:10.3758/s13423-013-0572-3.

Ioannidis, J. P. A. (2005). Why most published research findings are false. *PLoS Med.* 2, e124.
doi:10.1371/journal.pmed.0020124.

Jeffreys, H. (1961). *Theory of probability.* Oxford, UK: Oxford University Press.

Lakens, D. (2015). What p-hacking really looks like. Available at: https://osf.io/ycag9/ [Accessed
February 4, 2015].

Maier, M. A., Buechner, V. L., Kuhbandner, C., Pflitsch, M., Fernandez-Capo, M., and Gamiz-
Sanfeliu, M. (2014). Feeling the Future Again: Retroactive Avoidance of Negative Stimuli.
J. Conscious. Stud. 21, 121–152.

Masicampo, E. J., and Lalande, D. R. (2012). A peculiar prevalence of p values just below .05. *Q. J.*
Exp. Psychol. 2006 65, 2271–2279. doi:10.1080/17470218.2012.711335.

Morey, R. D., Rouder, J. N., Verhagen, J., and Wagenmakers, E.-J. (2014). Why Hypothesis Tests
Are Essential for Psychological Science: A Comment on Cumming (2014). *Psychol. Sci.*
doi:10.1177/0956797614525969.

Nuzzo, R. (2014). Scientific method: statistical errors. *Nature* 506, 150–152. doi:10.1038/506150a.

Pernet, C. R., Wilcox, R., and Rousselet, G. A. (2012). Robust correlation analyses: false positive
and power validation using a new open source matlab toolbox. *Front. Psychol.* 3, 606.
doi:10.3389/fpsyg.2012.00606.

Psychological Science (2014). 2014 Submission Guidelines - Association for Psychological Science.
Available at:
http://www.psychologicalscience.org/index.php/publications/journals/psychological_scie
nce/ps-submissions [Accessed April 2, 2014].

Rouder, J. N. (2014). Optional stopping: No problem for Bayesians. *Psychon. Bull. Rev.*
doi:10.3758/s13423-014-0595-4.

Rouder, J. N., Speckman, P. L., Sun, D., Morey, R. D., and Iverson, G. (2009). Bayesian t tests for
accepting and rejecting the null hypothesis. *Psychon. Bull. Rev.* 16, 225–237.
doi:10.3758/PBR.16.2.225.

Rousselet, G. A., and Pernet, C. R. (2012). Improving standards in brain-behaviour correlation
analyses. *Front. Hum. Neurosci.* 6:, 119. doi:10.3389/fnhum.2012.00119.

Savalei, V., and Dunn, E. (2015). Is the call to abandon p-values the red herring of the replicability
crisis? *Cognition* 6, 245. doi:10.3389/fpsyg.2015.00245.

Schwarzkopf, D. S., De Haas, B., and Rees, G. (2012). Better ways to improve standards in brain-
behavior correlation analysis. *Front. Hum. Neurosci.* 6, 200.
doi:10.3389/fnhum.2012.00200.

Schwarzkopf, D. S., and Rees, G. (2013). Subjective size perception depends on central visual
cortical magnification in human v1. *PloS One* 8, e60550.
doi:10.1371/journal.pone.0060550.

Schwarzkopf, D. S., Song, C., and Rees, G. (2011). The surface area of human V1 predicts the
subjective experience of object size. *Nat. Neurosci.* 14, 28–30. doi:10.1038/nn.2706.

Simmons, J. P., Nelson, L. D., and Simonsohn, U. (2011). False-positive psychology: undisclosed
flexibility in data collection and analysis allows presenting anything as significant. *Psychol.*
Sci. 22, 1359–1366. doi:10.1177/0956797611417632.

Trafimow, D., and Marks, M. (2015). Editorial. *Basic Appl. Soc. Psychol.* 37, 1–2.
doi:10.1080/01973533.2015.1012991.

Wagenmakers, E.-J. (2007). A practical solution to the pervasive problems of p values. *Psychon.*
Bull. Rev. 14, 779–804.

Wagenmakers, E.-J., Wetzels, R., Borsboom, D., Kievit, R., and van der Maas, H. L. J. (2011a). Yes,
psychologists must change the way they analyze their data: Clarifications for Bem, Utts, &
Johnson. Available at:
http://dl.dropbox.com/u/1018886/ClarificationsForBemUttsJohnson.pdf.

Wagenmakers, E.-J., Wetzels, R., Borsboom, D., and van der Maas, H. L. J. (2011b). Why
psychologists must change the way they analyze their data: the case of psi: comment on
Bem (2011). *J. Pers. Soc. Psychol.* 100, 426–432. doi:10.1037/a0022790.

Wagenmakers, E.-J., Wetzels, R., Borsboom, D., Maas, H. L. J. van der, and Kievit, R. A. (2012). An
Agenda for Purely Confirmatory Research. *Perspect. Psychol. Sci.* 7, 632–638.
doi:10.1177/1745691612463078.

Wetzels, R., and Wagenmakers, E.-J. (2012). A default Bayesian hypothesis test for correlations
and partial correlations. *Psychon. Bull. Rev.* 19, 1057–1064. doi:10.3758/s13423-012-
0295-x.

Wilcox, R. R. (2005). *Introduction to robust estimation and hypothesis testing*. Academic Press.

Wilcox, R. R., and Muska, J. (2001). Inferences about correlations when there is
heteroscedasticity. *Br. J. Math. Stat. Psychol.* 54, 39–47.

Young, F., Nightingale, S., and Temple, R. (1988). The preliminary report of the findings of the
aspirin component of the ongoing physicians' health study: The fda perspective on aspirin
for the primary prevention of myocardial infarction. *JAMA* 259, 3158–3160.
doi:10.1001/jama.1988.03720210048028.

Supplementary Figure 1. Distributions as shown in Figure 4 but for comparing the means of two samples. A.
No difference, i.e. normally distributed data with unit standard deviation and population mean of 0. B. A
weak effect with a population difference of 0.25 and unit standard deviation. All other conventions as in
Figure 4.

Supplementary Figure 2. Statistical evidence for H_1 plotted against sample size for a range of effect sizes

(see color code) when comparing the means of two samples. Individual panels show the bootstrapped

evidence ε (A) or the default Bayes factor[1] BF_{10} (B,C). Panel C replots the Bayes factor with y-axis zoomed in

on zero. The shaded grey region denotes "inconclusive" evidence (i.e. -0.5<ε<0.5 or ⅓<BF_{10}<3, respectively).

For the bootstrapped evidence (A) these data represent the mean across 100 simulations.