1 Testing for ancient selection using cross-population allele
2 frequency differentiation

3 Fernando Racimo[1,*]

4 **1 Department of Integrative Biology, University of California, Berkeley, CA, USA**

5 **∗ E-mail: fernandoracimo@gmail.com**

6 1 Abstract

A powerful way to detect selection in a population is by modeling local allele frequency changes in a

particular region of the genome under scenarios of selection and neutrality, and finding which model is

most compatible with the data. Chen et al. [1] developed a composite likelihood method called XP-CLR

that uses an outgroup population to detect departures from neutrality which could be compatible with

hard or soft sweeps, at linked sites near a beneficial allele. However, this method is most sensitive to recent

selection and may miss selective events that happened a long time ago. To overcome this, we developed

an extension of XP-CLR that jointly models the behavior of a selected allele in a three-population tree.

Our method - called 3P-CLR - outperforms XP-CLR when testing for selection that occurred before two

populations split from each other, and can distinguish between those events and events that occurred

specifically in each of the populations after the split. We applied our new test to population genomic

data from the 1000 Genomes Project, to search for selective sweeps that occurred before the split of

Africans and Eurasians, but after their split from Neanderthals, and that could have presumably led to

the fixation of modern-human-specific phenotypes. We also searched for sweep events that occurred in

East Asians, Europeans and the ancestors of both populations, after their split from Africans.

21 2 Introduction

22 Genetic hitchhiking will distort allele frequency patterns at regions of the genome linked to a beneficial

23 allele that is rising in frequency [2]. This is known as a selective sweep. If the sweep is restricted to a

24 particular population and does not affect other closely related populations, one can detect such an event

25 by looking for extreme patterns of localized population differentiation, like high values of F_{st} at a specific

26 locus [3]. This and other related statistics have been used to scan the genomes of present-day humans

from different populations, so as to detect signals of recent positive selection [4–7].

Once it became possible to sequence entire genomes of archaic humans (like Neanderthals) [8–10], researchers also began to search for selective sweeps that occurred in the ancestral population of all present-day humans. For example, ref. [8] searched for genomic regions with a depletion of derived alleles in a low-coverage Neanderthal genome, relative to what would be expected given the derived allele frequency in present-day humans. This is a pattern that would be consistent with a sweep in present-day humans. Later on, ref. [10] developed a hidden Markov model (HMM) that could identify regions where Neanderthals fall outside of all present-day human variation (also called "external regions"), and are therefore likely to have been affected by ancient sweeps in early modern humans. They applied their method to a high-coverage Neanderthal genome. Then, they ranked these regions by their genetic length, to find segments that were extremely long, and therefore highly compatible with a selective sweep. Finally, ref. [11] used summary statistics calculated in the neighborhood of sites that were ancestral in archaic humans but fixed derived in all or almost all present-day humans, to test if any of these sites could be compatible with a selective sweep model. While these methods harnessed different summaries of the patterns of differentiation left by sweeps, they did not attempt to explicitly model the process by which these patterns are generated over time.

Chen et al. [1] developed a method called XP-CLR, which is designed to test for selection in one population after its split from a second, outgroup, population t_{AB} generations ago. It does so by modeling the evolutionary trajectory of an allele under linked selection and under neutrality, and then comparing the likelihood of the data under each of the two models. The method detects local allele frequency differences that are compatible with the linked selection model [2], along windows of the genome.

XP-CLR is a powerful test for detecting selective events restricted to one population. However, it provides little information about when these events happened, as it models all sweeps as if they had immediately occurred in the present generation. Additionally, if one is interested in selective sweeps that took place before two populations a and b split from each other, one would have to run XP-CLR separately on each population, with a third outgroup population c that split from the ancestor of a and b t_{ABC} generations ago (with $t_{ABC} > t_{AB}$). Then, one would need to check that the signal of selection appears in both tests. This may miss important information about correlated allele frequency changes shared by a and b, but not by c, limiting the power to detect ancient events.

To overcome this, we developed an extension of XP-CLR that jointly models the behavior of an allele

in all 3 populations, to detect selective events that occurred before or after the closest two populations split from each other. Below we briefly review the modeling framework of XP-CLR and describe our new test, which we call 3P-CLR. In the Results, we show this method outperforms XP-CLR when testing for selection that occurred before the split of two populations, and can distinguish between those events and events that occurred after the split, unlike XP-CLR. We then apply the method to population genomic data from the 1000 Genomes Project [12], to search for selective sweep events that occurred before the split of Africans and Eurasians, but after their split from Neanderthals. We also use it to search for selective sweeps that occurred in the Eurasian ancestral population, and to distinguish those from events that occurred specifically in East Asians or specifically in Europeans.

3 Methods

3.1 XP-CLR

First, we review the procedure used by XP-CLR to model the evolution of allele frequency changes of two populations a and b that split from each other t_{AB} generations ago (Figure 1.A). For neutral SNPs, Chen et al. [1] use an approximation to the Wright-Fisher diffusion dynamics [13]. Namely, the frequency of a SNP in a population a (p_A) in the present is treated as a random variable governed by a normal distribution with mean equal to the frequency in the ancestral population (β) and variance proportional to the drift time ω from the ancestral to the present population:

$$p_A|\beta \sim N(\beta, \omega\beta(1-\beta)) \tag{1}$$

where $\omega = t_{AB}/(2N_e)$ and N_e is the effective size of population A.

If a SNP is segregating in both populations - i.e. has not hit the boundaries of fixation or extinction - this process is time-reversible. Thus, one can model the frequency of the SNP in population a with a normal distribution having mean equal to the frequency in population b and variance proportional to the sum of the drift time (ω) between a and the ancestral population, and the drift time between b and the ancestral population (ψ):

$$p_A|p_B \sim N(p_B, (\omega + \psi)p_B(1 - p_B)) \tag{2}$$

4

For SNPs that are linked to a beneficial allele that has undergone a sweep in population a only, Chen et al. [1] model the allele as evolving neutrally until the present and then apply a transformation to the normal distribution that depends on the distance to the selected allele r and the strength of selection s [14, 15]. Let $c = 1 - q_0^{r/2}$ where q_0 is the frequency of the beneficial allele in population a before the sweep begins. The frequency of a neutral allele is expected to increase from p to $1 - c + cp$ if the allele is linked to the beneficial allele, and this occurs with probability equal to the frequency of the neutral allele (p) before the sweep begins. Otherwise, the frequency of the neutral allele is expected to decrease from p to cp. This leads to the following transformation of the normal distribution:

$$f(p_A|p_B, r, s, \omega, \psi) = \frac{1}{\sqrt{2\pi}\sigma}\frac{p_A + c - 1}{c^2}e^{-\frac{(p_A + c - 1 - cp_B)^2}{2c^2\sigma^2}}I_{[1-c,1]}(p_A) + \frac{1}{\sqrt{2\pi}\sigma}\frac{c - p_A}{c^2}e^{-\frac{(p_A - cp_B)^2}{2c^2\sigma^2}}I_{[0,c]}(p_A) \tag{3}$$

where $\sigma^2 = (\omega + \psi)p_b(1 - p_b)$ and $I_{[x,y]}(z)$ is 1 on the interval $[x, y]$ and 0 otherwise.

For $s \to 0$ or $r >> s$, this distribution converges to the neutral case. Let $\mathbf{v}$ be the vector of all drift times that are relevant to the scenario we are studying. In this case, it will be equal to (ω, ψ) but in more complex cases below, it may include additional drift times. Let $\mathbf{r}$ be the vector of recombination fractions between the beneficial alleles and each of the SNPs within a window of arbitrary size. We can then calculate the product of likelihoods over all k SNPs in that window for either the neutral or the linked selection model, after binomial sampling of alleles from the population frequency, and conditioning on the event that the allele is segregating in the population:

$$CL_{XP-CLR}(\mathbf{r}, \mathbf{v}, s) = \prod_{j=1}^{k} \frac{\int_0^1 f(p_A^j|p_B^j, \mathbf{v}, s, r^j)\binom{n}{m_j}(p_A^j)^{m_j}(1 - p_A^j)^{n-m_j}dp_A^j}{\int_0^1 f(p_A^j|p_B^j, \mathbf{v}, s, r^j)dp_A^j} \tag{4}$$

We note that the denominator in the above equation is not explicitly stated in ref. [1] for ease of notation, but appears in the published online implementation of the method. Because we are ignoring the correlation in frequencies produced by linkage, this is a composite likelihood [16, 17]. Finally, we obtain a composite likelihood ratio statistic S_{XP-CLR} of the hypothesis of linked selection over the hypothesis of neutrality:

$$S_{XP-CLR} = 2[sup_{\mathbf{r},\mathbf{v},s}log(CL_{XP-CLR}(\mathbf{r}, \mathbf{v}, s)) - sup_{\mathbf{v}}log(CL_{XP-CLR}(\mathbf{r}, \mathbf{v}, s = 0))] \tag{5}$$

For ease of computation, Chen et al. [1] assume that $\mathbf{r}$ is given (via a recombination map) and we will do so too. Furthermore, they empirically estimate $\mathbf{v}$ using F_2 statistics [18] calculated over the whole genome, and assume selection is not strong or frequent enough to affect their genome-wide values. Because we are interested in selection over long time scales, the new methods we will present below are optimally run using drift times calculated from population split times and effective population sizes estimated using model-based demographic inference methods, like $\partial a \partial i$ [19] or fastsimcoal2 [20].

3.2   3P-CLR

We are interested in the case where a selective event occurred more anciently than the split of two populations (a and b) from each other, but more recently than their split from a third population c (Figure 1.B). We begin by modeling p_A and p_B as evolving from an unknown common ancestral frequency β:

$$p_A|\beta,\omega \sim N(\beta,\omega\beta(1-\beta)) \tag{6}$$

$$p_B|\beta,\psi \sim N(\beta,\psi\beta(1-\beta)) \tag{7}$$

Let χ be the drift time separating the most recent common ancestor of a and b from the most recent common ancestor of a, b and c. Additionally, let ν be the drift time separating population c in the present from the most recent common ancestor of a, b and c. Given these parameters, we can treat β as an additional random variable that either evolves neutrally or is linked to a selected allele that swept immediately more anciently than the split of a and b. In both cases, the distribution of β will depend on the frequency of the allele in population c (p_C) in the present. In the neutral case:

$$f_{neut}(\beta|p_C,\nu,\chi) = N(p_C,(\nu+\chi)p_C(1-p_C)) \tag{8}$$

In the linked selection case:

$$f_{sel}(\beta|p_C,\nu,\chi,r,s) = \frac{1}{\sqrt{2\pi}\kappa}\frac{\beta+c-1}{c^2}e^{-\frac{(\beta+c-1-cp_C)^2}{2c^2\kappa^2}}I_{[1-c,1]}(\beta) + \frac{1}{\sqrt{2\pi}\kappa}\frac{c-\beta}{c^2}e^{-\frac{(\beta-cp_C)^2}{2c^2\kappa^2}}I_{[0,c]}(\beta) \tag{9}$$

where $\kappa^2 = (\nu+\chi)p_C(1-p_C)$

The frequencies in a and b given the frequency in c can be obtained by integrating β out. This leads

to a density function that models selection in the ancestral population of a and b.

$$f(p_A, p_B|p_C, \mathbf{v}, r, s) = \int_0^1 f_{neut}(p_A|\beta, \omega) f_{neut}(p_B|\beta, \psi) f_{sel}(\beta|p_C, \nu, \chi, r, s) d\beta \tag{10}$$

Additionally, formula 10 can be modified to test for selection that occurred specifically in one of the terminal branches that lead to a or b (Figures 1.C and 1.D), rather than in the ancestral population of a and b. For example, the density of frequencies for a scenario of selection in the branch leading to a can be written as:

$$f(p_A, p_B|p_C, \mathbf{v}, r, s) = \int_0^1 f_{sel}(p_A|\beta, \omega, r, s) f_{neut}(p_B|\beta, \psi) f_{neut}(\beta|p_C, \nu, \chi) d\beta \tag{11}$$

We will henceforth refer to the version of 3P-CLR that is tailored to detect selection in the internal branch that is ancestral to a and b as 3P-CLR(Int). In turn, the versions of 3P-CLR that are designed to detect selection in each of the daughter populations a and b will be designated as 3P-CLR(A) and 3P-CLR(B), respectively.

We can now calculate the probability density of specific allele frequencies in populations a and b, given that we observe m_C derived alleles in a sample of size n_C from population c:

$$f(p_A, p_B|m_C, \mathbf{v}, r, s) = \int_0^1 f(p_A, p_B|p_C, \mathbf{v}, r, s) f(p_C|m_C) dp_C \tag{12}$$

and

$$f(p_C|m_C) = \frac{1}{B(m_C, n_C - m_C + 1)} p_C^{m_C-1} (1 - p_C)^{n_C - m_C} \tag{13}$$

where B(x,y) is the Beta function.

Conditioning on the event that the site is segregating in the population, we can then calculate the probability of observing m_A and m_B derived alleles in a sample of size n_A from population a and a sample of size n_B from population b, respectively, given that we observe m_C derived alleles in a sample of size n_C from population c, using binomial sampling:

$$P(m_A, m_B|m_C, \mathbf{v}, r, s) = \frac{\int_0^1 \int_0^1 P(m_A|p_A) P(m_B|p_B) f(p_A, p_B|m_C, \mathbf{v}, r, s) dp_A dp_B}{\int_0^1 \int_0^1 f(p_A, p_B|m_C, \mathbf{v}, r, s) dp_A dp_B} \tag{14}$$

138 where

$$P(m_A|p_A) = \binom{n_A}{m_A} p_A^{m_A}(1-p_A)^{n_A-m_A} \tag{15}$$

139 and

$$P(m_B|p_B) = \binom{n_B}{m_B} p_B^{m_B}(1-p_B)^{n_B-m_B} \tag{16}$$

140 This allows us to calculate a composite likelihood of the derived allele counts in a and b given the
141 derived allele counts in c:

$$CL_{3P-CLR}(\mathbf{r}, \mathbf{v}, s) = \prod_{j=1}^{k} P(m_A^j, m_B^j|m_C^j, \mathbf{v}, r^j, s) \tag{17}$$

As before, we can use this composite likelihood to produce a composite likelihood ratio statistic
that can be calculated over regions of the genome to test the hypothesis of linked selection centered on a
particular locus against the hypothesis of neutrality. Due to computational costs in numerical integration,
we skip the sampling step for population c (formula 13) in our implementation of 3P-CLR. In other
words, we assume $p_C = m_C/n_C$, but this is also assumed in XP-CLR when computing its corresponding
outgroup frequency. To perform the numerical integrations, we used the package Cubature (v.1.0.2). We
implemented our method in a freely available C++ program that can be downloaded from here:

`https://github.com/ferracimo` [WILL POST IT AFTER PUBLICATION]

150   4   Results

151   4.1   Simulations

We generated simulations in SLiM [21] to test the performance of XP-CLR and 3P-CLR in a three-
population scenario. We first focused on the performance of 3P-CLR(Int) in detecting ancient selective
events that occurred in the ancestral branch of two sister populations. We assumed that the population
history had been correctly estimated by the researcher (i.e. the drift parameters and population topology
were known). First, we simulated scenarios in which a beneficial mutation arose in the ancestor of
populations a and b, before their split from each other but after their split from c (Table 1). Although

158 both XP-CLR and 3P-CLR are sensitive to partial or soft sweeps (as they do not rely on extended

159 patterns of haplotype homozygosity [1]), we required the allele to have fixed before the split (at time t_{ab})

160 to ensure that the allele had not been lost before it, and also to ensure that the sweep was restricted to

161 the internal branch of the tree. We fixed the effective size of all three populations at $N_e = 10,000$. Each

162 simulation consisted in a 5 cM region and the beneficial mutation occurred in the center of this region.

163 The mutation rate was set at $2.5 * 10^{-8}$ per generation and the recombination rate was set at 10^{-8} per

164 generation.

165   To make a fair comparison to 3P-CLR(Int), and given that XP-CLR is a two-population test, we

166 applied XP-CLR in two ways. First, we pretended population b was not sampled, and so the "test"

167 panel consisted of individuals from a only, while the "outgroup" consisted of individuals from c. In the

168 second implementation (which we call "XP-CLR-avg"), we used the same outgroup panel, but pooled

169 the individulas from a and b into a single panel, and this pooled panel was the "test". The window size

170 was set at 0.5 cM and the space between the center of each window was set at 600 SNPs. To speed up

171 computation, and because we are largely interested in comparing the relative performance of the three

172 tests under different scenarios, we used only 20 randomly chosen SNPs per window in all tests. We note,

173 however, that the performance of all three tests can be improved by using more SNPs per window.

174   Figure 2 shows receiver operating characteristic (ROC) curves comparing the sensitivity and specificity

175 of 3P-CLR(Int), 3P-CLR(A), XP-CLR and XP-CLR-avg in the first six demographic scenarios described

176 in Table 1. Each ROC curve was made from 100 simulations under selection (with $s = 0.1$ for the central

177 mutation) and 100 simulations under neutrality (with $s = 0$ and no fixation required). In each simulation,

178 100 haploid individuals (or 50 diploids) were sampled from population a, 100 individuals from population

179 b and 100 individuals from the outgroup population c. For each simulation, we took the maximum value

180 at a region in the neighborhood of the central mutation $(+/- 0.5$ cM) and used those values to compute

181 ROC curves under the two models.

182   When the split times are recent or moderately ancient (models A to D), 3P-CLR(Int) outperforms

183 the two versions of XP-CLR. Furthermore, 3P-CLR(A) is the test that is least sensitive to selection in

184 the internal branch as it is only meant to detect selection in the terminal branch leading to population

185 a. When the split times are very ancient (models E and F), none of the tests perform well. The root

186 mean squared error (RMSE) of the genetic distance between the true selected site and the highest scored

187 window is comparable across tests in all six scenarios (Figure S2). 3P-CLR(Int) is the best test at finding

the true location of the selected site in almost all demographic scenarios.

We also simulated a situation in which only a few individuals (e.g. a small sample of archaic humans)
have been sequenced from the outgroup, while large numbers of sequences are available from the tests
(e.g. two populations of present-day humans). Figures S1 and S3 show the ROC curves and RMSE plots,
respectively, for a scenario in which 100 individuals were sampled from the test populations but only
10 individuals (5 diploids) were sampled from the outgroup. Unsurprisingly, all tests have less power to
detect selection when the split times and the selection events are recent to moderately ancient (models
A-D). Interestingly though, when the split times and the selective events are very ancient (models E-F),
both 3P-CLR and XP-CLR perform better when using a small ougroup panel (Figure S1) than when
using a large outgroup panel (Figure 2). This may be because both of these tests require the outgroup
sample at each site to be a segregating polymorphism, and sites that are polymorphic in a small panel
are, on average, more ancient than sites that are polymorphic in a large panel. Because the recent
polymorphisms in the outgroup carry little or no information about ancient selection in the tests, they
are less likely to contribute to differences in the likelihood functions for the selection and the neutrality
models, and so they make these tests less efficient at distinguishing these two models.

Importantly, the usefulness of 3P-CLR(Int) resides not just in its performance at detecting selective
sweeps in the ancestral population, but in its specific sensitivity to that particular type of events. Because
the test relies on correlated allele frequency differences in both population a and population b (relative
to the outgroup), selective sweeps that are specific to only one of the populations will not lead to high
3P-CLR(Int) scores, but will instead lead to high 3P-CLR(A) scores or 3P-CLR(B) scores, depending on
where selection took place. Figure 3 shows ROC curves in two scenarios in which a selective sweep event
occurred only in population a (models I and J in Table 1), using 100 sampled individuals from each of
the 3 populations. Here, XP-CLR performs well, but is outperformed by 3P-CLR(A). Furthermore, 3P-
CLR(Int) shows almost no sensitivity to the recent sweep. For example, in Model I, at a specificity of 90%,
3P-CLR(A) and XP-CLR(A) have 86% and 80% sensitivity, respectively, while at the same specificity,
3P-CLR(Int) only has 18% sensitivity. One can compare this to the same demographic scenario but with
selection occurring in the ancestral population of a and b (model C, Figure 2), where at 90% specificity,
3P-CLR(A) and XP-CLR(A) have 72% and 84% sensitivity, respectively, while 3P-CLR(Int) has 90%
sensitivity. We also observe that 3P-CLR(A) is the best test at finding the true location of the selected
site when selection occurs in the terminal branch leading to population a (Figure S4).

4.2   Selection in Eurasians

We first applied 3P-CLR to modern human data from the 1000 Genomes Project [12]. We used the African-American recombination map [22] to convert physical distances into genetic distances. We focused on Europeans and East Asians as the two sister populations, using Africans (excluding admixed African-Americans) as the outgroup population (Figure S5.A). We randomly sampled 100 individuals from each population and obtained sample derived allele frequencies every 10 SNPs in the genome. We then calculated likelihood ratio statistics by a sliding window approach, where we sampled a "central SNP" once every 20 SNPs. The central SNP in each window was the candidate beneficial SNP for that window. We set the window size to 0.25 cM, and randomly sampled 100 SNPs from each window, centered around the candidate beneficial SNP. In each window, we calculated 3P-CLR to test for selection at three different branches of the population tree: the terminal branch leading to Europeans (3P-CLR Europe), the terminal branch leading to East Asians (3P-CLR East Asia) and the ancestral branch of Europeans and East Asians (3P-CLR Eurasia). Results are shown in Figure 4. For each scan, we selected the windows in the top 99.9% quantile of scores and merged them together if they were contiguous. Tables 2, 3 and 4 show the top hits for Europeans, East Asians and the ancestral Eurasian branch, respectively

We observe several genes that have been identified in previous selection scans. In the East Asian branch, one of the top hits is *EDAR*. This gene codes for a protein involved in hair thickness and incisor tooth morphology [23,24]. It has been repeatedly identified in earlier selections scans as having undergone a sweep in East Asians [25,26].

Furthermore, 3P-CLR allows us to narrow down on the specific time at which selection occurred in the history of particular populations. For example, ref. [1] performed a scan of the genomes of East Asians using XP-CLR with Africans as the outgroup, and identified a number of genes as being under selection [1]. 3P-CLR confirms this signal in several of these loci when looking specifically at the East Asian branch: *CYP26B1, EMX1, SPR, SFXN5, PPARA, PKDREJ, GTSE1, TRMU, CELSR1, PINX1, XKR6, CD226, ACD, PARD6A, GFOD2, RANBP10, TSNAXIP1, CENPT, THAP11, NUTF2, CDH16, RRAD, FAM96B, CES2, CBFB, C16orf70, TRADD, FBXL8, HSF4, NOL3, EXOC3L1, E2F4, ELMO3, LRRC29, FHOD1, SLC9A5, PLEKHG4, LRRC36, ZDHHC1, HSD11B2, ATP6V0D1, AGRP, FAM65A, CTCF* and *RLTPR*. However, when applied to the ancestral Eurasian branch, 3P-CLR finds some genes that were previously found in the XP-CLR analysis of East Asians, but that are not among the top hits in 3P-CLR applied to the East Asian branch: *COMMD3, BMI1, SPAG6* and *ABCC12*. This suggests

selection in these regions occurred earlier, i.e. before the European-East Asian split. Figure 5.A shows a comparison between the 3P-CLR scores for the three branches in the region containing genes *BMI1* (a proto-oncogene [27]) and *SPAG6* (involved in sperm motility [28]). In that figure, the score within each window was standardized using its chromosome-wide mean and standard deviation, to make a fair comparison. One can observe that the signal of Eurasia-specific selection is evidently stronger than the other two signals.

When running 3P-CLR to look for selection specific to Europe, we find that *TYRP1* (Figure 5.B) and *MYO5A*, which play a role in human skin pigmentation [29–32], are among the top hits. Both of these genes have been previously found to be under strong selection in Europe [33], using a statistic called iHS, which measures extended patterns of haplotype homozygosity that are characteristic of selective sweeps. Interestingly, a change in the gene *TYRP1* has also been found to cause a blonde hair phenotype in Melanesians [34]. Another of our top hits is the region containing *SH2B3*, which was identified previously as a candidate for selection in Europe based on both iHS and Fst [35]. This gene contains a nonsynonymous SNP (rs3184504) segregating in Europeans. One of its alleles (the one in the selected haplotype) has been associated with celiac disease and type 1 diabetes [36, 37] but is also protective against bacterial infection [38].

We used Gowinda (v1.12) [39] to find enriched Gene Ontology (GO) categories among the regions in the 99% highest quantile for each branch score, relative to the rest of the genome (P < 0.05, FDR < 0.2). The significantly enriched categories are listed in Table 5. In the East Asian branch, we find categories related to pyruvate metabolism, cholesterol absorption and peroxisome proliferation. In the European branch, we find categories related to cuticle development, antioxidant activity and thyroid hormone generation, among others. In the Eurasian branch, we only find two categories that are related to the regulation of transcription by sequence-specific DNA-binding.

4.3 Selection in ancestral modern humans

We applied 3P-CLR to modern human data combined with recently sequenced archaic human data [10]. We sought to find selective events that occurred in modern humans after their spit from archaic groups. We used the combined Neanderthal and Denisovan high-coverage genomes [9, 10] as the outgroup population, and, for our two test populations, we randomly sampled 100 Eurasian genomes and 100 African genomes (excluding admixed African-Americans) from the 1000 Genomes data (Figure S5.B).

We used previously estimated drift times as fixed parameters [10], and tested for selective events that
occurred more anciently than the split of Africans and Eurasians, but more recently than the split from
Neanderthals. We ran 3P-CLR using 0.25 cM windows as above, but also verified that the density of
scores was robust to the choice of window size and spacing (Figure S6). As before, we selected the top
99.9% windows and merged them together if they were contiguous. Table 6 and Figure S7 show the
top hits. To find putative candidates for the beneficial variants in each region, we queried the catalogs
of modern human-specific high-frequency or fixed derived changes that are ancestral in the Neanderthal
and/or the Denisova genomes [10, 40].

We observe several genes that have been identified in previous scans that looked for selection in modern
humans after their split from archaic groups [8, 10]: *SIPA1L1, ANAPC10, ABCE1, RASA1, CCNH,*
KCNJ3, HBP1, COG5, GPR22, DUS4L, BCAP29, CADPS2, RNF133, RNF148, FAM172A, POU5F2,
FGF7, RABGAP1, GPR21, STRBP, SMURF1, GABRA2, ALMS1, PVRL3, EHBP1, VPS54, OTX1,
UGP2, HCN1, GTDC1, ZEB2, OIT3, USP54 and *MYOZ1.* One of our strongest candidate genes among
these is *ANAPC10.* This gene is a core subunit of the cyclosome, is involved in progression through the
cell cycle [41], and may play a role in oocyte maturation and human T-lymphotropic virus infection
(KEGG pathway [42]). *ANAPC10* is noteworthy because it was found to be significantly differentially
expressed in humans compared to other great apes and macaques: it is up-regulated in the testes [43].
The gene also contains two intronic changes that are fixed derived in modern humans, ancestral in both
Neanderthals and Denisovans and that have evidence for being highly disruptive, based on a composite
score that combines conservation and regulatory data (PHRED-scaled C-scores > 11 [10, 44]). The
changes, however, appear not to lie in any obvious regulatory region [45, 46].

We also find *ADSL* among the list of candidates. This gene is known to contain a nonsynonymous
change that is fixed in all present-day humans but homozygous ancestral in the Neanderthal genome,
the Denisova genome and two Neanderthal exomes [40] (Figure 6.A). It was previously identified as ly-
ing in a region with strong support for positive selection in modern humans, using summary statistics
implemented in an ABC method [11]. The gene is interesting because it is one of the members of the
Human Phenotype ontology category "aggression / hyperactivity" which is enriched for nonsynonymous
changes that occurred in the modern human lineage after the split from archaic humans [40, 47]. *ADSL*
codes for adenylosuccinase, an enzyme involved in purine metabolism [48]. A deficiency of adenylosucci-
nase can lead to apraxia, speech deficits, delays in development and abnormal behavioral features, like

hyperactivity and excessive laughter [49]. The nonsynonymous mutation (A429V) is in the C-terminal domain of the protein (Figure 6.B) and lies in a highly conserved position (primate PhastCons = 0.953; GERP score = 5.67 [44, 50, 51]). The ancestral amino acid is conserved across the tetrapod phylogeny, and the mutation is only three residues away from the most common causative SNP for severe adeny-losuccinase deficiency [52–56]. The change has the highest probability of being disruptive to protein function, out of all the nonsynonymous modern-human-specific changes that lie in the top-scoring regions (C-score = 17.69). While *ADSL* is an interesting candidate and lies in the center of the inferred selected region (Figure 6.A), there are other genes in the region too, including *TNRC6B* and *MKL1*. *TNRC6B* may be involved in miRNA-guided gene silencing [57], while *MKL1* may play a role in smooth muscle differentiation [58], and has been associated with acute megakaryocytic leukemia [59].

RASA1 was also a top hit in a previous scan for selection [8], and was additionally inferred to have a high Bayes factor in favor of selection in ref. [11]. The gene codes for a protein involved in the control of cellular differentiation [60], and has a modern human-specific fixed nonsynonymous change (G70E). Human diseases associated with *RASA1* include basal cell carcinoma [61] and arteriovenous malformation [62, 63].

The $GABA_A$ gene cluster in chromosome 4p12 is also among the top regions. The genes within the putatively selected region code for three of the subunits of the $GABA_A$ receptor (*GABRA2, GABRA4*), which codes for a ligand-gated ion channel that plays a key role in synaptic inhibtion in the central nervous system (see review by ref. [64]). *GABRA2* is significantly associated with the risk of alcohol dependence in humans [65], perception of pain [66] and asthma [67]. In turn, GABRA4 is associated with autism risk [68, 69].

Two other candidate genes that may be involved in brain development are *FOXG1* and *CADPS2*. *FOXG1* was not identified in any of the previous selection scans, and codes for a protein called forkhead box G1, which plays an important role during brain development. Mutations in this gene have been associated with a slow-down in brain growth during childhood resulting in microcephaly, which in turn causes various intellectual disabilities [70, 71]. *CADPS2* was identified in [8] as a candidate for selection, and has been associated with autism [72]. The gene has been suggested to be specifically important in the evolution of all modern humans, as it was not found to be selected earlier in great apes or later in particular modern human populations [73].

Finally, we find a signal of selection in a region containing the gene *EHBP1* and *OTX1*. This region

was identified in both of the two previous scans for modern human selection [8,10]. *EHBP1* codes for a protein involved in endocytic trafficking [74] and has been associated with prostate cancer [75]. *OTX1* is a homeobox family gene that may play a role in brain development [76]. Interestingly, *EHBP1* contains a single-nucleotide intronic change (chr2:63206488) that is almost fixed in all present-day humans and homozygous ancestral in Neanderthal and Denisova [10]. This change is also predicted to be highly disruptive (C-score = 13.1) and lies in a position that is extremely conserved across primates (PhastCons = 0.942), mammals (PhastCons = 1) and vertebrates (PhastCons = 1). The change is 18 bp away from the nearest splice site and overlaps a VISTA conserved enhancer region (element 1874) [77], which suggests a putative regulatory role for the change.

We again used Gowinda [39] to find enriched GO categories among the regions with high 3P-CLR scores in the Modern Human branch. This time, we used a stricter quantile cutoff (99.5%) to define candidate regions than we did when running the program in the Eurasian tree (99%) because the less strict cutoff yielded a very large number of enriched categories (114), though in both cases the enriched terms were very similar. The significantly enriched categories (P < 0.05, FDR < 0.2) are listed in Table 7. We find several GO terms related to the regulation of the cell cycle, cell proliferation in the bone marrow, lymphocyte chemotaxis and myeloid cell differentiation.

4.4 Modern human-specific high-frequency changes in GWAS catalog

We overlapped the genome-wide association studies (GWAS) database [78,79] with the list of fixed or high-frequency modern human-specific changes that are ancestral in archaic humans [10] and that are located within our top putatively selected regions in modern humans (Table 8). None of the resulting SNPs are completely fixed derived, because GWAS can only yield associations from sites that are segregating. Among these SNPs, the one with the highest probability of being disruptive (rs10003958, C-score = 16.58, Gerp score = 6.07) is located in a highly-conserved regulatory ("strong enhancer") region in the *RAB28* gene [45,46] (Primate PhastCons = 0.951), and is significantly associated with obesity [80] (Figure 7.A). Interestingly, the region containing *RAB28* is inferred to have been under positive selection in both the modern human and the Eurasian ancestral branches (Tables 4, 6). In line with this evidence, the derived allele of rs10003958 is absent in archaic humans, at very high frequencies in Eurasians (> 94%), and only at moderately high frequencies in Africans (74%) (Figure 7.B).

We also find a highly disruptive SNP (rs10171434, C-score = 8.358) associated with urinary metabo-

lites [81] and suicidal behavior in patients with mood disorders [82]. The SNP is located in an enhancer regulatory freature [45, 46] located between genes *PELI1* and *VPS54*, in the same putatively selected region as genes *EHBP1* and *OTX1* (see above). Finally, there is a highly disruptive SNP (rs731108, C-score = 10.31) that is associated with renal cell carcinoma [83]. This SNP is also located in an enhancer regulatory feature [45, 46], in an intron of *ZEB2*. In this last case, though, only the Neanderthal genome has the ancestral state, while the Denisova genome carries the modern human variant.

5 Discussion

We have developed a new method called 3P-CLR, which allows us to detect positive selection along the genome. The method is based on an earlier test (XP-CLR [1]) that uses linked allele frequency differences between two populations to detect population-specific selection. However, 3P-CLR can allow us to distinguish between selective events that occurred before and after the split of two populations. Our method also has some similiarities to an earlier method developed by [84], which used an F_{st}-like score to detect selection ancestral to two populations. In that case, though, the authors used summary statistics and did not explicitly model the process leading to allele frequency differentiation.

We used our method to confirm previously found candidate genes in particular human populations, like *EDAR, TYRP1, SH2B3* and *MYO5A*, and find some novel candidates too (Tables 2, 3, 4). Additionally, we can infer that certain genes, which were previously known to have been under selection in East Asians (like *SPAG6*), are more likely to have undergone a sweep in the population ancestral to both Europeans and East Asians than in East Asians only. We find that genes involved in pyruvate and cholesterol metabolism are particularly enriched among the East Asian candidate regions, which suggests these biological functions may have been subject to positive selection in recent times.

We also used 3P-CLR to detect selective events that occurred in the ancestors of modern humans, after their split from Neanderthals and Denisovans (Table 6). These events could perhaps have led to the spread of phenotypes that set modern humans apart from other hominin groups. We find several intersting candidates, like *SIPA1L1, ADSL, RASA1, OTX1, EHBP1, FOXG1, RAB28* and *ANAPC10*, some of which were previously detected using other types of methods [8, 10, 11]. We also find an enrichment for GO categories related to cell cycle regulation and lymphocyte chemotaxis among the candidate regions, suggesting that these biological processes might have been affected by positive selection after the split

394 from archaic humans.

395 An advantage of differentiation-based tests like XP-CLR and 3P-CLR is that, unlike other patterns **396** detected by tests of neutrality (like extended haplotype homozygostiy, [85]) that are exclusive to hard **397** sweeps, the patterns that both XP-CLR and 3P-CLR are tailored to find are based on regional allele **398** frequency differences between populations. These patterns can also be produced by soft sweeps from **399** standing variation or by partial sweeps [1], and there is some evidence that the latter phenomena may **400** have been more important than classic sweeps during human evolutionary history [86].

401 Another advantage of both XP-CLR and 3P-CLR is that they do not rely on an arbitrary division **402** of genomic space. Unlike other methods which require the partition of the genome into small windows **403** of fixed size, our composite likelihood ratios can theoretically be computed over windows that are as big **404** as each chromosome, while only switching the central candidate site at each window. This is because **405** the likelihood ratios use the genetic distance to the central SNP as input. SNPs that are very far away **406** from the central SNP will not contribute much to the likelihood function of both the neutral and the **407** selection models, while those that are close to it will. While we heuristically limit the window size in **408** our implementation in the interest of speed, this can be arbitrarily adjusted by the user as needed. **409** The use of genetic distance in the likelihood function also allows us to take advantage of the spatial **410** distribution of SNPs as an additional source of information, rather than only relying on patterns of **411** population differentiation restricted to tightly linked SNPs.

412 3P-CLR also has an advantage over HMM-based selection methods, like the one implemented in **413** ref. [10]. The likelihood ratio scores obtained from 3P-CLR can provide an idea of how credible a **414** selection model is for a particular region, relative to the rest of the genome. The HMM-based method **415** previously used to scan for selection in modern humans [10] can only rank putatively selected regions by **416** genetic distance, but cannot output a statistical measure that may indicate how likely each region is to **417** have been selected in ancient times. In contrast, 3P-CLR provides a composite likelihood ratio score, **418** which allows for a statistically rigorous way to compare the neutral model and a specific selection model **419** (for example, recent or ancient selection).

420 The score also gives an idea of how much fainter the signal of ancient selection in modern humans **421** is, relative to recent selection specific to a particular present-day population. For example, the outliers **422** from Figure 4 have much higher scores (relative to the rest of the genome) than the outliers from Figure **423** S7. This may be due to both the difference in time scales in the two sets of tests and to the uncertainty

that comes from estimating outgroup allele frequencies using only two archaic genomes. This pattern can

also be observed in Figure S8, where the densities of the scores looking for patterns of ancient selection

(3P-CLR Modern Human and 3P-CLR Eurasia) have much shorter tails than the densities of scores

looking for patterns of recent selection (3P-CLR Europe and 3P-CLR East Asia). Simulations show that

3P-CLR(Int) score distributions are naturally shorter than 3P-CLR(A) scores (Figure S9), which could

explain the short tail of the 3P-CLR Eurasia distribution. Additionally, the even shorter tail in the

distribution of 3P-CLR Modern Human scores may be a consequence of the fact that the split times of

the demographic history in that case are older than the split times in the Eurasian tree, as simulations

show that ancient split times tend to further shorten the tail of the 3P-CLR score distribution (Figure

S9).

A limitation of composite likelihood ratio tests is that the composite likelihood calculated for each

model under comparison is obtained from a product of individual likelihoods at each site, and so it

underestimates the correlation that exists between SNPs due to linkage effects [1, 16, 17, 87]. One way

to mitigate this problem is by using corrective weights based on linkage disequilibrium (LD) statistics

calculated on the outgroup population [1]. Our implementation of 3P-CLR allows the user to incorporate

such weights, if appropriate LD statistics are available from the outgroup. However, in cases where these

are unreliable, it may not be possible to fully correct for this (for example, when only a few unphased

genomes are available, as in the case of the Neanderthal and Denisova genomes).

While 3P-CLR relies on integrating over the possible allele frequencies in the ancestors of populations

a and b (formula 10), one could envision using ancient DNA to avoid this step. Thus, if enough genomes

could be sampled from that ancestral population that existed in the past, one could use the sample

frequency in the ancient set of genomes as a proxy for the ancestral population frequency. This may soon

be possible, as several early modern human genomes have already been sequenced in recent years [88–90].

Though we have limited ourselves to a three-population model in this manuscript, it should be straight-

forward to expand our model to a larger number of populations, albeit with additional costs in terms

of speed and memory. Our method relies on a similar framework to the demographic inference method

implemented in TreeMix [91], which can estimate complex population trees that include migration events,

using genome-wide data. With a more complex modeling framework, it may be possible to estimate the

time and strength of selective events with better resolution, and to incorporate additional demographic

forces, like continuous migration between populations or pulses of admixture.

Acknowledgments

We thank Montgomery Slatkin, Rasmus Nielsen, Joshua Schraiber, Nicolas Duforet-Frebourg, Emilia Huerta-Sánchez, Hua Chen, Nick Patterson, David Reich, Joachim Hermisson, Graham Coop and members of the Slatkin and Nielsen labs for helpful advice and discussions. This work was supported by NIH grant R01-GM40282 to Montgomery Slatkin.

References

1. Chen H, Patterson N, Reich D (2010) Population differentiation as a test for selective sweeps. Genome research 20: 393–402.

2. Smith JM, Haigh J (1974) The hitch-hiking effect of a favourable gene. Genetical research 23: 23–35.

3. Lewontin R, Krakauer J (1973) Distribution of gene frequency as a test of the theory of the selective neutrality of polymorphisms. Genetics 74: 175–195.

4. Akey JM, Zhang G, Zhang K, Jin L, Shriver MD (2002) Interrogating a high-density snp map for signatures of natural selection. Genome research 12: 1805–1814.

5. Weir BS, Cardon LR, Anderson AD, Nielsen DM, Hill WG (2005) Measures of human population structure show heterogeneity among genomic regions. Genome research 15: 1468–1476.

6. Oleksyk TK, Zhao K, Francisco M, Gilbert DA, O'Brien SJ, et al. (2008) Identifying selected regions from heterozygosity and divergence using a light-coverage genomic dataset from two human populations. PLoS One 3: e1712.

7. Yi X, Liang Y, Huerta-Sanchez E, Jin X, Cuo ZXP, et al. (2010) Sequencing of 50 human exomes reveals adaptation to high altitude. Science 329: 75–78.

8. Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, et al. (2010) A draft sequence of the neandertal genome. science 328: 710–722.

9. Meyer M, Kircher M, Gansauge MT, Li H, Racimo F, et al. (2012) A high-coverage genome sequence from an archaic denisovan individual. Science 338: 222–226.

10. Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, et al. (2014) The complete genome
sequence of a neanderthal from the altai mountains. Nature 505: 43–49.

11. Racimo F, Kuhlwilm M, Slatkin M (2014) A test for ancient selective sweeps and an application
to candidate sites in modern humans. Molecular biology and evolution 31: 3344–3358.

12. Consortium GP, et al. (2012) An integrated map of genetic variation from 1,092 human genomes.
Nature 491: 56–65.

13. Nicholson G, Smith AV, Jónsson F, Gústafsson Ó, Stefánsson K, et al. (2002) Assessing popula-
tion differentiation and isolation from single-nucleotide polymorphism data. Journal of the Royal
Statistical Society: Series B (Statistical Methodology) 64: 695–715.

14. Durrett R, Schweinsberg J (2004) Approximating selective sweeps. Theoretical population biology
66: 129–138.

15. Fay JC, Wu CI (2000) Hitchhiking under positive darwinian selection. Genetics 155: 1405–1413.

16. Lindsay BG (1988) Composite likelihood methods. Contemporary Mathematics 80: 221–39.

17. Varin C, Reid N, Firth D (2011) An overview of composite likelihood methods. Statistica Sinica
21: 5–42.

18. Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, et al. (2012) Ancient admixture in human
history. Genetics 192: 1065–1093.

19. Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD (2009) Inferring the joint demo-
graphic history of multiple populations from multidimensional snp frequency data. PLoS genetics
5: e1000695.

20. Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M (2013) Robust demographic infer-
ence from genomic and snp data. PLoS genetics 9: e1003905.

21. Messer PW (2013) Slim: simulating evolution with selection and linkage. Genetics 194: 1037–1039.

22. Hinch AG, Tandon A, Patterson N, Song Y, Rohland N, et al. (2011) The landscape of recombi-
nation in african americans. Nature 476: 170–175.

23. Fujimoto A, Kimura R, Ohashi J, Omi K, Yuliwulandari R, et al. (2008) A scan for genetic determinants of human hair morphology: Edar is associated with asian hair thickness. Human Molecular Genetics 17: 835–843.

24. Kimura R, Yamaguchi T, Takeda M, Kondo O, Toma T, et al. (2009) A common variation in edar is a genetic determinant of shovel-shaped incisors. The American Journal of Human Genetics 85: 528–535.

25. Sabeti PC, Varilly P, Fry B, Lohmueller J, Hostetter E, et al. (2007) Genome-wide detection and characterization of positive selection in human populations. Nature 449: 913–918.

26. Grossman SR, Shylakhter I, Karlsson EK, Byrne EH, Morales S, et al. (2010) A composite of multiple signals distinguishes causal variants in regions of positive selection. Science 327: 883–886.

27. Siddique HR, Saleem M (2012) Role of bmi1, a stem cell factor, in cancer recurrence and chemoresistance: preclinical and clinical evidences. Stem Cells 30: 372–378.

28. Sapiro R, Kostetskii I, Olds-Clarke P, Gerton GL, Radice GL, et al. (2002) Male infertility, impaired sperm motility, and hydrocephalus in mice deficient in sperm-associated antigen 6. Molecular and cellular biology 22: 6298–6305.

29. Pastural E, Ersoy F, Yalman N, Wulffraat N, Grillo E, et al. (2000) Two genes are responsible for griscelli syndrome at the same 15q21 locus. Genomics 63: 299–306.

30. Fukuda M, Kuroda T, Mikoshiba K (2002) Slac2-a/melanophilin, the missing link between rab27 and myosin va: implications of a tripartite protein complex for melanosome transport. The Journal of biological chemistry 277: 12432.

31. Halaban R, Moellmann G (1990) Murine and human b locus pigmentation genes encode a glycoprotein (gp75) with catalase activity. Proceedings of the National Academy of Sciences 87: 4809–4813.

32. Sulem P, Gudbjartsson DF, Stacey SN, Helgason A, Rafnar T, et al. (2008) Two newly identified genetic determinants of pigmentation in europeans. Nature genetics 40: 835–837.

33. Voight BF, Kudaravalli S, Wen X, Pritchard JK (2006) A map of recent positive selection in the
human genome. PLoS biology 4: e72.

34. Kenny EE, Timpson NJ, Sikora M, Yee MC, Moreno-Estrada A, et al. (2012) Melanesian blond
hair is caused by an amino acid change in tyrp1. Science 336: 554–554.

35. Pickrell JK, Coop G, Novembre J, Kudaravalli S, Li JZ, et al. (2009) Signals of recent positive
selection in a worldwide sample of human populations. Genome research 19: 826–837.

36. Todd JA, Walker NM, Cooper JD, Smyth DJ, Downes K, et al. (2007) Robust associations of
four new chromosome regions from genome-wide analyses of type 1 diabetes. Nature genetics 39:
857–864.

37. Hunt KA, Zhernakova A, Turner G, Heap GA, Franke L, et al. (2008) Newly identified genetic risk
variants for celiac disease related to the immune response. Nature genetics 40: 395–402.

38. Zhernakova A, Elbers CC, Ferwerda B, Romanos J, Trynka G, et al. (2010) Evolutionary and
functional analysis of celiac risk loci reveals sh2b3 as a protective factor against bacterial infection.
The American Journal of Human Genetics 86: 970–977.

39. Kofler R, Schlötterer C (2012) Gowinda: unbiased analysis of gene set enrichment for genome-wide
association studies. Bioinformatics 28: 2084–2085.

40. Castellano S, Parra G, Sánchez-Quinto FA, Racimo F, Kuhlwilm M, et al. (2014) Patterns of coding
variation in the complete exomes of three neandertals. Proceedings of the National Academy of
Sciences 111: 6666–6671.

41. Pravtcheva DD, Wise TL (2001) Disruption of apc10/doc1 in three alleles of oligosyndactylism.
Genomics 72: 78–87.

42. Kanehisa M, Goto S (2000) Kegg: kyoto encyclopedia of genes and genomes. Nucleic acids research
28: 27–30.

43. Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, et al. (2011) The evolution of gene
expression levels in mammalian organs. Nature 478: 343–348.

44. Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, et al. (2014) A general framework for estimating the relative pathogenicity of human genetic variants. Nature genetics 46: 310–315.

45. Consortium EP, et al. (2012) An integrated encyclopedia of dna elements in the human genome. Nature 489: 57–74.

46. Rosenbloom KR, Dreszer TR, Long JC, Malladi VS, Sloan CA, et al. (2011) Encode whole-genome data in the ucsc genome browser: update 2012. Nucleic acids research : gkr1012.

47. Robinson PN, Köhler S, Bauer S, Seelow D, Horn D, et al. (2008) The human phenotype ontology: a tool for annotating and analyzing human hereditary disease. The American Journal of Human Genetics 83: 610–615.

48. Van Keuren M, Hart I, Kao FT, Neve R, Bruns G, et al. (1987) A somatic cell hybrid with a single human chromosome 22 corrects the defect in the cho mutant (ade–i) lacking adenylosuccinase activity. Cytogenetic and Genome Research 44: 142–147.

49. Gitiaux C, Ceballos-Picot I, Marie S, Valayannopoulos V, Rio M, et al. (2009) Misleading behavioural phenotype with adenylosuccinate lyase deficiency. European Journal of Human Genetics 17: 133–136.

50. Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, et al. (2005) Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome research 15: 1034–1050.

51. Cooper GM, Goode DL, Ng SB, Sidow A, Bamshad MJ, et al. (2010) Single-nucleotide evolutionary constraint scores highlight disease-causing mutations. Nature methods 7: 250–251.

52. Kmoch S, Hartmannová H, Stibůrková B, Krijt J, Zikánová M, et al. (2000) Human adenylosuccinate lyase (adsl), cloning and characterization of full-length cdna and its isoform, gene structure and molecular basis for adsl deficiency in six patients. Human molecular genetics 9: 1501–1513.

53. Maaswinkel-Mooij P, Laan L, Onkenhout W, Brouwer O, Jaeken J, et al. (1997) Adenylosuccinase deficiency presenting with epilepsy in early infancy. Journal of inherited metabolic disease 20: 606–607.

54. Marie S, Cuppens H, Heuterspreute M, Jaspers M, Tola EZ, et al. (1999) Mutation analysis in adenylosuccinate lyase deficiency: Eight novel mutations in the re-evaluated full adsl coding sequence. Human mutation 13: 197–202.

55. Race V, Marie S, Vincent MF, Van den Berghe G (2000) Clinical, biochemical and molecular genetic correlations in adenylosuccinate lyase deficiency. Human molecular genetics 9: 2159–2165.

56. Edery P, Chabrier S, Ceballos-Picot I, Marie S, Vincent MF, et al. (2003) Intrafamilial variability in the phenotypic expression of adenylosuccinate lyase deficiency: a report on three patients. American Journal of Medical Genetics Part A 120: 185–190.

57. Meister G, Landthaler M, Peters L, Chen PY, Urlaub H, et al. (2005) Identification of novel argonaute-associated proteins. Current biology 15: 2149–2155.

58. Du KL, Chen M, Li J, Lepore JJ, Mericko P, et al. (2004) Megakaryoblastic leukemia factor-1 transduces cytoskeletal signals and induces smooth muscle cell differentiation from undifferentiated embryonic stem cells. Journal of Biological Chemistry 279: 17578–17586.

59. Mercher T, Busson-Le Coniat M, Monni R, Mauchauffé M, Khac FN, et al. (2001) Involvement of a human gene related to the drosophila spen gene in the recurrent t (1; 22) translocation of acute megakaryocytic leukemia. Proceedings of the National Academy of Sciences 98: 5776–5779.

60. Trahey M, Wong G, Halenbeck R, Rubinfeld B, Martin GA, et al. (1988) Molecular cloning of two types of gap complementary dna from human placenta. Science 242: 1697–1700.

61. Friedman E, Gejman PV, Martin GA, McCormick F (1993) Nonsense mutations in the c–terminal sh2 region of the gtpase activating protein (gap) gene in human tumours. Nature genetics 5: 242–247.

62. Eerola I, Boon LM, Mulliken JB, Burrows PE, Dompmartin A, et al. (2003) Capillary malformation–arteriovenous malformation, a new clinical and genetic disorder caused by rasa1 mutations. The American Journal of Human Genetics 73: 1240–1249.

63. Hershkovitz D, Bercovich D, Sprecher E, Lapidot M (2008) Rasa1 mutations may cause hereditary capillary malformations without arteriovenous malformations. British Journal of Dermatology 158: 1035–1040.

64. Whiting PJ, Bonnert TP, McKernan RM, Farrar S, Bourdelles BL, et al. (1999) Molecular and functional diversity of the expanding gaba-a receptor gene family. Annals of the New York Academy of Sciences 868: 645–653.

65. Edenberg HJ, Dick DM, Xuei X, Tian H, Almasy L, et al. (2004) Variations in gabra2, encoding the $\alpha2$ subunit of the gaba a receptor, are associated with alcohol dependence and with brain oscillations. The American Journal of Human Genetics 74: 705–714.

66. Knabl J, Witschi R, Hösl K, Reinold H, Zeilhofer UB, et al. (2008) Reversal of pathological pain through specific spinal gabaa receptor subtypes. Nature 451: 330–334.

67. Xiang YY, Wang S, Liu M, Hirota JA, Li J, et al. (2007) A gabaergic system in airway epithelium is essential for mucus overproduction in asthma. Nature medicine 13: 862–867.

68. Ma D, Whitehead P, Menold M, Martin E, Ashley-Koch A, et al. (2005) Identification of significant association and gene-gene interaction of gaba receptor subunit genes in autism. The American Journal of Human Genetics 77: 377–388.

69. Collins AL, Ma D, Whitehead PL, Martin ER, Wright HH, et al. (2006) Investigation of autism and gaba receptor subunit genes in multiple ethnic groups. Neurogenetics 7: 167–174.

70. Ariani F, Hayek G, Rondinella D, Artuso R, Mencarelli MA, et al. (2008) Foxg1 is responsible for the congenital variant of rett syndrome. The American Journal of Human Genetics 83: 89–93.

71. Mencarelli M, Spanhol-Rosseto A, Artuso R, Rondinella D, De Filippis R, et al. (2010) Novel foxg1 mutations associated with the congenital variant of rett syndrome. Journal of medical genetics 47: 49–53.

72. Sadakata T, Furuichi T (2010) Ca 2+-dependent activator protein for secretion 2 and autistic-like phenotypes. Neuroscience research 67: 197–202.

73. Crisci JL, Wong A, Good JM, Jensen JD (2011) On characterizing adaptive events unique to modern humans. Genome biology and evolution 3: 791–798.

74. Guilherme A, Soriano NA, Furcinitti PS, Czech MP (2004) Role of ehd1 and ehbp1 in perinuclear sorting and insulin-regulated glut4 recycling in 3t3-l1 adipocytes. Journal of Biological Chemistry 279: 40062–40075.

75. Gudmundsson J, Sulem P, Rafnar T, Bergthorsson JT, Manolescu A, et al. (2008) Common sequence variants on 2p15 and xp11. 22 confer susceptibility to prostate cancer. Nature genetics 40: 281–283.

76. Gong S, Zheng C, Doughty ML, Losos K, Didkovsky N, et al. (2003) A gene expression atlas of the central nervous system based on bacterial artificial chromosomes. Nature 425: 917–925.

77. Pennacchio LA, Ahituv N, Moses AM, Prabhakar S, Nobrega MA, et al. (2006) In vivo enhancer analysis of human conserved non-coding sequences. Nature 444: 499–502.

78. Li MJ, Wang P, Liu X, Lim EL, Wang Z, et al. (2011) Gwasdb: a database for human genetic variants identified by genome-wide association studies. Nucleic acids research : gkr1182.

79. Welter D, MacArthur J, Morales J, Burdett T, Hall P, et al. (2014) The nhgri gwas catalog, a curated resource of snp-trait associations. Nucleic acids research 42: D1001–D1006.

80. Paternoster L, Evans DM, Nohr EA, Holst C, Gaborieau V, et al. (2011) Genome-wide population-based association study of extremely overweight young adults–the goya study. PLoS One 6: e24303.

81. Suhre K, Wallaschofski H, Raffler J, Friedrich N, Haring R, et al. (2011) A genome-wide association study of metabolic traits in human urine. Nature genetics 43: 565–569.

82. Perlis RH, Huang J, Purcell S, Fava M, Rush AJ, et al. (2010) Genome-wide association study of suicide attempts in mood disorder patients. Genome 167.

83. Henrion M, Frampton M, Scelo G, Purdue M, Ye Y, et al. (2013) Common variation at 2q22. 3 (zeb2) influences the risk of renal cancer. Human molecular genetics 22: 825–831.

84. Schlebusch CM, Skoglund P, Sjödin P, Gattepaille LM, Hernandez D, et al. (2012) Genomic variation in seven khoe-san groups reveals adaptation and complex african history. Science 338: 374–379.

85. Sabeti PC, Reich DE, Higgins JM, Levine HZ, Richter DJ, et al. (2002) Detecting recent positive selection in the human genome from haplotype structure. Nature 419: 832–837.

86. Hernandez RD, Kelley JL, Elyashiv E, Melton SC, Auton A, et al. (2011) Classic selective sweeps were rare in recent human evolution. science 331: 920–924.

87. Pace L, Salvan A, Sartori N (2011) Adjusting composite likelihood ratio statistics. Statistica Sinica 21: 129.

88. Fu Q, Li H, Moorjani P, Jay F, Slepchenko SM, et al. (2014) Genome sequence of a 45,000-year-old modern human from western siberia. Nature 514: 445–449.

89. Seguin-Orlando A, Korneliussen TS, Sikora M, Malaspinas AS, Manica A, et al. (2014) Genomic structure in europeans dating back at least 36,200 years. Science 346: 1113–1118.

90. Lazaridis I, Patterson N, Mittnik A, Renaud G, Mallick S, et al. (2014) Ancient human genomes suggest three ancestral populations for present-day europeans. Nature 513: 409–413.

91. Pickrell JK, Pritchard JK (2012) Inference of population splits and mixtures from genome-wide allele frequency data. PLoS genetics 8: e1002967.

Tables

Table 1. Description of models tested. All times are in generations. Selection in the "ancestral population" refers to a selective sweep where the beneficial mutation and fixation occurred before the split time of the two most closely related populations. Selection in "daughter population A" refers to a selective sweep that occurred in one of the two most closely related populations (A), after their split from each other.

Model	Population where selection occurred	t_{AB}	t_{ABC}	t_M	s	N_e
A	Ancestral population	500	2,000	1,800	0.1	10,000
B	Ancestral population	1,000	4,000	2,500	0.1	10,000
C	Ancestral population	2,000	4,000	3,500	0.1	10,000
D	Ancestral population	3,000	8,000	5,000	0.1	10,000
E	Ancestral population	2,000	16,000	8,000	0.1	10,000
F	Ancestral population	4,000	16,000	8,000	0.1	10,000
I	Daughter population A	2,000	4,000	1,000	0.1	10,000
J	Daughter population A	3,000	8,000	2,000	0.1	10,000

Table 2. Top hits for 3P-CLR run on the European terminal branch, using Africans as the outgroup. We show the windows in the top 99.9% quantile of scores. Windows were merged together if they were contiguous. Win max = Location of window with maximum score. Win start = left-most end of left-most window for each region. Win end = right-most end of right-most window for each region. All positions were rounded to the nearest 100 bp. Score max = maximum score within region.

chr	Win max	Win start	Win end	Score max	Genes within region
17	19175100	18858600	19445800	213.864	SLC5A10,FAM83G,GRAP,GRAPL,EPN2,B9D1,MAPK7,MFAP4,RNF112,SLC47A1
15	29241200	29210600	29338200	180.648	APBA2
14	66765100	66417200	67923400	176.032	GPHN,FAM71D,MPP5,ATP6V1D,EIF2S1,PLEK2,TMEM229B
10	74736500	74007800	75402200	161.652	DDIT4,DNAJB12,MICU1,MCU,OIT3,PLA2G12B,P4HA1,NUDT13,ECD,FAM149B1, DNAJC9,MRPS16,TTC18,ANXA7,MSS51,PPP3CB,USP54,MYOZ1
1	35623900	35380800	36584500	150.189	DLGAP3,ZMYM6NB,ZMYM6,ZMYM1,SFPQ,ZMYM4,KIAA0319L, NCDN,TFAP2E,PSMB2,C1orf216,CLSPN,AGO4,AGO1,AGO3,TEKT2,ADPRHL2,COL8A2
11	64581000	64217100	64588600	148.952	RASGRP2,PYGM,SF1,MAP4K2,MEN1,SLC22A11,SLC22A12,NRXN2
12	113010000	111691000	113030000	145.64	BRAP,ACAD10,ALDH2,MAPKAPK5,TMEM116,ERP29,NAA25,TRAFD1,RPL6,PTPN11, RPH3A,CUX2,FAM109A,SH2B3,ATXN2
2	74507400	73404200	74970500	137.92	INO80B,WBP1,MOGS,MRPL53,CCDC142,TTC31,LBX2,PCGF1,TLX2,DQX1,AUP1, NOTO,SMYD5,PRADC1,CCT7,HTRA2,LOXL3,DOK1,M1AP,SEMA4F,FBXO41, EGR4,ALMS1,NAT8,TPRKB,DUSP11,C2orf78,STAMBP,ACTG2,DGUOK,TET3, BOLA3,MOB1A,MTHFD2,SLC4A5,DCTN1,WDR54,RTKN
15	45332100	45094600	45436600	137.885	C15orf43,SORD,DUOX2,DUOXA2,DUOXA1,DUOX1
7	98882800	98717700	99369400	135.106	ZSCAN25,CYP3A5,CYP3A7,CYP3A4,SMURF1,KPNA7,ARPC1A,ARPC1B,PDAP1, BUD31,PTCD1,ATP5J2- PTCD1,CPSF4,ATP5J2,ZNF789,ZNF394,ZKSCAN5,FAM200A,ZNF655
15	72654000	72057800	73138200	132.697	THSD4,MYO9A,SENP8,GRAMD2,PKM,PARP6,CELF6,HEXA,TMEM202,ARIH1, GOLGA6B,BBS4,ADPGK
9	91155000	90913100	91201600	125.597	SPIN1,NXNL2
10	83601100	83597800	83761400	120.455	NRG3
5	142116000	142074000	142194000	117.87	FGF1,ARHGAP26
10	31863100	31479100	31908500	113.751	ZEB1
18	66807000	66646700	66883000	108.186	CCDC102B
2	104933000	104749000	105027000	107.684	-
6	76751700	76636000	77261200	106.577	IMPG1
7	81142700	81087600	81298600	104.727	-
4	167411000	167094000	167644000	104.554	-
21	21424100	21378700	21643900	104.405	-
2	216600000	216551000	216628000	103.798	-
17	58512800	58075800	59174400	103.791	HEATR6,CA4,USP32,C17orf64,APPBP2,PPM1D,BCAS3
5	123509000	123370000	123604000	103.386	-
6	150686000	150637000	150738000	103.115	IYD
15	35551700	35444900	35727400	102.676	DPH6
6	121627000	121082000	121788000	102.59	TBC1D32,GJA1
4	60872200	60814500	61356600	98.9547	-
9	108572000	108412000	108755000	98.9047	TAL2,TMEM38B
1	204823000	204680000	204872000	98.6962	NFASC
20	53878400	53876100	54051800	95.6355	-
10	93143500	93060300	93325000	95.5239	HECTD2
1	162116000	162002000	162228000	95.4304	NOS1AP
9	12777200	12488900	12787600	95.1877	TYRP1,LURAP1L
18	7330950	7259810	7374120	93.1786	-
3	188661000	188641000	188840000	91.5491	TPRG1
15	52859500	52581800	52992200	91.5176	MYO5C,MYO5A,ARPP19,FAM214A
15	48211200	48153900	48308500	90.9933	-
12	80298900	80117100	80435100	90.0939	PPP1R12A
11	38229500	37879000	38607000	86.7175	-
5	82679100	82488400	82790300	85.7742	XRCC4,VCAN
6	43624100	43419100	43688500	85.5502	DLK2,TJAP1,LRRC73,POLR1C,YIPF3,XPO5,POLH,GTPBP2,MAD2L1BP,RSPH9, MRPS18A
13	48977600	48726500	49291500	85.2064	ITM2B,RB1,LPAR6,RCBTB2,CYSLTR2
1	53568300	53124700	53633500	84.8166	FAM159A,COA7,ZYG11B,ZYG11A,ECHDC2,SCP2,PODN,SLC1A7
10	54166700	54130800	54335700	84.1889	-
8	15994200	15838700	15997700	83.0166	MSR1
15	94107900	94022200	94185300	82.8563	-
11	129910000	129805000	130073000	82.6407	PRDM10,APLP2,ST14

Table 3. Top hits for 3P-CLR run on the East Asian terminal branch, using Africans as the outgroup. We show the windows in the top 99.9% quantile of scores. Windows were merged together if they were contiguous. Win max = Location of window with maximum score. Win start = left-most end of left-most window for each region. Win end = right-most end of right-most window for each region. All positions were rounded to the nearest 100 bp. Score max = maximum score within region.

chr	Win max	Win start	Win end	Score max	Genes within region
5	117510000	117345000	117716000	270.211	-
3	58238900	58104900	58557500	257.128	FLNB,DNASE1L3,ABHD6,RPP14,PXK,PDHB,KCTD6,ACOX2,FAM107A
10	94874300	94840100	95720400	235.225	MYOF,CEP55,FFAR4,RBP4,PDE6C,FRA10AC1,LGI1,SLC35G1
15	64166100	63723000	64339800	231.645	USP3,FBXL22,HERC1,DAPK2
4	42193900	41823000	42206800	231.205	TMEM33,DCAF4L1,SLC30A9,BEND4
2	72378700	72353700	73177300	221.359	CYP26B1,EXOC6B,SPR,EMX1,SFXN5
1	234347000	234207000	234380000	208.215	SLC35F3
11	25172400	25098500	25276200	206.418	LUZP2
4	158638000	158481000	158740000	190.705	-
17	61536300	60912100	61549600	185.021	TANC2,CYB561
20	24793800	24570100	25037800	177.801	SYNDIG1,CST7,APMAP,ACSS1
4	86504400	86438300	86602900	173.091	ARHGAP24
10	56026900	55868800	56209000	172.266	PCDH15
1	75622900	75277800	76729300	167.716	LHX8,SLC44A5,ACADM,RABGGTB,MSH4,ASB17,ST6GALNAC3
18	5299800	5203000	5314080	163.435	ZBTB14
7	112265000	112125000	112622000	157.219	LSMEM1,TMEM168,C7orf60
8	10836400	10559800	11126200	150.728	RP1L1,SOX7,PINX1,XKR6
1	172931000	172670000	172950000	149.67	-
4	135424000	134792000	135547000	149.303	-
10	53363100	53226200	53440300	147.8	PRKG1
7	1.09E+08	108741000	109226000	146.998	-
13	63542000	63261200	63971200	146.966	-
3	102005000	101902000	102242000	146.338	ZPLD1
6	69974500	69524500	70359500	144.236	BAI3
2	56096500	55929400	56198400	139.435	EFEMP1
2	109534000	108937000	109626000	139.089	LIMS1,RANBP2,CCDC138,EDAR,SULT1C4,GCC2
22	46760700	46594600	46831200	138.989	PPARA,CDPF1,PKDREJ,TTC38,GTSE1,TRMU,CELSR1
3	104826000	104604000	104910000	138.473	-
18	67572500	67533400	67877100	138.033	CD226,RTTN
2	26159900	25853900	26233500	135.742	KIF3C,DTNB
20	31604100	31304800	31614200	134.323	COMMD7,DNMT3B,MAPRE1,SUN5,BPIFB2
2	17456900	17247000	17564600	134.267	-
4	28858500	28537900	28879000	131.871	-
9	107052000	106657000	107058000	131.263	SMC2
12	93322200	92983200	93454700	129.603	C12orf74,PLEKHG7,EEA1
4	80074800	79878800	80250300	129.59	NAA11
5	153541000	153053000	153736000	129.577	GRIA1,FAM114A2,MFAP3,GALNT10
14	77987200	77720700	78086400	126.295	TMEM63C,NGB,POMT2,GSTZ1,TMED8,SAMD15,NOXRED1,VIPAS39,AHSA1,ISM2,SPTLC2
6	8002470	7975450	8112550	125.312	TXNDC5,BLOC1S5,EEF1E1
12	124021000	123925000	124275000	125.199	SNRNP35,RILPL1,TMED2,DDX55,EIF2B1,GTF2H3,TCTN2,ATP6V0A2,DNAH10
10	97039700	96682100	97059000	125.048	CYP2C9,CYP2C8,C10orf129,PDLIM1
16	67607200	66947800	68430200	124.191	ACD,PARD6A,ENKD1,C16orf86,GFOD2,RANBP10,TSNAXIP1,CENPT,THAP11,NUTF2,EDC4,NRN1L,PSKH1,CTRL,PSMB10,LCAT,SLC12A4,DPEP3,DPEP2,DUS2,DDX28,NFATC3,ESRP2,PLA2G15,SLC7A6,SLC7A6OS,PRMT7,SMPD3,CDH16,RRAD,FAM96B,CES2,CES3,CES4A,CBFB,C16orf70,B3GNT9,TRADD,FBXL8,HSF4,NOL3,KIAA0895L,EXOC3L1,E2F4,ELMO3,LRRC29,TMEM208,FHOD1,SLC9A5,PLEKHG4,KCTD19,LRRC36,TPPP3,ZDHHC1,HSD11B2,ATP6V0D1,AGRP,FAM65A,CTCF,RLTPR
12	103350000	103178000	103439000	123.87	PAH,ASCL1
5	111988000	111981000	112344000	123.442	APC,SRP19,REEP5,DCP2
14	69489100	69424000	69782800	122.252	ACTN1,DCAF5,EXD2,GALNT16
3	17197100	17188900	17897600	122.173	TBC1D5

Table 4. Top hits for 3P-CLR run on the Eurasian ancestral branch, using Africans as the outgroup. We show the windows in the top 99.9% quantile of scores. Windows were merged together if they were contiguous. Win max = Location of window with maximum score. Win start = left-most end of left-most window for each region. Win end = right-most end of right-most window for each region. All positions were rounded to the nearest 100 bp. Score max = maximum score within region.

chr	Win max	Win start	Win end	Score max	Genes within region
17	58778100	58075800	59308200	549.661	HEATR6,CA4,USP32,C17orf64,APPBP2,PPM1D,BCAS3
10	22705100	22428900	22798800	500.245	EBLN1,COMMD3,COMMD3-BMI1,BMI1,SPAG6
4	41834200	41452400	42195900	496.603	LIMCH1,PHOX2B,TMEM33,DCAF4L1,SLC30A9,BEND4
18	67572500	67533400	67881500	493.711	CD226,RTTN
17	62870600	62655400	63061700	484.401	SMURF2,LRRC37A3,GNA13
7	99227800	98717700	99374100	446.801	ZSCAN25,CYP3A5,CYP3A7,CYP3A4,SMURF1,KPNA7,ARPC1A,ARPC1B,PDAP1, BUD31,PTCD1,ATP5J2- PTCD1,CPSF4,ATP5J2,ZNF789,ZNF394,ZKSCAN5,FAM200A,ZNF655
2	73545700	72370500	74117400	444.484	CYP26B1,EXOC6B,SPR,EMX1,SFXN5,RAB11FIP5,NOTO,SMYD5,PRADC1,CCT7, FBXO41,EGR4,ALMS1,NAT8,TPRKB,DUSP11,C2orf78,STAMBP
10	93143500	93049100	93325000	440.141	HECTD2
1	230018000	229912000	230132000	437.859	-
2	22420000	22187700	22469200	436.745	-
17	61536300	60888200	61549600	435.653	TANC2,CYB561
20	54054100	53877600	54056600	429.932	-
9	90946300	90908100	91200000	426.111	SPIN1,NXNL2
8	30625000	30515900	30891400	416.16	GSR,PPP2CB,TEX15,PURG,WRN
11	39699100	39604100	39937900	413.515	-
6	10644100	10578900	10784300	408.918	GCNT2,C6orf52,PAK1IP1,TMEM14C,TMEM14B,SYCP2L,MAK
3	188751000	188646000	188859000	400.376	TPRG1
1	64483200	64340800	64538400	398.325	ROR1
10	31863100	31479100	31908500	397.626	ZEB1
4	177625000	177608000	177889000	395.393	VEGFC
14	57824300	57607700	58048400	392.93	EXOC5,AP5M1,NAA30,C14orf105
10	66018600	65795200	66311900	389.492	-
11	19609000	19591300	19731200	387.843	NAV2
13	49136300	48726500	49293400	386.345	ITM2B,RB1,LPAR6,RCBTB2,CYSLTR2
4	13424000	13143500	13535500	379.283	RAB28
8	52617300	52362200	52930500	377.6	PXDNL,PCMTD1
6	3149410	3073260	3204820	376.252	RIPK1,BPHL,TUBB2A
1	25592800	25517600	25869600	372.377	SYF2,C1orf63,RHD,TMEM50A,RHCE,TMEM57
3	97346000	96453200	97364600	371.806	EPHA6
6	105946000	105800000	105954000	371.534	PREP
15	45332100	45094600	45436600	369.82	C15orf43,SORD,DUOX2,DUOXA2,DUOXA1,DUOX1
12	111447000	111331000	111655000	368.231	CCDC63,MYL2,CUX2
6	131952000	131736000	132060000	366.757	ARG1,MED23,ENPP3,OR2A4,CTAGE9
5	11741300	11640500	11850200	365.997	CTNND2
1	116880000	116723000	117028000	365.955	ATP1A1
8	43401200	42499600	49036000	364.188	CHRNB3,CHRNA6,THAP1,RNF170,HOOK3,FNTA,POMK,HGSNAT,SPIDR,CEBPD, MCM4,UBE2V2
7	30270500	30178800	30471600	360.954	MTURN,ZNRF2,NOD1
4	33576600	33301000	33643000	359.425	-

Table 5. Enriched GO categories in the European, East Asian and Eurasian branches. We tested for ontology enrichment among the regions in the 99% quantile of the 3P-CLR scores for each population branch (P < 0.05, FDR < 0.2).

Population Branch	Raw p-value	FDR	GO category
Europe	0.00001	0.009243333	protein localization to membrane
Europe	0.00001	0.009243333	cellular protein localization
Europe	0.00001	0.009243333	cellular macromolecule localization
Europe	0.00003	0.016986	single-organism cellular localization
Europe	0.00003	0.016986	single-organism localization
Europe	0.00005	0.019915556	NAD(P)H oxidase activity
Europe	0.00005	0.019915556	cellular localization
Europe	0.00006	0.019915556	thyroid hormone generation
Europe	0.00006	0.019915556	cuticle development
Europe	0.0001	0.029226923	protein localization
Europe	0.00012	0.029226923	thyroid hormone metabolic process
Europe	0.00013	0.029226923	antioxidant activity
Europe	0.00013	0.029226923	macromolecule localization
Europe	0.00018	0.041946429	cellular response to reactive oxygen species
Europe	0.00022	0.047034444	protein localization to plasma membrane
Europe	0.00024	0.047034444	hydrogen peroxide catabolic process
Europe	0.00026	0.047034444	oxidoreductase activity, acting on peroxide as acceptor
Europe	0.00026	0.047034444	peroxidase activity
Europe	0.00034	0.061767895	cellular component assembly involved in morphogenesis
Europe	0.0004	0.069824	cellular response to hydrogen peroxide
Europe	0.00052	0.085701905	bicarbonate transport
Europe	0.00055	0.087832273	oxidoreductase activity, acting on NAD(P)H, oxygen as acceptor
Europe	0.00073	0.112074348	cell-cell junction
Europe	0.00085	0.127015	recycling endosome membrane
Europe	0.00101	0.144204	MLL5-L complex
Europe	0.00108	0.146987037	single-organism membrane organization
Europe	0.00112	0.146987037	interleukin-2 biosynthetic process
Europe	0.00143	0.171870313	plasma membrane organization
Europe	0.00145	0.171870313	PTW/PP1 phosphatase complex
Europe	0.00148	0.171870313	localization
Europe	0.0015	0.171870313	response to reactive oxygen species
Europe	0.0015	0.171870313	determination of left/right symmetry
Europe	0.00174	0.192166471	cleavage furrow
Europe	0.00178	0.192166471	hydrogen peroxide metabolic process
East Asia	0.00004	0.11411	regulation of peroxisome proliferator activated receptor signaling pathway
East Asia	0.00013	0.151273333	pyruvate dehydrogenase (acetyl-transferring) kinase activity
East Asia	0.00025	0.151273333	negative regulation of intestinal cholesterol absorption
East Asia	0.00025	0.151273333	regulation of intestinal cholesterol absorption
East Asia	0.00025	0.151273333	negative regulation of intestinal phytosterol absorption
East Asia	0.00028	0.151273333	negative regulation of peroxisome proliferator activated receptor signaling pathway
Eurasia	0.00001	0.0272	RNA polymerase II core promoter proximal region sequence-specific DNA binding transcription factor activity involved in negative regulation of transcription
Eurasia	0.00003	0.042205	RNA polymerase II transcription regulatory region sequence-specific DNA binding transcription factor activity involved in negative regulation of transcription

Table 6. Top hits for 3P-CLR run on the ancestral branch to Eurasians and Africans, using archaic humans as the outgroup. We show the windows in the top 99.9% quantile of scores. Windows were merged together if they were contiguous. Win max = Location of window with maximum score. Win start = left-most end of left-most window for each region. Win end = right-most end of right-most window for each region. All positions were rounded to the nearest 100 bp. Score max = maximum score within region.

chr	Win max	Win start	Win end	Score max	Genes within region
21	34916200	34737300	35222100	868.062	IFNGR2,TMEM50B,DNAJC28,GART,SON,DONSON,CRYZL1,ITSN1
17	56595700	56373200	57404800	846.821	BZRAP1,SUPT4H1,RNF43,HSF5,MTMR4,SEPT4,C17orf47,TEX14,RAD51C,PPM1E, TRIM37,SKA2,PRR11,SMG8,GDPD1
12	79919700	79756800	80109400	840.098	SYT1,PAWR
14	71790600	71658900	72283600	835.723	SIPA1L1
12	116589000	116366000	116760000	826.65	MED13L
2	37989300	37917400	38021500	823.495	CDC42EP3
3	36941700	36836900	37517500	819.361	TRANK1,EPM2AIP1,MLH1,LRRFIP2,GOLGA4,C3orf35,ITGA9
14	29635300	29222200	29696100	813.978	FOXG1
5	86911000	86463700	87101400	811.741	RASA1,CCNH
1	213498000	213145000	213561000	810.015	VASH2,ANGEL2,RPS6KC1
2	156468000	155639000	156764000	809.474	KCNJ3
4	146155000	145355000	146222000	806.198	HHIP,ANAPC10,ABCE1,OTUD4
7	121700000	121620000	122369000	794.997	PTPRZ1,AASS,FEZF1,CADPS2,RNF133,RNF148
17	61237300	60906000	61544500	793.984	TANC2,CYB561
7	107229000	106639000	107308000	784.544	PRKAR2B,HBP1,COG5,GPR22,DUS4L,BCAP29,SLC26A4
3	50576000	50177100	51929300	778.276	SEMA3F,GNAT1,GNAI2,LSMEM2,IFRD2,HYAL3,NAT6,HYAL1,HYAL2,TUSC2, RASSF1,ZMYND10,NPRL2,CYB561D2,TMEM115,CACNA2D2,C3orf18,HEMK1, CISH,MAPKAPK3,DOCK3,MANF,RBM15B,RAD54L2,TEX264,GRM2,IQCF6, IQCF3,IQCF2,IQCF5,IQCF1
5	93214400	92677500	93645500	772.879	NR2F1,FAM172A,POU5F2,KIAA0825
7	99167000	98722100	99375300	764.946	ZSCAN25,CYP3A5,CYP3A7,CYP3A4,SMURF1,KPNA7,ARPC1A,ARPC1B,PDAP1, BUD31,PTCD1,ATP5J2- PTCD1,CPSF4,ATP5J2,ZNF789,ZNF394,ZKSCAN5,FAM200A,ZNF655
21	36769600	36689700	36842100	764.846	RUNX1
13	96782600	96180700	97420500	759.296	CLDN10,DZIP1,DNAJC3,UGGT2,HS6ST3
1	176411000	175890000	176437000	758.886	RFWD2,PAPPA2
15	49269100	49247500	50036600	758.119	SECISBP2L,COPS2,GALK2,FAM227B,FGF7,DTWD1,SHC4
22	40521100	40350900	41223300	757.637	GRAP2,FAM83F,TNRC6B,ADSL,SGSM3,MKL1,MCHR1,SLC25A17,ST13
9	125562000	125505000	126059000	755.506	ZBTB26,RABGAP1,GPR21,STRBP,OR1L6,OR5C1,PDCL,OR1K1,RC3H2,ZBTB6
5	89579400	89413800	89654700	755.236	-
19	19313800	19100800	19788800	755.007	SUGP2,ARMC6,SLC25A42,TMEM161A,MEF2BNB- MEF2B,MEF2B,MEF2BNB,RFXANK,NR2C2AP,NCAN,HAPLN4,TM6SF2,SUGP1, MAU2,GATAD2A,TSSK6,NDUFA13,YJEFN3,CILP2,PBX4,LPAR2,GMIP,ATP13A1,ZNF101
2	63889800	62767900	64394800	753.877	EHBP1,OTX1,WDPCP,MDH1,UGP2,VPS54,PELI1
2	73506800	73482800	74054300	749.705	FBXO41,EGR4,ALMS1,NAT8,TPRKB,DUSP11,C2orf78
1	66833100	66772600	66952600	745.941	PDE4B
4	46634200	46361400	46994100	743.893	GABRA2,COX7B2,GABRA4
3	110709000	110513000	110932000	739.819	PVRL3
5	50148300	44575900	50411900	737.793	MRPS30,HCN1,EMB,PARP8
2	145109000	144689000	145219000	736.476	GTDC1,ZEB2
13	44935700	44887400	45237200	736.039	SERP2,TSC22D1
11	65212900	65129200	65379700	734.241	KCNK7,MAP3K11,SLC25A45,FRMD8,SCYL1,LTBP3,SSSCA1,FAM89B,EHBP1L1
8	79211300	78656900	79554100	734.157	PKIA
4	22923300	22826000	23196800	733.814	-
14	76101700	76054700	76450300	733.155	FLVCR2,TTLL5,C14orf1,IFT43,TGFB3
4	13346700	13142500	13533100	732.271	RAB28
12	97334600	96828100	97424100	729.578	NEDD1
5	135849000	135588000	136038000	724.441	TRPC7
1	21083000	21012100	21629100	724.414	KIF17,SH2D5,HP1BP3,EIF4G3,ECE1
10	74730800	74007800	75399900	723.785	DDIT4,DNAJB12,MICU1,MCU,OIT3,PLA2G12B,P4HA1,NUDT13,ECD,FAM149B1, DNAJC9,MRPS16,TTC18,ANXA7,MSS51,PPP3CB,USP54,MYOZ1
2	201156000	200636000	201355000	720.454	C2orf69,TYW5,C2orf47,SPATS2L,KCTD18
3	147223000	146941000	147360000	719.916	ZIC4,ZIC1
5	127571000	127254000	127734000	712.643	SLC12A2,FBN2
9	23839000	23717700	24073800	711.977	ELAVL2

Table 7. Enriched GO categories in the Modern Human branch. We tested for ontology enrichment among the regions in the 99.5% quantile of the 3P-CLR scores corresponding to the Modern Human branch after the split from archaic humans (P < 0.05, FDR < 0.2).

Population Branch	Raw p-value	FDR	GO category
Modern Human	0.00006	0.100843	B cell chemotaxis
Modern Human	0.00011	0.100843	histone H4 acetylation
Modern Human	0.00012	0.100843	transcription, DNA-templated
Modern Human	0.00015	0.100843	RNA biosynthetic process
Modern Human	0.00025	0.100843	heterocycle biosynthetic process
Modern Human	0.00028	0.100843	aromatic compound biosynthetic process
Modern Human	0.00028	0.100843	regulation of cell proliferation in bone marrow
Modern Human	0.00028	0.100843	positive regulation of cell proliferation in bone marrow
Modern Human	0.0003	0.100843	cellular nitrogen compound biosynthetic process
Modern Human	0.00031	0.100843	nucleobase-containing compound biosynthetic process
Modern Human	0.00049	0.138157692	organic cyclic compound biosynthetic process
Modern Human	0.00052	0.138157692	lymphocyte chemotaxis
Modern Human	0.00055	0.138157692	cellular macromolecule biosynthetic process
Modern Human	0.00063	0.144245625	RNA metabolic process
Modern Human	0.00069	0.144245625	regulation of cell cycle
Modern Human	0.0007	0.144245625	activation of Rap GTPase activity
Modern Human	0.00078	0.146167222	negative regulation of cell cycle process
Modern Human	0.0008	0.146167222	regulation of organelle organization
Modern Human	0.00099	0.149766452	germinal center formation
Modern Human	0.00103	0.149766452	regulation of cytoskeleton organization
Modern Human	0.00106	0.149766452	mitotic spindle assembly checkpoint
Modern Human	0.0011	0.149766452	nuclear division
Modern Human	0.00114	0.149766452	negative regulation of mitotic metaphase/anaphase transition
Modern Human	0.00114	0.149766452	negative regulation of metaphase/anaphase transition of cell cycle
Modern Human	0.00115	0.149766452	leucine zipper domain binding
Modern Human	0.00124	0.149766452	mitotic spindle checkpoint
Modern Human	0.00126	0.149766452	spindle assembly checkpoint
Modern Human	0.00135	0.149766452	cell cycle process
Modern Human	0.00139	0.149766452	negative regulation of molecular function
Modern Human	0.0014	0.149766452	anaphase-promoting complex
Modern Human	0.00142	0.149766452	mitosis
Modern Human	0.00151	0.1552175	spindle checkpoint
Modern Human	0.00162	0.158729118	lymphocyte migration
Modern Human	0.00162	0.158729118	intra-Golgi vesicle-mediated transport
Modern Human	0.00181	0.167352222	establishment or maintenance of monopolar cell polarity
Modern Human	0.00181	0.167352222	establishment of monopolar cell polarity
Modern Human	0.00192	0.171855135	mitotic anaphase
Modern Human	0.00208	0.177203721	mitotic cell cycle
Modern Human	0.00216	0.177203721	negative regulation of cell cycle
Modern Human	0.00222	0.177203721	establishment of protein localization to Golgi
Modern Human	0.00222	0.177203721	protein targeting to Golgi
Modern Human	0.00231	0.177203721	anaphase
Modern Human	0.00234	0.177203721	myeloid cell differentiation
Modern Human	0.00255	0.179768044	negative regulation of mitosis
Modern Human	0.00257	0.179768044	histone H4-K5 acetylation
Modern Human	0.00257	0.179768044	histone H4-K8 acetylation
Modern Human	0.00279	0.188911458	organelle fission
Modern Human	0.00284	0.188911458	nucleobase-containing compound metabolic process
Modern Human	0.003	0.1893844	regulation of metaphase/anaphase transition of cell cycle
Modern Human	0.003	0.1893844	regulation of mitotic metaphase/anaphase transition

Table 8. Overlap between GWAS catalog and catalog of modern human-specific high-frequency changes in the top modern human selected regions. Chr = chromosome. Pos = position (hg19). ID = SNP rs ID. Hum = Present-day human major allele. Anc = Human-Chimpanzee ancestor allele. Arch = Archaic human allele states (Altai Neanderthal, Denisova) where H=human-like allele and A=ancestral allele. Freq = present-day human derived frequency. Cons = consequence. C = C-score. PubMed = PubMed article ID for GWAS study.

Chr	Pos	ID	Hum	Anc	Arch	Freq	Gene	Cons	C	GWAS trait	PubMed
2	64279606	rs10171434	C	T	A/A,A/A	0.92	NA	regulatory	8.358	Suicide attempts in bipolar disorder	21041247
2	64279606	rs10171434	C	T	A/A,A/A	0.92	NA	regulatory	8.358	Urinary metabolites	21572414
2	144783214	rs16823411	T	C	A/A,A/A	0.93	GTDC1	intron	4.112	Body mass index	21701565
2	144783214	rs16823411	T	C	A/A,A/A	0.93	GTDC1	intron	4.112	Body mass index	21701565
2	145213638	rs731108	G	C	A/A,H/H	0.92	ZEB2	regulatory	10.31	Renal cell carcinoma	23184150
2	156506516	rs4407211	C	T	A/A,A/A	0.92	NA	intergenic	1.348	Alcohol consumption	23953852
3	51142359	rs4286453	T	C	A/A,A/A	0.91	DOCK3	intron	4.96	Multiple complex diseases	17554300
3	51824167	rs6796373	G	C	A/A,A/A	0.94	NA	intergenic	1.381	Response to taxane treatment (placlitaxel)	23006423
3	147200492	rs9876193	G	A	H/H,A/A	0.95	ZIC1	intron,nc	6.856	Type 2 diabetes	17463246
4	13325741	rs2867467	G	C	A/A,A/A	0.91	NA	intergenic	0.476	Obesity (extreme)	21935397
4	13328373	rs6842438	C	T	A/A,A/A	0.92	NA	intergenic	5.241	Obesity (extreme)	21935397
4	13330095	rs10019897	C	T	A/A,A/A	0.92	NA	upstream	1.472	Multiple complex diseases	17554300
4	13330095	rs10019897	C	T	A/A,A/A	0.92	NA	upstream	1.472	Obesity (extreme)	21935397
4	13333413	rs9996364	A	G	A/A,A/A	0.92	HSP90AB2P	upstream	5.865	Obesity (extreme)	21935397
4	13338465	rs11945340	C	T	A/A,A/A	0.92	HSP90AB2P	non coding exon	12.04	Obesity (extreme)	21935397
4	13340249	rs6839621	T	C	A/A,A/A	0.92	HSP90AB2P	non coding exon	0.074	Obesity (extreme)	21935397
4	13346602	rs11930614	C	T	A/A,A/A	0.92	NA	intergenic	0.587	Obesity (extreme)	21935397
4	13350973	rs10021881	T	C	A/A,A/A	0.92	NA	regulatory	3.032	Obesity (extreme)	21935397
4	13356393	rs16888596	G	A	A/A,A/A	0.94	NA	intergenic	2.344	Obesity (extreme)	21935397
4	13357274	rs11732938	A	G	A/A,A/A	0.94	NA	intergenic	15.45	Obesity (extreme)	21935397
4	13360622	rs11947529	T	A	A/A,A/A	0.93	RAB28	downstream	4.356	Obesity (extreme)	21935397
4	13363958	rs12331157	A	G	A/A,A/A	0.97	RAB28	intron	1.3	Obesity (extreme)	21935397
4	13363974	rs12332023	C	T	A/A,A/A	0.97	RAB28	intron	0.75	Obesity (extreme)	21935397
4	13366481	rs7673680	C	T	A/A,A/A	0.93	RAB28	downstream	4.16	Obesity (extreme)	21935397
4	13370308	rs10003958	T	C	A/A,A/A	0.93	RAB28	regulatory	16.58	Obesity (extreme)	21935397
4	13373583	rs9999851	C	T	A/A,A/A	0.97	RAB28	intron	1.305	Obesity (extreme)	21935397
4	13374462	rs9291610	G	A	A/A,A/A	0.93	RAB28	intron	3.264	Obesity (extreme)	21935397
4	13393897	rs9998914	A	T	A/A,A/A	0.96	RAB28	intron	0.414	Obesity (extreme)	21935397
4	13403855	rs11943295	G	A	A/A,A/A	0.94	RAB28	intron	1.702	Multiple complex diseases	17554300
4	13403855	rs11943295	G	A	A/A,A/A	0.94	RAB28	intron	1.702	Obesity (extreme)	21935397
4	13403998	rs11943330	G	A	A/A,A/A	0.93	RAB28	intron	3.295	Obesity (extreme)	21935397
4	13404130	rs7677336	G	T	A/A,A/A	0.94	RAB28	intron	0.752	Obesity (extreme)	21935397
4	13404717	rs7673732	A	C	A/A,A/A	0.93	RAB28	intron	0.702	Obesity (extreme)	21935397
4	13440031	rs11737264	C	G	A/A,A/A	0.93	RAB28	intron	1.159	Obesity (extreme)	21935397
4	13440271	rs11737360	C	T	A/A,A/A	0.94	RAB28	intron	2.745	Obesity (extreme)	21935397
4	13449532	rs16888654	A	C	A/A,A/A	0.94	RAB28	intron	0.46	Obesity (extreme)	21935397
4	13452022	rs16888661	C	A	A/A,A/A	0.91	RAB28	intron	5.359	Obesity (extreme)	21935397
4	13463991	rs11933841	T	C	A/A,A/A	0.93	RAB28	intron	4.193	Obesity (extreme)	21935397
4	13465710	rs11947665	T	A	A/A,A/A	0.93	RAB28	intron	4.41	Obesity (extreme)	21935397
4	23095293	rs6825402	C	T	A/A,A/A	0.96	NA	intergenic	2.599	Multiple complex diseases	17554300
5	45393261	rs6874279	G	A	A/A,A/A	0.93	HCN1	intron	1.47	Alcohol dependence	20201924
5	45393261	rs6874279	G	A	A/A,A/A	0.93	HCN1	intron	1.47	Alcoholism	pha002891
5	89540468	rs2935504	C	T	A/A,A/A	0.97	RP11-61G23.1	non coding exon	4.52	Multiple complex diseases	17554300
7	106720932	rs12154324	G	A	A/A,A/A	0.93	NA	regulatory	5.411	Multiple complex diseases	17554300
13	44978167	rs9525954	C	A	A/A,A/A	0.95	RP11-269C23.3	intron	2.731	Type 2 diabetes	17463246
13	45034814	rs9533862	G	C	A/A,A/A	0.93	FILIP1LP1	intron	2.026	Suicide attempts in bipolar disorder	21041247
13	45055091	rs17065868	T	C	A/A,A/A	0.92	FILIP1LP1	intron	3.214	Antineutrophil cytoplasmic antibody-associated vasculitis	22808956

 # Figures

Figure 1. Schematic tree of selective sweeps detected by XP-CLR and 3P-CLR. While XP-CLR can only use two populations (an outgroup and a test) to detect selection (panel A), 3P-CLR can detect selection in the ancestral branch of two populations (3P-CLR(Int), panel B) or on the branches specific to each population (3P-CLR(A) and 3P-CLR(B), panels C and D, respectively). The greek letters denote the known drift times for each branch of the population tree.

Figure 2. ROC curves for performance of 3P-CLR(Int), 3P-CLR(A) and two variants of XP-CLR in detecting selective sweeps that occurred before the split of two populations a **and** b**, under different demographic models**. In this case, the outgroup panel from population c contained 100 haploid genomes. The two sister population panels (from a and b) also have 100 haploid genomes each.

Figure 3. 3P-CLR(Int) is tailored to detect selective events that happened before the split t_{ab}, so it is largely insensitive to sweeps that occurred after the split. ROC curves show performance of 3P-CLR(Int) and two variants of XP-CLR for models where selection occurred in population a after its split from b.

Figure 4. 3P-CLR scan of Europeans (upper panel), East Asians (middle panel) and the ancestral population to Europeans and East Asians (lower panel), using Africans as the outgroup in all 3 cases. The red line denotes the 99.9% quantile cutoff.

Figure 5. 3P-CLR scan of Europeans (blue), East Asians (black) and the ancestral Eurasian population (red) reveals regions under selection in different branches of the population tree. To make a fair comparison, all 3P-CLR scores were standardized by substracting the chromosome-wide mean from each window and dividing the resulting score by the chromosome-wide standard deviation. A) The region containing genes *SPAG6* and *BMI1* is a candidate for selection in the ancestral population of Europeans and East Asians. B) The region containing *TYRPP1* is a candidate for selection in the European population. The image was built using the GenomeGraphs package in Bioconductor.

Figure 6. ADSL is a candidate for selection in the modern human lineage, after the split from Neanderthal and Denisova. A) One of the top-scoring regions when running 3P-CLR on the modern human lineage contains genes *TNRC6B*, *ADSL*, *MKL1*, *MCHR1*, *SGSM3* and *GRAP2*. The most disruptive nonsynonymous modern-human-specific change in the entire list of top regions is in an exon of *ADSL* and is fixed derived in all present-day humans but ancestral in archaic humans. It is highly conserved across tetrapods and lies only 3 residues away from the most common mutation leading to severe adenylosuccinase deficiency. B) The *ADSL* gene codes for a tetrameric protein. The mutation is in the C-terminal domain of each tetramer (red arrows), which are near the active sites (light blue arrows). Scores in panel A were standardized using the chromosome-wide mean and standard deviation. Vertebrate alignments were obtained from the UCSC genome browser (Vertebrate Multiz Alignment and Conservation track) and the image was built using the GenomeGraphs package in Bioconductor and Cn3D.

Figure 7. *RAB28* **is a candidate for selection in both the Eurasian and the modern human ancestral lineages.** A) The gene lies in the middle of a 3P-CLR peak for both ancestral populations. The putatively selected region also contains several SNPs that are significantly associated with obesity and that are high-frequency derived in present-day humans ($> 93\%$) but ancestral in archaic humans (red dots). The SNP with the highest C-score among these (rs10003958, pink circle) lies in a highly conserved strong enhancer region adjacent to the last exon of the gene. Color code for ChromHMM segmentation regions in UCSC genome browser: red = promoter, orange = strong enhancer, yellow = weak enhancer, green = weak transcription, blue = insulator. The image was built using the GenomeGraphs package in Bioconductor and the UCSC Genome Browser. B) Derived allele frequencies of SNP rs10003958 in the Denisova and Neanderthal genomes, and in different 1000 Genomes continental populations. AFR = Africans. AMR = Native Americans. SAS = South Asians. EUR = Europeans. EAS = East Asians.

671 Supplementary Figures

Figure S1. ROC curves for performance of 3P-CLR(Int), 3P-CLR(A) and two variants of XP-CLR in detecting selective sweeps that occurred before the split of two populations a **and** b**, under different demographic models**. In this case, the outgroup panel from population c contained 10 haploid genomes. The two sister population panels (from a and b) have 100 haploid genomes each.

Figure S2. Root-mean squared error for the location of sweeps inferred by 3P-CLR(Int), 3P-CLR(A) and two variants of XP-CLR under different demographic scenarios, when the sweeps occurred before the split of populations a **and** b**.** In this case, the outgroup panel from population c contained 100 haploid genomes and the two sister population panels (from a and b) have 100 haploid genomes each.

Figure S3. Root-mean squared error for the location of the sweep inferred by 3P-CLR(Int), 3P-CLR(A) and two variants of XP-CLR under different demographic scenarios, when the sweeps occurred before the split of populations a **and** b**.** the outgroup panel from population c contained 10 haploid genomes and the two sister population panels (from a and b) have 100 haploid genomes each.

Figure S4. Root-mean squared error for the location of the sweep inferred by 3P-CLR(Int), 3P-CLR(A) and two variants of XP-CLR under different demographic scenarios, when the sweeps occurred in the terminal population branch leading to population a, after the split of populations a and b. In this case, the outgroup panel from population c contained 100 haploid genomes and the two sister population panels (from a and b) have 100 haploid genomes each.

Figure S5. A. Three-population tree separating Europeans, East Asians and Africans. B. Three-population tree separating Eurasians, Africans and archaic humans (Neanderthal+Denisova).

Figure S6. Comparison of 3P-CLR on the modern human ancestral branch under different window sizes and central SNP spacing. The red density is the density of standardized scores for 3P-CLR run using 0.25 cM windows, 100 SNPs per window and a spacing of 20 SNPs between each central SNP. The blue dashed density is the density of standardized scores for 3P-CLR run using 1 cM windows, 200 SNPs per window and a spacing of 80 SNPs between each central SNP.

Figure S7. 3P-CLR scan of the ancestral branch to Africans and Eurasians, using the Denisovan and Neanderthal genomes as the outgroup. The red line denotes the 99.9% quantile cutoff.

Figure S8. Genome-wide densities of each of the 3P-CLR scores described in this work.
The distributions of scores testing for recent selection (Europeans and East Asians) have much longer
tails than the distributions of scores testing for more ancient selection (Modern Humans and
Eurasians). All scores were standardized using their genome-wide means and standard deviations.

Figure S9. Distribution of 3P-CLR(Int) and 3P-CLR(A) scores under different demographic histories. We combined all scores obtained from 100 neutral simulations and 100 simulations with a selective sweep under different demographic and selection regimes. We then plotted the densities of the resulting scores. Top panel: Model A; Middle panel: Model C; Bottom panel: Model I. See Table 1 for details about each model.