Cis-regulatory changes associated with a recent mating system shift and floral adaptation in *Capsella*

Kim A. Steige[1], Johan Reimegård[2], Daniel Koenig[3], Douglas G. Scofield[1], Tanja Slotte[1,4,]*

[1]Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden

[2]Science for Life Laboratory, Department of Cell and Molecular Biology, Uppsala University,

Uppsala, Sweden

[3]Max Planck Institute for Developmental Biology, Tübingen, Germany

[4]Department of Ecology, Environment and Plant Sciences, Science for Life Laboratory, Stockholm

University, Stockholm, Sweden

*Corresponding author:

Tanja Slotte

Email: tanja.slotte@su.se (TS)

Abstract

The selfing syndrome constitutes a suite of floral and reproductive trait changes that have evolved repeatedly across many evolutionary lineages in response to the shift to selfing. Convergent evolution of the selfing syndrome suggests that these changes are adaptive, yet our understanding of the detailed molecular genetic basis of the selfing syndrome remains limited. Here, we investigate the role of *cis*-regulatory changes during the recent evolution of the selfing syndrome in *Capsella rubella*, which split from the outcrosser *Capsella grandiflora* less than 200 kya. We assess allele-specific expression (ASE) in leaves and flower buds at a total of 18,452 genes in three interspecific F1 *C. grandiflora* x *C. rubella* hybrids. Using a hierarchical Bayesian approach that accounts for technical variation using genomic reads, we find evidence for extensive *cis*-regulatory changes. On average, 44% of the assayed genes show evidence of ASE, however only 6% show strong allelic expression biases. Flower buds, but not leaves, show an enrichment of *cis*-regulatory changes in genomic regions responsible for floral and reproductive trait divergence between *C. rubella* and *C. grandiflora*. We further detected an excess of heterozygous transposable element (TE) insertions near genes with ASE, and TE insertions targeted by uniquely mapping 24-nt small RNAs were associated with reduced expression of nearby genes. Our results suggest that *cis*-regulatory changes have been important during the recent adaptive floral evolution in *Capsella* and that differences in TE dynamics between selfing and outcrossing species could be important for rapid regulatory divergence in association with mating system shifts.

Introduction

The transition from outcrossing to predominant self-fertilization has occurred repeatedly in flowering plants (Stebbins 1950). In association with this shift, marked changes in floral and reproductive traits have occurred independently in many different lineages (Barrett 2002). In general, selfers tend to show reduced allocation of resources to traits involved in pollinator attraction and reward (e.g. smaller petals, less nectar per flower, less scent), exhibit changes in floral morphology that may improve the efficacy of autonomous self-pollination (e.g. reduced separation between stigma and anthers), and show reduced allocation of resources to male function (reduced ratio of pollen to ovules) (reviewed in Sicard and Lenhard 2011). Together, this combination of floral and reproductive traits is termed "the selfing syndrome" (Ornduff 1969).

Despite the striking pattern of convergent floral evolution in association with the shift to selfing, we currently have a limited understanding of the molecular genetic basis of the selfing syndrome. Quantitative trait loci (QTL) for the selfing syndrome have been identified in a handful of systems (e.g. *Capsella*; Sicard et al 2011; Slotte et al 2012; *Leptosiphon*; Goodwillie et al 2006; *Mimulus*; Fishman et al 2002; Fishman et al 2015; *Oryza*; Grillo et al 2009; *Solanum*; Bernacchi and Tanksley 1997). In domesticated tomatoes, *cis*-regulatory changes at the *Style2.1* gene have been implicated in reduced stigma exsertion (Chen et al 2007), but in most other systems, the molecular basis of the selfing syndrome is not known. A major unresolved question thus concerns the general importance of *cis*-regulatory changes vs. other types of molecular changes for the evolution of the selfing syndrome.

Cis-regulatory changes have long been hypothesized to be important for organismal adaptation (Doebley and Lukens 1998; Carroll 2000; Wray 2007; Carroll 2008; Stern and Orgogozo 2008; but see Hoekstra and Coyne 2007), due to their potentially limited negative pleiotropic effects (Wray 2007). The prospects for identifying *cis*-regulatory changes on a transcriptome-wide scale have greatly improved due to the advent of massively parallel sequencing (Fraser 2011). In particular, methods for assessing allele-specific expression (ASE) that contrast the relative levels of expression of two alleles in an individual allow for transcriptome-scale assessment of *cis*-regulatory changes. ASE studies require the presence of transcribed polymorphisms as well as rigorous bioinformatic approaches, but have benefits over mapping approaches (e.g. eQTL mapping) in terms of cost and resolution, and can identify individual genes with *cis*-regulatory changes (Pastinen 2010).

As part of our broad goal to examine molecular genetic changes associated with the selfing syndrome, we examine the influence of *cis*-regulatory changes on the evolution of the selfing syndrome in *Capsella rubella*. We further test whether silencing of TEs through the RNA-directed methylation pathway is important for global *cis*-regulatory divergence in association with the shift to selfing. The crucifer genus *Capsella* is a promising system for assessing the role of *cis*-regulatory changes in association with plant mating system shifts and adaptation, because of the availability of a sequenced genome of *C. rubella* (Slotte et al 2013) and because it is possible to generate viable

offspring from crosses between *Capsella* species that differ in their mating system (e.g. Slotte et al

2012, Rebernig et al 2015).

In *C. rubella*, the transition to selfing occurred relatively recently (<200 kya), and was

associated with speciation from an outcrossing progenitor similar to present-day *C. grandiflora* (Slotte

et al. 2013, Foxe et al. 2009, Guo et al. 2009, St Onge et al. 2011, Brandvain et al. 2013). Despite the

recent shift to selfing, *C. rubella* already exhibits a derived reduction in petal size and a reduced

pollen-ovule ratio, as well as a reduction of the degree of flower opening (Sicard et al 2011, Slotte et

al. 2012). *C. rubella* therefore exhibits floral and reproductive characters typical of a selfing

syndrome. The selfing syndrome of *C. rubella* is associated with improved efficacy of autonomous

self-pollination (Sicard et al. 2011), and regions with quantitative trait loci (QTL) for floral divergence

between *C. rubella* and *C. grandiflora* exhibit an excess of fixed differences and reduced

polymorphism in *C. rubella* (Slotte et al. 2012). Together, these observations suggest that the rapid

evolution of the selfing syndrome in *C. rubella* was driven by positive selection.

While the molecular genetic basis of the selfing syndrome in *C. rubella* has not been

identified, it has been suggested that *cis*-regulatory changes could be involved, and a previous study

found many flower and pollen development genes to be differentially expressed in flower buds of *C.*

grandiflora and *C. rubella* (Slotte et al. 2013). However, these results could be confounded by

differences in floral organ sizes and pollen number between *C. rubella* and *C. grandiflora,* and Slotte

et al. (2013) did not directly assess *cis*-regulatory changes or investigate possible causes of *cis*-

regulatory divergence. There is reason to believe that *cis*-regulatory changes could be partly caused by

differences in TE abundance between selfers and outcrossers, as TE silencing can affect nearby gene

expression in plants (Hollister and Gaut 2009; Hollister et al. 2011). As *C. rubella* harbors fewer TEs

close to genes than *C. grandiflora* (Ågren et al. 2014), this system offers an opportunity to investigate

the role of TEs for *cis*-regulatory evolution and for the evolution of floral and reproductive traits in

association with the shift to selfing.

In this study we directly assessed *cis*-regulatory divergence by analyzing allele-specific

expression in F1 hybrids of *C. grandiflora* and *C. rubella*, and investigated the role of *cis*-regulatory

changes for the selfing syndrome in *C. rubella*. We conducted deep sequencing of transcriptomes,

small RNAs, as well as genomes of *C. grandiflora* x *C. rubella* hybrids to identify genes with *cis*-

regulatory divergence in flower buds and leaves, and tested whether *cis*-regulatory changes in flowers

were overrepresented in genomic regions responsible for adaptive phenotypic divergence. We further

identifed TEs in *C. rubella* and *C. grandiflora* and tested whether TE insertions targeted by uniquely

mapping 24-nt siRNAs were associated with *cis*-regulatory divergence. Our results provide insight

into the role of *cis*-regulatory changes in association with the shift to selfing in a wild plant system.

Results

Many genes exhibit allele-specific expression in interspecific F1 hybrids

In order to quantify ASE between *C. grandiflora* and *C. rubella*, we generated deep whole

transcriptome RNAseq data from flower buds and leaves of three *C. grandiflora* x *C. rubella* F1

hybrids (total 52.1 vs 41.8 Gbp with Q≥30 for flower buds and leaves, respectively). We included

three technical replicates for one F1 in order to examine the reliability of our expression data. For all

F1s and their *C. rubella* parents, we also generated deep (38-68x) whole genome resequencing data in

order to reconstruct parental haplotypes and account for read mapping biases.

F1 RNAseq reads were mapped with high stringency to reconstructed parental haplotypes

specific for each F1, i.e. reconstructed reference genomes containing whole-genome haplotypes for

both the *C. grandiflora* and the *C. rubella* parent of each F1 (see Material and Methods). We

conducted stringent filtering of genomic regions where SNPs were deemed unreliable for ASE

analyses due to e.g. high repeat content, copy number variation, or a high proportion of heterozygous

genotypes in an inbred *C. rubella* line (for details, see Material and Methods and S1 text); this mainly

resulted in removal of pericentromeric regions (S2 Fig - S5 Fig). After filtering, we identified ~18,200

genes with ~274,000 transcribed heterozygous SNPs that were amenable to ASE analysis in each F1

(Table 1). The mean allelic ratio of genomic read counts at these SNPs was 0.5 (S6 Fig), suggesting

that our bioinformatic procedures efficiently minimized read mapping biases. Furthermore, technical

reliability of our RNAseq data was high, as indicated by a mean Spearman's ρ between replicates of

0.98 (range 0.94-0.99).

We assessed ASE using a Bayesian statistical method with a reduced false positive rate

compared to the standard binomial test (Skelly et al. 2011). The method uses genomic read counts to

model technical variation in ASE and estimates the global proportion of genes with ASE, independent

of specific significance cutoffs, and also yields gene-specific estimates of the ASE ratio and the

posterior probability of ASE. The model also allows for and estimates the degree of variability in ASE

along the gene, through the inclusion of a dispersion parameter.

Based on this method, we estimate that on average, the proportion of assayed genes with ASE

is 44.6% (Table 1; S8 Table). In general, most allelic expression biases were moderate, and only 5.9%

of assayed genes showed ASE ratios greater than 0.8 or less than 0.2 (Figs. 1 and 2). There was little

variation in ASE ratios along genes, as indicated by the distribution of the dispersion parameter

estimates having a mode close to zero and a narrow range (Figs. 1 and 2). This suggests that unequal

expression of differentially spliced transcripts is not a major contributor to regulatory divergence

between *C. rubella* and *C. grandiflora* (Figs. 1 and 2).

For genes with evidence for ASE (hereafter defined as posterior probability of ASE ≥ 0.95),

there was a moderate shift toward higher expression of the *C. rubella* allele (mean ratio *C.*

rubella/total=0.56; Figs. 1 and 2). This shift was present for all F1s, for both leaves and flowers (Figs.

1 and 2). No such shift was apparent for genomic reads, and ratios of genomic read counts for SNPs in

genes with ASE were very close to 0.5 (mean ratio *C. rubella*/total=0.51; Figs 1 and 2). Furthermore,

qPCR with allele-specific probes for five genes validated our ASE results empirically (S9 Table).

Thus, *C. rubella* alleles appear to be on average expressed at a higher level than *C. grandiflora* alleles

in our F1s.

The mean ASE proportion, as well as the absolute number of genes with ASE was greater for

leaves (49%; 6010 genes) than for flower buds (40%; 5216 genes), although this difference was

largely driven by leaf samples from one of our F1s (Table 1). Most instances of ASE were specific to

either leaves or flower buds, and on average, only 15% of genes expressed in both leaves and flower

buds showed consistent ASE in both organs (Fig. 3). Many cases of ASE were also specific to a

particular F1, and across all three F1s, there were 1305 genes that showed consistent ASE in flower

buds, and 1663 in leaves (Fig. 3).

Enrichment of *cis*-regulatory changes in genomic regions responsible for phenotypic divergence

We used permutation tests to check for an excess of genes showing ASE within five previously-

identified narrow (<2 Mb) QTL regions responsible for floral and reproductive trait divergence (Slotte

et al. 2012). These genomic regions harbor major QTL for petal size and flowering time, but also

encompass part of the confidence intervals for QTL for sepal size, stamen length and ovule number, as

QTL for different floral and reproductive traits are highly overlapping (Slotte et al 2012). As the

selfing syndrome has a shared genetic basis in independent *C. rubella* accessions (Sicard et al. 2011,

Slotte et al. 2012), we reasoned that genes with consistent ASE across all F1s would be most likely to

represent candidate *cis*-regulatory changes underlying QTL. Out of the 1305 genes with ASE in flower

buds of all F1s, 85 were found in narrow QTL regions, and this overlap was significantly greater than

expected by chance (permutation test, P=0.03; Fig. 4; see Material and Methods for details). In

contrast, for leaves, there was no significant excess of genes showing ASE in narrow QTL

(permutation test, P=1; Fig. 4). Thus, the association between QTL and ASE in flower buds is unlikely

to be an artifact of locally elevated heterozygosity facilitating both ASE and QTL detection, which

should affect analyses of both leaf and flower samples.

List enrichment analyses reveal floral candidate genes with ASE

We conducted list enrichment analyses to characterize the functions of genes showing ASE relative to

all genes amenable to analysis of ASE (i.e. harboring heterozygous transcribed SNPs and expressed at

detectable levels). There was an enrichment of Gene Ontology (GO) terms involved in defense and

stress responses for genes with ASE in flower buds and in leaves (S10 Table). GO terms related to

hormonal responses, including brassinosteroid and auxin biosynthetic processes, were specifically

enriched among genes with ASE in flower buds (S10 Table). Genes with nearby heterozygous TE

insertions were also enriched for a number of GO terms related to reproduction and defense (S11-S12

Table), suggesting that heterozygous TE insertions could be important for patterns of GO term

enrichment for genes with ASE

We further identified nineteen genes involved in floral and reproductive development in *A. thaliana*, which are located in QTL regions (see above), and show ASE in flower buds (Table 2). These genes are of special interest as candidate genes for detailed studies of the genetic basis of the selfing syndrome in *C. rubella*.

Intergenic divergence is elevated near genes with ASE

To investigate the role of polymorphisms in regulatory regions for ASE, we assessed levels of heterozygosity in intergenic regions 1 kb upstream of genes, and in previously identified conserved noncoding regions (Williamson et al. 2014) within 5 kb and 10 kb of genes. Genes with ASE were not significantly more likely to be associated with conserved noncoding regions with heterozygous SNPs than genes without ASE. However, levels of intergenic heterozygosity 1 kb upstream of genes were slightly but significantly higher for genes with ASE than for those without ASE (median heterozygosity of 0.016 vs. 0.014, respectively in leaves (Wilcoxon rank sum test, W = 295692325, p-value = $2.26*10^{-115}$), median heterozygosity of 0.017 vs. 0.014, respectively in flowers (Wilcoxon rank sum test, W = 297625040, p-value = $6.16*10^{-142}$), S13 Table), suggesting that polymorphisms in regulatory regions upstream of genes might contribute to *cis*-regulatory divergence.

Enrichment of TEs near genes with ASE

To test whether differences in TE content might contribute to *cis*-regulatory divergence between *C. rubella* and *C. grandiflora*, we examined whether heterozygous TE insertions near genes were associated with ASE. We identified TE insertions specific to the *C. grandiflora* or *C. rubella* parents of our F1s using genomic read data, as in Ågren et al. (2014) (Table 3; see Material and Methods). Overall, we found that *C. rubella* harbored fewer TE insertions close to genes than *C. grandiflora* (on average, 482 vs 1154 insertions within 1 kb of genes in *C. rubella* and *C. grandiflora*, respectively). Among heterozygous TE insertions, *Gypsy* insertions were the most frequent (Table 3); they were also the most frequent genome-wide (Table 3). There was a significant association between heterozygous TE insertions within 1 kb of genes and ASE, for both leaves and flower buds, and the strength of the association was greater for TE insertions closer to genes (Table 4; Fig. 5). This was true for individual F1s, as well as for all F1s collectively (Table 4; Fig. 5; S14 Table).

TEs targeted by uniquely mapping 24-nt small RNAs are associated with reduced expression of nearby genes

To test whether siRNA-based silencing of TEs might be responsible for the association between TE insertions and ASE in *Capsella*, we analyzed data for flower buds from one of our F1s, for which we

had matching small RNA data (see Material and Methods). We selected only those 24-nt siRNA reads

that mapped uniquely, without mismatch, to one site within each of our F1s, because uniquely

mapping siRNAs have been shown to have a more marked association with gene expression in

Arabidopsis (Hollister et al. 2009). For each gene, we then assessed the ASE ratio of the allele on the

same chromosome as a TE insertion (i.e. ASE ratios were polarized such that relative ASE was equal

to the ratio of the expression of the allele with a TE insertion on the same chromosome over the total

expression of both alleles), and then further examined the influence of nearby siRNAs.

Overall, the mean relative ASE was reduced for genes with nearby TE insertions (Fig. 6) with

a more pronounced effect for TE insertions within 1 kb (within the gene: Wilcoxon rank sum test, W =

1392103, p-value = $8.76*10^{-3}$; within 200 bp: Wilcoxon rank sum test, W = 1903047, p-value =

$7.17*10^{-3}$; within 1 kb: Wilcoxon rank sum test, W = 3687972, p-value = $8.19*10^{-3}$). The magnitude

of the effect on ASE was more pronounced for genes near TE insertions targeted by uniquely mapping

24-nt siRNAs (Fig. 6; for genes with a TE insertion within the gene: Wilcoxon rank sum test, W =

423369, p-value = $1.36*10^{-4}$; within 200 bp: W = 540926, p-value = $1.82*10^{-5}$; within 1 kb: W =

983938, p-value = $3.13*10^{-3}$). In contrast, no significant effect on ASE was apparent for genes near TE

insertions that were not targeted by uniquely mapping 24-nt siRNAs (Fig. 6). Thus, uniquely mapping

siRNAs targeting TE insertions appear to be responsible for the association we observe between ASE

and TE insertions. Globally, Gypsy and hAT insertions made up a greater proportion of the TE

insertions that were targeted by siRNA, compared to those that were not (Chi-squared test, $\chi=35.9468$,

$P=1.796*10^{-5}$, Supplementary Figure S7). However, for heterozygous TE insertions within 1 kb of

genes there were no significant differences in the composition of TEs that were vs. were not targeted

by uniquely mapping siRNAs.

Discussion

In this study, we have quantified allele-specific expression in order to understand the role of *cis*-

regulatory changes in association with a recent plant mating system shift. Our results indicate that

many genes, on average over 40%, harbor *cis*-regulatory differences between *C. rubella* and *C.*

grandiflora. The proportion of genes with ASE may seem high given the recent divergence (~100 kya)

between *C. rubella* and *C. grandiflora* (Brandvain et al. 2013, Slotte et al. 2013). However, the

majority of genes with ASE showed relatively mild allelic expression biases, and while our estimates

are higher than those in a recent microarray-based study of interspecific *Arabidopsis* hybrids (<10%)

(He et al. 2012a), our results are consistent with recent analyses of RNAseq data from intraspecific F1

hybrids of *Arabidopsis* accessions (~30%) (Cubillos et al. 2014). Somewhat higher levels of ASE

were found in a recent study of maize and teosinte (~70% of genes showed ASE in at least one tissue

and F1 individual (Lemmon et al. 2014), and using RNAseq data and the same hierarchical Bayesian

analysis that we employed, Skelly et al. (2011) estimated that a substantially higher proportion, >70%

of assayed genes, showed ASE among two strains of *Saccharomyces cerevisiae.* Thus, our estimates

of the proportion of genes with ASE fall within the range commonly observed for recently diverged

accessions or lines based on RNAseq data.

Two lines of evidence suggest that *cis*-regulatory changes have contributed to floral and

reproductive adaptation to selfing in *C. rubella*. First, we find an excess of genes with ASE in flower

buds within previously identified narrow QTL regions for floral and reproductive traits that harbor a

signature of selection (Slotte et al. 2012). This suggests either that multiple *cis*-regulatory changes

were involved in the evolution of the selfing syndrome in *C. rubella*, or that these regions harbor an

excess of *cis*-regulatory changes for other reasons, for instance due to hitchhiking of *cis*-regulatory

variants with causal variants for the selfing syndrome. Distinguishing between these hypotheses will

require identification of causal genetic changes for the selfing syndrome in *C. rubella*. In contrast, no

such excess is present for genes with ASE in leaves, suggesting that this observation is not simply a

product of higher levels of divergence among *C. rubella* and *C. grandiflora* in certain genomic regions

facilitating both QTL delimitation and ASE analysis. Second, we find that genes involved in hormonal

responses, including brassinosteroid biosynthesis, are overrepresented among genes with ASE in

flower buds, but not in leaves. Based on a study of differential expression and functional information

from *Arabidopsis thaliana*, regulatory changes in this pathway were previously suggested to be

important for the selfing syndrome in *C. rubella* (Slotte et al. 2013). While we do not identify ASE at

the same genes as in Slotte et al. 2013, our work nonetheless provides support for *cis*-regulatory

changes at other genes in the brassinosteroid pathway contributing to the selfing syndrome of *C.*

rubella. Future studies should conduct fine-scale mapping and functional validation to fully explore

this hypothesis. To facilitate this work, we have identified a set of candidate genes with ASE that are

located in genomic regions harboring QTL for floral and reproductive trait divergence between *C.*

rubella and *C. grandiflora*. Of particular interest in this list is the gene *JAGGED* (*JAG*), which is

involved in determining petal growth and shape by promoting cell proliferation in *A. thaliana* (Sauret-

Güeto et al. 2013, Schiessl et al. 2014). As *C. rubella* has reduced petal size due to a shortened period

of proliferative growth (Sicard et al. 2011), and the *C. rubella* allele is expressed at a lower level than

the *C. grandiflora* allele, this gene is a very promising candidate gene for the selfing syndrome.

Our work also provides general insights into the nature of *cis*-regulatory divergence. Indeed,

many instances of ASE were specific to a particular individual or tissue, an observation also supported

by recent studies (e.g. Lemmon et al. 2014, He et al. 2012a). This suggests that there is substantial

variation in ASE depending on genotype and developmental stage, consistent with the reasoning that

cis-regulatory changes can have very specific effects, but expression noise is probably also a

contributing factor. It is also difficult to completely rule out the possibility that some cases of subtle

ASE may not represent biologically meaningful *cis*-regulatory variation. However, in our analyses, we

took several steps to model and account for technical variation in order to reduce the incidence of false

positives. We also cannot fully rule out imprinting effects as potential causes of ASE, because

generating reciprocal F1 hybrids was not possible due to seed abortion in *C. rubella* x *C. grandiflora*

crosses. However, we do not expect these effects to make a major contribution to the patterns we

observed; in *Arabidopsis*, imprinting effects are only prevalent in endosperm tissue, and are rare in

more advanced stage tissues such as those analyzed here (Scott et al. 1998, Wolff et al. 2011, Cubillos

et al. 2014), which suggests that imprinting is not likely to be responsible for the patterns we observe.

One somewhat unexpected finding was the global shift in expression levels toward higher

relative expression of the *C. rubella* allele in the F1 hybrids. No marked bias was present for the same

SNPs and genes in our genomic data, suggesting that if systematic bioinformatic biases are the cause,

the effect is specific to transcriptomic reads. This seems unlikely to completely explain the shift in

expression that we observe, as we made considerable effort to avoid reference mapping bias, including

high stringency mapping of transcriptomic reads to reconstructed parental haplotypes specific to each

F1. Similar global shifts toward higher expression of the alleles from one parent have also been

observed in F1s of maize and teosinte (Lemmon et al. 2014) and *Drosophila* (McManus et al. 2010).

An even stronger bias toward higher expression of the *A. lyrata* allele was recently observed in F1s of

A. thaliana and *A. lyrata* (He et al. 2012a), and was attributed to interspecific differences in gene

silencing. Our results mirror those seen in some allopolyploids, where homeologs from one parental

species can be expressed at a markedly higher level than those from the other parental species (e.g.

Chang et al 2010; Flagel & Wendel 2010; Schnable et al 2011; Yoo et al. 2013).

To investigate potential mechanisms for *cis*-regulatory divergence, we first examined

heterozygosity in regulatory regions and conserved noncoding regions close to genes. While genes

with ASE in general showed slightly elevated levels of heterozygosity upstream of genes, there was no

enrichment of conserved noncoding regions with heterozygous SNPs close to genes with ASE. It thus

seems likely that divergence in regulatory regions in the proximity of genes, but not specifically in

conserved noncoding regions, has contributed to global *cis*-regulatory divergence between *C. rubella*

and *C. grandiflora*.

To examine biological explanations for the shift toward a higher relative expression of *C.*

rubella alleles, we examined the relationship between TE insertions and ASE. As *C. rubella* harbors a

lower number of TE insertions near genes than *C. grandiflora,* we reasoned that TE silencing might

contribute to the global shift in expression toward higher relative expression of the *C. rubella* allele,

with *C. grandiflora* alleles being preferentially silenced due to targeted methylation of nearby TEs,

through transcriptional gene silencing mediated by 24-nt siRNAs. Our results are consistent with this

hypothesis. Not only is there is an association between genes with TEs and heterozygous TE insertions

in our F1s, there is also reduced expression of alleles that reside on the same haplotype as a nearby TE

insertion, and the reduction is particularly strong for TEs that are targeted by uniquely mapping

siRNAs. In contrast, no effect on ASE is apparent for TEs that are not targeted by uniquely mapping

siRNAs. Moreover, the relatively limited spatial scale over which siRNA-targeted TE insertions are

associated with reduced expression of nearby genes (<1 kb) is consistent with previous results from

Arabidopsis (Hollister et al. 2009, Hollister et al. 2011, Wang et al. 2013). Our findings therefore

suggest that silencing of TE insertions close to genes is important for global *cis*-regulatory divergence

between *C. rubella* and *C. grandiflora*.

Why then do *C. rubella* and *C. grandiflora* differ with respect to silenced TEs near genes? In

Arabidopsis, methylated TE insertions near genes appear to be predominantly deleterious, and exhibit

a signature of purifying selection (Hollister et al. 2009). The reduced prevalence of TE insertions near

genes in *C. rubella* could be caused by rapid purging of recessive deleterious alleles due to increased

homozygosity as a result of self-fertilization (Arunkumar et al. 2014). However, we prefer the

alternative interpretation that deleterious alleles that were rare in the outcrossing ancestor were

preferentially lost in *C. rubella*, mainly as a consequence of the reduced effective population size

associated with the shift to selfing. This is in line with analyses of polymorphism and divergence at

nonsynonymous sites, for which *C. rubella* exhibits patterns consistent with a general relaxation of

purifying selection (Slotte et al. 2013).

If TE dynamics are generally important for *cis*-regulatory divergence in association with plant

mating system shifts, we might expect different effects on *cis*-regulatory divergence depending not

only on the genome-wide distribution of TEs, but also on the efficacy of silencing mechanisms in the

host (Hollister et al. 2009, Hollister et al. 2011, Ågren et al. 2015). For instance, He et al. (2012a)

found a shift toward higher relative expression of alleles from the outcrosser *A. lyrata,* which harbors

a higher TE content, a fact which they attributed to differences in silencing efficacy between *A.*

thaliana and *A. lyrata*; indeed, TEs also showed upregulation of the *A. lyrata* allele (He et al. 2012b)

and *A. lyrata* TEs were targeted by a lower fraction of uniquely mapping siRNAs (Hollister et al.

2011). In contrast, we found no evidence for a difference in silencing efficacy between *C. rubella* and

C. grandiflora, which harbor similar fractions of uniquely mapping siRNAs (12% vs 10% uniquely

mapping/total 24-nt RNA reads for *C. rubella* and *C. grandiflora*, respectively). Thus, in the absence

of strong divergence in silencing efficacy, differences in the spatial distribution of TEs, such as those

we observe between *C. rubella* and *C. grandiflora*, might be more important for *cis*-regulatory

divergence. More studies of ASE in F1s of selfers of different ages and their outcrossing relatives are

needed to assess the general contribution of differences in silencing efficacy versus genomic

distribution of TE insertions for *cis*-regulatory divergence in association with mating system shifts.

Conclusions

We have shown that many genes exhibit *cis*-regulatory changes between *C. rubella* and *C. grandiflora*

and that there is an enrichment of genes with floral ASE in genomic regions responsible for

phenotypic divergence. In combination with analyses of the function of genes with floral ASE, this

suggests that *cis*-regulatory changes have contributed to the evolution of the selfing syndrome in *C.*

rubella. We further observe a general shift toward higher relative expression of the *C. rubella* allele,

an observation that can in part be explained by elevated TE content close to genes in *C. grandiflora*

and reduced expression of *C. grandiflora* alleles due to silencing of nearby TEs. These results support

the idea that TE dynamics and silencing are of general importance for *cis*-regulatory divergence in

association with plant mating system shifts.

Material and Methods

Plant material

We generated three interspecific *C. grandiflora* x *C. rubella* F1s by crossing two accessions of the

selfer *C. rubella* as pollen donor with three accessions of the outcrosser *C. grandiflora* as seed parent

(S16 Table). No viable seeds were obtained from reciprocal crosses. Seeds from F1s and their *C.*

rubella parental lines were surface-sterilized and germinated on 0.5 x Murashige-Skoog medium. We

transferred one-week old seedlings to soil in pots that were placed in randomized order in a growth

chamber (16 h light: 8 h dark; 20° C: 14° C). After four weeks, but prior to bolting, we sampled young

leaves for RNA sequencing. Mixed-stage flower buds were sampled 3 weeks later, when all F1s were

flowering. To assess data reliability, we collected three separate samples of leaves and flower buds

from one F1 individual, and three biological replicates of one *C. rubella* parental line. For genomic

DNA extraction, we sampled leaves from all three F1 individuals as well as from their *C. rubella*

parents. For small RNA sequencing, we germinated six F2 offspring from one of our F1 individuals

and sampled flower buds as described above.

Sample preparation and sequencing

We extracted total RNA for whole transcriptome sequencing with the RNEasy Plant Mini Kit (Qiagen,

Hilden, Germany). For small RNA sequencing, we extracted total RNA using the mirVana kit (Life

Technologies). For whole genome sequencing, we used a modified CTAB DNA extraction (Doyle and

Doyle 1987) to obtain predominantly nuclear DNA. RNA sequencing libraries were prepared using

the TruSeq RNA v2 protocol (Illumina, San Diego, CA, USA). DNA sequencing libraries were

prepared using the TruSeq DNA v2 protocol. Small RNA libraries were prepared from 1 μg of total

RNA using the TruSeq SmallRNA SamplePrep fom Illumina according to the manufacturer's protocol

(#15004197 rev E; Illumina, San Diego, CA, USA). Sequencing was performed on an Illumina HiSeq

2000 instrument (Illumina, San Diego, CA, USA) to gain 100bp paired end reads, except for small

RNA samples for which single end 50 bp reads were obtained. Sequencing was done at the Uppsala

SNP & SEQ Technology Platform, Uppsala University, except for accession *C. rubella* Cr39.1 where

genomic DNA sequencing was done at the Max Planck Institute of Developmental Biology, Tübingen.

In total, we obtained 93.9 Gbp (Q≥30) of RNAseq data, with an average of 9.3 Gbp per sample. In

addition we obtained 45.6 Gbp (Q≥30) of DNAseq data, corresponding to a mean expected coverage

per individual of 52x, and 106,110,000 high-quality (Q≥30) 50 bp small RNA reads. All sequence data

has been submitted to the European Bioinformatics Institute (www.ebi.ac.uk), with study accession

number: PRJEB9020.

Sequence quality and trimming

We merged read pairs from fragment spanning less than 185 nt (this also removes potential adapter

sequences) in SeqPrep (https://github.com/jstjohn/SeqPrep) and trimmed reads based on sequence

quality (phred cutoff of 30) in CutAdapt 1.3 (Martin 2011). For DNA and RNAseq reads, we removed

all read pairs where either of the reads was shorter than 50 nt. We then analyzed each sample

individually using fastQC v. 0.10.1 (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) to

identify potential errors that could have occurred in the process of amplifying DNA and RNA. We

assessed RNA integrity by analyzing the overall depth of coverage over annotated coding genes, using

geneBody_coverage.py that is part of the RSeQC package v. 2.3.3 (Wang et al. 2012). For DNA reads

we analyzed the genome coverage using bedtools v.2.17.0 (Quinlan and Hall 2010) and removed all

potential PCR duplicates using Picard v.1.92 (http://picard.sourceforge.net). Small RNA reads were

trimmed using custom scripts and CutAdapt 1.3 and filtered to retain only reads of 24 nt length.

Read mapping and variant calling

We mapped both genomic reads and RNAseq reads to the v1.0 reference *C. rubella* assembly (Slotte

et al. 2013) (http://www.phytozome.net/capsella). For RNAseq reads we used STAR v.2.3.0.1 (Dobin

et al. 2013) with default parameters. For genomic reads we modified the default STAR settings to

avoid splitting up reads, and for mapping 24-nt small RNA we used STAR with settings modified to

require perfect matches to the parental haplotypes of the F1s as well as to a TE library based on

multiple Brassicaceae species and previously used in Slotte et al. (2013).

Variant calling was done in GATK v. 2.5-2 (McKenna et al. 2010) according to GATK best

practices (DePristo et al. 2011, Van der Auwera et al. 2013). Briefly, after duplicate marking, local

realignment around indels was undertaken, and base quality scores were recalibrated, using a set of

1,538,085 SNPs identified in *C. grandiflora* (Williamson et al. 2014) as known variants. Only SNPs

considered high quality by GATK were kept for further analysis. Variant discovery was done jointly

on all samples using the UnifiedGenotyper, and for each F1, genotypes were phased by transmission,

by reference to the genotype of its highly inbred *C. rubella* parental accession.

We validated our procedure for calling variants in genomic data by comparing our calls for the

inbred line *C. rubella* 1GR1 at 176,670 sites sequenced in a different individual from the same line by

Sanger sequencing (Slotte et al. 2010). Overall, we found 29 calls that differed among the two sets,

resulting in an error rate of 0.00016, considerably lower than the level of divergence among *C. rubella*

and *C. grandiflora* (0.02; Brandvain et al. 2013).

Reconstruction of parental haplotypes of interspecific F1s

We reconstructed genome-wide parental haplotype sequences for each interspecific F1 and used these

as a reference sequence for mapping genomic and transcriptomic reads for ASE analyses. This was

done to reduce effects of read mapping biases on our analyses of ASE by increasing the number of

mapped reads and reducing mismapping that can result when masking heterozygous SNPs in F1s

(Degner et al. 2009).

To reconstruct parental genomes for each F1, we first conducted genomic read mapping,

variant calling and phasing by reference to the inbred *C. rubella* parent as described in the section

"Read Mapping and Variant Calling" above. The resulting phased vcf files were used in conjunction

with the *C. rubella* reference genome sequence to create a new reference for each F1, containing both

of its parental genome-wide haplotypes. Read mapping of both genomic and RNA reads from each F1

was then redone to its specific parental haplotype reference genome, and read counts at all reliable

SNPs (see section "Filtering" below) were obtained using Samtools mpileup and a custom software

written in javascript by Johan Reimegård. The resulting files with allele counts for genomic and

transcriptomic data were used in all downstream analyses of allelic expression biases (see section

"Analysis of Allele-Specific Expression" below).

Filtering

We used two approaches to filter the genome assembly to identify regions where we have high

confidence in our SNP calls. Genomic regions with evidence for large-scale copy number variation

were identified using Control-FREEC (Boeva et al. 2011), and repeats and selfish genetic elements

were identified using RepeatMasker 4.0.1 (http://www.repeatmasker.org). Additionally, we identified

genomic regions with unusually high proportions of heterozygous genotype calls in a lab-inbred *C.*

rubella line, which is expected to be highly homozygous. Regions with evidence for high proportions

of repeats, copy number variation or high proportion of heterozygous calls in the inbred line mainly

corresponded to centromeric and pericentromeric regions, and these were removed from consideration

in further analyses of allele-specific expression (S2 Fig. - S5 Fig.).

Analysis of allele-specific expression

Analyses of allele-specific expression (ASE) were done using a hierarchical Bayesian method

developed by Skelly et al. (2011). The method requires read counts at heterozygous coding SNPs for

both genomic and transcriptomic data. Genomic read counts are used to fit the parameters of a beta-

binomial distribution, in order to obtain an empirical estimate of the distribution of variation in allelic

ratios due to technical variation (as there is no true ASE for genomic data on read counts for

heterozygous SNPs). This distribution is then used in analyses of RNAseq data where genes are

assigned posterior probabilities of exhibiting ASE.

We conducted ASE analyses using the method of Skelly et al. (2011) for each of our three F1

individuals. Prior to analyses, we filtered the genomic data to only retain read counts for heterozygous

SNPs in coding regions that did not overlap with neighboring genes, and following Skelly et al.

(2011), we also removed SNPs that were the most strongly biased in the genomic data (specifically, in

the 1% tails of a beta-binomial distribution fit to all heterozygous SNPs in each sample), as such

highly biased SNPs may result in false inference of variable ASE if retained. The resulting data set

showed very little evidence for read mapping bias affecting allelic ratios: the mean ratio of *C. rubella*

alleles to total was 0.507 (S6 Fig).

All analyses were run in triplicate and MCMC convergence was checked by comparing

parameter estimates across independent runs from different starting points, and by assessing the

degree of mixing of chains. For all analyses of RNA counts, we used median estimates of the

parameters of the beta-binomial distribution from analyses of genomic data for all three F1s (S8

Table). Runs were completed on a high-performance computing cluster at Uppsala University

(UPPMAX) using the pqR implementation of R (http://www.pqr-project.org), for 200,000 generations

or a maximum runtime of 10 days. We discarded the first 10% of each run as burn-in prior to

obtaining parameter estimates.

ASE validation by qPCR

We validated ASE results by performing qPCR with TaqMan® Reverse Transcription Reagents

(LifeTechnologies, Carlsbad, CA, USA) using oligo(dT)$_{16}$s to convert mRNA into cDNA using the

manufacturers protocol and performed qPCR with the Custom TaqMan® Gene Expression Assay

(LifeTechnologies, Carlsbad, CA, USA) with the colors FAM and VIC using manufacturers protocol.

The qPCR for both alleles was multiplexed in one well to directly compare the two alleles using a Bio-

Rad CFX96 Touch™ Real-Time PCR Detection System (Bio-Rad, Hercules, CA, USA). To exclude

color bias, we used reciprocal probes with VIC and FAM colorant (S15 Table). The expression

difference between the *C. rubella* and *C. grandiflora* allele was quantified using the difference in

relative expression between the two alleles, as well as the Quantification Cycle (Cq value). A lower

Cq value correlates with a higher amount of starting material in the sample. If the direction of allelic

imbalance inferred by qPCR was the same as for ASE inferred by the method by Skelly et al. (2011),

we considered that the qPCR supported the ASE results. For further details see S1 Text.

Enrichment of genes with ASE in genomic regions responsible for phenotypic divergence

We tested whether there was an excess of genes with evidence for ASE (posterior probability of ASE

≥ 0.95 in all three F1 hybrids) in previously identified genomic regions harboring QTL for phenotypic

divergence between *C. rubella* and *C. grandiflora* (Slotte et al. 2012). For this purpose, we

concentrated on narrow QTL regions, defined as in a previous study (Slotte et al. 2012) (i.e. QTL

regions with 1.5-LOD confidence intervals <2 Mb). The five QTL regions that met our criteria for

inclusion as narrow QTL were non-overlapping and corresponded to previously identified QTL for

floral and reproductive traits (on scaffolds 2 and 7 for petal width, on scaffold 7 for petal length and

on scaffolds 1 and 3 for flowering time). As QTL for floral and reproductive traits are generally highly

overlapping these genomic regions also encompass part of the confidence intervals for other QTL,

including a major QTL for petal length on scaffold 2, and QTL for sepal length, stamen length and

ovule number on scaffold 7). Significance was based on a permutation test (1000 permutations) in R

3.1.2.

List enrichment tests of GO terms

We tested for enrichment of GO biological process terms using Fisher exact tests in the R package

TopGO (Alexa et al. 2006). GO terms were downloaded from TAIR (http://www.arabidopsis.org) on

September 3rd, 2013, for all *A. thaliana* genes that have orthologs in the *C. rubella* v1.0 annotation,

and we only considered GO terms with at least two annotated members in the background set.

We tested for enrichment of GO biological process terms among genes with ASE in all of our

F1s Separate tests were conducted for leaf and flower bud samples, and background sets consisted of

all genes where we could assess ASE in either leaves or flower buds.

We used the same approach to test for enrichment of GO biological process terms among

genes within 1 kb and 2 kb of heterozygous TE insertions in F1 Inter4.1, for which we had matching

small RNA data. For this purpose, separate tests were done for all heterozygous TE insertions,

heterozygous TE insertions targeted by uniquely mapping siRNAs, and heterozygous TE insertions

not targeted by siRNAs. For these tests, the background sets consisted of all annotated *C. rubella*

genes.

Intergenic heterozygosity in regulatory and conserved noncoding regions

We quantified intergenic heterozygosity 1 kb upstream of genes using VCFTools (Danecek et al.

2011), and compared levels of polymorphism among genes with and without ASE using a Wilcoxon

rank sum test. We further assessed whether there was an enrichment of conserved noncoding elements

(identified in Williamson et al. (2014)) with heterozygous SNPs within 5 kb of genes with ASE, using

Fisher exact tests. Separate tests were conducted for each F1.

Identification of TE insertions and association with ASE

We used PoPoolationTE (Kofler et al. 2012) to identify transposable elements in our F1s. While

intended for pooled datasets, this method can also be used on genomic reads from single individuals

(Ågren et al. 2014). For this purpose we used a library of TE sequences based on several Brassicaceae

species (Slotte et al. 2013). We used the default pipeline for PoPoolationTE, modified to require a

minimum of 5 reads to call a TE insertion, and the procedure in Ågren et al. (2014) to determine

heterozygosity or homozygosity of TE insertions. Parental origins of TE insertions were inferred by

combining information from runs on F1s and their *C. rubella* parents. We used chi-square tests to

assess tested whether the composition of heterozygous TE insertions targeted by uniquely mapping

siRNAs differed from those not targeted by siRNAs.

We tested whether heterozygous TE insertions within a range of different window sizes close

to genes (200 bp, 1 kbp, 2 kbp, 5 kbp, and 10 kbp) were associated with ASE by performing Fisher

exact tests. We tested whether the expression of the allele on the same chromosome as a nearby

(within 1 kbp) TE insertion was reduced compared to ASE at against genes without nearby TE

insertions using a Wilcoxon rank sum test. Similar tests were conducted to test for an effect on relative

ASE of TE insertions with uniquely mapping siRNAs.

Acknowledgements

The authors thank Daniel Skelly, Duke University, for helpful advice on ASE analyses, Emily

Josephs, University of Toronto and Adrian Platts, McGill University for information on conserved

noncoding regions in *Capsella*, and Michael Nowak, Stockholm University, for valuable comments on

the manuscript. Sequencing was performed by the SNP&SEQ Technology Platform in Uppsala. The

facility is part of the National Genomics Infrastructure (NGI) Sweden and Science for Life

Laboratory. The SNP&SEQ Platform is also supported by the Swedish Research Council and the Knut

and Alice Wallenberg Foundation. The computations were performed on resources provided by SNIC

through Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX) under

Project b2012122. This work was supported by grants from the Swedish Research Council, the Erik

Philip-Sörensen foundation, the Nilsson-Ehle foundation, the Magnus Bergvall foundation, and the

Royal Swedish Academy of Sciences to T.S. D.K. acknowledges funding from the Human Frontier

Science Program (LT000783) and the German Research Foundation Priority Program 1529 –

'Adaptomics' (WE 2897).

References

Alexa A, Rahnenführer J, Lengauer T. 2006. Improved scoring of functional groups from gene

expression data by decorrelating GO graph structure. Bioinformatics. 22:1600–1607.

Arunkumar R, Ness RW, Wright SI, Barrett SCH. 2014. The Evolution of Selfing Is Accompanied by

Reduced Efficacy of Selection and Purging of Deleterious Mutations. Genetics. 199(3):817-829

Barrett SC. 2002. The evolution of plant sexual diversity. Nature Reviews Genetics. 3:274-284

Bernacchi D, Tanksley SD. 1997. An interspecific backcross of *Lycopersicon esculentum* x *L. hirsutum*:

linkage analysis and a QTL study of sexual compatibility factors and floral traits. Genetics.

147:861-877

Boeva V, Zinovyev A, Bleakley K, Vert J-P, Janoueix-Lerosey I, Delattre O, Barillot E. 2011. Control-

free calling of copy number alterations in deep-sequencing data using GC-content normalization.

Bioinformatics. 27:268–269.

Brandvain Y, Slotte T, Hazzouri KM, Wright SI, Coop G. 2013. Genomic Identification of Founding
Haplotypes Reveals the History of the Selfing Species *Capsella rubella*. PLoS Genet.
9:e1003754.

Carroll SB. 2000. Endless forms: the evolution of gene regulation and morphological diversity. Cell.
101:577–580.

Carroll SB. 2008. Evo-devo and an expanding evolutionary synthesis: a genetic theory of morphological
evolution. Cell. 134:25–36.

Chang PL, Dilkes BP, McMahon M, Comai L, Nuzhdin SV. 2010. Homoeolog-specific retention and
use in allotetraploid *Arabidopsis suecica* depends on parent of origin and network partners.
Genome Biology. 11:R125

Chen K-Y, Cong B, Wing R, Vrebalov J, Tanksley SD. 2007. Changes in regulation of a transcription
factor lead to autogamy in cultivated tomatoes. Science. 318:643-645

Cubillos FA, Stegle O, Grondin C, Canut M, Tisné S, Gy I, Loudet O. 2014. Extensive *cis*-regulatory
variation robust to environmental perturbation in *Arabidopsis*. Plant Cell. 26:4298–4310.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth
GT, Sherry ST et al. 2011. The variant call format and VCFtools. Bioinformatics. 27:2156–2158.

Degner JF, Marioni JC, Pai AA, Pickrell JK, Nkadori E, Gilad Y, Pritchard JK. 2009. Effect of read-
mapping biases on detecting allele-specific expression from RNA-sequencing data.
Bioinformatics. 25:3207–3212.

DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del Angel G,
Rivas MA, Hanna M et al. 2011. A framework for variation discovery and genotyping using next-
generation DNA sequencing data. Nat Genet. 43:491–498.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.
2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29:15–21.

Doebley J, Lukens L. 1998. Transcriptional regulators and the evolution of plant form. Plant Cell.
10:1075–1082.

Doyle JJ, Doyle JL. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue.
Phytochem bull. 19: 11-15.

Fishman L, Kelly AJ, Willis JH. 2002. Minor quantitative trait loci underlie floral traits associated with
mating system divergence in Mimulus. Evolution. 56:2138–2155.

Fishman L, Beardsley PM, Stathos A, Williams CF, Hill JP. 2015. The genetic architecture of traits
associated with the evolution of self-pollination in Mimulus. New Phytologist. 205:907-917

Flagel LE, Wendel JF 2010. Evolutionary rate variation, genomic dominance and duplicate gene
expression evolution during allotetraploid cotton speciation. New Phytologist. 186:184-193

Foxe JP, Slotte T, Stahl EA, Neuffer B, Hurka H, Wright SI. 2009. Recent speciation associated with
the evolution of selfing in *Capsella*. Proceedings of the National Academy of Sciences.
106:5241–5245.

Fraser HB. 2011. Genome-wide approaches to the study of adaptive gene expression evolution:
systematic studies of evolutionary adaptations involving gene expression will allow many
fundamental questions in evolutionary biology to be addressed. Bioessays. 33:469–477.
Goodwillie C, Ritland C, Ritland K. 2006 The genetic basis of floral traits associated with mating
system evolution in Leptosiphon (Polemoniaceae): an analysis of quantitative trait loci. Evolution.
60:491-504
Grillo MA, Changbao L, Fowlkes AM, Briggeman TM, Zhou A. Schemske DW, Sang T. 2009. Genetic
architecture for the adaptive origin of wild rice, *Oryza nivara*. Evolution. 63:870-883
Guo Y-L, Bechsgaard JS, Slotte T, Neuffer B, Lascoux M, Weigel D Schierup MH. 2009. Recent
speciation of *Capsella rubella* from *Capsella grandiflora*, associated with loss of self-
incompatibility and an extreme bottleneck. Proceedings of the National Academy of Sciences.
106:5246–5251.
He F, Zhang X, Hu J, Turck F, Dong X, Goebel U, Borevitz J, de Meaux J. 2012a. Genome-wide
analysis of *cis*-regulatory divergence between species in the *Arabidopsis* genus. Mol Biol Evol.
29:3385–3395.
He F, Zhang X, Hu J-Y, Turck F, Dong X, Goebel U, Borevitz JO, de Meaux J. 2012b. Widespread
interspecific divergence in cis-regulation of transposable elements in the *Arabidopsis* genus. Mol
Biol Evol. 29:1081–1091.
Hoekstra HE, Coyne JA. 2007. The locus of evolution: evo devo and the genetics of adaptation.
Evolution. 61:995–1016.
Hollister JD, Gaut BS. 2009. Epigenetic silencing of transposable elements: a trade-off between reduced
transposition and deleterious effects on neighboring gene expression. Genome Res. 19:1419–
1428.
Hollister JD, Smith LM, Guo Y-L, Ott F, Weigel D, Gaut BS. 2011. Transposable elements and small
RNAs contribute to gene expression divergence between *Arabidopsis thaliana* and *Arabidopsis*
lyrata. Proceedings of the National Academy of Sciences. 108:2322–2327.
Kofler R, Betancourt AJ, Schlötterer C. 2012. Sequencing of pooled DNA samples (Pool-Seq) uncovers
complex dynamics of transposable element insertions in *Drosophila melanogaster*. PLoS Genet.
8:e1002487.
Lemmon ZH, Bukowski R, Sun Q, Doebley JF. 2014. The Role of cis Regulatory Evolution in Maize
Domestication. PLoS Genet. 10:e1004745.
Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal. 17:10–12.
McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D,
Gabriel S, Daly M et al. 2010. The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data. Genome Res. 20:1297–1303.
McManus CJ, Coolon JD, O'Duff M, Eipper-Mains J, Graveley BR, Wittkopp PJ. 2010. Regulatory

divergence in *Drosophila* revealed by mRNA-seq. Genome Res. 20: 816-825

Ornduff R. 1969. Reproductive Biology in Relation to Systematics. Taxon. 18(2):121-133

Pastinen T. 2010. Genome-wide allele-specific analysis: insights into regulatory variation. Nat Rev
Genet. 11:533–538.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 26:841–842.

Rebernig CA, Lafon-Placette C, Hatorangan MR, Slotte T, Köhler C. 2015. Non-reciprocal interspecies
hybridization barriers in the Capsella genus are established in the endosperm. PLoS Genetics. 11:
e1005295

Sauret-Güeto S, Schiessl K, Bangham A, Sablowski R, Coen E. 2013. JAGGED controls *Arabidopsis*
petal growth and shape by interacting with a divergent polarity field. Plos Biol. 11:e1001550.

Schiessl K, Muiño JM, Sablowski R. 2014. *Arabidopsis* JAGGED links floral organ patterning to tissue
growth by repressing Kip-related cell cycle inhibitors. Proceedings of the National Academy of
Sciences. 111:2830–2835.

Schnable JC, Springer NM, Freeling M. 2011 Differentiation of the maize subgenomes by genome
dominance and both ancient and ongoing gene loss. Proceedings of the National Academy of
Sciences. 108:4069-4074

Scott RJ, Spielman M, Bailey J, Dickinson HG. 1998. Parent-of-origin effects on seed development in
Arabidopsis thaliana. Development. 125:3329–3341.

Sicard A, Lenhard M. 2011. The selfing syndrome: a model for studying the genetic and evolutionary
basis of morphological adaptation in plants. Annals of Botany. 107(9):1433-1443

Sicard A, Stacey N, Hermann K, Dessoly J, Neuffer B, Bäurle I, Lenhard M. 2011. Genetics, evolution,
and adaptive significance of the selfing syndrome in the genus *Capsella*. Plant Cell. 23:3156–
3171.

Skelly DA, Johansson M, Madeoy J, Wakefield J, Akey JM. 2011. A powerful and flexible statistical
framework for testing hypotheses of allele-specific gene expression from RNA-seq data. Genome
Res. 21:1728–1737.

Slotte T, Foxe JP, Hazzouri KM, Wright SI. 2010. Genome-wide evidence for efficient positive and
purifying selection in *Capsella grandiflora*, a plant species with a large effective population size.
Mol Biol Evol. 27:1813-1821.

Slotte T, Hazzouri KM, Stern D, Andolfatto P, Wright SI. 2012. Genetic architecture and adaptive
significance of the selfing syndrome in *Capsella*. Evolution. 66:1360–1374.

Slotte T, Hazzouri KM, Ågren JA, Koenig D, Maumus F, Guo YL, Steige K, Platts AE, Escobar JS,
Newman LK et al. 2013. The *Capsella rubella* genome and the genomic consequences of rapid
mating system evolution. Nat Genet. 2013; 45(7):831-835

St Onge KR, Källman T, Slotte T, Lascoux M, Palmé AE. 2011. Contrasting demographic history and
population structure in *Capsella rubella* and *Capsella grandiflora*, two closely related species

with different mating systems. Mol Ecol. 20:3306–3320.

Stebbins GL. 1950. Variation and Evolution in Plants. Columbia Univ. Press, New York.

Stern DL, Orgogozo V. 2008. The loci of evolution: how predictable is genetic evolution? Evolution.

62(9):2155–2177.

Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, Jordan T,

Shakir K, Roazen D, Thibault J et al. 2013. From FastQ data to high confidence variant calls: the

Genome Analysis Toolkit best practices pipeline. Curr Protoc Bioinformatics. 11:11.10.1–

11.10.33.

Wang L, Wang S, Li W. 2012. RSeQC: quality control of RNA-seq experiments. Bioinformatics.

28:2184–2185.

Wang X, Weigel D, Smith LM. 2013. Transposon variants and their effects on gene expression in

Arabidopsis. PLoS Genet. 9:e1003255.

Williamson RJ, Josephs EB, Platts AE, Hazzouri KM, Haudry A, Blanchette M, Wright SI. 2014.

Evidence for widespread positive and negative selection in coding and conserved noncoding

regions of *Capsella grandiflora*. PLoS Genet. 10:e1004622.

Wolff P, Weinhofer I, Seguin J, Roszak P, Beisel C, Donoghue MT, Spillane C, Nordborg M,

Rehmsmeier M, Köhler C. 2011. High-resolution analysis of parent-of-origin allelic expression in

the *Arabidopsis* Endosperm. PLoS Genet. 7:e1002126.

Wray GA. 2007. The evolutionary significance of *cis*-regulatory mutations. Nat Rev Genet. 8:206–216

Yoo MJ, Szadkowski E., Wendel JF. 2013 Homoeolog expression bias and expression level dominance

in allopolyploid cotton. Heredity. 110:171-180

Ågren JA, Wang W, Koenig D, Neuffer B, Weigel D, Wright SI. 2014. Mating system shifts and

transposable element evolution in the plant genus *Capsella*. BMC Genomics. 15:602.

Ågren JA, Wright SI. 2015. Selfish genetic elements and plant genome size evolution. Trends Plant Sci.

doi: 10.1016/j.tplants.2015.03.007

Supporting Information

S1 Information: S1 Text containing detailed procedures for filtering genomic regions, qPCR details,

Supporting Figures (S2-S7) and Tables (S8-S16).

Tables

Table 1. Genes amenable to analysis of ASE in flower bud and leaf samples from the three *C.*

grandiflora x *C. rubella* F1s, counts of genes with evidence for ASE and the estimated false discovery

rate (FDR) and proportion of genes with ASE.

F1 designation	Sample	Genes amenable to ASE analysis[a]	Analyzed genes[b]	Heterozygous SNPs in analyzed genes	Genes with ASE PP ≥ 0.95[c]	FDR	ASE proportion[d]
Inter3.1	Flower buds	18299	16857	262120	4728	0.0013	0.38
Inter4.1		18270	17837	272126	5744	0.0022	0.42
Inter5.1		18144	17448	262696	5176	0.0020	0.40
Inter3.1	Leaves	18299	14877	238786	5105	0.0012	0.44
Inter4.1		18270	15784	249181	8129	0.0024	0.62
Inter5.1		18144	15478	240653	4795	0.0018	0.41

[a]Total number of genes with heterozygous SNPs in coding regions remaining after filtering.

[b]Number of genes amenable to ASE analyses with expression data in at least one of the replicates of

the sample.

[c]Number of genes with evidence for ASE (posterior probability ≥ 0.95).

[d]Direct estimate of the ASE proportion independent of significance cutoffs.

Table 2. Selfing syndrome candidate genes identified based on ASE, QTL information, and Arabidopsis annotation.

C. rubella ortholog	Arabidopsis ortholog	Arabidopsis annotation	GO biological process terms related to floral and reproductive development
Carubv10012851m[a,b]	AT3G24340	CHR40	regulation of flower development
Carubv10016094m[a,b]	AT3G24650	ATABI3, ABI3, SIS10	embryo development, cotyledon development
Carubv10007602m[a,b]	AT4G21600	ENDO5	brassinosteroid biosynthetic process
Carubv10000655m[b,d]	AT5G08130	BIM1	brassinosteroid mediated signaling pathway, primary shoot apical meristem specification
Carubv10006681m[b,d]	AT4G28720	YUC8	brassinosteroid mediated signaling pathway
Carubv10021883m[a,c]	AT1G68480	JAG	sepal formation, flower development, abaxial cell fate specification, anther development, carpel development, stamen development, petal formation, specification of floral organ identity
Carubv10021345m[a,c]	AT1G68640	PAN, TGA8	petal formation, sepal formation, regulation of flower development
Carubv10013321m[a,c]	AT3G22420	ATWNK2, WNK2, ZIK3	photoperiodism, flowering
Carubv10016406m[a,c]	AT3G23270	-	pollen tube growth
Carubv10014951m[a,c]	AT3G23440	EDA6, MEE37	megagametogenesis
Carubv10014152m[a,c]	AT3G23630	ATIPT7, IPT7	pollen tube growth, reciprocal meiotic recombination
Carubv10010238m[a,c]	AT3G62210	EDA32	polar nucleus fusion
Carubv10004312m[a,c]	AT4G16760	ATACX1, ACX1	pollen development
Carubv10005585m[a,c]	AT4G17030	AT-EXPR, EXPR,	sexual reproduction

		ATEXLB1, ATEXPR1, EXLB1	
Carubv10007441m[a,c]	AT4G20370	TSF	regulation of flower development, photoperiodism, flowering, positive regulation of flower development
Carubv10004229m[a,c]	AT4G20910	CRM2, HEN1	specification of floral organ identity, floral organ formation, petal formation, regulation of flower development, sepal formation, meristem initiation, meristem development, ovule development
Carubv10015623m[a,c]	AT4G21380	ARK3, RK3	recognition of pollen
Carubv10007227m[a,c]	AT4G21530	APC4	ovule development
Carubv10007633m[a,c]	AT4G21590	ENDO3	petal development, stamen development, pollen tube growth, ovule development

[a]located within narrow QTL regions

[b]ASE in all three F1s

[c]ASE in the F1 with data for three replicates, but not in all three F1s

[d]located within QTL regions, but not narrow QTL regions

Table 3. Mean number of TE insertions in three interspecific F1s. The table shows the overall number,

as well as heterozygous insertions with parent of origin information.

TE family	Mean copy number	Heterozygous insertions	Insertions specific to the *C. rubella* parental genome	Insertions specific to the *C. grandiflora* parental genome
CACTA	84	40	10	30
Copia	710	483	144	339
Gypsy	1124	602	153	449
Harbinger	176	109	26	83
hAT	83	55	16	40
Helitron	236	127	30	97
LINE	229	165	38	128
MuDR	203	109	28	81
SINE	113	92	9	83
Total	2958	1782	454	1330

745   Table 4. Enrichment of heterozygous TEs near genes with ASE. The table shows mean counts over all

746   three F1s, and Fisher exact test *P*-values. The four categories of counts correspond to numbers of

747   genes with ASE (posterior probability of ASE $\geq$ 0.95) and TE insertions within a specific window size

748   near the gene (+ASE,+TE), with ASE but without TEs (+ASE,-TE), without ASE but with TE

749   insertions (-ASE,+TE), and with neither ASE nor TEs (-ASE,-TE). NS indicates not significant.

Sample	Window size (bp)	+ASE,+TE	+ASE-TE	-ASE, +TE	-ASE,-TE	*P*
Flower buds	200	113	5103	136	12029	$4.32*10^{-19}$
	1000	218	4998	339	11826	$5.07*10^{-16}$
	2000	307	4909	540	11624	$6.53*10^{-12}$
	5000	566	4650	1108	11057	$8.22*10^{-10}$
	10000	958	4258	2006	10159	$2.32*10^{-7}$
Leaves	200	108	5902	115	9255	$8.52*10^{-7}$
	1000	216	5793	277	9093	$1.49*10^{-4}$
	2000	317	5693	435	8935	$2.25*10^{-3}$
	5000	595	5415	877	8493	NS
	10000	1027	4983	1576	7795	NS

Figure 1

Fig. 1. ASE in flower buds. Distributions of ASE ratios (*C. rubella*/Total) for all assayed genes (A, B,

C), and for genes with at least 0.95 posterior probability of ASE (D, E, F). Ratio of *C. rubella* to total

for genomic reads, for genes with significant ASE (G, H, I), and the distribution of the dispersion

parameter that quantifies variability in ASE across genes (J, K, L). All distributions are shown for

each of the three interspecific F1s inter 3.1 (left), inter4.1 (middle) and inter5.1 (right).

Figure 2

Fig. 2. ASE in leaves. Distributions of ASE ratios (*C. rubella*/Total) for all assayed genes (A, B, C),

and for genes with at least 0.95 posterior probability of ASE (D, E, F). Ratio of *C. rubella* to total for

genomic reads, for genes with significant ASE (G, H, I), and the distribution of the dispersion

parameter that quantifies variability in ASE across genes (J, K, L). All distributions are shown for

each of the three interspecific F1s inter 3.1 (left), inter4.1 (middle) and inter5.1 (right).

Figure 3

767
Fig. 3. Many cases of ASE are specific to individuals or samples. Venn diagrams showing
intersections of genes with ASE in flower buds (A) and leaves (B) of the three F1 individuals, and (C)
in all leaf and flower samples, for the set of genes assayed in all F1s.

Figure 4

Fig. 4. Enrichment of genes with ASE in narrow QTL regions. There is an excess of genes with ASE

in narrow QTL regions for flower buds (A) but not for leaves (B). Histograms show the distribution of

numbers of genes with ASE that fall within narrow QTL regions, based on 1000 random permutations

of the observed number of genes with ASE among all genes where we could assess ASE. Arrows

indicate the observed number of genes with ASE that are located in narrow QTL regions.

Figure 5

Fig. 5. Enrichment of TEs near genes with ASE. The Figure shows odds ratios (OR) of the association

between genes with ASE and TEs, with TE insertions scored in four different window sizes (within a

distance of 0 bp, 1 kbp, 2 kbp, 5 kbp, and 10 kbp of each gene) . Odds ratios for flower buds are

shown for all three F1s studies, with values for flower buds in black and leaves in grey.

Fig. 6. The effect of TE insertions on relative allelic expression. Boxplots show the relative allelic
expression (expression of the allele on same haplotype as TE insertion relative to expression of both
alleles) for genes near heterozygous TE insertions, scored in a range of window sizes ranging from 0
bp (within the gene) to 10 kbp from the gene. A. The relative allelic expression is reduced for genes
with nearby TE insertions. B. The degree of reduction of relative allelic expression is stronger for

791   genes near TE insertions targeted by uniquely mapping siRNA. C. There is no reduction of relative

792   allelic expression for genes near TE insertions that are not targeted by uniquely mapping siRNA.