Title:

Brain composition in *Heliconius* butterflies, post-eclosion growth and experience

dependent neuropil plasticity

Authors:

Stephen H. Montgomery[1], Richard M. Merrill[2,3], Swidbert R. Ott[4]

Institutional affiliations:

[1] Dept. Genetics, Evolution & Environment, University College London, Gower

Street, London, UK, WC1E 6BT

[2] Dept. Zoology, University of Cambridge, Downing Street, Cambridge, UK, CB2

3EJ

[3] Smithsonian Tropical Research Institute, MRC 0580-12, Unit 9100 Box 0948, DPO

AA 34002-9998, Panama

[4] Dept. Neuroscience, Psychology and Behaviour, University of Leicester, Adrian

Building, University Road, Leicester, UK, LE1 7RH

Corresponding author: Stephen H. Montgomery: Stephen.Montgomery@cantab.net

Running head: *Anatomy and plasticity of* Heliconius *brains*

Key words: adaptive brain evolution, comparative neuroanatomy, *Heliconius*,

Lepidoptera, mushroom bodies, plasticity

Financial support: This research was supported by research fellowships from the

Royal Commission of the Exhibition of 1851 and Leverhulme Trust, a Royal Society

Research Grant (RG110466) and a British Ecological Society Early Career Project

Grant awarded to SHM. RMM was supported by a Junior Research Fellowship from

King's College, Cambridge and an Ernyst Mayr Fellowship from STRI. SRO was

supported by a University Research Fellowship from the Royal Society, London

(UK).

ABSTRACT

Behavioral and sensory adaptations are often based in the differential expansion of brain components. These volumetric differences represent changes in investment, processing capacity and/or connectivity, and can be used to investigate functional and evolutionary relationships between different brain regions, and between brain composition and behavioral ecology. Here, we describe the brain composition of two species of *Heliconius* butterflies, a long-standing study system for investigating ecological adaptation and speciation. We confirm a previous report of striking mushroom body expansion, and explore patterns of post-eclosion growth and experience-dependent plasticity in neural development. This analysis uncovers age- and experience-dependent post-emergence mushroom body growth comparable to that in foraging hymenoptera, but also identifies plasticity in several other neuropil. An interspecific analysis indicates that *Heliconius* display remarkable levels of investment in mushroom bodies for a lepidopteran, and indeed rank highly compared to other insects. Our analyses lay the foundation for future comparative and experimental analyses that will establish *Heliconius* as a useful case study in evolutionary neurobiology.

INTRODUCTION

Behavioral adaptations are largely based in changes in brain function. In some cases this includes differential expansion of individual brain structures, or functionally related systems, that betray underlying changes in neuron number or circuitry. These provide an opportunity to study the neural basis of adaptive behavior, particularly in clades with known ecological specializations. The Neotropical genus *Heliconius* (Heliconiinae, Nymphalidae) display a number of striking behavioral adaptations including a dietary adaptation unique among Lepidoptera; adult pollen feeding (Gilbert, 1972, 1975). With the exception of four species formerly ascribed to the genus *Neruda* (Beltrán et al., 2007; Kozak et al., 2015), all *Heliconius* actively collect and ingest pollen as adults. This provides a source of amino acids and permits a greatly extended lifespan of up to six months without reproductive senescence (Gilbert, 1972; Benson, 1972; Ehrlich and Gilbert, 1973). Without access to pollen *Heliconius* suffer a major reduction in longevity and reproductive success (Gilbert, 1972; Dunlap-Pianka et al., 1977; O'Brien et al., 2003).

Several lines of evidence suggest selection for pollen feeding has shaped *Heliconius* foraging behavior. Pollen is collected from a restricted range of mostly Cucurbitaceous plants (Estrada and Jiggins, 2002), which occur at low densities (Gilbert, 1975). Individuals inhabit home ranges of typically less than 1 km^2, within which they repeatedly utilize a small number of roosting sites that they return to with high fidelity (Turner, 1971; Benson, 1972; Gilbert, 1975; Mallet, 1986; Murawski and Gilbert, 1986; Finkbeiner, 2014). On leaving the roost individuals visit feeding sites with a level of consistency in time and space that strongly suggests 'trap-lining' behavior (Ehrlich and Gilbert, 1973; Gilbert, 1975, 1993; Mallet, 1986), analogous to that observed in foraging bees (Janzen, 1971; Heinrich, 1979). Roosts themselves are located visually (Jones, 1930; Gilbert, 1972; Ehrlich and Gilbert, 1973; Mallet, 1986), and older individuals tend to be more efficient foragers (Boggs et al., 1981; Gilbert, 1993). Together these observations suggest the evolution of pollen feeding in *Heliconius* was facilitated by an enhanced, visually-orientated and time-compensated memory that utilizes long distance landmarks (Gilbert, 1975). The evolution of this behavior must involve "some elaboration of the nervous system" (Turner, 1981). This elaboration is suggested to occur in the mushroom bodies, which Sivinski (1989)

reported are 3–4 times larger in *Heliconius charithonia* than in six other species of
butterfly, including two non-pollen feeding Heliconiini.

Insect mushroom bodies have a variety of roles in olfactory associative
learning, sensory integration, filtering and attention (Zars, 2000; Farris, 2005, 2013;
Menzel, 2014). Direct experimental evidence suggests the mushroom bodies mediate
place memory in *Periplaneta americana* (Mizunami et al., 1998; Lent et al., 2007).
Two lines of indirect evidence further implicate the mushroom bodies in allocentric
memory in other insects. First, comparisons across species suggest that extreme
evolutionary expansion of the mushroom body may commonly be associated with
changes in foraging behavior that depend on spatial memory or the complexity of
sensory information utilized by the species (Farris, 2005, 2013). For example,
phylogenetic comparisons across Hymenoptera demonstrate the volumetric expansion
and elaboration of the Euhymenopteran mushroom body occurs coincidently with the
origin of parasitoidism (Farris and Schulmeister, 2011), a behavioral adaptation that
involves place-centered foraging and spatial memory for host location (Rosenheim,
1987; van Nouhuys and Kaartinen, 2008). Second, mushroom bodies are
ontogenetically plastic, and this plasticity has been linked to foraging behavior.
Again, the trap-lining Hymenoptera illustrate the link between foraging behavior and
the mushroom bodies (Withers et al., 1993; Durst et al., 1994; Capaldi et al., 1999;
Farris et al., 2001). Honeybees show two forms of post-eclosion growth in mushroom
body volume; age dependent growth, which occurs regardless of environmental
variation, and experience dependent growth which increases with foraging or social
experience (Withers et al., 1993; Durst et al., 1994; Fahrbach et al., 1998, 2003; Farris
et al., 2001; Maleszka et al., 2009). In other Hymenoptera there is close
correspondence between the rate and timing of mushroom body growth and the onset
of foraging behavior (Gronenberg et al., 1996; Kühn-Bühlmann and Wehner, 2006;
Withers et al., 2008; Jones et al., 2013). Whether the trap-lining behavior observed in
Heliconius is associated with similar ontogenetic plasticity is not known.

Here we confirm Sivinksi's (1989) observation of a phylogenetic expansion of
the mushroom bodies in *Heliconius*. We further demonstrate ontogenetic and
environmentally induced plasticity comparable in size to trap-lining Hymenoptera.
Together these analyses suggest the mushroom bodies may have a role in the
allocentric spatial foraging observed in *Heliconius*, and lay the groundwork for

comparative analyses across Heliconiini examining the origin and timing of
mushroom body expansion.
135

MATERIALS & METHODS

Animals

We collected five males and five females of two species of *Heliconius*, *H. hecale*
melicerta and *H. erato demophoon* from wild populations around Gamboa (9°7.4′ N,
79°42.2′ W, elevation 60 m) and the nearby Soberanía National Park, República de
Panamá. We assume all wild-caught individuals were sexually mature, and that the
age range is not biased between species or sexes. Wild individuals were compared to
individuals from first or second-generation insectary-reared stock populations,
descended from wild caught parents from the same sampling localities. Stock
populations were kept in controlled conditions in cages (c. 1 × 2 × 2 m) of mixed sex
at roughly equal densities. Cages were housed at the *Heliconius* insectaries at the
Smithsonian Tropical Research Institute's (STRI) facility in Gamboa. Stocks had
access to their preferred host plant (*Passiflora biflora* and *P. vitifolia* respectively for
H. erato and *H. hecale*), a pollen source (*Psychotria elata*) and feeders containing c.
20% sugar solution with an additional bee-pollen supplement to ensure an excess of
pollen. Larvae were allowed to feed naturally on the host plant.

After emergence from the pupae insectary-reared individuals were collected
for two age groups, a recently emerged 'young' group (1–3 days post emergence) and
an 'old' group (2–3 weeks post emergence). *Heliconius* undergo a "callow" period of
general inactivity immediately after emergence that lasts about 5 days, during which
flight behavior is weak and males are sexually inactive (Mallet, 1980). These age
groups therefore represent behaviorally immature and mature individuals. For *H.*
hecale 5 males and 5 females were sampled for both age groups, in *H. erato* 4 males
and 6 females were sampled for the 'young' group and 5 males and 4 females were
sampled for the 'old' group. In samples for which the exact time of emergence was
known there was no significant difference between *H. hecale* and *H. erato* in age
structure of the old (*H. erato*: mean = 22.6 days, SD = 8.6; *H. hecale*: mean = 26.4
days, SD = 5.5; t_{13} = -0.899, p = 0.385) or young (*H. erato*: mean = 1.7 days, SD =
0.8; *H. hecale*: mean = 1.3 days, SD = 1.1; t_{17} = 0.829, p = 0.419) insectary-reared
groups. Three body size measurements were taken for each individual: body mass,

weighted to 0.01 g using a OHAUS pocket balance (model YA102), body length, and wingspan, measured using FreeLOGIX digital calipers. Samples were collected and exported under permits SEX/A-3-12 and SE/A-7-13 obtained from the Autoridad Nacional del Ambiente, República de Panamá in conjunction with STRI.

Antibodies and sera for neuropil staining

We used indirect immunofluorescence staining against synapsin to reveal the neuropil structure of the brain under a confocal microscope (Ott, 2008). This technique exploits the abundant expression of synapsin, a vesicle-associated protein, at presynaptic sites. Monoclonal mouse anti-synapsin antibody 3C11 (anti-SYNORF1; (Klagges et al., 1996) was obtained from the Developmental Studies Hybridoma Bank (DSHB), University of Iowa, Department of Biological Sciences, Iowa City, IA 52242, USA (RRID: AB_2315424). The 3C11 antibody was raised against a bacterially expressed fusion protein generated by adding a glutathione S-transferase (GST)-tag to cDNA comprised of most of the 5′ open reading frame 1 of the *Drosophila melanogaster* synapsin gene (*Syn,* CG3985). The binding specificity of this antibody was characterised in *D. melanogaster* (Klagges et al., 1996) and confirmed in synapsin null mutants by Godenschwege et al. (2004). The epitope was later narrowed down to within LFGGMEVCGL in the C domain (Hofbauer et al., 2009). Bioinformatic analysis has confirmed the presence of this motif in lepidopteran genomes, and demonstrated that it is highly conserved across Lepidoptera (Montgomery and Ott, 2015). Binding specificity in *M. sexta* has been confirmed by western blot analysis (Utz et al., 2008) and 3C11 immunostaining has been used as an anatomical marker of synaptic neuropil in a wide range of arthropod species including several Lepidoptera: *D. plexippus* (Heinze and Reppert, 2012), *G. zavaleta* (Montgomery and Ott, 2015), *H. virescens* (Kvello et al., 2009) and *M. sexta* (El Jundi et al., 2009). Cy2-conjugated affinity-purified polyclonal goat anti-mouse IgG (H+L) antibody (Jackson ImmunoResearch Laboratories, West Grove, PA) was obtained from Stratech Scientific Ltd., Newmarket, Suffolk, UK (Jackson ImmunoResearch Cat No. 115-225-146, RRID: AB_2307343).

Immunocytochemistry

Brains were fixed and stained following a published protocol (Ott, 2008). The protocol was divided into two stages, the first of which was performed at the STRI

Gamboa Field Station. The brain was exposed under HEPES-buffered saline (HBS;

150 mM NaCl; 5 mM KCl; 5 mM $CaCl_2$; 25 mM sucrose; 10 mM HEPES; pH 7.4)

and fixed in situ for 16–20 hours at room temperature (RT) in zinc-formaldehyde

solution (ZnFA; 0.25% (18.4 mM) $ZnCl_2$; 0.788% (135 mM) NaCl; 1.2% (35 mM)

sucrose; 1% formaldehyde) under agitation. The brain was subsequently dissected out

under HBS, washed (3 × in HBS), placed into 80% methanol/20% DMSO for 2 hours

under agitation, transferred to 100% methanol and stored at RT. After transportation

to the UK samples were stored at -20˚C.

In the second stage of the protocol the samples were brought to RT and

rehydrated in a decreasing methanol series (90%, 70%, 50%, 30%, 0% in 0.1 M Tris

buffer, pH 7.4, 10 minutes each). Normal goat serum (NGS; New England BioLabs,

Hitchin, Hertfordshire, UK) and antibodies were diluted in 0.1 M phosphate-buffered

saline (PBS; pH 7.4) containing 1% DMSO and 0.005% NaN_3 (PBSd). After a pre-

incubation in 5% NGS (PBSd-NGS) for 2 hours at RT, antibody 3C11 was applied at

a 1:30 dilution in PBSd-NGS for 3.5 days at 4˚C under agitation. The brains were

rinsed in PBSd (3 × 2 hours) before applying the Cy2-conjugated anti-mouse antibody

1:100 in PBSd-NGS for 2.5 days at 4˚C under agitation. This was followed by

increasing concentrations of glycerol (1%, 2%, 4% for 2 hours each, 8%, 15%, 30%,

50%, 60%, 70% and 80% for 1 hour each) in 0.1 M Tris buffer with DMSO to 1%.

The brains were then passed in a drop of 80% glycerol directly into 100% ethanol.

After agitation for 30 minutes the ethanol was refreshed (3 × 30 minute incubations),

before being underlain with methyl salicylate. The brain was allowed to sink, before

the methyl salicylate was refreshed (2 × 30 minute incubations).

Confocal imaging

All imaging was performed on a confocal laser-scanning microscope (Leica TCS SP8,

Leica Microsystem, Mannheim, Germany) using a 10× dry objective with a numerical

aperture of 0.4 (Leica Material No. 11506511), a mechanical z-step of 2 μm and an x-

y resolution of 512 × 512 pixels. Imaging the whole brain required capturing 3×2 tiled

stacks in the x-y dimensions (20% overlap) that were automatically merged in Leica

Applications Suite Advanced Fluorescence software. Each brain was scanned from

the posterior and anterior side to span the full z-dimension of the brain. These image

stacks were then merged in Amira 3D analysis software 5.5 (FEI Visualization

Sciences Group; custom module 'Advanced Merge'). The *z*-dimension was scaled

1.52× to correct the artifactual shortening associated with the 10× air objective

(Heinze and Reppert, 2012; Montgomery and Ott, 2015). Images that illustrate key

morphological details were captured separately as single confocal sections with an *x-y*

resolution of 1024 × 1024 pixels.

Neuropil segmentations and volumetric reconstructions

We assigned image regions to anatomical structures in the Amira 5.5 *labelfield*

module by defining outlines based on the brightness of the synapsin

immunofluorescence. Within each stack, every forth or fifth image was manually

segmented and interpolated in the *z*-dimension across all images that contain the

neuropil of interest. The *measure statistics* module was used to determine volumes (in

μm^3) for each neuropil. 3D polygonal surface models of the neuropils were

constructed from the smoothed labelfield outlines (*SurfaceGen* module). The color

code used for the neuropils in the 3D models is consistent with previous

neuroanatomical studies of insect brains (Brandt et al., 2005; Kurylas et al., 2008; El

Jundi et al., 2009a, b; Dreyer et al., 2010; Heinze and Reppert, 2012; Montgomery

and Ott, 2015).

The whole-brain composite stacks were used to reconstruct and measure six

paired neuropils in the optic lobes, and seven paired and two unpaired neuropils in the

midbrain where distinct margins in staining intensity delineate their margins. All

paired neuropils were measured on both sides of the brain in wild-caught individuals

to permit tests of asymmetry, yielding two paired measurements per brain (*i.e. N* = 10

× 2) for each structure. We found no evidence of volumetric asymmetry for either

species (p > 0.05 for each neuropil in a paired t-tests) and therefore summed the

volumes of paired neuropil to calculate the total volume of that structure. In insectary-

reared individuals we subsequently measured the volume of paired neuropil from one

hemisphere, chosen at random, and multiplied the measured volume by two. We

measured the total neuropil volume of the midbrain to permit statistical analyses that

control for allometric scaling. In keeping with the earlier lepidopteran literature, we

use the term 'midbrain' for the fused central mass that comprises the protocerebral

neuromere excluding the optic lobes, the deuto- and tritocerebral neuromeres, and the

sub-esophageal neuromeres. For the following statistical analyses we analyzed the

central body as a single structure and, unless otherwise stated, summed the volumes

of the mushroom body lobes and peduncles.

Intraspecific statistical analyses

In all statistical analyses continuous variables were log_{10}-transformed. Unpaired two-

tailed two-sample *t*-tests were used to test for volumetric differences between sexes or

groups. We found no evidence of sexual dimorphism in neuropil volume of wild

caught individuals that could not be explained by allometric scaling and therefore

combined male and female data.

All statistical analyses were performed in R v.3.1 (R Development Core

Team, 2008). Our analyses focused on two intra-specific comparisons: i) we

compared 'young' and 'old' insectary-reared individuals and interpret significant

differences as evidence for post-eclosion growth; and ii) we compared wild-caught

individuals with 'old' insectary-reared individuals and interpret significant differences

as evidence for environmentally induced, or experience dependent plasticity. These

comparisons were made by estimating the allometric relationship between each

neuropil and a measure of overall brain size (total volume of the midbrain minus the

combined volume of all segmented neuropil in the midbrain: 'rest of midbrain', rMid)

using the standard allometric scaling relationship: $\log y = \beta \log x + \alpha$. We used

standard major axis regressions in the SMATR v.3.4-3 (Warton et al., 2012) to test for

significant shifts in the allometric slope (β). Where we identified no heterogeneity in

β we performed two further tests: 1) for differences in α that suggest discrete 'grade-

shifts' in the relationship between two variables, 2) for major axis-shifts along a

common slope. Patterns of brain:body allometry were explored in a similar manner,

using total neuropil volume as the dependent variable (summed volumes of all optic

lobes neuropil plus the total midbrain volume), and comparing the results obtained

using alternative body size measurements as the independent variable. We also

present the effect size, measured by the correlation coefficient (*r*). Effect sizes of

$0.1 < r < 0.3$ are interpreted as 'small' effects, $0.3 < r < 0.5$ 'medium' effects, and $r < 0.5$

'large' effects (Cohen, 1988).

Interspecific statistical analyses

To analyze interspecific patterns of divergence in brain composition we collected

published data for neuropil volumes of four other Lepidoptera; *D. plexippus*; (Heinze

and Reppert, 2012), *G. zavaleta* (Montgomery and Ott, 2015), *M. sexta*; (El Jundi et al., 2009a) and *H. virescens* (Kvello et al., 2009). Data were available for eight neuropils across all four species. Relative size was measured by calculating the residuals from a phylogenetically-corrected least squares (PGLS) linear regression between each structure and the rest of the brain (whole brain or midbrain as indicated) performed in BayesTraits (freely available from www.evolution.rdg.ac.uk; Pagel, 1999). For this analysis, a phylogeny of the six species was created using data on two loci, *COI* and *EF1a* (GenBank Accession IDs, *COI*: EU069042.1, GU365908.1, JQ569251.1, JN798958.1, JQ539220.1, HM416492.1; *EF1a*: EU069147.1, DQ157894.1, U20135.1, KC893204.1, AY748017.1, AY748000.1). The data were aligned and concatenated using MUSCLE (Edgar, 2004), before constructing a maximum likelihood tree in MEGA v.5 (Tamura et al., 2011). Differences in brain composition across species were analyzed by Principal Component analysis of these data, and visualized as biplots (Greenacre, 2010) in R package *ggbiplot* (V.Q. Vu, https://github.com/vqv/ggbiplot). Finally, we extended our phylogenetic analysis across insects using a similar approach. We restricted this analysis to volumetric data collected with similar methodology (Rein et al., 2002; Brandt et al., 2005; Kurylas et al., 2008; Dreyer et al., 2010; Ott and Rogers, 2010; Wei et al., 2010). The phylogenetic relationship of these insects was taken from Trautwein et al. (2012).

RESULTS

General layout of the *Heliconius* brain

The overall layout and morphology of the *Heliconius* brain (Fig. 1) is similar to that of other Lepidoptera (El Jundi et al., 2009; Kvello et al., 2009; Heinze and Reppert, 2012a; Montgomery and Ott, 2015). The midbrain forms a single medial mass, containing the supra-esophageal ganglion to which the sub-esophageal ganglion is fused. Together with the rest of the midbrain (rMid), which lacks distinct internal boundaries and was therefore unsuitable for further segmentation in the current analysis, we measured the volumes of six paired neuropils in the optic lobes, and eight paired and two unpaired neuropils in the midbrain in 59 individuals across both species (Table 1).

Sensory neuropil

The large optic lobes (OL; Fig. 2) account for approximately 64% of the total brain

volume. As is the case in both *D. plexippus* and *G. zavaleta* the lamina (La), two-

layered medulla (Me) (Fig. 2E), accessory medulla (aMe), lobula (Lob) and lobula

plate (Lop) are well defined and positioned in the OL as nested structures from lateral

to medial (Fig. 2A). The La has a distinct, brightly stained inner rim (iRim; Fig. 2E), a

feature common to all diurnal butterflies analyzed thus far (Heinze and Reppert, 2012;

Montgomery and Ott, 2015). In common with *D. plexippus* we identify a thin strip of

irregularly shaped neuropil running ventrally from the aME to the Me (Fig. 2G–H).

We also identify a sixth neuropil in the OL that we believe to be homologous

to the optic glomerulus (OG; Fig. 2B,F) identified in *D. plexippus* (Heinze and

Reppert, 2012)*,* which is absent in other lepidopteran brains described to date and was

postulated to be Monarch-specific. As in *D. plexippus* this neuropil is a multi-lobed,

irregularly shaped structure positioned to the medial margin of the Lob with which it

appears to be connected. In *Heliconius* the OG is not as extended in the anterior

margin as in *D. plexippus* and is subsequently confined to the OL, without protrusion

into the optic stalk or midbrain (Fig. 2A,B,F). The position of the OG in *Heliconius*

is also similar to that of a much smaller neuropil observed in *G. zavaleta*

(Montgomery and Ott, 2015) that may be homologous.

The midbrain contains further neuropils with primary functions in processing

visual information that include the anterior optic tubercule (AOTu). We identify the

same four components of the AOTu previously described in *D. plexippus* and *G.*

zavaleta butterflies (Heinze and Reppert, 2012; Montgomery and Ott, 2015): the

small, closely clustered nodular unit (NU), strap (SP) and lower unit (LU), and the

much larger upper unit (UU) (Fig. 2C). As in other butterflies, the UU is expanded

compared with nocturnal moths (El Jundi et al., 2009; Kvello et al., 2009). The

proportion of total neuropil comprised of the AOTu is, however, larger in *D.*

plexippus (0.74%) than *Heliconius* (0.40% in *H. hecale* and 0.37% in *H. erato*).

The antennal lobes (AL), the primary olfactory neuropil, are comprised of

small, round glomeruli that are innervated by axons from olfactory receptor neurons

in the antennae. These glomeruli are arranged around a Central Fibrous Neuropil

(CFN) (Figure 3A,B). In *Heliconius* the AL comprises 2% of the total brain neuropil

volume, and contains approximately 68 glomeruli (estimated in one individual of each

sex: *H. erato* ♂ = 69, ♀ = 68; *H. hecale* ♂ = 68, ♀ = 67) which matches closely the

number of olfactory receptor genes (70) identified in the *H. melpomene* genome

(Dasmahapatra et al., 2012). We found no expanded macro-glomeruli complex
(MGC) or obvious candidates for sexually dimorphic glomeruli. This is in keeping
with all diurnal butterflies described to date (Rospars, 1983; Heinze and Reppert,
2012; Carlsson et al., 2013), with the exception of the more olfactorily orientated *G.*
zavaleta (Montgomery and Ott, 2015).

We took advantage of comparable datasets for *H. erato, H. hecale* and *G.*
zavaleta to investigate whether changes in relative AL volume are due to an increased
volume of glomeruli or CFN. Both glomerular and CFN volume are larger in *G.*
zavaleta relative to midbrain volume, as indicated by significant grade-shifts in
allometric scaling in *G. zavaleta* and *Heliconius* (glomerular, *H. erato*: Wald $\chi 2$ =
10.709, p = 0.001; *H. hecale*: Wald $\chi 2$ = 9.139, p = 0.003; CFN, *H. erato*: Wald $\chi 2$ =
30.282, p < 0.001; *H. hecale*: Wald $\chi 2$ = 26.638, p < 0.001). However, CFN
expansion in *G. zavaleta* is disproportionately large, driving a grade-shift in the
scaling relationship between glomerular and CFN volume in *G. zavaleta* when
compared with either *Heliconius* (*H. erato*: Wald $\chi 2$ = 19.680, p < 0.001; *H. hecale*:
Wald $\chi 2$ = 31.663, p < 0.001; Fig. 3D).

Central complex
The central complex is a multimodal integration center linked to a range of functions
from locomotor control to memory (Pfeiffer and Homberg, 2014). Within the
limitations of the current analysis, the anatomy of the *Heliconius* central complex
shows strong conservation with *D. plexippus* and *G. zavaleta* (Heinze and Reppert,
2012; Montgomery and Ott, 2015). The central body (CB) is positioned along the
midline of the central brain and is formed of two neuropils, the upper (CBU) and
lower (CBL) divisions, which are associated with small paired neuropils, the noduli
(No), located ventrally to the CB (Fig. 4A–D,G). Two further paired neuropils, the
protocerebral bridge (PB; Fig. 4A,E) and posterior optic tubercles (POTu; Fig. 4A,F),
are positioned towards the posterior margin of the brain.

Mushroom bodies
The most striking aspect of *Heliconius* brain morphology is the hugely expanded
mushroom bodies which span the depth of the brain along the anterior-posterior axis
(Fig. 5). On the anterior side, the mushroom body lobes (MB-lo) lie above the AL. As
in *D. plexippus* (Heinze and Reppert, 2012) the distinct lobe structure observed in

moths (El Jundi et al., 2009; Kvello et al., 2009) is lost, possibly due to extensive expansion. The only identifiable feature is a lobe curving round the medial margin, likely to be part of the vertical lobe (Fig. 5D,F). The MB-lo merges with the cylindrical pedunculus (MB-pe) that extends to the posterior midbrain. The boundary between the MB-lo and MB-pe is not distinct. The combined volume of the MBlo+pe accounts for 12.2% of total midbrain volume in *H. hecale* and 14.6% of total midbrain volume in *H. erato*, at least twice that reported for other Lepidoptera (Sjöholm et al., 2005; El Jundi et al., 2009; Kvello et al., 2009; Heinze and Reppert, 2012a; Montgomery and Ott, 2015). At the posterior end, the MB-pe splits into two roots that are encircled by the mushroom body calyx (MB-ca; Fig. 5A,H,K). A Y-tract runs parallel to the MB-pe from the posterior boundary of the MB-lo to the junction between the MB-pe and MB-ca. The Y-tract ventral loblets seen in other Lepidoptera (El Jundi et al., 2009; Kvello et al., 2009) are not distinct, having merged with the MB-lo (Fig. 5A,J,N).

Heliconius have an un-fused double MB-ca with a deeply cupped morphology (Fig. 5A,C). Two concentric zones can be identified (Fig. 5E), though the boundary is not distinct throughout the depth of the neuropil. The MB-ca comprises 20.7% and 23.9% of total midbrain volume in *H. hecale* and *H. erato* respectively, at least three times greater than reported in other Lepidoptera (Sjöholm et al., 2005; El Jundi et al., 2009; Kvello et al., 2009; Heinze and Reppert, 2012a; Montgomery and Ott, 2015). In some individuals the MB-ca is so large that it protrudes into the OL resulting in a distortion of shape caused by constriction around the optic stalk (Fig. 5H). We also observe some degree of pitting in the posterior surface of the MB-ca (Fig. 5I). This pitting is related to radially arranged columnar domains that are apparent within the calycal neuropil (Fig. 5J,K). We do not observe any structure clearly identifiable as an accessory calyx. We do see a brightly stained globular neuropil below the MB-ca/pe junction but it is quite some distance away from the junction and lacks the 'spotty' appearance of the accessory calyx in *D. plexippus* (Heinze and Reppert, 2012). It seems more likely that this structure is a 'satellite' neuropil that is not part of the MB (Farris, 2005). Its position corresponds roughly to the medial end of the expanded OG in *D. plexippus*. In some preparations one can follow a narrow faint fiber tract from here to an area of more intense staining in the optic stalk and on to the medial margin of the OG. If this is a functional connection, it is conceivable that the medial expansion of the OG in *D. plexippus* occurred along this pre-existing pathway.

Interspecific divergence in brain composition and mushroom body expansion in
Heliconius

After correcting for allometric scaling by phylogenetically-corrected regressions
against total neuropil volume, the six lepidopteran species can be separated along the
first two principal components that together explain 90.7% of variance. PC1 (65.9%
of Var) is heavily loaded by sensory neuropil in one direction, and the MB-ca and
MB-lo+ped in the other (Table 2). PC2 (24.8% of Var) is heavily loaded by the Me in
one direction and the AL and CB in the other. This roughly separates the six species
into three pairs, representing (i) *H. hecale* and *H. erato*; (ii) the other diurnal
butterflies, *D. plexippus* and *G. zavaleta*; and (iii) the night-flying moths, *H. virescens*
and *M. sexta* (Fig. 6B). When midbrain neuropil are analyzed separately, PC1 (68.7%
of Var) marks an axis dominated by AL, CB and MB, and PC2 (23.3% of Var) is
strongly loaded by the AOTu (Fig. 6C). This leads to two clusters grouping (i) *H.
hecale* and *H. erato*, which invest heavily in the mushroom body neuropil, and (ii) the
night-flying moths and *G. zavaleta*, which invest heavily in olfactory neuropil;
leaving *D. plexippus* isolated by its large AOTu volume.

The combined volume of the calyx, pedunculus and lobes account for 13.7%
of total brain neuropil volume in *H. erato*, and 11.9% in *H. hecale*. This is much
larger than reported for any other Lepidoptera measured with similar methods (range
2.3–5.1%). Expressed as a percentage of the midbrain, to remove the effects of
variation in the large OL, *H. erato* (38.5%) and *H. hecale* (32.9%) again exceed other
Lepidoptera (4.8–13.5%) by 3–7 fold. These figures are also much larger than
reported for *H. charithonia* (4.2% of total brain size) by Sivinski (1989), whose
figures for other Lepidoptera are also much lower suggesting the difference is
explained by variation in methodology.

Beyond Lepidoptera, the most comparable data available are from *Apis
mellifera* (Brandt et al., 2005) and *Schistocerca gregaria* (Kurylas et al., 2008) for
which mushroom body volume and midbrain volume are reported (Fig. 6D). In terms
of raw volume (Table 1) *Heliconius* mushroom bodies are roughly equal in size to *A.
mellifera*. However, in *A. mellifera* the mushroom bodies comprise 65.4% of the
midbrain, (40.6% MB-ca, 24.8% MB-lo+ped) (Brandt et al., 2005), in gregarious-
phase *S. gregaria* they comprise 15.1% (8.2% MB-ca including the accessory calyx,
6.3% MB-lo+ped) (Kurylas et al., 2008). Further comparisons can be made

expressing mushroom body size as a percentage of segmented neuropil (Me+Lobula

system, CB, MB and AL) that were labeled across a wider range species. As a ratio of

percentage mushroom body volume to the percentage of the two other midbrain

neuropil (AL and CB), *Heliconius* have ratios (*H. erato*: 6.4; *H. hecale*: 6.7) that

exceed even *A. mellifera* (3.8). To account of the dominant effect of OL size on

scaling with overall brain size we also analysed residual variance from a PGLS

regression (Fig. 6E) between percentage OL and percentage MB volume. This shows

Heliconius (*H. erato* +8.2; *H. hecale* +7.5) have the second largest residual MB size

following *A. mellifera* (+11.9).

Brain : body allometry

In both species, larger wild individuals have larger brains when using total neuropil

volume and either body length or wingspan as measures of brain and body size (log-

log SMA regression, *H. hecale,* body length $p = 0.020$; wingspan $p = 0.019$; *H. erato,*

body length $p = 0.011$; wingspan $p = 0.010$). The brain size : body *mass* relationship

is not significant in wild individuals (*H. hecale, $p = 0.055$*; *H. erato, $p = 0.863$*), most

likely because body mass varies much with reproductive state and feeding condition.

We therefore used body length as a proxy for body size to analyze the effect of age

and experience on the relative size of the brain.

Both species showed a clear grade-shift with age towards increased relative

brain size (*H. hecale:* Wald $\chi^2 = 5.780$, $p = 0.016$; *H. erato:* Wald $\chi^2 = 10.124$, $p =$

0.001). Body length was very similar in old and young individuals (*H. hecale* $t_{18} = -$

0.918, $p = 0.371$; *H. erato* $t_{17} = 0.581$, $p = 0.568$) suggesting the effect reflects an

increase in absolute neuropil volume. Indeed, old individuals had significantly larger

absolute midbrain volumes in both species (*H. erato:* $t_{17} = 4.192$, $p = 0.001$, $r = 0.713$;

H. hecale: $t_{18} = 3.054$, $p = 0.007$, $r = 0.595$; Fig. 7A,D). An absolute increase in OL

and total brain volume, however, was strongly supported only in *H. erato* (OL: $t_{17} =$

5.076, $p < 0.001$, $r = 0.776$; total, $t_{17} = 5.153$, $p < 0.001$, $r = 0.708$) and not evident in

H. hecale (OL, $t_{18} = 0.280$, $p = 0.783$; total, $t_{18} = 1.082$, $p = 0.293$).

Only *H. hecale* showed a clear response in overall brain size to experience.

The total neuropil was 40% larger in wild-caught than in old insectary-reared

individuals ($t_{17} = 2.553$, $p = 0.020$, $r = 0.526$) driven by a significant difference in

midbrain volume ($t_{17} = 3.658$, $p = 0.002$, $r = 0.664$), but not OL volume ($t_{18} = 1.728$, p

$= 0.101$; Fig. 7D). Although there was no matching difference in body length ($t_{18} =$

0.983, p = 0.436), a grade-shift towards larger relative brain size in wild *hecale* was

not supported (Wald χ^2 = 2.058, p = 0.151). However, we do observe a grade-shift

when the midbrain is analyzed separately (Wald χ^2 = 4.725, p = 0.030). No significant

brain or body size differences were found between wild and old insectary-reared

individuals in *H. erato* (total neuropil: t_{17} = -0.432, p = 0.671; midbrain: t_{17} = -0.732, p

=0.474; OL: t_{17} = -0.123, p = 0.904; body length: t_{17} = 1.009, p = 0.327; Fig. 7A).

Post-eclosion growth in the volume of individual neuropil regions

The age-related increase in overall absolute brain size in *H. erato* was reflected in

volumetric increases in nearly all brain regions, with only the OG failing to show a

significant expansion in old individuals (Table 3A). There was some evidence for

age-related differences in the allometric scaling coefficients for aMe and PB, and for

grade-shifts in OG and POTu, but these were weak relative to the strong major axis

shifts observed for all neuropils investigated (Table 3A). The largest shifts were

observed for the POTu (difference in fitted-axis mean, Δ_{FA} = 0.604), aME (Δ_{FA} =

0.536), MB-ca (Δ_{FA} = 0.496) and MB-lo+ped (Δ_{FA} = 0.393; Fig. 8A-C).

In contrast, in *H. hecale,* age-related size increases in volume were confined to

the midbrain and not all segmented midbrain regions showed the same pattern of

expansion; the rMid, components of the mushroom body complex, central complex

and AL were all significantly larger in old individuals, but the AOTu, POTu and all

optic lobe neuropil were not (Table 3B). Neuropil expansion appears to occur in a co-

ordinated manner, such that the allometric relationship between each neuropil and

rMid is maintained (Table 3B). The only exceptions were the La, Me and OG, which

showed significant grade-shifts towards a reduced volume relative to rMid in old

individuals. All other segmented neuropil showed major-axis shifts along a common

slope towards higher values in old individuals (Table 3B). The largest shifts were

observed in the mushroom body (MB-ca, Δ_{FA} = 0.279; MB-lo+ped, Δ_{FA} = 0.250;

Fig. 8A1–C1).

Experience-dependent plasticity in neuropil volume

Although wild *H. erato* do not have significantly larger absolute volumes for any

measured neuropil (Table 4A), differences in allometric scaling or grade-shifts

between wild and old insectary-reared individuals are nevertheless evident. Altered

scaling affects the MB-ca, Lop, and PB, all of which show shallower scaling

relationships (smaller β) with rMid in wild-caught individuals (Table 4A; Figure 7B,C). The MB-lo+ped shows an unambiguous grade-shift towards larger size in wild whilst maintaining a common slope, and also shows a major axis shift ($\Delta_{FA} = 0.250$; Fig. 8B1).

In *H. hecale* wild individuals have significantly larger total midbrains ($t_{18} = 3.658$, p = 0.002). The only segmented neuropil to reflect this difference, however, are the MB-ca and MB-lo+ped (Table 4B; Fig. 8A2,C2), while the rMid is also larger in wild individuals ($t_{18} = 3.417$, p = 0.003). The average MB-ca volume of old insectary-reared individuals is only 68.3% of the average wild MB-ca volume, for the young insectary-reared individuals it is 49.3% (Figure 8A2,C2). For MB-lo+pe these figures are 76.9% and 58.7% respectively (Figure 8A2,B2). For comparison, in *H. erato* the average MB-ca volume of old insectary-reared individuals is 96.2% of the average wild MB-ca volume, for the young insectary-reared individuals it is 59.7% (Fig. 8A1–C1). For MB-lo+pe these figures are 96.9% and 63.9% respectively (Fig. 8A1–C1).

The only neuropil in the optic lobes to differ significantly in volume in *H. hecale* is the Me. The allometric relationship between neuropil volumes and rMid differs for all neuropil either in the allometric scaling coefficient or the intercept, except for the mushroom body components and aMe (Table 4A; Figure 7E,F). However, for aME this pattern is caused by a lack of allometric scaling in insectary-reared individuals (SMA p = 0.552). The mushroom bodies show evidence of a major axis shift along a common slope (MB-ca, $\Delta_{FA} = 0.355$; MB-lo+ped, $\Delta_{FA} = 0.299$; Fig 8B2, C2). Given all grade-shifts result in smaller neuropil volumes relative to rMid volume (Fig. 7E,F) we interpret this as indicating the rMid and mushroom bodies show coordinated environment-dependent increases in volume whilst other neuropil volumes remain largely constant, but with subsequently altered allometric relationships with rMid.

Allometric scaling of mushroom body components

We further explored the allometric scaling relationships between the three main mushroom body components, the MB-lo and MB-pe (analyzed separately), and the MB-ca. Within wild caught individuals, pairwise comparisons between these structures do not reveal any significant deviation from isometric scaling relationships (test $\beta \neq 1$, p > 0.05). However, the ontogenetic growth we observe between the

young and old groups of both species occur through concerted expansion of the MBlo
and MBca (i.e. a major axis shift), both of which show grade-shifts in their allometric
scaling with the MBpe between the young and old groups (Table 5A). A similar
pattern is found comparing *H. hecale* wild and old groups, but there are no significant
differences between wild and old *H. erato* with the exception of a narrowly
significant difference in the scaling coefficient suggesting MB-lo becomes
disproportionally larger as MB-ca increases in wild individuals compared to insectary
reared individuals (Table 5B).

DISCUSSION

We have described the layout and volume of the major brain neuropils in two species
of *Heliconius* butterflies. Our analyses illustrate the role ecology plays in shaping
brain structure, and confirm the substantial evolutionary expansion of the *Heliconius*
mushroom body first noted by Sivinski (1989). Indeed, our data suggest this previous
work underestimated their size. We have further identified neuropil-specific patterns
of volumetric variation across young and old insectary-reared and wild individuals
that indicate significant post-eclosion growth and experience-dependent plasticity. In
the mushroom body, the timing and extent of this ontogenetic plasticity is comparable
to that found in insects that strongly rely on spatial memory for foraging (e.g. Withers
et al., 1993, 2008; Gronenberg et al., 1996; Fahrbach et al., 1998, 2003; Maleszka et
al., 2009; Jones et al., 2013).

Interspecific divergence and mushroom body expansion in **Heliconius**

Our interspecific analyses across Lepidoptera reveal an unambiguously mosaic
pattern of brain evolution (Barton and Harvey, 2000), where the size of individual
neuropils deviate from the allometric expectation. Mosaic patterns in mammals, fishes
and ants have been interpreted as strong evidence for evolutionary responses to a
species' particular ecological needs (Barton et al., 1995; Huber et al., 1997;
Gronenberg and Hölldobler, 1999). In Lepidoptera, this is particularly noticeable in
the sensory neuropils (Fig. 6B). The relative volume of the visual neuropils closely
reflects diel activity patterns, and the size of the antennal lobes also appears to be
strongly associated with a nocturnal or low-light diurnal niche. This is illustrated in a
PCA of midbrain neuropil (Fig. 6C) that clusters the olfactorily driven butterfly *G.*

zavaleta with night-flying moths (Montgomery and Ott, 2015). Our data further
indicate that much of the divergence in AL size among Lepidoptera reflects changes
in CFN volume rather than total glomerular volume, implying that changes in the
number or branching complexity of AL interneurons dominate over numerical
differences in afferent sensory neuron supply, and associated sensitivity. Similarly,
the relative constancy in AL glomeruli number indicates that the dimensionality of the
afferent coding space is comparable across Lepidoptera with divergent diel patterns
(Boeckh and Boeckh, 1979; Rospars, 1983; Berg et al., 2002; Huetteroth and
Schachtner, 2005; Masante-Roca et al., 2005; Skiri et al., 2005; Kazawa et al., 2009;
Heinze and Reppert, 2012; Carlsson et al., 2013; Montgomery and Ott, 2015).

In contrast with other species-differences that are dominated by changes in the
sensory neuropils, *Heliconius* are clearly set apart in our multivariate analysis along
an axis heavily loaded by the mushroom bodies. As a percentage of total brain
volume, or indeed as a raw volume, *Heliconius* have the largest mushroom body so
far reported in Lepidoptera (Sivinski, 1989; Sjöholm et al., 2005; Rø et al., 2007;
Kvello et al., 2009; Snell-Rood et al., 2009; Dreyer et al., 2010; Heinze and Reppert,
2012b; Montgomery and Ott, 2015) and one of the largest across insects. This
phylogenetic expansion must reflect adaptive change in mushroom body function in
response to ecological selection pressures. The derived pollen-feeding behavior of
Heliconius provides a likely source of this selection (Sivinski, 1989). Several studies
have reported this behavior to entail spatially and temporally faithful foraging
patterns, guided by visual landmarks (Ehrlich and Gilbert, 1973; Gilbert, 1975, 1993;
Mallet, 1986) comparable with the landmark-based trap-lining behavior of foraging
bees. Experimental interventions (Mizunami et al., 1998) and comparative neuro-
ecological studies (Farris and Schulmeister, 2011) implicate mushroom bodies in
visually based spatial memory.

Comparisons across *Heliconius* and non-pollen feeding Heliconiini may
provide a test of this spatial memory hypothesis. Sivinski (1989) reported that two
individuals of *Dione juno* and *Dryas iulia*, both non-pollen feeding allies to
Heliconius, had mushroom bodies within the size range of other Lepidoptera. This
provides preliminary support that mushroom body expansion coincided with a single
origin of pollen feeding at the base of *Heliconius*. However, sampling in a wider
range of genera, including the specious *Eueides* which is most closely related to

Heliconius (Beltrán et al., 2007; Kozak et al., 2015), is required to confirm this
conclusion.

Alternative selection pressures also need to be considered, including the
degree of host-plant specialization (Brown, 1981) and the evolution of social roosting
(Benson, 1972; Mallet, 1986). These factors may well be inter-related, as visits to
Passiflora may be incorporated into trap-lines between pollen plants (Gilbert, 1975,
1993), and the sedentary home-range behavior required for trap-lining may predispose
Heliconius to sociality (Mallet, 1986). The latter scenario would parallel the
hypothesized origin of sociality in hymenoptera and primates in exaptations of an
expanded brain that may have first evolved to support specialization in foraging
behavior (Barton, 1998; Farris and Schulmeister, 2011). Regardless of whether pollen
feeding provided the initial selection pressure for mushroom body expansion, it is
likely that it contributes to meeting the energetic costs of increased neural investment.
649

Age- and experience-dependent growth in neuropil volume
In both *H. erato and H. hecale*, the mushroom bodies are significantly larger in aged
individuals. Volume increases of 38.0% for the calyx and 34.0% for the lobe system
in *H. erato,* and 27.9% for the calyx and 23.7% for the lobes in *H. hecale* are
comparable to, if not greater than, the ontogenetic changes seen in Hymenoptera (e.g.
c. 30% in *Camponotus floridanus* (Gronenberg et al., 1996); c. 20% in *Bombus*
impatiens (Jones et al., 2013)). Our comparisons between aged insectary-reared and
wild caught individuals also identify experience-dependent plasticity. This
'experience' in the wild likely includes greater range of movement, greater challenges
in foraging, and more variable environmental conditions and social interactions.

Our data suggest experience-dependent plasticity particularly affects
mushroom body maturation, though the pattern differs between species. In *H. hecale* a
strong volumetric difference is found between old insectary-reared and wild caught
individuals for both the calyx (32%) and lobes (24%). A concomitant expansion of
the unsegmented midbrain results in a pronounced major-axis shift. This is not simply
the result of an increased total brain size, however: no other neuropil region shows a
comparable increase in wild caught individuals, resulting in widespread grade-shifts
in these other neuropils towards smaller size relative to the unsegmented midbrain.
This may reflect a coordinated growth between the mushroom bodies and
unsegmented midbrain areas or, alternatively, coincident independent expansions. In

H. erato old insectary-reared and wild-caught individuals have mushroom bodies of

similar absolute size, but allometric grade-shifts over the unsegmented midbrain result

in greater relative volumes in wild compared to insectary-reared individuals. The

cause of this species difference is unclear, but warrants further investigation.

Finally, it is also notable that plasticity, in particularly age-related growth, is

not restricted to the mushroom bodies. Several visual and olfactory neuropils show

age- and experience-dependent expansions in *Heliconius,* as they do in other insects

(Kühn-Bühlmann and Wehner, 2006; Snell-Rood et al., 2009; Ott and Rogers, 2010;

Smith et al., 2010; Heinze and Florman, 2013; Jones et al., 2013). We also find

evidence of plasticity in components of the central complex. In *D. plexippus*, size

plasticity in the central body and protocerebral bridge has been linked to migratory

experience and, by association, the sky compass navigation that supports it (Heinze et

al., 2013). The occurrence of similar plasticity in non-migratory butterfly species

implies that it may be associated with foraging or locomotor experience more

generally, even at much smaller spatial scales.

Functional relevance of phylogenetic mushroom body expansion

Phylogenetic trends towards larger mushroom bodies involve increases in Kenyon

Cell (KC) numbers, clustered into larger numbers of functional sub-units (Farris,

2008). Farris and Roberts (2005) suggest that increasing KC number may provide

greater computational capacity by facilitating the processing of more complex

combinatorial inputs from afferent projection neurons (Sivan and Kopell, 2004), or

through integration across increasingly specialized sub-units (Strausfeld, 2002).

Novel pathways between such specialized KC sub-populations may play an

important role in the origin of derived behaviors that require the integration of

different sensory modalities (Chittka and Niven, 2009; Strausfeld et al., 2009).

Examples of this are provided by Hymenoptera and phytophagous scarab beetles

where, in addition to olfactory inputs, the mushroom body calyx receives direct input

from the optic lobes (Gronenberg, 2001; Farris and Roberts, 2005; Farris and

Schulmeister, 2011). This additional input is reflected in the subdivision of the calyx

into the lip, which processes olfactory information, and the collar and basal ring,

which process visual information (Gronenberg and Hölldobler, 1999). Visual input to

the mushroom bodies has also been demonstrated in some butterflies (Snell-Rood et

al., 2009) and moths (Sjöholm et al., 2005) but it has yet to be investigated in

Heliconius. The *Heliconius* calyx lacks the clear zonation observed in *D. plexippus*
that has been suggested to be analogous to the *A. mellifera* lip, collar and basal ring
(Heinze and Reppert, 2012). We do not interpret the lack of distinct zonation in
Heliconius as evidence against functional sub-division, as *Spodoptera littoralis*
displays localization of visual processing in the calyx that is not apparent without
labeling individual neurons. Given the implied role for visual landmark learning in
Heliconius foraging behavior (Jones, 1930; Gilbert, 1972, 1975; Mallet, 1986), we
hypothesise that their massively expanded mushroom body supports an integration of
visual information.

In other species the mushroom body also receives gustatory and
mechanosensory input (Schildberger, 1983; Homberg, 1984; Li and Strausfeld, 1999;
Farris, 2008). These may also be of relevance in *Heliconius* given the importance of
gustatory and mechanosensory reception in host-plant identification (Schoonhoven,
1968; Renwick and Chew, 1994; Briscoe et al., 2013) and pollen loading (Krenn and
Penz, 1998; Penz and Krenn, 2000), although it should be noted that there is currently
no evidence these behaviors are learnt (Kerpel and Moreira, 2005; Salcedo, 2011;
Silva et al., 2014).

Potential cellular changes associated with ontogenetic mushroom body expansion
The cellular basis of ontogenetic and environmentally induced plasticity may provide
further clues as to the functional changes associated with mushroom body expansion
during *Heliconius* evolution. The volumetric changes we observe must reflect
differences in cell numbers and/or branching and connectivity. It is unknown whether
KC neurogenesis is restricted to the larval and pupal stages in Lepidoptera, as it is in
Hymenoptera (Fahrbach et al., 1995) where post-eclosion expansion results solely
from increased neurite branching (Gronenberg et al., 1996; Farris et al., 2001). In
Hymenoptera, age-dependent expansion of the MB-ca accompanies growth of
extrinsic neuron processes, whilst increased branching complexity of KCs is
associated experience-dependent expansion and foraging specialization in social
castes (Farris et al., 2001; Jones et al., 2009). This suggests that changes in the calyx
circuitry involving increased synaptic connections onto KC dendrites may be
responsible for the volumetric changes associated with behavioral experience.
Although we have not yet measured KC number or dendritic branch length, the grade-
shifts of MB-ca and MB-lo over MB-pe, that are uncovered by our allometric

analyses, indicate that post-eclosion growth is not solely (if at all) due to additional KCs. This is because each additional KC will necessarily contribute volume to all three major MB compartments, MB-ca, MB-pe and MB-lo. Indeed, Ott and Rogers (2010) proposed that wiring overheads might increase non-linearly with increasing KC numbers to explain the positive allometric scaling of MB-ca over MB-lo observed in some other insects. The resultant scaling need not be isometric, however, this effect will always produce a constant allometric scaling relationship. In contrast, the grade-shifts we observe are most likely explained by increased dendritic growth and connectivity in the MB-ca and MB-lo. Confirming this interpretation, and understanding its functional relevance, may provide some insight into how environmental information is stored during post-eclosion development.

Conclusions

Our volumetric analyses uncover extensive phylogenetic expansion and ontogenetic plasticity of *Heliconius* mushroom bodies. Both processes may be linked to the derived foraging behavior or this genus, which relies on allocentric memory of pollen resources (Gilbert, 1975; Sivinski, 1989). This hypothesis must now be confirmed in wider comparative analyses and tested explicitly in behavioral experiments. Our phenotypic observations furthermore provide the necessary framework for analyses of the underpinning neuronal mechanisms regulating neuropil size, and of the consequences for circuit function.

Acknowledgments

The authors are grateful to Adriana Tapia, Moises Abanto, William Wcislo, Owen McMillan, Chris Jiggins, and the Smithsonian Tropical Research Institute for assistance, advice, and the use of the *Heliconius* insectaries at Gamboa, Panama, and the Ministerio del Ambiente for permission to collect butterflies in Panama. We also thank Judith Mank's research group at UCL for helpful advice and feedback, and the UCL Imaging Facility for help with confocal microscopy.

Conflict of interest statement

The authors declare no conflict of interest.

Role of authors

All authors read and approved the final manuscript. Study conception: SHM. Study design and preliminary experiments: SHM, RMM, SRO. Fieldwork and insectary rearing: SHM, RMM. Acquisition of data, analysis, interpretation and initial manuscript draft: SHM. Final interpretation and drafting: SHM, RMM, SRO.

Literature cited

Barton RA. 1998. Visual specialization and brain evolution in primates. Proc Biol Sci 265:1933–1937.

Barton RA, Purvis A, Harvey PH. 1995. Evolutionary radiation of visual and olfactory brain systems in primates, bats and insectivores. Philos Trans R Soc Lond B Biol Sci 348:381–92.

Beltrán M, Jiggins CD, Brower AVZ, Bermingham E, Mallet J. 2007. Do pollen feeding , pupal-mating and larval gregariousness have a single origin in *Heliconius* butterflies: Inferences from multilocus DNA sequence data. :221–239.

Benson WW. 1972. Natural selection for Miillerian mimicry in *Heliconius erato* in Costa Rica. Science (80) 176:936–939.

Berg BG, Galizia CG, Brandt R, Mustaparta H. 2002. Digital atlases of the antennal lobe in two species of tobacco budworm moths, the Oriental *Helicoverpa assulta* (male) and the American *Heliothis virescens* (male and female). J Comp Neurol 446:123–134.

Boeckh J, Boeckh V. 1979. Threshold and odor specificity of pheromone-sensitive neurons in the deutocerebrum of *Antheraea pernyi* and *A. polyphemus* (Saturnidae). J Comp Physiol A 132:235–242.

Boggs CL, Smiley JT, Gilbert LE. 1981. Patterns of pollen exploitation by *Heliconius* butterflies. Oecologia 48:284–289.

Brandt R, Rohlfing T, Rybak J, Krofczik S, Maye A, Westerhoff M, Hege H-C, Menzel R. 2005. Three-dimensional average-shape atlas of the honeybee brain and its applications. J Comp Neurol 492:1–19.

Briscoe AD, Macias-Muñoz A, Kozak KM, Walters JR, Yuan F, Jamie GA, Martin SH, Dasmahapatra KK, Ferguson LC, Mallet J, Jacquin-Joly E, Jiggins CD. 2013. Female behaviour drives expression and evolution of gustatory receptors in butterflies. PLoS Genet 9:e1003620.

Brown KS. 1981. The biology of *Heliconius*. Annu Rev Entomol 26:427–457.

Capaldi EA, Robinson GE, Fahrback SE. 1999. Neuroethology of spatial learning: the
birds and the bees. Annu Rev Psychol 50:651–682.

Cardoso MZ, Gilbert LE. 2013. Pollen feeding, resource allocation and the evolution
of chemical defence in passion vine butterflies. J Evol Biol 26:1254–60.

Carlsson MA, Schäpers A, Nässel DR, Janz N. 2013. Organization of the olfactory
system of nymphalidae butterflies. Chem Senses 38:355–67.

Chittka L, Niven J. 2009. Are Bigger Brains Better? Curr Biol 19:R995–R1008.

Cohen J. 1988. Statistical power analysis for the behavioral sciences. 2nd ed.
Hillsdale, NJ.: Lawrence Earlbaum Associates.

Dasmahapatra KK, Walters JR, Briscoe AD, Davey JW, Whibley A, Nadeau NJ,
Zimin AV, Hughes DST, Ferguson LC, Martin SH, Salazar C, Lewis JJ, Adler S,
Ahn S-J, Baker DA, Baxter SW, Chamberlain NL, Chauhan R, Counterman BA,
Dalmay T, Gilbert LE, Gordon K, Heckel DG, Hines HM, Hoff KJ, Holland
PWH, Jacquin-Joly E, Jiggins FM, Jones RT, Kapan DD, Kersey P, Lamas G,
Lawson D, Mapleson D, Maroja LS, Martin A, Moxon S, Palmer WJ, Papa R,
Papanicolaou A, Pauchet Y, Ray DA, Rosser N, Salzberg SL, Supple MA,
Surridge A, Tenger-Trolander A, Vogel H, Wilkinson PA, Wilson D, Yorke JA,
Yuan F, Balmuth AL, Eland C, Gharbi K, Thomson M, Gibbs RA, Han Y,
Jayaseelan JC, Kovar C, Mathew T, Muzny DM, Ongeri F, Pu L-L, Qu J,
Thornton RL, Worley KC, Wu Y-Q, Linares M, Blaxter ML, Ffrench-Constant
RH, Joron M, Kronforst MR, Mullen SP, Reed RD, Scherer SE, Richards S,
Mallet J, Owen McMillan W, Jiggins CD. 2012. Butterfly genome reveals
promiscuous exchange of mimicry adaptations among species. Nature 487:94–8.

Dreyer D, Vitt H, Dippel S, Goetz B, El Jundi B, Kollmann M, Huetteroth W,
Schachtner J. 2010. 3D standard brain of the Red Flour Beetle *Tribolium*
castaneum: A tool to study metamorphic development and adult plasticity. Front
Syst Neurosci 4:3.

Dunlap-Pianka H, Boggs CL, Gilbert LE. 1977. Ovarian dynamics in heliconiine
butterflies: programmed senescence versus eternal youth. Science (80) 197:487–
490.

Durst C, Eichmüller S, Menzel R. 1994. Development and experience lead to
increased volume of subcompartments of the honeybee mushroom body. Behav
Neural Biol 62:259–263.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and
high throughput. Nucleic Acids Res 32:1792–7.

Ehrlich PR, Gilbert LE. 1973. Population Structure and dynamics of the Tropical
butterfly *Heliconius ethilla*. Biotropica 5:69–82.

Estrada C, Jiggins CD. 2002. Patterns of pollen feeding and habitat preference among
Heliconius species. Ecol Entomol 27:448–456.

Fahrbach SE, Farris SM, Sullivan JP, Robinson GE. 2003. Limits on volume changes
in the mushroom bodies of the honeybee brain. J Neurobiol 57:141–151.

Fahrbach SE, Giray T, Robinson GE. 1995. Volume changes in the mushroom bodies
of adult honey bee queens. Neurobiol Learn Mem 63:181–191.

Fahrbach SE, Moore D, Capaldi E a., Farris SM, Robinson GE. 1998. Experience-
expectant plasticity in the mushroom bodies of the honeybee. Learn Mem 5:115–
123.

Farris SM, Roberts NS. 2005. Coevolution of generalist feeding ecologies and
gyrencephalic mushroom bodies in insects. Proc Natl Acad Sci U S A
102:17394–17399.

Farris SM, Robinson GE, Fahrbach SE. 2001. Experience- and age-related outgrowth
of intrinsic neurons in the mushroom bodies of the adult worker honeybee. J
Neurosci 21:6395–6404.

Farris SM, Schulmeister S. 2011a. Parasitoidism, not sociality, is associated with the
evolution of elaborate mushroom bodies in the brains of hymenopteran insects.
Proc Biol Sci 278:940–951.

Farris SM. 2005. Evolution of insect mushroom bodies: Old clues, new insights.
Arthropod Struct Dev 34:211–234.

Farris SM. 2008. Tritocerebral tract input to the insect mushroom bodies. Arthropod
Struct Dev 37:492–503.

Farris SM. 2013. Evolution of complex higher brain centers and behaviors: behavioral
correlates of mushroom body elaboration in insects. Brain Behav Evol 82:9–18.

Finkbeiner SD. 2014. Communal roosting in heliconius butterflies (Nymphalidae):
roost recruitment, establishment, fidelity, and resource use trends based on age
and sex. J Lepid Soc 68:10–16.

Gilbert LE. 1972. Pollen feeding and reproductive biology of *Heliconius* butterflies.
Proc Natl Acad Sci U S A 69:1403–1407.

Gilbert LE. 1975. Ecological consequences of a coevolved mutualism between
butterflies and plants. In: Gilbert LE, Raven P, editors. Coevolution of animals
and plants. p 210–240.

Gilbert LE. 1993. An evolutionary food web and its relationship to Neotropical
biodiversity. In: Barthlott W, Naumann CM, Schmidt-Loske K, Schuchmann
KL, editors. Animal-Plant Interactions in Tropical Environments. Zoologisches
Forschungsinstitut und Museum Alexander Koenig, Bonn, Germany. p 17–28.

Godenschwege T a., Reisch D, Diegelmann S, Eberle K, Funk N, Heisenberg M,
Hoppe V, Hoppe J, Klagges BRE, Martin JR, Nikitina E a., Putz G, Reifegerste
R, Reisch N, Rister J, Schaupp M, Scholz H, Schwärzel M, Werner U, Zars TD,

Buchner S, Buchner E. 2004. Flies lacking all synapsins are unexpectedly
healthy but are impaired in complex behaviour. Eur J Neurosci 20:611–622.

Greenacre MJ. 2010. Biplots in practice. Fundacion BBVA.

Gronenberg W, Heeren S, Hölldobler B. 1996. Age-dependent and task-related
morphological changes in the brain and the mushroom bodies of the ant
Camponotus floridanus. J Exp Biol 199:2011–9.

Gronenberg W, Hölldobler B. 1999. Morphologic representation of visual and
antennal information in the ant brain. J Comp Neurol 412:229–240.

Gronenberg W. 2001. Subdivisions of hymenopteran mushroom body calyces by their
afferent supply. J Comp Neurol 435:474–489.

Heinrich B. 1979. Resource heterogeneity and patterns of movement in foraging
bumblebees. Oecologia 40:235–245.

Heinze S, Florman J, Asokaraj S, El Jundi B, Reppert SM. 2013. Anatomical basis of
sun compass navigation II: the neuronal composition of the central complex of
the monarch butterfly. J Comp Neurol 521:267–98.

Heinze S, Reppert SM. 2012b. Anatomical basis of sun compass navigation I: the
general layout of the monarch butterfly brain. J Comp Neurol 520:1599–628.

Hofbauer A, Ebel T, Waltenspiel B, Oswald P, Chen Y, Halder P, Biskup S,
Lewandrowski U, Winkler C, Sickmann A, Buchner S, Buchner E. 2009. The
Wuerzburg hybridoma library against *Drosophila* brain. J Neurogenet 23:78–91.

Homberg U. 1984. Processing of antennal information in extrinsic mushroom body
neurons of the bee brain. J Comp Physiol A Neuroethol Sens Neural Behav
Physiol:825–836.

Huber R, van Staaden MJ, S K Les, Liem KF. 1997. Microhabitat use, trophic
patterns, and the evolution of brain structure in African Cichlids. Brain Behav
Evol 50:167–182.

Huetteroth W, Schachtner J. 2005. Standard three-dimensional glomeruli of the
Manduca sexta antennal lobe: a tool to study both developmental and adult
neuronal plasticity. Cell Tissue Res 319:513–24.

Janzen ADH. 1971. Euglossine bees as long-distance pollinators of Tropical plants.
Science (80) 171:203–205.

Jones BM, Leonard AS, Papaj DR, Gronenberg W. 2013. Plasticity of the worker
bumblebee brain in relation to age and rearing environment. Brain Behav Evol
82:250–261.

Jones F. 1930. The sleeping heliconias of Florida. Nat Hist 30:635–644.

Jones TA, Donlan NA, O'Donnell S. 2009. Growth and pruning of mushroom body
Kenyon cell dendrites during worker behavioral development in the paper wasp,
Polybia aequatorialis (Hymenoptera: Vespidae). Neurobiol Learn Mem 92:485–
495.

El Jundi B, Huetteroth W, Kurylas AE, Schachtner J. 2009. Anisometric brain
dimorphism revisited: Implementation of a volumetric 3D standard brain in
Manduca sexta. J Comp Neurol 517:210–25.

Kazawa T, Namiki S, Fukushima R, Terada M, Soo K, Kanzaki R. 2009. Constancy
and variability of glomerular organization in the antennal lobe of the silkmoth.
Cell Tissue Res 336:119–36.

Kerpel SM, Moreira GRP. 2005. Absence of learning and local specialization on host
plant selection by *Heliconius erato*. J Insect Behav 18:433–452.

Klagges BR, Heimbeck G, Godenschwege TA, Hofbauer A, Pflugfelder GO,
Reifegerste R, Reisch D, Schaupp M, Buchner S, Buchner E. 1996. Invertebrate
synapsins: a single gene codes for several isoforms in *Drosophila*. J Neurosci
16:3154–65.

Kozak KM, Wahlberg N, Dasmahapatra KK, Mallet J, Jiggins CD. 2015. Multilocus
species trees show the recent adaptive radiation of the mimetic *Heliconius*
butterflies. Syst Biol. 64 (3): 505-524.

Krenn HW, Penz CM. 1998. Mouthparts of *Heliconius* butterflies (Lepidoptera :
Nymphalidae): A search for anatomical adaptations to pollen-feeding behavior.
Int J Insect Morphol Embryol 27:301–309.

Kühn-Bühlmann S, Wehner R. 2006. Age-dependent and task-related volume changes
in the mushroom bodies of visually guided desert ants, *Cataglyphis bicolor*. J
Neurobiol 66:511–521.

Kurylas AE, Rohlfing T, Krofczik S, Jenett A, Homberg U. 2008. Standardized atlas
of the brain of the desert locust, *Schistocerca gregaria*. Cell Tissue Res
333:125–45.

Kvello P, Løfaldli BB, Rybak J, Menzel R, Mustaparta H. 2009. Digital, three-
dimensional average shaped atlas of the *Heliothis virescens* brain with integrated
gustatory and olfactory neurons. Front Syst Neurosci 3:14.

Lent DD, Pinter M, Strausfeld NJ. 2007. Learning with half a brain. Dev Neurobiol
67:740–751.

Li Y, Strausfeld NJ. 1999. Multimodal efferent and recurrent neurons in the medial
lobes of cockroach mushroom bodies. J Comp Neurol 409:647–63.

Maleszka J, Barron AB, Helliwell PG, Maleszka R. 2009. Effect of age, behaviour
and social environment on honey bee brain plasticity. J Comp Physiol A
Neuroethol Sensory, Neural, Behav Physiol 195:733–740.

Mallet J. 1980. A laboratory study of gregarious roosting in the butterfly *Heliconius*
melpomene. MSc Thesis, University of Newcastle-upon-Tyne.

Mallet J. 1986. Gregarious roosting and home range in *Heliconius* butterflies. Natl
Geogr Res. 2(2): 198-215

Masante-Roca I, Gadenne C, Anton S. 2005. Three-dimensional antennal lobe atlas of
male and female moths, *Lobesia botrana* (Lepidoptera: Tortricidae) and
glomerular representation of plant volatiles in females. J Exp Biol 208:1147–59.

Menzel R. 2014. The insect mushroom body, an experience-dependent recoding
device. J Physiol Paris 108:84–95.

Mizunami M, Weibrecht JM, Strausfeld NJ. 1998. Mushroom bodies of the
cockroach: Their participation in place memory. J Comp Neurol 402:520–537.

Montgomery SH, Ott SR. 2015. Brain composition in *Godyris zavaleta* , a diurnal
butterfly, Reflects an increased reliance on olfactory information. J Comp Neurol
523:869–891.

Murawski DA, Gilbert LE. 1986. Pollen flow in *Psiguria warscewiczii*: a comparison
of *Heliconius* butterflies and hummingbirds. Oecologia 68:161–167.

Van Nouhuys S, Kaartinen R. 2008. A parasitoid wasp uses landmarks while
monitoring potential resources. Proc Biol Sci 275:377–385.

O'Brien DM, Boggs CL, Fogel ML. 2003. Pollen feeding in the butterfly *Heliconius*
charitonia: isotopic evidence for essential amino acid transfer from pollen to
eggs. Proc Biol Sci 270:2631–2636.

Ott SR, Rogers SM. 2010. Gregarious desert locusts have substantially larger brains
with altered proportions compared with the solitarious phase. Proc Biol Sci
277:3087–96.

Ott SR. 2008. Confocal microscopy in large insect brains: zinc-formaldehyde fixation
improves synapsin immunostaining and preservation of morphology in whole-
mounts. J Neurosci Methods 172:220–30.

Pagel M. 1999. Inferring the historical patterns of biological evolution. Nature
401:877–84.

Penz CM, Krenn HW. 2000. Behavioral adaptations to pollen-Feeding in *Heliconius*
butterflies (Nymphalinae, Heliconiinae): An experiment using *Lantana* flowers.
J Insect Behav 13 :865–880.

Pfeiffer K, Homberg U. 2014. Organization and functional roles of the central
complex in the insect brain. Annu Rev Entomol 59:165–84.

Rein K, Zöckler M, Mader MT, Grübel C, Heisenberg M. 2002. The *Drosophila*
standard brain. Curr Biol 12:227–31.

Renwick JAA, Chew FS. 1994. Oviposition behavior in Lepidoptera. Annu Rev
Entomol 39:377–400.

Rø H, Müller D, Mustaparta H. 2007. Anatomical organization of antennal lobe
projection neurons in the moth *Heliothis virescen*s. J Comp Neurol 500:658–75.

Rosenheim J a. 1987. Host location and exploitation by the cleptoparasitic wasp
Argochrysis armilla: the role of learning (Hymenoptera: Chrysididae). Behav
Ecol Sociobiol 21:401–406.

Rospars JP. 1983. Invariance and sex-specific variations of the glomerular
organization in the antennal lobes of a moth, *Mamestra brassicae*, and a
butterfly, Pieris brassicae. J Comp Neurol 220:80–96.

Salcedo C. 2011. Pollen preference for *Psychotria* sp. is not learned in the passion
flower butterfly, *Heliconius erato*. J Insect Sci 11:25.

Schildberger K. 1983. Local interneurons associated with the mushroom bodies and
the central body in the brain of *Acheta domesticus*. Cell Tissue Res 230:573–
586.

Schoonhoven LM. 1968. Chemosensory bases of host plant eelection. Annu Rev
Entomol 13:115–136.

Silva AK, Gonçalves GL, Moreira GRP. 2014. Larval feeding choices in heliconians:
Induced preferences are not constrained by performance and host plant
phylogeny. Anim Behav 89:155–162.

Sivan E, Kopell N. 2004. Mechanism and circuitry for clustering and fine
discrimination of odors in insects. Proc Natl Acad Sci U S A 101:17861–17866.

Sivinski J. 1989. Mushroom body development in nymphalid butterflies: A correlate
of learning? J Insect Behav 2:277–283.

Sjöholm M, Sinakevitch I, Ignell R, Strausfeld NJ, Hansson BS. 2005. Organization
of Kenyon cells in subdivisions of the mushroom bodies of a lepidopteran insect.
J Comp Neurol 491:290–304.

Skiri HT, Rø H, Berg BG, Mustaparta H. 2005. Consistent organization of glomeruli
in the antennal lobes of related species of heliothine moths. J Comp Neurol
491:367–80.

Smith AR, Seid M a, Jiménez LC, Wcislo WT. 2010. Socially induced brain
development in a facultatively eusocial sweat bee *Megalopta genalis*
(Halictidae). Proc Biol Sci 277:2157–2163.

Snell-Rood EC, Papaj DR, Gronenberg W. 2009. Brain size: a global or induced cost
of learning? Brain Behav Evol 73:111–28.

Strausfeld NJ, Sinakevitch I, Brown SM, Farris SM. 2009. Ground plan of the insect
mushroom body: functional and evolutionary implications. J Comp Neurol
513:265–91.

Strausfeld NJ. 2002. Organization of the honey bee mushroom body: Representation
of the calyx within the vertical and gamma lobes. J Comp Neurol 450:4–33.

Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011. MEGA5:
molecular evolutionary genetics analysis using maximum likelihood,
evolutionary distance, and maximum parsimony methods. Mol Biol Evol
28:2731–9.

Trautwein MD, Wiegmann BM, Beutel R, Kjer KM, Yeates DK. 2012. Advances in
insect phylogeny at the dawn of the postgenomic era. Annu Rev Entomol
57:449–468.

Turner JRG. 1971. Experiments on the demography of tropical butterflies. II.
Longevity and home-range behaviour in *Heliconius erato*. Biotropica 3:21–31.

Turner JRG. 1981. Adaptation and evolution in *Heliconius*: A defense of
NeoDarwinism. Annu Rev Ecol Syst 12:99–121.

Utz S, Huetteroth W, Vomel M, Schachtner J. 2008. Mas-Allatotropin in the
developing antennal lobe of the Sphinx Moth *Manduca sexta*: Distribution, time
course, developmental regulation, and colocalization with other neuropeptides.
Dev Neurobiol 68:123–142.

Warren AD, Davis KJ, Strangeland EM, Pelham JP, Grishin NV. 2013. Illustrated
lists of American butterflies. http://butterfliesofamerica.com

Warton DI, Duursma RA, Falster DS, Taskinen S. 2012. smatr 3: an R package for
estimation and inference about allometric lines. Methods Ecol Evol 3:257–259.

Wei H, el Jundi B, Homberg U, Stengl M. 2010. Implementation of pigment-
dispersing factor-immunoreactive neurons in a standardized atlas of the brain of
the cockroach *Leucophaea maderae*. J Comp Neurol 518:4113–33.

Withers GS, Day NF, Talbot EF, Dobson HEM, Wallace CS. 2008. Experience-
dependent plasticity in the mushroom bodies of the solitary bee *Osmia lignaria*
(Megachilidae). Dev Neurobiol 68:73–82.

Withers GS, Fahrbach SE, Robinson GE. 1993. Selective neuroanatomical plasticity
and division of labour in the honeybee. Nature 364:238–40.

Zars T. 2000. Behavioral functions of the insect mushroom bodies. Curr Opin
Neurobiol 10:790–795.

1060

1061

1062 ## Abbreviations

AL	antennal lobe
aMe	accessory medulla
AN	antennal nerve
AOTu	anterior optic tubercule
CB	central body
CBL	lower central body
CBU	upper central body
CFN	central fibrous neuropil of AL
DMSO	dimethyl suphoxide
Glom	glomeruli
HBS	HEPES-buffered saline
iMe	inner medulla
iRim	inner rim of the lamina
KC	kenyon cell
La	lamina
LAL	lateral accessory lobes
Lo	lobula
LoP	lobula plate
LU	lower unit of AOTu
MB	mushroom body
MB-ca	mushroom body calyx
MB-lo	mushroom body lobes
MB-pe	mushroom body peduncle
MB-lo+pe	mushroom body lobes and peduncle combined
MBr	midbrain
Me	medulla
MGC	macro-glomeruli complex
NGS	normal goat serum
no	noduli
NU	nodule unit of AOTu
oMe	outer medulla
OR	olfactory receptor
OGC	optic glomerular complex
PA	pyrrolizidine alkaloids
PB	protocerebral bridge
PC	principal component
POTu	posterior optic tubercle
rMid	rest of midbrain
SP	strap of AOTu
UU	upper unit of AOTu
ZnFA	Zinc-Formaldehyde solution

1063

1064

1065 **Tables**

1066

1067 **Table 1:** Neuropil volumes and body size of A) *H. erato* and B) *H. hecale*

1068

1069 **Table 2:** Loadings on Principal Components Analysis of the relative size of brain

1070 components across six Lepidoptera.

1071

1072 **Table 3:** Comparisons between old (O) and young (Y) insectary-reared individuals

1073 for A) *H. erato* and B) *H. hecale. r* is the effect size. DI indicates the group with a

1074 higher value of α, β or fitted axis mean.

1075

1076 **Table 4:** Comparisons between wild caught (W) and old insectary-reared individuals

1077 for A) *H. erato* and B) *H. hecale. r* is the effect size. DI indicates the group with a

1078 higher value of α, β or fitted axis mean.

1079

1080 **Table 5:** Effects of age (A) and environmental experience (B) on scaling relationships

1081 between mushroom body components

1082

1083

1084

1085

1086

1087

1088

Table 1

A) *H. erato*

	wild caught				old insectary reared		young insectary reared	
	mean ($n = 10$)	SD	Rel. SD (%)	% total neuropil	mean ($n = 10$)	SD	mean ($n = 10$)	SD
Body mass (g)	0.093	0.017	19.999	-	0.074	0.014	0.088	0.019
Body length (mm)	23.833	1.426	5.983	-	23.095	1.773	22.671	0.951
Wing span (mm)	71.408	3.278	4.591	-	69.744	4.12	68.786	2.55
La	7.409E+07	1.052E+07	14.192	13.459	6.95E+07	1.61E+07	5.49E+07	1.25E+07
Me	2.396E+08	3.617E+07	15.094	43.523	2.45E+08	2.76E+07	1.90E+08	3.32E+07
aMe	1.633E+05	3.609E+04	22.094	0.030	1.59E+05	4.61E+04	9.77E+04	1.93E+04
Lon	2.630E+07	4.203E+06	15.984	4.777	2.79E+07	2.89E+06	2.07E+07	4.32E+06
LoP	1.393E+07	2.083E+06	14.952	2.531	1.35E+07	2.22E+06	1.04E+07	2.07E+06
OG	1.054E+06	2.400E+05	22.769	0.191	1.05E+06	2.42E+05	8.85E+05	2.26E+05
AL	1.185E+07	2.450E+06	20.671	2.153	1.19E+07	2.49E+06	7.72E+06	1.10E+06
AOTu	2.199E+06	4.535E+05	20.618	0.400	2.26E+06	3.28E+05	1.52E+06	3.27E+05
MB-ca	4.672E+07	9.290E+06	19.886	8.486	4.50E+07	1.22E+07	2.79E+07	5.75E+06
MB-pe	6.043E+06	1.109E+06	18.343	1.098	6.15E+06	1.35E+06	5.57E+06	1.58E+06
MB-lo	2.267E+07	5.812E+06	25.641	4.118	2.17E+07	4.26E+06	1.28E+07	2.31E+06
CBL	3.017E+05	5.189E+04	17.198	0.055	2.83E+05	6.00E+04	2.24E+05	3.81E+04
CBU	1.180E+06	1.788E+05	15.153	0.214	1.17E+06	2.57E+05	8.90E+05	1.36E+05
Nod	2.966E+04	1.146E+04	38.631	0.005	3.09E+04	1.64E+04	3.16E+04	8.46E+03
PB	2.120E+05	4.804E+04	22.658	0.039	1.96E+05	5.04E+04	1.39E+05	2.02E+04
POTu	4.213E+04	9.976E+03	23.681	0.008	4.20E+04	1.43E+04	2.73E+04	7.93E+03
Total midbrain	1.954E+08	3.365E+07	17.222	35.490	2.04E+08	2.70E+07	1.39E+08	2.28E+07

B) *H. hecale*

	wild caught				old insectary reared		young insectary reared	
	mean (*n* = 10)	SD	Rel. SD (%)	% total neuropil	mean (*n* = 9)	SD	mean (*n* = 10)	SD
Body mass (g)	0.163	0.025	15.317	-	0.154	0.046	0.171	0.047
Body length (mm)	29.693	3.097	10.431	-	28.189	3.0631	29.206	2.75
Wing span (mm)	88.129	8.004	9.082	-	80.6	7.134	86.34	8.012
La	9.751E+07	1.826E+07	18.721	13.939	9.39E+07	2.17E+07	9.64E+07	1.50E+07
Me	2.986E+08	5.342E+07	17.888	42.689	2.48E+08	3.81E+07	2.42E+08	3.66E+07
aMe	1.660E+05	2.951E+04	17.782	0.024	1.40E+05	2.80E+04	1.38E+05	3.67E+04
Lon	3.056E+07	5.630E+06	18.422	4.369	2.80E+07	4.64E+06	2.45E+07	5.06E+06
LoP	1.648E+07	2.972E+06	18.031	2.356	1.45E+07	2.45E+06	1.27E+07	2.53E+06
OG	1.099E+06	3.396E+05	30.894	0.157	9.93E+05	2.12E+05	9.24E+05	2.10E+05
AL	1.216E+07	2.056E+06	16.905	1.739	1.09E+07	1.34E+06	9.36E+06	1.59E+06
AOTu	2.572E+06	6.144E+05	23.891	0.368	2.30E+06	4.46E+05	2.02E+06	3.76E+05
MB-ca	5.271E+07	1.611E+07	30.569	7.534	3.60E+07	7.49E+06	2.60E+07	7.48E+06
MB-pe	6.680E+06	1.525E+06	22.834	0.955	5.92E+06	1.30E+06	4.91E+06	1.39E+06
MB-lo	2.421E+07	6.279E+06	25.930	3.461	1.79E+07	3.56E+06	1.32E+07	3.51E+06
CBL	3.109E+05	6.362E+04	20.467	0.044	2.91E+05	7.15E+04	2.47E+05	3.74E+04
CBU	1.093E+06	2.026E+05	18.541	0.156	1.16E+06	2.05E+05	9.65E+05	1.79E+05
Nod	4.207E+04	1.713E+04	40.730	0.006	3.34E+04	8.35E+03	3.06E+04	1.28E+04
PB	2.424E+05	5.657E+04	23.335	0.035	2.00E+05	3.09E+04	1.64E+05	1.75E+04
POTu	4.183E+04	1.257E+04	30.057	0.006	3.74E+04	8.47E+03	3.20E+04	8.27E+03
Total midbrain	2.551E+08	6.253E+07	24.513	36.465	1.82E+08	2.28E+07	1.50E+08	2.25E+07

Table 2

A) Midbrain only

| Neuropil | Loadings | |
| | Residuals | |
	PC1	PC2
AL	-0.981	-0.045
CB L+U	-0.798	0.406
MB-ca	0.962	0.110
MB-lo+pe	0.952	0.231
AOTu	-0.047	0.966

B) Whole neuropil

| Neuropil | Loadings | |
| | Residuals | |
	PC1	PC2
AL	0.761	0.619
CB L+U	0.671	0.670
MB-ca	-0.961	0.212
MB-lo+pe	-0.942	0.222
AOTu	0.811	0.024
Me	0.042	-0.949
Lob	0.920	-0.354
LoP	0.962	-0.167

Table 3

A) *H. erato*

	Volume			Scaling coefficient (β)			Intercept (α)			Major Axis Shift		
	t_{17}	p	r (DI)	LR	p	r (DI)	Wald χ2	p	r (DI)	Wald χ2	p	r (DI)
La	2.432	0.026	0.508 (O)	2.019	0.155	-	2.895	0.089	-	13.196	0.000	0.833 (O)
Me	4.118	0.001	0.707 (O)	0.090	0.765	-	1.127	0.288	-	19.405	0.000	1.000 (O)
aME	3.802	0.001	0.678 (O)	3.976	0.046	0.458 (O)	-	-	-	-	-	-
Lob	4.173	0.001	0.711 (O)	0.246	0.620	-	1.587	0.208	-	20.284	0.000	1.000 (O)
LoP	3.266	0.005	0.621 (O)	0.523	0.470	-	3.802	0.051	-	19.034	0.000	1.000 (O)
OG	1.412	0.176	-	0.385	0.535	-	5.694	0.017	0.547 (Y)	10.622	0.001	0.748 (O)
AL	5.080	0.000	0.776 (O)	4.169	0.041	-	0.214	0.644	-	27.584	0.000	1.000 (O)
AOT	5.192	0.000	0.783 (O)	0.109	0.741	-	0.123	0.726	-	26.321	0.000	1.000 (O)
MB-ca	4.050	0.001	0.701 (O)	3.679	0.055	-	1.607	0.205	-	19.177	0.000	1.000 (O)
MB-lo+pe	4.806	0.000	0.759 (O)	0.963	0.326	-	0.373	0.541	-	23.250	0.000	1.000 (O)
CB L+U	3.272	0.004	0.622 (O)	2.364	0.124	-	1.807	0.179	-	16.530	0.000	0.933 (O)
PB	3.169	0.006	0.609 (O)	5.996	0.014	0.562 (O)	-	-	-	-	-	-
POTu	2.772	0.013	0.558 (O)	1.539	0.215	-	4.124	0.042	0.466 (Y)	14.953	0.000	0.887 (O)
Total Mid	4.192	0.001	0.713 (O)	-	-	-	-	-	-	-	-	-
rMid	5.771	0.000	0.814 (O)	-	-	-	-	-	-	-	-	-
Total OL	5.076	0.000	0.776 (O)	-	-	-	-	-	-	-	-	-
Total neuropil	5.153	0.000	0.781 (O)	-	-	-	-	-	-	-	-	-

B) *H. hecale*

	Volume			Scaling coefficient (β)			Intercept (α)			Major Axis Shift		
	t_{18}	p	r (DI)	LR	p	r (DI)	Wald χ2	p	r (DI)	Wald χ2	p	r (DI)
La	-0.424	0.677	-	1.866	0.172	-	11.902	0.001	0.771 (Y)	1.043	0.307	-
Me	0.333	0.743	-	0.494	0.482	-	21.674	0.000	1.000 (Y)	1.971	0.160	-
aME	0.238	0.814	-	0.094	0.759	-	3.044	0.081	-	3.088	0.079	-
Lob	1.538	0.141	-	0.002	0.961	-	3.544	0.060	-	4.501	0.034	0.474 (O)
LoP	1.683	0.110	-	0.066	0.797	-	1.577	0.209	-	5.031	0.025	0.502 (O)
OG	0.617	0.545	-	0.266	0.606	-	4.408	0.036	0.470 (Y)	3.045	0.081	-
				-								
AL	2.418	0.026	-	1.795	0.180	-	2.396	0.122	-	6.451	0.011	0.570 (O)
AOT	1.496	0.152	-	0.101	0.751	-	2.166	0.141	-	4.656	0.031	0.483 (O)
MB-ca	3.177	0.005	0.599 (O)	0.283	0.595	-	0.104	0.747	-	9.166	0.002	0.677 (O)
MB-lo+pe	2.707	0.014	0.538 (O)	0.147	0.702	-	0.015	0.902	-	7.594	0.006	0.616 (O)
CB L+U	2.218	0.040	0.463 (O)	3.291	0.070	-	0.859	0.354	-	7.221	0.007	0.601 (O)
PB	3.291	0.004	0.613 (O)	1.043	0.307	-	0.172	0.678	-	9.448	0.002	0.687 (O)
POTu	1.494	0.153	-	0.078	0.780	-	0.736	0.391	-	6.292	0.012	0.561 (O)
Total Mid	3.054	0.007	0.584 (O)	-	-	-	-	-	-	-	-	-
rMid	2.854	0.011	0.558 (O)	-	-	-	-	-	-	-	-	-
Total OL	0.280	0.783	-	-	-	-	-	-	-	-	-	-
Total neuropil	1.082	0.293	-	-	-	-	-	-	-	-	-	-

Table 4

A) *H. erato*

	Volume			Scaling coefficient (β)			Intercept (α)			Major Axis Shift		
	t_{17}	p	r (DI)	LR	p	r (DI)	Wald $\chi2$	p	r (DI)	Wald $\chi2$	p	r (DI)
La	0.892	0.385	-	3.685	0.055	-	3.605	0.058	-	0.210	0.646	-
Me	-0.426	0.676	-	0.269	0.604	-	2.056	0.152	-	1.269	0.260	-
aME	0.359	0.724	-	5.150	0.023	0.521 (O)	-	-	-	-	-	-
Lob	-1.056	0.306	-	0.283	0.595	-	1.004	0.316	-	1.999	0.157	-
LoP	0.430	0.673	-	4.963	0.026	0.511 (O)	-	-	-	-	-	-
OG	0.116	0.909	-	2.148	0.143	-	2.055	0.152	-	0.848	0.357	-
AL	-0.035	0.972	-	1.695	0.193	-	2.269	0.132	-	0.899	0.343	-
AOT	-0.490	0.631	-	0.483	0.487	-	1.318	0.251	-	1.456	0.227	-
MB-ca	0.511	0.616	-	5.833	0.016	0.554 (O)	-	-	-	-	-	-
MB-lo+pe	0.239	0.814	-	0.714	0.398	-	4.418	0.036	0.482 (W)	7.594	0.006	0.632 (W)
CB L+U	0.394	0.699	-	4.272	0.039	0.474 (O)	-	-	-	-	-	-
PB	0.845	0.410	-	4.413	0.036	0.482 (O)	-	-	-	-	-	-
POTu	0.196	0.847	-	3.726	0.054	-	2.730	0.098	-	0.905	0.341	-
Total Mid	-0.732	0.474	-	-	-	-	-	-	-	-	-	-
rMid	-1.787	0.092	-	-	-	-	-	-	-	-	-	-
Total OL	-0.123	0.904	-	-	-	-	-	-	-	-	-	-
Total neuropil	-0.432	0.671	-	-	-	-	-	-	-	-	-	-

B) *H. hecale*

	Volume			Scaling coefficient (β)			Intercept (α)			Major Axis Shift		
	t_{18}	p	r (DI)	LR	p	r (DI)	Wald χ^2	p	r (DI)	Wald χ^2	p	r (DI)
La	0.437	0.667	-	6.725	0.010	0.580 (O)	-	-	-	-	-	-
Me	2.293	0.034	0.475 (W)	9.165	0.002	0.677 (O)	-	-	-	-	-	-
aME	1.898	0.074	-	3.728	0.054	-	1.463	0.227	-	8.056	0.005	0.688 (W)
Lob	1.017	0.322	-	9.760	0.002	0.699 (O)	-	-	-	-	-	-
LoP	1.609	0.125	-	6.081	0.014	0.551 (O)	-	-	-	-	-	-
OG	0.614	0.547	-	4.262	0.039	0.462 (O)	-	-	-	-	-	-
AL	1.519	0.146	-	7.095	0.008	0.596 (O)	-	-	-	-	-	-
AOT	1.088	0.291	-	3.938	0.047	0.444 (O)	-	-	-	-	-	-
MB-ca	3.126	0.006	0.593 (W)	1.657	0.198	-	0.395	0.530	-	10.432	0.001	0.722 (W)
MB-lo+pe	2.536	0.021	0.513 (W)	3.759	0.053	-	1.603	0.205	-	8.811	0.003	0.664 (W)
CB L+U	-0.446	0.661	-	1.665	0.197	-	11.013	0.001	0.742 (O)	2.385	0.122	-
PB	1.919	0.071	-	5.043	0.025	0.502 (O)	-	-	-	-	-	-
POTu	0.551	0.588	-	5.420	0.020	0.521 (O)	-	-	-	-	-	-
Total Mid	3.658	0.002	0.653 (W)	-	-	-	-	-	-	-	-	-
rMid	3.417	0.003	0.627 (W)	-	-	-	-	-	-	-	-	-
Total OL	1.728	0.101	-	-	-	-	-	-	-	-	-	-
Total neuropil	2.553	0.020	0.516 (W)	-	-	-	-	-	-	-	-	-

Table 5

A) *Old vs young insectary reared*

	components	Scaling coefficient (β)			Intercept (α)			Major Axis Shift		
		Likelihood Ratio	p	r	Wald $\chi2$	p	r	Wald $\chi2$	p	r (DI)
H. erato	MB ca + lo	0.627	0.428	-	2.249	0.134		16.987	0.000	0.946 (O)
	MB ca + pe	1.224	0.269	-	12.457	0.000	0.810	-	-	-
	MB lo + pe	0.206	0.650	-	29.286	0.000	1.000	-	-	-
H. hecale	MB ca + lo	0.100	0.752	-	0.058	0.810		8.771	0.003	0.662 (O)
	MB ca + pe	0.376	0.540	-	6.422	0.011	0.567	-	-	-
	MB lo + pe	0.118	0.731	-	5.462	0.019	0.523	-	-	-

B) *Wild vs. old insectary reared*

	components	Scaling coefficient (β)			Intercept (α)			Major Axis Shift		
		Likelihood Ratio	p	r	Wald $\chi2$	p	r	Wald $\chi2$	p	r (DI)
H. erato	MB ca + lo	4.083	0.043	0.464	0.139	0.709	-	0.186	0.667	-
	MB ca + pe	0.311	0.577	-	0.732	0.392	-	0.044	0.834	-
	MB lo + pe	1.296	0.255	-	0.213	0.645	-	0.011	0.916	-
H. hecale	MB ca + lo	0.307	0.580	-	0.398	0.528	-	7.901	0.005	0.629 (W)
	MB ca + pe	2.942	0.086	-	7.340	0.007	0.606	-	-	-
	MB lo + pe	1.553	0.213	-	4.086	0.043	0.452	-	-	-

Figure 1: Overview of the anatomy of the Heliconius brain.

3D models of *H. erato* (A1–G1) and *H. hecale* (A2–G2). B1–D1 and B2–D2: Volume rendering of synapsin immunofluorescence showing the surface morphology of the brain neuropil from the anterior (A1/A2), posterior (B1/B2), and dorsal (C1/C2) view. E1–G1 and E2–G2: Surface reconstructions of the major neuropil compartments from the anterior (D1/D2), posterior (E1/E2), and dorsal (F1/F2) view. Neuropil in yellow-orange: visual neuropil, green: central complex, blue: antennal lobes, red: mushroom bodies. See Figures 2–4 for further anatomical detail. The individuals displayed are male. Images in A1/A2 are from Warren et al. (2013). Scale bars = 25 mm in A1/A2; 500 µm in B1–D1/B2–D2.

Figure 2: Anatomy of the major visual neuropils.

A: Surface reconstructions of the optic lobe neuropils viewed from anterior (left image) and posterior (right image). They comprise the lamina (La), the medulla (Me), and accessory medulla (aMe), the lobula (Lo), the lobula plate (LoP) and the optic glomerulus (OG). B: Surface reconstruction of the optic glomerulus (OG) viewed along the anterior-posterior axis (top) and an anterior view (bottom). C: Surface reconstruction of the anteriot optic tubercle (AOTu). D–J: Synapsin immunofluorescence in single confocal sections of the optic lobe of H. hecale. D: Horizontal section showing all four major optic lobe neuropils (La, Me, Lo, LoP). E: Frontal section showing the inner rim (iRim) of the lamina, a thin layer on its inner surface that is defined by intense synapsin immunofluorescence. Synapsin immunostaining also reveals the laminated structure of the medulla with two main subdivisions, the outer and inner medulla (oMe, iMe). F: The OG is located medially to the Lo; frontal section, the midbrain (MBr) occupies the left half of the frame. G,H: Frontal sections showing a small, irregular neuropil (ir) observed running from the anterior-ventral boundary of the aME as in *D. plexippus* (Heinze and Reppert, 2012). All images are from male *H. hecale*.

Figure 3: Anatomy of the antennal lobe

A: 3D reconstruction of individual antennal lobe (AL) glomeruli superimposed on a volume rendering of the anterior surface of the midbrain. B: Synapsin immunofluorescence in a single frontal confocal section showing the glomeruli (Glom) surrounding the central fibrous neuropil (CFN). Images A–B are from male *H. hecale*. C,D: Allometric grade-shifts between Glomerular (circles) or CFN (triangles) volume and unsegmented midbrain volume (C), and between Glomerular and CFN volume (D) in *G. zavaleta* (solid blue), *H. erato* (black filled with red) and *H. hecale* (orange filled with yellow). Scale bars = 500 μm in A; 50 μm in B,C,G,H; 100 μm in B–F, J; 200 μm in I.

Figure 4: Anatomy of the central complex

A1/A2: Surface reconstruction of the central complex from an anterolateral (A1) and oblique posteroventral (A2) view, showing the upper and lower subunit of the central body (CBU, CBL), the noduli (No), the protocerebral bridge (PB) and posterior optic tubercles (POTu). B–G: Synapsin immunofluorescence in single confocal sections. B: Horizontal section showing the upper and lower subunit of the CB in relation to the antennal lobes (AL) and the calyx (MB-ca) and pedunculus (MB-pe) of the mushroom body. C,D: Frontal confocal sections at the level of the CBL (C) and CBU (D); the CB subunits are flanked by the profiles of the vertically running MB-pe on either side. E: Frontal section showing the location of the PB ventrally to the MB-ca. F: POTu positioned ventrally to the MB-ca in a frontal section. G: Frontal section showing position of the paired No ventrally to CBL and CBU. All images are from a male H. hecale. Scale bars = 100 μm in B–D, G; 50 μm in E,F.

Figure 5: Anatomy of the mushroom body

A–C: Surface reconstruction of the mushroom body viewed orthogonal to the anterior-posterior axis from a medial vantage point level with the peduncle (A); from anterior (B); and from posterior (C). The main components are the calyx (MB-ca) shown in dark red, and the peduncule (MB-pe) and lobes (MB-lo) shown in bright red. A Y-tract, shown in magenta, runs parallel and slightly medial to the MB-pe. D–O: Synapsin immunofluorescence in individual confocal sections. D: anterior view of the midbrain showing the MB-lo, an asterik indicates what is most likely the ventral lobe, otherwise the individual lobes and loblets of the MB-lo are fused. E: Frontal section at a posterior level near the end of the MB-pe, showing the profiles of the MB-ca with its zonation into an outer and a medial ring. F,G and J,K: Horizontal confocal sections through the midbrain at increasing depths from dorsal towards ventral, showing MB structure in relation to neighboring neuropil: the anterior optic tubercle (AOTu in F,G); the antennal lobe (AL in G,J); and the central body upper division (CBU in K). H: An example of a female *H. erato* where the MB-ca is deformed due expansion into the optic lobe and constriction (labeled C) at the optic stalk by the neural sheath surrounding the brain. I: Pitted surface of the MB-ca in a very posterior tangential horizontal section. The pitting is related to what appear to be columnar domains within the calyx neuropil (cf. MB-ca in J,K,M). L: Areas of intense synapsin staining in the optic stalk (OS*); Lo, lobula; OG, optic glomerulus. M: Frontal section near the base of the calyx (MB-ca) showing a satellite neuropil (sat.) located near to the MB-pe. N: A Y-tract runs parallel with, and dorsally and slightly medially to the MB-pe; both are seen in profile in this frontal section. O: A fiber bundle (fb) connected to the AOTu running near the junction between the MB-pe and MB-lo. With the exception of I, all images are from a male *H. hecale*. Scale bars A-G, J-K = 200 µm, H-I, L-O = 100 µm.

Figure 6: Divergence in brain structure across Lepidoptera, and in mushroom body size across insects.

A: Phylogenetic relationships of Lepidoptera (red branches) and other insects (grey branches) for which directly comparable data are available. Branches are not drawn proportional to divergence dates, numbers refer to labels in panel E. B,C: Principal Component analysis of segmented neuropil volumes, corrected for allometric scaling with the unsegmented midbrain and phylogeny. B shows the results of an analysis using all neuropil. C shows the results of an analysis excluding the optic lobe neuropil. Species data points are indicated by the first letter of their genus and species name: D.p = *Danaus plexippus*; H.e = *Heliconius erato*; H.h = *H. hecale*; G.z = *Godyris zavaleta*; H.v = *Heliothis virescens*; M.s = *Manduca sexta*. D: The proportion of the midbrain occupied by MB-ca (dark red) and MB-lo+pe (light red) in four butterflies, and two other insects with fully comparable data. E: Across a wider sample of insects (shown in A), when expressed as a percentage of total volume of OL, AL, CB and MB, *Apis mellifera* (solid blue) and *Heliconius* (solid red) stand out as having expanded mushroom bodies, correcting for the size of the optic neuropil, compared to other Lepidoptera (unfilled red circles) and other insects (unfilled blue circles). The line was fitted by PGLS. All insect images in A are from Wikimedia commons and were released under the Creative Commons License, except *Heliconius* (see Fig. 1).

Figure 7: Age and environment dependent growth of brain components

A1,A2: Comparisons of raw volumes of total neuropil, total OL neuropil, total midbrain neuropil between wild-caught, old and young insectary-reared individuals of *H. erato* (A1) and *H. hecale* (A2). Significance of pair-wise comparisons are shown along the x-axis (young-old = orange; old-wild = dark red; n.s. = p>0.05, * = p<0.05, ** = p <0.01, *** = p < 0.001). B: Allometric scaling of LoP in *H. erato*. C: Allometric scaling of PB in *H. erato*. D: Allometric scaling of OG in *H. hecale*. E: Allometric scaling of CB in *H. hecale*. Note in E and F the shifts in allometry occur along the x-axis, this is explained by the large difference in unsegmented midbrain volume observed between wild-caught and old insectary-reared individuals in *H. hecale* as displayed in D.

Figure 8: Age and environment dependent growth of the mushroom bodies

Surface reconstruction of the mushroom body viewed along the anterior-posterior axis for wild-caught, old and young insectary-reared individuals of H. erato (A1) and H. hecale (A2). Representative individuals were chosen as those closest to the group mean volume. Scale bar = 200 μm. B1-C1/B2-C2: allometric relationships between MB-lo+pe (B1/B2), or MB-ca (C1/C2),and the volume of the unsegmented midbrain (rMid) for *H. erato* (B1/C1) and *H. hecale* (B2/C2). Data for wild caught individuals are in green, data for old insectary-reared individuals in dark blue, and data for young insectary-reared individuals are in light blue. Allometric slopes for each group are shown, the slope, intercepts and major-axis means are compared in Table 3,4.