Invasive plants as novel food resources, the pollinators' perspective

Ignasi Bartomeus[1], Jochen Fründ[2,3] and Neal M. Williams[3]

[1] Estación Biólogica de Doñana - Consejo Superior de Investigaciones Científicas (EBD-CSIC), Dept. Integrative Ecology. Sevilla, Spain. <nacho.bartomeus@gmail.com>
[2] University of Guelph, Dept. Integrative Biology, Guelph, ON, Canada
[3] University of California, Dept. Entomology and Nematology, Davis, CA, USA

Biological invasions are one of the main drivers of global change and have negatively impacted all biomes and trophic levels (Hobbs 2000, Vilà et al. 2011). While most introduced species fail to establish, or establish small naturalized populations (hereafter exotic species), a few become invasive and rapidly increase in abundance and/or range (hereafter invasives or invaders; Pysek et al. 2004). It is these invader species that are most often linked to negative impacts on native / endemic communities. Although most interactions between invasive and native species at the same trophic level result in negative direct impacts (e.g. plant-plant competitive interactions, Vilà et al. 2011), when the invasive plant species can be used as a resource those interactions can also be positive for consumers such as native herbivores, predators or mutualists, at least for some species (Heleno et al. 2009, Bezemer et al. 2004). Entomophilous exotic plant species, for example, not only compete directly for space and light with other plants, but also offer resource opportunities for the native pollinator community (Stouffer et al. 2014). Most research on this topic to date has taken the plant perspective, focusing on how successful plant invaders integrate into the native plant-pollinator interaction networks (Vilà et al. 2009), and how this integration in turn impacts the native plant community (Morales and Traveset 2009). However, species specific responses of pollinators to the addition of exotic plants are rarely taken into account. This represent an important gap in our knowledge as pollinator foraging choices determine the structure of interactions within communities, which in turn have important implications for the community stability (Thébault and Fontain 2010) and functioning (Thomson et al. 2012). How different pollinators respond to the changed composition of floral species within the community that results from exotic plant invaders deserves more attention.

From the pollinators' point of view, exotic plants are novel food resources, and as such their relative abundance, attractiveness, rewards (i.e. nutritional value) and distinctiveness partly

38　determine their use by various pollinators (Carvalheiro et al. 2014). Although exotics as a

group are not preferred or avoided within their new communities (Williams et al. 2010), it

40　might be that particular exotic species are preferred by some pollinators while avoided by

others. The intrinsic preferences for different plant hosts is an important factor determining

42　host use. Hence, the direct benefits or costs of a novel resource use will differ among

pollinator species. Moreover, in a community context, the preferences of each pollinator

44　affect the other pollinators' choices, potentially leading to indirect effects. For example, some

pollinator species may obtain indirect benefits if the invasive plants distract other pollinators

46　from natives, reducing competition. Alternatively, pollinators may pay indirect costs if

competition is increased or if invasive plants reduce the availability of a preferred native

48　plant.

50　We review the evidence on direct and indirect benefits and costs of invasive plants on

pollinators and re-analyze Williams et al. (2010) dataset on pollinator specific preferences so

52　as to identify species that prefer some exotic plants over native plants and vice-versa. This

information is crucial to understanding the consequence for the pollinator community

54　because if only some pollinators take advantage of alien plants this can favour populations of

some pollinator species (winners) over others (losers). By using an approach that takes into

56　account both pollinator behavioral responses and interaction network structure, we can better

understand the invasion process, with important implications for conservation actions.

Effects of plant invasions on pollinator populations and community structure.

The impact of invasive plants interacting with native pollinators has received considerable

62　attention for its potential to disrupt native mutualisms (Traveset and Richardson 2006).

However, most work to date has focused solely on how invasive plant interactions with

64　native pollinators changes the pollination success of native plants. Interestingly, existing data

show that invasive plants can have positive, neutral, or negative effects on pollination of

66　native plants (Bjerknes et al. 2007). The contrasting results may reflect invasive plant density

(Muñoz and Cavieres 2008, Dietzsch et al. 2011), spatial aggregation (Cariveau and Norton

68　2009) or flower morphology and attractiveness (Morales and Traveset 2009, Carvalheiro et

al. 2014).

In contrast, the effects of exotic or invasive species on the pollinator populations and

communities have received far less attention (Stout and Morales 2008). Because most plant-pollinator systems are generalized (Waser et al. 1996), invasive plants are usually well

integrated in the plant–pollinator network of interactions (Vilà et al 2009). Hence, we might expect overall effects on pollinators to mirror the changes (positive or negative) in floral

resources offered by the newly invaded community. If the entomophilous exotic or invasive plants increase the resources present in the community, this should also allow the increase of

most pollinator populations (Tepedino et al. 2008). Stout and Morales (2008) cite indirect evidence that some social pollinators (e.g., bumblebees) can be favoured by non-native mass

flowering crops (Westphal et al. 2003, Herrmann et al. 2007), which may be analogous to the effect of abundant invasive species. However, the same authors note examples where exotic

plants are not used by native pollinators due to flower morphology or chemistry. Despite Stout and Morales' (2008) call for more research on this topic, few additional studies have

been published since then.

Current evidence suggests that food resource availability may indeed regulate pollinator's populations, at least those of bees (reviewed in, Roulston and Goodell 2011, Williams and

Kremen 2007, Crone 2013, but see Steffan-Dewenter and Schiele 2008 for potential regulation by nesting resources). However, studies of the effect of exotic plants on

pollinators' population dynamics are extremely rare, particularly for non-invasive exotic plants that establish small populations. Palladini and Maron (2014) provide one of the few

examples of effects of exotic plants (mainly *Euphorbia esula*) on the reproduction of a solitary bee species (*Osmia lignaria*). For this species, the number of nests established and

offspring production per female was positively related to native plant abundance and negatively related to exotic plant species. This may be because although *Osmia lignaria*

foraged on exotics for nectar, the species did not use *Euphorbia esula* exotic pollen to provision nests. Thus, the specific ability to use the invader resources emerges as a key factor

affecting the potential impacts of the invader on pollinators. The only other evidence to date for direct effects of invasive plants through resource augmentation is for bumblebees, whose

annual social life history allows demographic responses to be measured within a single season. Within-season abundance can increase almost four times in sites invaded by the plant

Lupinus polyphyllus, compared to in non-invaded sites (Jakobsson and Padrón 2014). In addition, the foraging season of *Bombus terrestris* in the United Kingdom can be extended

into winter through its use of exotic plants that fill a late season phenological niche (Stelzer et

        al. 2010). Such within-season demographic responses are likely to have longer-term

106   population effects, although no study has quantified such effect to date.

108   A second group of studies provides indirect evidence that exotic plants affect

        pollinators' by using community approaches to compare invaded with non-invaded sites.

110   These studies show a variety of pollinator responses, including increased abundance and

        species richness in invaded sites (Lopezaraiza-mikel et al. 2007), lower abundance and

112   diversity of pollinators in invaded areas (Moron et al. 2009), or no difference in abundance

        between invaded and non-invaded sites (Nienhuis et al. 2009). It is therefore not surprising

114   that a recent meta-analysis reported no changes in overall pollinator abundance in invaded

        sites (Montero-castaño and Vilà 2012). However, the studies included in the meta-analysis

116   were not designed to infer population changes, and the result should be interpreted with

        caution. Moreover, most of the examples concern abundant invasive plants. Plant abundance

118   can strongly influence pollinators decision to incorporate a new plant into its diet

        (Valdovinos et al. 2010), and thus the results may differ when examining non-invasive exotic

120   plants. Likewise, given the wide array of pollinators ranging from birds and bats to bees and

        hoverflies, it is unlikely to find a consistent overall response.

        The importance of behaviour

        While exotic plants can represent new resource opportunities for native animals, evidence

126   suggests that only a minority of pollinator species can take advantage of these new

        opportunities. For example, generalist bees more commonly forage on invasive exotic plants

128   than specialists (Lopezaraiza-Mikel et al., 2007; Tepedino et al., 2008; Padrón et al. 2009), or

        that social bumblebees are more common in invaded sites than solitary bees (Nienhuis et al.

130   2009). Bartomeus et al. (2009), for example, report that bees were more often recorded on

        native species than on the invader *Carpobrotus* aff. *acinaciformis*, except for the social

132   bumblebee *Bombus terrestris* and for most beetles, which visited *Carpobrotus* flowers almost

        exclusively. Hence, pollinators can discriminate between native and exotic plants, and the

134   decision of exploiting one or another can vary across species. Analyzing the invasion process

        as a novel resource availability for pollinators may give us a framework to predict which

136   pollinators can beneficiate from the invasion process.

138   Incorporating any novel resource into the diet requires a series of conditions to be met. First,
       the pollinator must recognize the novel resource as a host, second, the visitor must be able to

140   use this new resource and third, the resource must be profitable (i.e., a net benefit) for the
       visitor to exploit it. Hence, exotic plant use depends on intrinsic traits of the plant and

142   pollinator. We cannot assume that for native pollinators exotic flowering species are always
       fundamentally different than the native flowering species; nevertheless, plants presenting new

144   colors, shapes or chemical compounds may not be easily used by all pollinators in the
       community (Stout and Morales 2008). Furthermore many pollinators have innate preferences

146   for certain colors and/or shapes (Gumbert 2000, Riffell et al. 2008), which may make novel
       flowers less attractive than the natives which have coevolved in the community. Neophobic

148   responses also may make some pollinators unlikely to approach and explore novel food
       opportunities (Forrest and Thomson 2009). However, some pollinators may learn how to

150   exploit new resources if their behaviour is flexible enough (Chittka et al. 2009, Forrest and
       Thomson 2009). In particular, bees, which are the main pollinator group both in numbers and

152   in effectiveness (reviewed in Winfree et al. 2011), have a powerful neuronal system able to
       learn new tasks (Chittka et al. 2009) and their behavior flexibility has been suggested to be

154   linked to their ability to persist in disturbed environments (Kevan and Menzel 2012).
       However the role of learning abilities in incorporating new foraging plants is little explored

156   (Dukas and Real 1993) and most information comes from a handful of species (mostly
       social).

158
       Even if the exotic plant is recognized as a potential host, pollinators might not be able to

160   exploit the new plant species if they are unable to handle its flowers (Parker 1997, Corbet et
       al. 2001), or to digest its nectar (Adler and Irwin, 2005) or pollen (Sedivy et al. 2011,

162   Palladini and Maron 2014). Morphological matching between flowers and pollinators may
       thus be important in determining pollinator visitation patterns (Gibson et al. 2012, Bartomeus

164   2013, Stang et al. 2009, see also Pearse et al. 2014 for antagonist insect-plant interactions).
       Second, even in the cases where pollinators recognize and can use the novel resource, their

166   decision to include it in its diet will depend on its quality and abundance relative to others in
       the community. The thresholds for switching to a resource based on its quality or abundance

168   have been show to be variable among different species in birds (Carnicer et al. 2008). Insect
       pollinators switch between foraging plants depending on the resource availability (Inouye

170   1978, Chittka et al 1997). Like for birds, thresholds for switching are likely to be different for

        different species, however, the experimental evidence that insect pollinators can discriminate

172   between different resources and learn to forage on the preferred one is limited to a handful of

        bee species, and the switching strategies among species is mostly unknown.

176   **The importance of the community context**.

        Pollinator-plant interactions do not occur in isolation, but form part of a complex network of

178   interactions (Bascompte and Jordano 2007). For example, competitive exclusion between

        pollinator species can drive foraging behavior patterns (Jhonson and Hubell 1995). Invasive

180   plant species can modify the native network structure not only by creating new interactions

        with native pollinators but also by modifying the existing plant-pollinator interactions among

182   the native species (Bartomeus et al. 2008, Albrecht et al. 2014). Such changes are especially

        likely when invaders are abundant and provide accessible resources (i.e. nectar and pollen),

184   thus potentially interacting with a large proportion of the resident pollinator species (Albrecht

        et al. 2014). In any case, modifications to pairwise interactions can have cascading effects

186   throughout the community. For example, dynamic pollination network models have been

        used to show that removal of well-established invasive plants negatively affected the

188   persistence of pollinator interactions through the network (Valdovinos et al. 2009,

        Carvalheiro et al. 2008). Detailed empirical studies, also show that co-flowering neighboring

190   invasive species affect pollinator choices (Cariveau and Norton 2009, Waters et al. 2014).

192   The behavioural switching (also called interaction rewiring) between resources has been

        recently studied at the community level using network theory. There is increased evidence

194   that pollinators can rewire their community-wide interactions depending on the context

        (Kaiser-Bunbury et al. 2010), such as the addition of a plant invasion. Jackobson and Padrón

196   (2014) speculate that the attraction of bumblebees to the invasive plant *Lupinus polyphyllus*,

        reduced competition for the native plants, allowing an increase of solitary bee visits to

198   natives. Similarly, Montero-Castaño (2014) showed that monopolization of the invasive

        species *Hedysarum coronarium* by honeybees allowed other bee species to establish

200   interactions with natives that are not realized when the honeybee is present. Context

        dependent rewiring is supported by findings that despite for consistent species-specific

202   preferences for certain flowers across communities, there is also important variation and

        flexibility in preferences among different contexts (Fründ et al. 2010). Hence, both direct

 effects and indirect effects on pollinators are expected after a plant invasion, and those can only be understood in a community-wide framework.

Figure 1. Simplified plant-pollinator networks before the invasion process (A) in two distinctive scenarios: (B) The exotic plant (orange square) adds a novel resource without affecting the rest of the community, and (C) a superabundant invasive plant (orange square) adds an abundant novel resource, while reducing native plants resource availability. In the schemes we can see the interactions established in each case (lines connecting pollinators and plants). The color of links depicts whether their frequency is increased (blue) or decreased (red) with respect to A. The changes in size of the plants or pollinators depict the winners (blue border) and losers (red border) of the invasion process. When the effect is indirect is noted with dashed lines. See text for further explanation.

In Figure 1, we illustrate a simplified plant-pollinator network with two distinctive scenarios. In the first one (Fig. 1.B), we add an entomophilous exotic plant that does not reduce the abundances of other plants. Pollinator species able to use this plant (identified as p1 and p2) will establish new links with this exotic plant (blue links). Species p1 will have more food resources and can potentially increase its population over time (bigger blue circle denotes population size increase) whereas species p2 will experience a neutral effect because it changes from foraging on natives to foraging on the exotic. These are the direct (often neutral or positive) effects of the exotic plant on pollinators. Other pollinators may not be able or may not choose to visit the new invasive plant (p3), but as the competition for their preferred resources is changed, they may receive indirect benefits (p3). Experiments removing dominant pollinators have shown that a relaxed competition for resources may lead to diet

expansion of some species (Brosi et al 2013), supporting our example with species p3.

In the second scenario (Fig 1. C), the exotic plant is an abundant invader that also reduces native plant abundance by direct competition (e.g. for space). In this scenario, only a few pollinators (p1) may benefit, while all others will experience increased competition for resources (p2, p3). This is an oversimplification, and of course the net benefit for pollinators will depend not only on the number of visits, but on the quality of those visits (e.g. reward uptake, nutritional content of the exotic species, etc.). Moreover, some species will require a variety of pollen sources to complete larvae development (Roulston and Cane 2000) highlighting the importance of maintaining plant diversity. The magnitude of the indirect and direct effects will depend also on the time-scale at which it is evaluated, with functional responses and local switching occurring faster than numerical responses (i.e. population growth). Moreover, the relative phenological timing of plants and bees can modify their mutual influence. All in all, the net costs and benefits are likely to depend on many factors, but this framework supports the scarce information presented above, where some social generalized species tend to increase their abundance after invasion by highly attractive species, but other pollinators have mixed responses.

Case study: bee preferences in California

Can we predict which pollinators will be winners or losers of the invasion process? Measuring population responses or fecundity is a daunting task, especially at the community level; however, we can gain indirect evidence by looking at pollinator preferences. Within a plant community, pollinators do not prefer exotic plants as a group (Williams et al. 2010) or even prefer natives (Chrobock et al. 2013), but individual pollinator preferences have not been explored yet. Preference is defined as using a resource more than expected given its abundance. Conversely, avoidance occurs when a resource is underused relative to its abundance. The null model of no preferences is the case when pollinators visit flowers in proportion to their abundance in the community. Deviations from this null model can help us identify pollinator species that prefer exotic species (hence receiving a potential direct benefit) and species that avoid the exotics (hence, receiving negative, neutral or positive indirect effects in some cases). We recognize that we cannot infer direct fitness consequences, or predict indirect effects from a static network. Ours, nonetheless, is the first

262   attempt to identify direct effects and serve as a proxy for identifying pollinator winners after

the invasion process. Most importantly, this way we can emphasize that pollinators differ in

264   their behavior, acknowledging that the effects on specific pollinators cannot be generalized.

Furthermore, in the future, we can explore what determines pollinator preferences. Are they

266   driven by plant traits, such as abundance or morphology, by pollinator traits, or a

combination of both?

To explore this preference-based proxy, we used the same dataset used in Williams et al.

270   (2010). For simplicity we show here only 7 sites from semi-natural habitats in California.

This system is especially suitable to test our questions, because it contains several exotic

272   plants, ranging from abundant invaders to naturalized exotic plants, as well as a variety of

pollinator species. We calculated preferences pooling all sites, but we separate our analysis in

274   three sampling periods (early, mid and late season). We treated periods separately because

plant turnover was substantial over the season and otherwise might have masked the

276   preference relationships.

278   First we re-evaluated that pollinators do not prefer exotic plants as a group within a

quantitative framework, where expected (E) visitation values are calculated based on plant

280   mean abundance across sites, and observed (O) visitation values are the sums of pollinator

visitation to each plant across sites. Chi statistics were used for each of the three tests (i.e. one

282   per season) to assess if there is an overall preference. The Pearson residuals of the Chi tests

$(O-E/\sqrt{E})$ estimate the magnitude of preference or avoidance for a given plant based on

284   deviation from expected values and its significance was assessed by building Bonferroni

confidence intervals (see Neu et al. 1974 and Byers and Steinhorst 1984). In order to test for

286   differences between exotics and natives we compared the *Electivity* values of exotic and

native plants using linear models. *Electivity* values (E'-O'/E'+O', where E' and O' are the

288   proportional expected and observed values, Ivlev 1964) are bound between -1 and +1, easier

to interpret and highly correlated with Pearson residuals. R code to calculate these indexes

290   can be found at (https://gist.github.com/ibartomeus/cdddca21d5dbff26a25e).

292   We show that when pooling visits for all pollinator species, some plants are preferred over

others (Chi square test p-values for early, mid and late seasons < 0.001; *Electivity* values

294   range from -0.84 to 0.82 indicating we find both over-preferred plants and under-preferred

plants; Fig. 2). In accordance with the results reported in Williams et al. (2010), pooled

296   pollinators showed no preferences for exotic plants in any season (Fig. 2; early season: $F_{1,25} =$

1.29, p-value: 0.27; mid season: $F_{1,22} = 0.06$, p-value: 0.81; late season: $F_{1,15} = 0.46$, p-value:

298   0.51), but this general trend does not contradict the fact that specific native or exotic plants

are indeed preferred (in red in Fig. 2).

302   Figure 2: Boxplots of the *electivity* indices per plant species separated by plant origin (exotic,

native) and season (early, mid, late). Each plant value is plotted in the background in grey

304   when preference is not significant and in red when significant (significance based on Chi

square Pearson residuals test). Positive values indicate plants that are preferred and negative

306   values those that are avoided.

308   Second, we shift the focus of our analysis to analyze pollinator species-specific preferences.

We excluded pollinator species with less than 20 visits recorded per season in order to

310   prevent confounding rarity with specialization (Blüthgen 2010). We end up with 16 pollinator

species, some of them present in several seasons, making 22 pollinator-season combinations.

312   Again, each of the 22 pollinator-season combination was evaluated using Chi square tests and

electivity values were compared between exotic and native plants for each pollinator species

314   using mixed models with season as random effect. All pollinators showed significant

preferences for certain plant species (chi square p-values < 0.05). When analyzed individually

316   most pollinators do not show a consistent preference for exotic or native plants, with the

exception of three species (Fig. 3; all species p-values > 0.1, except *Bombus melanopygus* =

318   0.05, *Dialictus incompletum* = 0.07 and *Halictus ligatus* = 0.03), all of them preferring

exotics over natives. However, some pollinators of the 16 analyzed do prefer only one or

320   more native plants (4 species: *Evylaeus sp, Osmia nemoris, Ceratina arizonensis, Calliopsis*

fracta), others prefer only one or more exotic plants (7 species: *Synhalonia actuosa,*

322   *Synhalonia frater, Bombus vosnesenskii, Halictus ligatus, Halictus tripartitus, Megachile*

apicalis, Bombus californicus) and some prefer a mix of exotic and natives (4 species:

324   *Bombus melanopygus, Ashmeadiella aridula, Dialictus incompletum, Dialictus*

tegulariforme). The differential preferences regarding the exotic status create the basis for

326   expecting winners and losers after an invasion process (see Fig. 1).

328   Figure 3. Boxplot of the effect sizes (i.e. model estimates) indicating the difference between

electivity to exotics and to natives for each pollinator species. Positive values indicate overall

330   preferences for natives, and negative values to exotics. Data points for each pollinator species

are indicated in grey when not significant, in orange when $p < 0.1$ and in red when $p < 0.05$.

In conclusion, although the overall pattern is no preference for exotic plants, some particular

334   exotic (and native) plants are overall preferred. Similarly, most pollinators do not have

overall preferences for exotics, but a few species do favor them. Those are social species,

336   usually common and sometimes even considered species typical of disturbed areas (e.g.

Halictus ligatus, Dialictus incompletum). Interestingly, even within the species with no

338   overall preference for exotics, we identify pollinators that prefer particular exotic plants.

These pollinators are more likely to be positively affected by the invasion process, the others

340   negatively affected, as their preferred resources will potentially diminish through

displacement by invasive plant species.

Relevance for conservation

We highlighted that pollinator species vary in response to plant invasions, including pollinators use, preference, and in some cases population dynamic consequences. Assessing the winners and losers in front of the rapid rate of invasive species introductions is crucial for understanding the responses of species groups performing important ecosystem functions, like pollination (Ollerton et al. 2011, Klein et al. 2007). There has been a recent awareness of pollinator declines globally (Potts et al. 2010, Gonzalez-Varo et al. 2013). However, biological invasions, especially by plant species, have received little attention as a threat (but see Stout and Morales 2008, and Morales et al. 2013 for effects of animal invasions). We already know that not all pollinators are equally affected by global change, a few are winners and many are losers (Bartomeus et al. 2013). Interestingly, among the winner pollinators we found species that are able to use flowering crops and tolerate new human-modified habitats (Bartomeus and Winfree 2013). Gaining information about which species are able also to exploit new exotic plants will be a way forward to understand which species will be flexible enough to survive in novel ecosystems, often dominated by exotic plants. In a changing world, species able to adapt their foraging strategies to use new resources may be the better suited to survive. For example, bumble bees that use the widespread plant invader *I. glandulifera* in the EU are thriving, while endangered bumble bee species do not use it (Kleijn and Raemaker 2008).

If we are going to manage emerging novel ecosystems, we need to incorporate pollinator specific responses to different global change drivers, including plant invasions. Some bumble bees and other trophic generalist bees can benefit from exotic plant invasions, as shown by the fact that those can use and even prefer to forage on new exotic plants. This behavioral flexibility may be the key to persisting in a changing world, and maintaining an important ecosystem function. More research is needed on the degree that plant invasions negatively affect those species in comparison with other disturbances that are occurring simultaneously. We need to implement better population monitoring programs at the community level (so indirect responses can be accounted for), but overall, understanding better which role play the pollinator behavior flexibility and cognitive capabilities in the process of adapting to novel environments is a promising line of research.

376   **Acknowledgments:**

We thank the book editors and an anonymous reviewer for their comments on the manuscript.

378   IB acknowledge funding from the project SURVIVE_HIREC (CGL2013-47448-P)

380   **References:**

Adler, L. S., & Irwin, R. E. (2005). Ecological costs and benefits of defenses in

382   nectar. Ecology, 86(11), 2968-2978.

384   Albrecht, M., Padrón, B., Bartomeus, I., & Traveset, A. (2014). Consequences of plant

invasions on compartmentalization and species' roles in plant–pollinator

386   networks. Proceedings of the Royal Society B: Biological Sciences, 281(1788), 20140773.

388   Bartomeus, I. (2013). Understanding linkage rules in plant-pollinator networks by using

hierarchical models that incorporate pollinator detectability and plant traits. PloS one, 8(7),

390   e69200.

392   Bartomeus, I., & Winfree, R. (2013). Pollinator declines: reconciling scales and implications

for ecosystem services. F1000Research, 2.

Bartomeus, I., Ascher, J. S., Gibbs, J., Danforth, B. N., Wagner, D. L., Hedtke, S. M., &

396   Winfree, R. (2013). Historical changes in northeastern US bee pollinators related to shared

ecological traits. Proceedings of the National Academy of Sciences, 110(12), 4656-4660.

Bartomeus, I., Bosch, J., & Vilà, M. (2008). High invasive pollen transfer, yet low deposition

400   on native stigmas in a Carpobrotus-invaded community. Annals of Botany, 102(3), 417-424.

402   Bartomeus, I., Vilà, M., & Santamaría, L. (2008). Contrasting effects of invasive plants in

plant–pollinator networks. Oecologia, 155(4), 761-770.

Bascompte, J., & Jordano, P. (2007). Plant-animal mutualistic networks: the architecture of

406   biodiversity. Annual Review of Ecology, Evolution, and Systematics, 567-593.

408   Bezemer, T. M., Harvey, J. A., & Cronin, J. T. (2014). Response of native insect communities to invasive plants. Annual review of entomology, 59, 119-141.

Bjerknes, A. L., Totland, Ø., Hegland, S. J., & Nielsen, A. (2007). Do alien plant invasions

412   really affect pollination success in native plant species?. Biological Conservation, 138(1), 1-12.

Blüthgen, N. (2010). Why network analysis is often disconnected from community ecology: a

416   critique and an ecologist's guide. Basic and Applied Ecology, 11(3), 185-195.

418   Brosi, B. J., & Briggs, H. M. (2013). Single pollinator species losses reduce floral fidelity and plant reproductive function. Proceedings of the National Academy of Sciences, 110(32),

420   13044-13048.

422   Byers, C. R., Steinhorst, R. K., & Krausman, P. R. (1984). Clarification of a technique for analysis of utilization-availability data. The Journal of Wildlife Management, 1050-1053.

Cariveau, D. P., & Norton, A. P. (2009). Spatially contingent interactions between an exotic

and native plant mediated through flower visitors. Oikos,118(1), 107-114.

Carnicer, J., Abrams, P.A. & Jordano, P. (2008). Switching behavior, coexistence and diversification: comparing empirical community-wide evidence with theoretical predictions

Ecology Letters, 11: 802–808

Carvalheiro, L. G., Barbosa, E. R. M., & Memmott, J. (2008). Pollinator networks, alien species and the conservation of rare plants: Trinia glauca as a case study. Journal of Applied

Ecology, 45(5), 1419-1427.

Carvalheiro, L. G., Biesmeijer, J. C., Benadi, G., Fründ, J., Stang, M., Bartomeus, I., ... & Kunin, W. E. (2014). The potential for indirect effects between co-flowering plants via

shared pollinators depends on resource abundance, accessibility and relatedness. Ecology letters, 17(11), 1389-1399.

Chittka, L. Gumbert, A. & Kunze, J. (1997) Foraging dynamics of bumble bees: correlates of

442   movements within and between plant species. Behavioral Ecology 8(3):239-249.

444   Chittka, L., & Niven, J. (2009). Are bigger brains better? Current Biology,19(21), R995-

R1008.

Chrobock, T., Winiger, P., Fischer, M., & van Kleunen, M. (2013). The cobblers stick to their

448   lasts: pollinators prefer native over alien plant species in a multi-species

experiment. Biological invasions, 15(11), 2577-2588.

Corbet, S. A., Bee, J., Dasmahapatra, K., Gale, S., Gorringe, E., La Ferla, B., ... &

452   Vorontsova, M. (2001). Native or exotic? Double or single? Evaluating plants for pollinator-

friendly gardens. Annals of Botany, 87(2), 219-232.

Crone, E. E. (2013). Responses of social and solitary bees to pulsed floral resources. The

456   American Naturalist, 182(4), 465-473.

458   Dietzsch, A. C., Stanley, D. A., & Stout, J. C. (2011). Relative abundance of an invasive

alien plant affects native pollination processes. Oecologia, 167(2), 469-479.

Forrest, J., & Thomson, J. D. (2009). Pollinator experience, neophobia and the evolution of

462   flowering time. Proceedings of the Royal Society B: Biological Sciences, 276(1658), 935-

943.

Fründ, J., Linsenmair, K. E., & Blüthgen, N. (2010). Pollinator diversity and specialization in

466   relation to flower diversity. Oikos, 119(10), 1581-1590.

468   Gibson, M. R., Richardson, D. M., & Pauw, A. (2012). Can floral traits predict an invasive

plant's impact on native plant–pollinator communities? Journal of Ecology, 100(5), 1216-

470   1223.

472 González-Varo, J. P., Biesmeijer, J. C., Bommarco, R., Potts, S. G., Schweiger, O., Smith, H. G., ... & Vila, M. (2013). Combined effects of global change pressures on animal-mediated

474 pollination. Trends in ecology & evolution, 28(9), 524-530.

476 Gumbert, A. (2000). Color choices by bumble bees (*Bombus terrestris*): innate preferences and generalization after learning. Behavioral Ecology and Sociobiology, 48(1), 36-43.

Heleno, R. H., Ceia, R.S., Ramos, J.A., & Memmott, J. (2009) Effects of Alien Plants on

480 Insect Abundance and Biomass: a Food-Web Approach. Conservation Biology, 23(2), 410-419.

Herrmann, F., Westphal, C., Moritz, R. F., & Steffan-Dewenter, I. (2007). Genetic diversity

484 and mass resources promote colony size and forager densities of a social bee (*Bombus pascuorum*) in agricultural landscapes. Molecular Ecology, 16(6), 1167-1178.

Hobbs, R. J. (Ed.). (2000). Invasive species in a changing world. Island Press.

Inouye, D.W. (1978). Resource partitioning in bumblebees: Experimental Studies of Foraging

490 Behavior. Ecology 59(4), 672-678

492 Ivlev, V. S. (1964). Experimental ecology of the feeding of fishes.

494 Jakobsson, A., & Padrón, B. (2014). Does the invasive *Lupinus polyphyllus* increase pollinator visitation to a native herb through effects on pollinator population

496 sizes?. Oecologia, 174(1), 217-226.

498 Jhonson, L.K. & Hubell, P. (1975) Contrasting Foraging Strategies and Coexistence of Two Bee Species on a Single Resource. Ecology 56 (6), 1398-1406

502 Kaiser-Bunbury, C. N., Muff, S., Memmott, J., Müller, C. B., & Caflisch, A. (2010). The robustness of pollination networks to the loss of species and interactions: a quantitative

504 approach incorporating pollinator behaviour. Ecology Letters, 13(4), 442-452.

506 Kevan, P. G., & Menzel, R. (2012). The plight of pollination and the interface of neurobiology, ecology and food security. The Environmentalist, 32(3), 300-310.

Kleijn, D., & Raemakers, I. (2008). A retrospective analysis of pollen host plant use by stable

and declining bumble bee species. Ecology, 89(7), 1811-1823.

Klein, A. M., Vaissiere, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., & Tscharntke, T. (2007). Importance of pollinators in changing landscapes for world

crops. Proceedings of the Royal Society B: Biological Sciences, 274(1608), 303-313.

Lopezaraiza–Mikel, M. E., Hayes, R. B., Whalley, M. R., & Memmott, J. (2007). The impact of an alien plant on a native plant–pollinator network: an experimental approach. Ecology

Letters, 10(7), 539-550.

Montero-Castaño, A. (2014) Interacciones entre polinizadores y la planta exótica *Hedysarum coronarium* a distintas escalas espaciales. PhD Thesis

Montero-Castaño, A., & Vila, M. (2012). Impact of landscape alteration and invasions on

pollinators: a meta-analysis. Journal of Ecology, 100(4), 884-893.

Morales, C. L., & Traveset, A. (2009). A meta-analysis of impacts of alien vs. native plants on pollinator visitation and reproductive success of co-flowering native plants. Ecology

Letters, 12(7), 716-728.

Morales, C. L., Arbetman, M. P., Cameron, S. A., & Aizen, M. A. (2013). Rapid ecological replacement of a native bumble bee by invasive species. Frontiers in Ecology and the

Environment, 11(10), 529-534.

Moroń, D., Lenda, M., Skórka, P., Szentgyörgyi, H., Settele, J., & Woyciechowski, M. (2009). Wild pollinator communities are negatively affected by invasion of alien goldenrods

in grassland landscapes. Biological Conservation, 142(7), 1322-1332.

Muñoz, A. A., & Cavieres, L. A. (2008). The presence of a showy invasive plant disrupts

pollinator service and reproductive output in native alpine species only at high

540   densities. Journal of Ecology, 96(3), 459-467.

542   Neu, C. W., Byers, C. R., & Peek, J. M. (1974). A technique for analysis of utilization-

availability data. The Journal of Wildlife Management, 541-545.

Nienhuis, C. M., Dietzsch, A. C., & Stout, J. C. (2009). The impacts of an invasive alien

546   plant and its removal on native bees. Apidologie, 40(4), 450-463.

548   Ollerton, J., Winfree, R., & Tarrant, S. (2011). How many flowering plants are pollinated by

animals?. Oikos, 120(3), 321-326.

Padrón, B., Traveset, A., Biedenweg, T., Díaz, D., Nogales, M., & Olesen, J. M. (2009).

552   Impact of alien plant invaders on pollination networks in two archipelagos. PLoS One, 4(7),

e6275.

Palladini, J. D., & Maron, J. L. (2014). Reproduction and survival of a solitary bee along

556   native and exotic floral resource gradients. Oecologia, 176(3), 789-798.

558   Parker, I. M. (1997). Pollinator limitation of *Cytisus scoparius* (Scotch broom), an invasive

exotic shrub. Ecology, 78(5), 1457-1470.

Pearse, I. S., Harris, D. J., Karban, R., & Sih, A. (2013). Predicting novel herbivore–plant

562   interactions. Oikos, 122(11), 1554-1564.

564   Potts, S. G., Biesmeijer, J. C., Kremen, C., Neumann, P., Schweiger, O., & Kunin, W. E.

(2010). Global pollinator declines: trends, impacts and drivers. Trends in ecology &

566   evolution, 25(6), 345-353.

568   Pyšek, P., Richardson, D. M., Rejmánek, M., Webster, G. L., Williamson, M., & Kirschner,

J. (2004). Alien plants in checklists and floras: towards better communication between

570   taxonomists and ecologists. Taxon, 131-143.

572   Riffell, J. A., Alarcón, R., Abrell, L., Davidowitz, G., Bronstein, J. L., & Hildebrand, J. G.
(2008). Behavioral consequences of innate preferences and olfactory learning in hawkmoth–
574   flower interactions. Proceedings of the National Academy of Sciences, 105(9), 3404-3409.

576   Roulston, T. A. H., & Goodell, K. (2011). The role of resources and risks in regulating wild
bee populations. Annual review of entomology, 56, 293-312.
578

Roulston, T. H., & Cane, J. H. (2000). Pollen nutritional content and digestibility for
580   animals. Plant Systematics and Evolution, 222(1-4), 187-209.

582   Sedivy, C., Müller, A., & Dorn, S. (2011). Closely related pollen generalist bees differ in
their ability to develop on the same pollen diet: evidence for physiological adaptations to
584   digest pollen. Functional Ecology, 25(3), 718-725.

586   Stang, M., Klinkhamer, P. G., Waser, N. M., Stang, I., & van der Meijden, E. (2009). Size-
specific interaction patterns and size matching in a plant–pollinator interaction web. Annals
588   of Botany, 103(9), 1459-1469.

590   Steffan-Dewenter, I., & Schiele, S. (2008). Do resources or natural enemies drive bee
population dynamics in fragmented habitats. Ecology, 89(5), 1375-1387.
592

Stelzer, R. J., Chittka, L., Carlton, M., & Ings, T. C. (2010). Winter active bumblebees
594   (*Bombus terrestris*) achieve high foraging rates in urban Britain. PLoS One, 5(3), e9559.

596   Stouffer, D. B., Cirtwill, A. R., & Bascompte, J. (2014). How exotic plants integrate into
pollination networks. Journal of Ecology, 102(6), 1442-1450.
598

Stout, J. C., & Morales, C. L. (2009). Ecological impacts of invasive alien species on
600   bees. Apidologie, 40(3), 388-409.

602   Tepedino, V. J., Bradley, B. A., & Griswold, T. L. (2008). Might flowers of invasive plants

increase native bee carrying capacity? Intimations from Capitol Reef National Park,

604   Utah. Natural Areas Journal, 28(1), 44-50.

606   Thébault, E., & Fontaine, C. (2010). Stability of ecological communities and the architecture

of mutualistic and trophic networks. Science, 329(5993), 853-856.

Thompson, R. M., Brose, U., Dunne, J. A., Hall Jr, R. O., Hladyz, S., Kitching, R. L., ... &

610   Tylianakis, J. M. (2012). Food webs: reconciling the structure and function of

biodiversity. Trends in ecology & evolution, 27(12), 689-697.

Traveset, A., & Richardson, D. M. (2006). Biological invasions as disruptors of plant

614   reproductive mutualisms. Trends in Ecology & Evolution, 21(4), 208-216.

616   Valdovinos, F. S., Ramos-Jiliberto, R., Flores, J. D., Espinoza, C., & López, G. (2009).

Structure and dynamics of pollination networks: the role of alien plants. Oikos, 118(8), 1190-

618   1200.

620   Valdovinos, F. S., Ramos-Jiliberto, R., Garay-Narváez, L., Urbani, P., & Dunne, J. A. (2010).

Consequences of adaptive behaviour for the structure and dynamics of food webs. Ecology

622   letters, 13(12), 1546-1559.

624   Vilà, M., Bartomeus, I., Dietzsch, A. C., Petanidou, T., Steffan-Dewenter, I., Stout, J. C., &

Tscheulin, T. (2009). Invasive plant integration into native plant–pollinator networks across

626   Europe. Proceedings of the Royal Society B: Biological Sciences, 276(1674), 3887-3893.

628   Vilà, M., Espinar, J. L., Hejda, M., Hulme, P. E., Jarošík, V., Maron, J. L., ... & Pyšek, P.

(2011). Ecological impacts of invasive alien plants: a meta-analysis of their effects on

630   species, communities and ecosystems. Ecology Letters, 14(7), 702-708.

632   Waser, N. M., Chittka, L., Price, M. V., Williams, N. M., & Ollerton, J. (1996).

Generalization in pollination systems, and why it matters. Ecology, 77(4), 1043-1060.

Waters, S. M., Fisher, S. E., & Hille Ris Lambers, J. (2014). Neighborhood-contingent indirect interactions between native and exotic plants: multiple shared pollinators mediate reproductive success during invasions. Oikos,123(4), 433-440.

Westphal, C., Steffan-Dewenter, I., & Tscharntke, T. (2003). Mass flowering crops enhance pollinator densities at a landscape scale. Ecology Letters, 6(11), 961-965.

Williams, N. M., & Kremen, C. (2007). Resource distributions among habitats determine solitary bee offspring production in a mosaic landscape. Ecological applications, 17(3), 910-
921.

Williams, N. M., Cariveau, D., Winfree, R., & Kremen, C. (2011). Bees in disturbed habitats use, but do not prefer, alien plants. Basic and Applied Ecology, 12(4), 332-341.

Winfree, R., Bartomeus, I., & Cariveau, D. P. (2011). Native pollinators in anthropogenic habitats. Annual Review of Ecology, Evolution, and Systematics,42(1), 1.