The Cf-4 receptor-like protein associates with the BAK1 receptor-like kinase to initiate

receptor endocytosis and plant immunity

Jelle Postma[1], Thomas W. H. Liebrand[2,3], Guozhi Bi[2,4], Alexandre Evrard[1,5], Ruby R.

Bye[2], Malick Mbengue[1,6], Matthieu H. A. J. Joosten[2,*], Silke Robatzek[1,*]

[1]The Sainsbury Laboratory, Norwich Research Park, Norwich, NR4 7UH, United Kingdom

[2]Laboratory of Phytopathology, Wageningen University, Droevendaalsesteeg 1, 6708 PB

Wageningen, The Netherlands

[3]Present address: Department of Plant Pathology, 576 Hutchison Hall, University of

California, Davis, USA

[4]College of Horticulture, Northeast Agricultural University, Harbin 150030, China

[5]Present address: Vilmorin SA. Route du Manoir, 49250 La Ménitré, France

[6]Present address: French National Institute for Agricultural Research**,** Centre de Recherche de

Toulouse**,** France

[*]Corresponding authors:

E-mail: Matthieu.Joosten@wur.nl

Telephone: 0031 317 483411

E-mail: Robatzek@TSL.ac.uk

Telephone: 0044 1603 450408

Manuscript information:

Word count:

Main body: 5519

Introduction: 1180

Materials and Methods: 1069

Results: 2148

Discussion: 1087

Acknowledgements   35

Figures: 7 (all in colour)

Supporting Information: 10 figures, 2 videos

Summary

• The first layer of plant immunity is activated by cell surface receptor-like kinases (RLKs)

and proteins (RLPs) that detect infectious pathogens. Constitutive interaction with the

RLK SUPPRESSOR OF BIR1 (SOBIR1) contributes to RLP stability and kinase activity.

As RLK activation requires transphosphorylation with a second associated RLK, it

remains elusive how RLPs initiate downstream signaling. To address this, we investigated

functioning of Cf RLPs that mediate immunity of tomato against *Cladosporium fulvum*.

• We employed live-cell imaging and co-immunoprecipitation in tomato and *Nicotiana*

benthamiana to investigate the requirement of associated kinases for Cf activity and

ligand-induced subcellular trafficking of Cf-4.

• Upon elicitation with the matching effector ligands Avr4 and Avr9, BRI1-ASSOCIATED

KINASE 1 (BAK1) associates with Cf-4 and Cf-9. Furthermore, Cf-4 that interacts with

SOBIR1 at the plasma membrane, is recruited to late endosomes after elicitation.

Significantly, BAK1 is required for Avr4-triggered endocytosis, effector-triggered

defenses in Cf-4 plants and resistance of tomato against *C. fulvum*.

• Our observations indicate that RLP-mediated immune signaling and endocytosis require

ligand-induced recruitment of BAK1, reminiscent of BAK1 interaction and subcellular

fate of the FLAGELLIN SENSING 2 RLK. This reveals that diverse classes of cell

surface immune receptors share common requirements for signaling initiation and

endocytosis.

Introduction

The innate immune system of higher eukaryotes senses a wide range of pathogens through

distinct pattern recognition receptors (PRRs). The engagement of these receptors leads to the

activation of signaling pathways resulting in the induction of defense responses. Plants rely

on plasma membrane-resident receptor-like kinases (RLKs) and receptor-like proteins (RLPs)

as a first layer of the immune system against phytopathogenic microbes, many of them

exploiting the extracellular space of plant tissues for their growth (Faulkner & Robatzek,

2012). Therefore, decoding the mechanisms underlying the perception of pathogen-derived

signals by RLKs and RLPs at the host cell surface and subsequent activation of defense

signaling is an important aspect for understanding how plant immunity is initiated. The *Cf*

resistance genes encode leucine-rich repeat (LRR)-RLPs conferring immunity to specific

races of the pathogenic fungus *Cladosporium fulvum* causing leaf mold disease of tomato

(*Solanum lycopersicum*) (Joosten & de Wit, 1999; Rivas & Thomas, 2005; Stergiopoulos &

de Wit, 2009). Distinct members of the Cf receptor family are activated upon recognition of

their matching ligands, which are the so-called avirulence (Avr) proteins that are secreted by

the fungus upon invasion of tomato leaves. The founding member of the Cf proteins and

LRR-RLPs in general is Cf-9, which mediates resistance to Avr9-producing strains of *C.*

fulvum, whereas for example Cf-2 and Cf-4 recognize Avr2 and Avr4, respectively (Rivas &

Thomas, 2005). The Cf receptors mediate race-specific immunity to *C. fulvum*, which is

typically associated with the hypersensitive response (HR), a form of programmed cell death.

Well-characterized PRRs are FLAGELLIN SENSING 2 (FLS2) and ELONGATION

FACTOR-TU RECEPTOR (EFR), which are LRR-RLKs initiating broad-spectrum immunity

against bacterial infection upon detection of the pathogen-associated molecular patterns

(PAMPs) flagellin (flg22) and EF-Tu, respectively (Boller & Felix, 2009). To activate

immune signaling, FLS2 and EFR form ligand-induced complexes with members of the

SOMATIC EMBRYOGENESIS RECEPTOR KINASE (SERK) LRR-RLK family, of which

BRI1-ASSOCIATED KINASE 1 (BAK1)/SERK3 plays a prominent role in FLS2-mediated

immunity (Heese *et al.*, 2007; Roux *et al.*, 2011). BAK1 was initially found to interact with

the LRR-RLK brassinosteroid receptor BRASSINOSTEROID INSENSITIVE 1 (BRI1), and

is involved in a number of immune pathways and developmental processes, in line with its

broader function as a co-receptor (Santiago *et al.*, 2013; Sun *et al.*, 2013; Liebrand *et al.*,

2014). All receptors mentioned above are plasma membrane-localized proteins allowing them

to detect extracellular microbe-derived ligands. In order to reach their destination after their

maturation in the endoplasmic reticulum (ER), they have to be targeted to the plasma

membrane by secretory trafficking (Beck *et al.*, 2012a). As part of the immune response,

plasma membrane-resident FLS2, when activated by flg22, is recruited to ARA7/RabF2b-

and ARA6/RabF1-positive endosomes, which are compartments of the late endosomal

trafficking pathway (Robatzek *et al.*, 2006; Beck *et al.*, 2012c). This process depends on

BAK1/SERK3 (Chinchilla *et al.*, 2007; Beck *et al.*, 2012c), which itself undergoes

constitutive endocytosis (Russinova *et al.*, 2004).

In contrast to RLKs, RLPs lack an intracellular signaling domain and therefore it has

been long suggested that these receptors require the interaction with a protein kinase to

initiate signaling (Jones *et al.*, 1994; Joosten & de Wit, 1999; Liebrand *et al.*, 2014). For

example, the Toll-like receptors (TLRs) of the mammalian innate immune system require

interaction with the MyD88 adaptor protein to recruit the kinase IRAK and subsequently

activate immune signaling (Moresco *et al.*, 2011; Broz & Monack, 2013). Following the

paradigm of the involvement of BAK1/SERK3 in RLK-mediated signaling, there is genetic

evidence for a role of BAK1/SERK3 also in plant defense responses mediated by RLPs. Both

resistance to the fungal pathogen *Verticillium dahliae*, conferred by the tomato LRR-RLP Ve1

and immunity elicited by *Sclerotinia sclerotiorum* through *Arabidopsis thaliana* RLP30, are

dependent on BAK1/SERK3 (Fradin *et al.*, 2009; Fradin *et al.*, 2011; Zhang *et al.*, 2013).

Furthermore, another SERK family member, SERK1, was also shown to be genetically

required for Ve1 and Cf-4 antifungal immunity in *Arabidopsis* and tomato, respectively

(Fradin *et al.*, 2011). However, no molecular interaction between these LRR-RLPs and the

SERKs has been reported, leaving the possibility that the observed phenotypes are indirect,

for example through BAK1/SERK3 function in cell death control or damage-induced

immunity (Kemmerling *et al.*, 2007; Gao *et al.*, 2009; Schwessinger *et al.*, 2011; Liu *et al.*,

2013; Tintor *et al.*, 2013). In addition to its positive regulatory role, BAK1/SERK3 was also

described to negatively regulate immunity in tomato (*Sl*) mediated by *Sl*Eix2, an LRR-RLP

recognizing fungal xylanase (Bar *et al.*, 2010). In a BAK1/SERK3-dependent manner, the

close homologue *Sl*Eix1 impairs *Sl*Eix2 endocytosis and HR (Bar *et al.*, 2010).

Besides genetic evidence for a role of SERK1, no direct involvement in Cf-mediated

immunity of the SERKs has been reported to date. Instead, another LRR-RLK, referred to as

SUPPRESSOR OF BIR1-1/EVERSHED (SOBIR1/EVR), has recently been identified as a

critical component in LRR-RLP-mediated immunity (Jehle *et al.*, 2013; Liebrand *et al.*, 2013;

Zhang *et al.*, 2013; Zhang *et al.*, 2014). *Sl*SOBIR1 interacts specifically with RLPs, including

the Cf-2, Cf-4 and Cf-9 proteins, and this interaction occurred independently of the presence

of their matching Avr ligands. The constitutive association of these LRR-RLPs with

*Sl*SOBIR1 is required for RLP stability and has been proposed to allow activation of a

cytoplasmic signaling cascade upon perception of an Avr by the interacting Cf protein

(Liebrand *et al.*, 2013; Liebrand *et al.*, 2014). It however remains to be elucidated how

ligand-dependent activation of LRR-RLP signaling actually occurs.

Colonization of tomato leaf tissue by *C. fulvum* is fully confined to the apoplast and

the fungus does not form specialized feeding structures like haustoria. Consistent with this

life-style, the Avr proteins secreted by the fungus appear to solely accumulate in the apoplast

(Joosten & de Wit, 1999; Thomma *et al.*, 2005; Stergiopoulos & de Wit, 2009; Joosten, 2012).

This suggests that the Cf receptors perceive the various secreted Avr proteins of *C. fulvum* at

the plasma membrane of the host cells. However, the exact subcellular localization of Cf

proteins has remained unclear ever since their first identification (Jones *et al.*, 1994). Initially

it was shown that Cf-9 is present both at the ER and at the plasma membrane (Piedras *et al.*,

2000). A subsequent study demonstrated a role of a C-terminal dilysine motif of Cf-9 in ER

targeting (Benghezal *et al.*, 2000), which however was later found to be dispensable for Cf-9

function (van der Hoorn *et al.*, 2001). More recently, ER-resident chaperones were identified

as interacting proteins of Cf-4 and were shown to be important for Avr4-triggered immunity

(Liebrand *et al.*, 2012), which provided further evidence for ER localization of these RLPs.

146   Here, we have used live-cell imaging of transiently and stably expressed fluorescent

protein fusions in *N. benthamiana* to investigate the subcellular localization and dynamics of

Cf-4 *in planta*. Our data suggest that Cf-4, together with *Sl*SOBIR1, perceives Avr4 at the

plasma membrane of the host cells, a process that subsequently induces the association of Cf-

4 with BAK1/SERK3, and likewise with SERK1. We furthermore demonstrate a role of these

SERK family members in Cf-mediated immunity and show that the Cf-4 receptor undergoes

BAK1/SERK3-dependent endocytosis upon Avr4 perception.

Materials and Methods

Plant materials and constructs

Nicotiana benthamiana plants were grown under 16 hours of light at 24°C and 45 - 65 %

humidity. Various constructs used in this study have been described before (in all cases GFP

refers to enhanced (e)GFP); Cf-4-GFP, Cf-9-GFP, *Sl*SOBIR1-GFP, *Sl*SOBIR1-like-GFP,

*At*SOBIR1-GFP, *At*SOBIR1-Myc and *At*SOBIR1^{D489N}-Myc (Liebrand *et al.*, 2012; Liebrand

et al., 2013); FLS2-YFPn/c (Frei dit Frey *et al.*, 2012); MEMB12-mCherry (Geldner *et al.*,

2009); VHA-a1-RFP, RFP-ARA7 and ARA6-RFP (Dettmer *et al.*, 2006); AtBAK1

(Schwessinger *et al.*, 2011). Cf-4-YFPn/c and *Sl*SOBIR1-YFPn/c were obtained by PCR-

amplifying the coding regions of the respective genes, which were cloned into the pENTR/D-

TOPO$^{®}$ system (Invitrogen) and subsequently recombined into *pAM-PAT-GWPro$_{35S}$*

(GenBank AY436765). The *35S::ACA8-mCherry* construct was generated by Gateway

recombination (Invitrogen) using pDONR201 containing the coding sequences of *ACA8*

devoid of stop codon (Frei dit Frey *et al.*, 2012) and a *35S::GW-mCherry* destination vector

(provided by R. Panstruga, RWTH Aachen University, Germany). *Sl*SOBIR1-HA was

obtained by recombining pENTR/D-TOPO$^{®}$-*Sl*SOBIR1 (Liebrand *et al.*, 2013) to pGWB14,

carrying the HA coding sequence (Nakagawa *et al.*, 2007). *Sl*SERK1-Myc and *Sl*SERK3a-

Myc were obtained by recombining pENTR/D-TOPO®-*Sl*SERK1 and pDONR201®-

*Sl*SERK3a (Liebrand *et al.*, 2013) to pGWB20 carrying the Myc coding sequence (Nakagawa

et al., 2007), respectively. pUB::*At*BAK1^{D416N} was obtained by PCR amplification from

pGWB14-BAK1^{D416N}-HA (Schwessinger *et al.*, 2011) using primers BAK1-TOPO-F (5'-

CAC CAT GGA ACG AAG ATT AAT GAT C-3') and BAK1-TOPO-R (5'-TTA TCT TGG

ACC CGA GGG GTA TTC-3') introducing a stop codon, directional cloning into pENTR/D-

TOPO® and subsequent recombination of pENTR/D-TOPO-BAK1^{D416N} with pUB-DEST

(Grefen *et al.*, 2010) All recombinations were performed using the classical Gateway LR

clonase reaction (Invitrogen). The correctness of all constructs was confirmed by sequencing.

Transient, stable transformation, and Virus-Induced Gene Silencing (VIGS)

Transient transformation of *N. benthamiana* epidermal cells was done as described before

(Choi *et al.*, 2013). Briefly, two-day cultures of *Agrobacterium tumefaciens* GV3101 carrying

the respective expression constructs in liquid LB medium supplemented with antibiotics were

washed in water prior to *N. benthamiana* leaf infiltrations. For single protein localization and

co-localization purposes, final OD$_{600}$=0.25 and OD$_{600}$=0.5, respectively, were used for agro-

infiltrations. Microscopy was performed at 2-3 dpi. *N. benthamiana* lines stably expressing

Cf-4-GFP were obtained by incubating *N. benthamiana* leaf explants with *Agrobacterium*

strain GV3101 carrying plasmid pBIN-KS-35S::Cf-4-GFP (Liebrand *et al.*, 2012). Selection

of transformed plants was done as described before (Horsch *et al.*, 1985; Gabriëls *et al.*,

2006). Using segregation analysis based on kanamycin resistance, a single locus insertion line

was selected. Tobacco Rattle Virus (TRV)-mediated Virus-Induced Gene Silencing (VIGS) in

N. benthamiana was performed as described before (Liebrand *et al.*, 2012). Briefly, *A.*

tumefaciens GV3101 carrying *TRV-RNA1* at OD$_{600}$ = 0.4 and GV3101 carrying *TRV-RNA2*

containing the respective target sequences at OD$_{600}$ = 0.2 were mixed and co-infiltrated into

N. benthamiana leaves of two-weeks old plants, and three weeks later leaves were used for

further analysis. *TRV* alone was used as a control. The effects on Cf-4-GFP accumulation of

VIGS targeting various endogenous *N. benthamiana* genes in combination with co-

expression of *A. thaliana* genes were observed via immunoblotting and RT-PCR (Supporting

Information Figs. 6c, 9c,d).

Bioassay for Avr4-induced immunity

TRV   constructs   targeting   *NbSERK3a/b*   (Chaparro-Garcia   *et   al.*,   2011),

TRV2::*NbSOBIR1/NbSOBIR1-like* (Liebrand *et al.*, 2013) and TRV2::*Cf-4* (Gabriëls *et al.*,

2006), alongside with TRV2::*GUS* (Tameling & Baulcombe, 2007) controls, were agro-

infiltrated into leaves of one-week-old *N. benthamiana:Cf-4-GFP* plants at O.D.600 = 0.5.

After three weeks, Avr4 (O.D.600 = 0.03, twice) (Gabriëls *et al.*, 2007), RxD460V (O.D.600

= 0.1) (Bendahmane *et al.*, 2002) and BAX (O.D.600 = 0.5) (Lacomme & Santa Cruz, 1999),

were agro-infiltrated into mature, fully expanded leaves of the TRV-inoculated plants. HR

scores were recorded three days after agro-infiltration. In addition, three-week old

NbSERK3a/b-silenced leaves were transiently transformed with Cf-4-GFP and 3 days later

infiltrated with 300μM Avr4 protein. HR was observed 6 days after Avr4 infiltration. VIGS in

tomato, followed by inoculation with conidia of *C. fulvum* race 5-pGPD:*GUS* and subsequent

GUS staining and quantification were performed as described before (Liebrand *et al.*, 2013).

Immunoblot analysis and co-immunoprecipitation

Immunoblot analysis with anti-GFP and anti-Myc antibodies and pull-down experiments

were carried out as previously reported (Liebrand *et al.*, 2012; Liebrand *et al.*, 2013), with the

following modifications: Transiently transformed *N. benthamiana* leaves were infiltrated with

Avr4, Avr2, or Avr9 proteins or flg22 peptide, at the indicated concentrations, and total

proteins were extracted. For detection of the HA-epitope tag the anti-HA-Biotin High

Affinity Antibody (clone 3F10; Roche Applied Science) was used. In contrast to the

theoretical mass of Cf-4-GFP (app. 123 kDa) Immunoblot analysis revealed a specific band at

about 140 kDa in accordance with what was observed in previous studies (Liebrand *et al.*,

2012).

qRT-PCR analysis

For qRT-PCR, total RNA was isolated from *N. benthamiana:Cf-4-GFP* leaf material, at 2

weeks after agro-inoculation with the various VIGS constructs. RNA extraction, cDNA

synthesis and qRT-PCR were performed as described (Liebrand *et al.*, 2012). *NbSERK3a/b*

expression was investigated using primers rbo16 (5'-TGC GCT GAA GAC CAA CTT GGC

T-3') and rbo17 (5'-CTG AAG CTT GCC CAA TGT GTC G-3'). *NbSERK1* expression was

investigated using primers rbo20 (5'-ATT GCA CAG TCT GCG T-3') and rbo21 (5'-CGA

AGG AAT CTC AAT TTA GTC-3'). Expression of *NbSOBIR1* was investigated using

primers to266 and to267 (Liebrand *et al.*, 2013). Expression of endogenous actin was used to

calibrate the expression level of the query genes, as previously described (Liebrand *et al.*,

2012). qRT-PCRs to determine the expression levels of *AtBAK1* and *AtBAK1*^{D416N} were

performed using primers *At*BAK1-F (5'-TGG ACT TGC AAA ACT CAT GG-3') and

*At*BAK1-R (5'-GAT CAA AAG CCC TTT GTC CA-3').

Confocal microscopy and image analysis

Confocal laser microscopy was performed using a Leica SP5 laser point scanning microscope

(Leica, Germany) mounted with hybrid detectors (HyD™) as described previously (Beck *et*

al., 2012c). Briefly, GFP and RFP/mCherry fluorophores were excited using the 488-nm

argon laser and the 561 nm diode, respectively, and fluorescence emission was captured

between 500 and 550 nm for GFP and between 580 and 620 nm for RFP/mCherry. For GFP

only images, chloroplast autofluorescence was captured between 700 and 800 nm. Sequential

scan mode was used for simultaneous imaging of GFP and RFP/mCherry. Abaxial sides of *N.*

benthamiana leaf discs were imaged. Images were taken using a 63X water immersion

objective and processed using the Leica LAS-AF and FIJI (ImageJ) software packages. Spot

detection and quantification on confocal micrographs were performed using EndoQuant,

which is a modification of EndomembraneQuantifier suitable for standard confocal images

(Beck *et al.*, 2012c).

Results

Cf-4 interacts with SlSOBIR1 at the plasma membrane

To address the localization and dynamics of the Cf-4 receptor in antifungal immunity, we

used functional fluorescently-tagged Cf-4 (Cf-4-GFP) mediating Avr4-triggered HR in *N.*

benthamiana (Supporting Information Fig. S1; (Liebrand *et al.*, 2012)). We monitored Cf-4-

GFP subcellular localization and revealed its presence at the plasma membrane of leaf

epidermal cells by co-localizing Cf-4-GFP with the plasma membrane autoinhibitory calcium

ATPase ACA8, fused to mCherry (Fig. 1, Supporting Information Fig. S2; (Frei dit Frey *et al.*,

2012)). The cell surface localization of this Cf protein is in agreement with the described

apoplastic localization of its ligand, Avr4, (Joosten *et al.*, 1994; van den Burg *et al.*, 2006)

and the plasma membrane-localization of its constitutive interactor, SOBIR1 (Fig. 2.; (Leslie

et al., 2010; Liebrand *et al.*, 2013)). *Sl*SOBIR1-GFP, its close homolog *Sl*SOBIR1-like fused

to GFP and also *At*SOBIR1-GFP were all detected at the plasma membrane, co-localizing

with ACA8-mCherry, and internal vesicles (Figs. 1, 2, Supporting Information Figs. S2, S3,

S4).

To investigate the subcellular localization of the Cf-4-*Sl*SOBIR1 complex, we used

bimolecular fluorescence complementation (BiFC). We observed reconstitution of the YFP

molecule by the detection of fluorescence when transiently co-expressing Cf-4 and

*Sl*SOBIR1 that were C-terminally fused to respectively the C- and N-terminal halves of YFP

(respectively YFPc and YFPn), indicative of Cf-4-*Sl*SOBIR1 heterodimerization (Fig. 1). Cf-

4-*Sl*SOBIR1 heterodimerization was found at the plasma membrane, which was confirmed by

ACA8-mCherry co-localization (Fig. 1). In this assay, we did not detect positive BiFC when

co-expressing FLS2 and *Sl*SOBIR1 fused to the YFP halves, whereas BiFC did occur when

co-expressing FLS2-YFPc and FLS2-YFPn (Supporting Information Fig. S5). These

observations indicate homodimerization of FLS2 but no heterodimerization between FLS2

and *Sl*SOBIR1, which is consistent with previous findings and thus supporting the specificity

of BiFC in this system (Frei dit Frey *et al.*, 2012; Sun *et al.*, 2012; Liebrand *et al.*, 2013). All

these results show that Cf-4 localizes at the plasma membrane, where this LRR-RLP interacts

with the RLK *Sl*SOBIR1.

SOBIR1 localizes to endosomes

In addition to its localization at the plasma membrane, and in accordance to what was shown

in previous studies, *At*SOBIR1 was also observed at internal vesicles that showed co-labeling

with the endocytic tracer FM4-64, indicative of endosomal localization (Leslie *et al.*, 2010).

To examine the identity of these SOBIR1-positive vesicles, we transiently co-expressed

*Sl*SOBIR1-GFP, *Sl*SOBIR1-like-GFP and *At*SOBIR1-GFP with described fluorescent

markers of the endomembrane trafficking pathways. These include MEMB12-mCherry for

labeling Golgi compartments, VHA-a1-RFP for labeling the *trans*-Golgi network (TGN) and

the Rab5 GTPases RFP-ARA7/RabF2b and ARA6/RabF1-RFP for labeling endosomes

(Geldner *et al.*, 2009; Beck *et al.*, 2012c). No co-localization between the SOBIR1-positive

vesicles and MEMB12- and VHA-a1-labeled compartments was detected (Fig. 2, Supporting

Information Fig. S3a,b). By contrast, we found clear localization of the SOBIR1 receptors to

RFP-ARA7/RabF2b- and ARA6/RabF1-RFP-positive endosomes (Fig. 2, Supporting

Information Fig. S3c,d). These observations are in agreement with the described FM4-64-

positive endosomal localization of *At*SOBIR1-YFP in *Arabidopsis* (Leslie *et al.*, 2010), and

suggest that constitutive endocytosis of SOBIR1 receptors takes place. Furthermore, co-

localization with ARA6/RabF1 indicates that the SOBIR1 receptors might enter the late

endosomal pathway (Beck *et al.*, 2012c).

Avr4 triggers endocytosis of the Cf-4/SOBIR1 complex

The observation that Cf-4 interacts with *Sl*SOBIR1 at the plasma membrane, together with

the finding that SOBIR1 receptors are present at endosomes, prompted us to investigate

whether Cf-4 is endocytosed when activated by Avr4. To test this, we treated *N. benthamiana*

leaves transiently and stably expressing Cf-4-GFP with purified Avr4 and Avr2 proteins, of

which the latter is specifically recognized by the Cf-2 receptor. When elicited with Avr4, this

triggered internalization of Cf-4-GFP (Fig. 3a, Supporting Information Fig. S6). Co-

expression with ARA6/RabF1-RFP showed a strong overlap of this marker with the Avr4-

induced Cf-4-GFP-positive vesicles, thereby revealing endosomal localization of activated

Cf-4 (Fig. 3, Supporting Information Videos S1, S2). Cf-4-As expected, GFP maintained

plasma membrane localization upon control treatment with Avr2, indicating that endocytosis

of Cf-4 is ligand-specific and depends on the activation of the Cf-4 receptor (Fig. 3a,

Supporting Information Fig. S6). Consistent with SOBIR1 localizing to endosomes (Fig. 2,

Supporting Information Fig. S3), we observed co-localization of Avr4-induced Cf-4-GFP-

positive endosomes and endosomal *Sl*SOBIR1-mCherry (Fig. 3).

As both, SOBIR1 and Cf-4 underwent endocytosis and localized at late endosomal

compartments we questioned whether they might get co-internalized. Therefore we

investigated whether Cf-4 is endocytosed together with *Sl*SOBIR1. Transient co-expression

of Cf-4 and *Sl*SOBIR1, C-terminally fused to the C- and N-terminal halves of YFP,

respectively, results in positive BiFC at the plasma membrane (Fig. 1) and when triggering

with Avr4, BiFC was also observed at internal vesicles (Fig. 3b). These vesicles also showed

co-localization with ARA6/RabF1-RFP, demonstrating endosomal localization of the

reconstituted Cf-4-*Sl*SOBIR1 BiFC heterodimer (Fig. 3b), and in agreement with Avr4-

induced co-localization of Cf-4-GFP and *Sl*SOBIR1-mCherry at endosomes (Fig. 3a).

Treatment with Avr2 did not trigger re-localization of the plasma membrane-resident

reconstituted Cf-4-*Sl*SOBIR1 BiFC heterodimer (Fig. 3b), further supporting ligand-specific

internalization of this Cf-4 together with *Sl*SOBIR1. These results indicate that the receptors

can be endocytosed as heterodimers, an observation that was previously made for the BRI1-

BAK1 complex in *Arabidopsis* protoplasts (Russinova *et al.*, 2004). In contrast to this finding,

we did not observe flg22-induced endocytosis of the FLS2 homodimer as the reconstituted

BiFC signal was maintained at the plasma membrane after elicitation (Supporting

Information Fig. S5b). Endocytosis of activated FLS2 depends on BAK1/SERK3 (Chinchilla

et al., 2007; Beck *et al.*, 2012c), and this may indicate that the reconstituted FLS2 BiFC

homodimer could be affected in complex formation with BAK1.

Cf-4 endocytosis requires BAK1/SERK3

Because flg22-activated FLS2 traffics via the ARA6/RabF1-positive late endosomal pathway and depends on functional BAK1/SERK3 (Beck *et al.*, 2012c; Choi *et al.*, 2013), this raises the possibility that Avr4-induced endocytosis of Cf-4 also involves BAK1/SERK3. To test this, we applied virus-induced gene silencing (VIGS), using a tobacco rattle virus (TRV) construct targeting the two *NbSERK3* homologues (*NbSERK3a* and *b*) in *N. benthamiana* (Chaparro-Garcia *et al.*, 2011). Monitoring Cf-4-GFP localization in leaves of *TRV::NbSERK3a/b*-inoculated *N. benthamiana* revealed a strongly reduced amount of Avr4-induced Cf-4-GFP-positive endosomes, as compared to the situation in control leaves from plants that had been inoculated with *TRV* only (Fig. 4, Supporting Information Fig. S10b), while Cf-4 protein levels were unaltered (Supporting Information Fig. S10c). By using a heterologous functional complementation approach, Avr4-induced endocytosis of Cf-4-GFP was partially restored when *AtBAK1* was transiently expressed in *NbSERK3a/b*-silenced leaves. No reduction was observed in the amount of ARA6/RabF1-labelled endosomes when *NbSERK3a/b* was silenced (Fig. 4, Supporting Information Fig. S10b,c). Importantly, this suggests that, as is the case for the RLK FLS2, ligand-induced endocytosis of the RLP Cf-4 is BAK1/SERK3-dependent, and that this RLK plays a role in Cf-4 function. This finding is further strengthened by expression of a kinase-inactive *AtBAK1* variant (*AtBAK1-KD*) in *NbSERK3a/b*-silenced leaves that did not restore Cf-4-GFP endocytosis upon triggering with Avr4 (Fig. 4, Supporting Information Fig. S10b,c).

As a next step, we investigated whether SOBIR1 is required for Avr4-induced Cf-4 endocytosis using a similar heterologous functional complementation approach. In agreement with previous findings revealing that Cf-4 abundance is dependent on SOBIR1 (Liebrand *et al.*, 2013), Cf-4-GFP levels in *N. benthamiana* were increased by transient expression of *At*SOBIR1-Myc and its kinase-inactive variant (Supporting Information Fig. S7c). Importantly, transient expression of *At*SOBIR1-Myc but not kinase-inactive *At*SOBIR1-KD-Myc restored Avr4-induced endocytosis of Cf-4-GFP in *NbSOBIR1/-like*-silenced leaves (Supporting Information Fig. S7a,b.). Altogether, these data indicate that the kinase activities of both BAK1 and SOBIR1 are required for endocytosis of Cf-4 upon Avr4 recognition, suggesting the active removal of triggered receptors from the plasma membrane.

Cf-4 associates with SERK members in a ligand-depending manner

We addressed a possible role of BAK1/SERK3 in Cf-4-mediated immunity by co-

immunoprecipitation experiments. Cf-4-GFP was purified from *N. benthamiana* leaves

transiently co-expressing Cf-4-GFP, *Sl*SOBIR1-HA and *Sl*SERK3a-Myc, that had either been

mock-treated or challenged with Avr4, Avr2 or flg22 at two days after agro-infiltration of the

three constructs. While C-terminally tagged BAK1/SERK3 fusion proteins are not signalling

competent after flg22 and elf18 trigger during immunity their recruitment behavior to the

FLS2 receptor complex is still intact, which suggests suitability for interaction studies with

other receptors (Ntoukakis *et al.*, 2011). In all cases, *Sl*SOBIR1-HA was detected upon

immunoprecipitation of Cf-4-GFP, independent of the treatment conditions (Fig. 5,

Supporting Information Fig. S8). This confirms the constitutive interaction between these two

receptors as shown by BiFC (Fig. 1) and as was previously reported (Liebrand *et al.*, 2013).

By contrast, *Sl*SERK3a-Myc was only revealed upon immunoprecipitation of Cf-4-GFP from

agro-infiltrated leaves that had first been infiltrated with the matching ligand Avr4, but not

when infiltrated with Avr2 or flg22, which was included as control treatment (Fig. 5,

Supporting Information Fig. S8). Importantly, this demonstrates that ligand-dependent hetero-

complex formation of RLPs as the Cf-4 receptor with *Sl*SERK3a takes place, in a way that is

similar to what has been shown for RLK-type PRRs (Monaghan & Zipfel, 2012; Liebrand *et*

al., 2014).

Because of the previous genetic evidence for a role of *Sl*SERK1 in Cf-mediated immunity

(Fradin *et al.*, 2011), we examined whether Cf-4 also interacts with this co-receptor by co-

immunoprecipitation experiments after transient co-expression of the various receptors in

N. benthamiana leaves. Unlike what is the case for *Sl*SERK3a-Myc, *Sl*SERK1-Myc was

revealed after immunoprecipitation of Cf-4-GFP, independent of elicitation of Cf-4 by Avr4

(Supporting Information Fig. S8). However, the amount of co-immunoprecipitated *Sl*SERK1-

Myc was strongly enhanced in the Avr4-treated leaves, as compared to the leaves treated with

Avr2. We cannot exclude the possibility that this differential pattern between Cf-4 interaction

with *Sl*SERK1 and *Sl*SERK3a is caused by the higher steady state expression levels of

*Sl*SERK1-Myc compared to those of *Sl*SERK3a-Myc (Supporting Information Fig. S8; see

input). However, following the recent notion of the presence of BRI1-BAK1/SERK3

preassembled heterodimers (Bücherl *et al.*, 2013), our data could also indicate that Cf-4 exists

in a preformed complex with *Sl*SOBIR1 and *Sl*SERK1, which becomes stabilized and/or

recruits increased amounts of *Sl*SERK1 upon elicitation with Avr4.

407   We next explored the possibility that ligand-induced recruitment of *Sl*SERK1 and -3

could exist as a more general mechanism of Cf receptor complex activation. To this end, we

purified Cf-9-GFP from *N. benthamiana* leaves transiently co-expressing this RLP together

with *Sl*SOBIR1-HA and either *Sl*SERK1-Myc or *Sl*SERK3a-Myc. Similar to what we

observed for activated Cf-4, both *Sl*SERK1-Myc and *Sl*SERK3a-Myc were strongly co-

immunoprecipitated with Cf-9-GFP when elicited with its matching ligand effector Avr9 but,

as expected, not significantly with Avr4 that was included as control treatment (Supporting

Information Fig. S9). This supports the idea that additional SOBIR1-dependent RLPs, such as

Ve1, RLP30 and ReMax (Fradin *et al.*, 2009; Fradin *et al.*, 2011; Zhang *et al.*, 2013), also

recruit members of the SERK family to form active receptor complexes.

SERK members mediate Avr4-triggered immunity

Avr4 triggers HR in Cf-4-expressing *N. benthamiana*, which can be diminished by silencing

of the gene encoding the Cf-4 receptor itself or *NbSOBIR1/-like* ((Liebrand *et al.*, 2013); Fig.

6a). To determine whether BAK1/SERK3 is involved in Cf-4-mediated immunity, we

examined the Avr4-triggered HR in Cf-4-GFP-expressing *N. benthamiana* leaves (transiently

and stably), without and with silencing of *NbSERK3a/b*. We consistently found that the Avr4-

triggered HR was significantly reduced in *NbSERK3a/b*-silenced Cf-4-GFP-expressing leaves,

as compared to the leaves of *TRV* control plants (Fig. 6a, Supporting Information Fig. S10a).

NbSERK3a/b-silencing did not generally affect programmed cell death, as this was still

induced by the auto-active variant of the nucleotide binding-LRR immune receptor RxD460V

and the pro-apoptotic factor BCL2-ASSOCIATED PROTEIN X (BAX) (Lacomme & Santa

Cruz, 1999; Bendahmane *et al.*, 2002). The capability of the *NbSERK3a/b*-VIGS construct to

knock down the expression of both *NbSERK3a/b* gene homologues was previously shown

(Chaparro-Garcia *et al.*, 2011). We further investigated the specificity of *NbSERK3a/b*-

silencing within the SERK family and found that also a likely *NbSERK1* homologue was

knocked down upon expression of the *NbSERK3a/b*-VIGS construct (Supporting Information

Fig. S10b). Thus, it could be that Avr4-triggered HR in *N. benthamiana* requires not only

BAK1/SERK3 but also SERK1, an observation that is consistent with the earlier finding that

VIGS of *SlSERK1* compromises Cf-4-mediated immunity to *C. fulvum* in tomato (Fradin *et*

al., 2011).

To examine whether BAK1/SERK3, in addition to SERK1, is also involved in Cf-4-mediated

resistance to Avr4-secreting *C. fulvum* strains, their encoding genes were silenced in Cf-4-

expressing tomato plants. A strain secreting Avr4 and expressing the *GUS* reporter gene

exhibited strong colonization of tomato leaves lacking the Cf-4 receptor (MM-Cf-0) and

when the *Cf-4* gene was silenced (Fig. 6b). Colonization by this fungus was significantly

more abundant in Cf-4-expressing tomato leaves silenced for *SlSERK3*, as increased numbers

of successful colonization attempts were observed compared to the *GUS*-silenced negative

control. This phenotype is reminiscent of the effect of silencing of *SlSOBIR1(-like)* on Cf-4-

mediated resistance to *C. fulvum* (Liebrand *et al.*, 2013). We furthermore silenced *SlSERK3*

together with *SlSERK1*, which also compromised resistance to the pathogen. Taken together,

these results show that BAK1/SERK3 is a key positive regulator of full Cf-4-mediated

immunity induced upon Avr4 perception.

Discussion

The Cf signaling pathway is essential for the immune response of tomato to *C. fulvum* (Rivas

& Thomas, 2005; Stergiopoulos & de Wit, 2009). A number of Cf signaling components were

identified through genetic and proteomic approaches, but the mechanism by which Cf-4

initiates downstream signaling remained unclear (Liebrand *et al.*, 2014). Here, we show that

both SERK1 and BAK1/SERK3 are recruited to the Cf-4 receptor in an Avr4-dependent

manner, a mechanism we additionally confirmed for Avr9-induced activation of Cf-9.

Consistently, we show that Cf-4 requires these RLKs for its function. Our observations imply

that Avr4 induces the formation of a complex of Cf-4 and *Sl*SERK1/3 to induce Cf-4

signaling. Furthermore, we confirm that Cf-4 interacts with SOBIR1 at the plasma membrane

and, more importantly demonstrate ligand-induced SERK3 dependent late endocytic

trafficking of the Cf-4 RLP together with *Sl*SOBIR1 as a novel pathway in Cf-mediated

downstream events.

465   The requirement of BAK1/SERK3 hints at clear similarities between the Cf-4/9

effector receptors and FLS2 MAMP receptor pathways, further evidenced by overlapping

transcriptional reprogramming upon activating FLS2- and Cf-mediated immune responses

(Navarro *et al.*, 2004). This overlap potentially involves the regulation of downstream

responses through similar components which, in addition to the SERK family members,

include E3 ligases (PUB12/13 for FLS2 and CMPG1 for Cf-4 immunity; (Gilroy *et al.*, 2011;

Lu *et al.*, 2011)) and receptor-like cytoplasmic kinases (BIK1/PBL1 for FLS2, ACIK1 for Cf-

4 and CAST AWAY for SOBIR1 (EVR) signaling; (Burr *et al.*, 2011; Monaghan & Zipfel,

2012; Liebrand *et al.*, 2014)). Given these similarities, it is conceivable that the constitutive

association between the RLP Cf-4 and the RLK *Sl*SOBIR1 represents a PRR complex, in

which the Cf-4 ectodomain mediates specific ligand recognition and the *Sl*SOBIR1 kinase

domain is regarded as the signaling part. This is in contrast to PRRs exemplified by FLS2, in

which both functions are present within the same molecule. Ligand-induced interaction of

PRRs with SERK member RLKs is subsequently required to trigger downstream signaling by

both RLP- and RLK-type PRRs (Fig. 7).

Regarding Cf-4 as a PRR is in agreement with the observation that Cf-4 not only recognizes

the Avr4 protein secreted by *C. fulvum*, but is also activated by the Avr4 homologue produced

by the banana pathogen *Mycosphaerella fijiensis* (Stergiopoulos *et al.*, 2010). Both *C. fulvum*

and *M. fijiensis* Avr4 proteins bind to chitin (van den Burg *et al.*, 2003; Stergiopoulos *et al.*,

2010) and it was suggested that, reminiscent of PAMP recognition, the chitin-binding motif

of Avr4 is the pattern that is recognized through interaction with the LRRs of Cf-4 (Thomma

et al., 2011). As elicitation with Avr4 appears to maintain the Cf-4 association with

*Sl*SOBIR1 ((Liebrand *et al.*, 2013); Fig. 5) and induces the interaction of Cf-4 with

*Sl*SERK1/3, this could suggest the possibility that a complex consisting of Cf-4, *Sl*SOBIR1

and *Sl*SERK3a is formed. In turn, this could result in increased phosphorylation events

between the *Sl*SOBIR1 and *Sl*SERK1/3 kinases, which then trigger downstream signaling

(Fig. 7), similar to what has been proposed for the FLS2-BAK1 model (Schwessinger *et al.*,

2011). Furthermore, our data show that *Sl*SERK1 and BAK1/*Sl*SERK3 are also recruited to

the Cf-9 receptor upon its activation with Avr9, supporting the idea of a general mechanism

by which SOBIR1-dependent RLPs form active receptor complexes. Although no molecular

interactions have been reported so far, a similar scenario could be proposed for RLP30, which

also genetically depends on SOBIR1 and BAK1/SERK3 (Zhang *et al.*, 2013). A role for

SERK receptors is less clear for *Sl*Eix2, because in this case BAK1 was found to negatively

regulate this receptor through interaction with its close homologue *Sl*Eix1 (Bar *et al.*, 2010).

Future studies should determine whether, similar to the trans-phosphorylation of FLS2 by

BAK1/SERK3 (Schwessinger *et al.*, 2011), the phosphorylation status of the kinase domain

of SOBIR1 also alters upon SERK1/3 recruitment.

Receptor-mediated endocytosis is part of the eukaryotic immune response and for

example is found for the FLS2 receptor (Husebye *et al.*, 2006; Robatzek *et al.*, 2006; Spallek

et al., 2013). An important role of receptor-mediated endocytosis is to control the abundance

of receptor (complexes) at the plasma membrane, a process that is well established in animals

and involves lysosomal/vacuolar degradation (Lemmon & Schlessinger, 2010). Activated

FLS2 traffics into the late endosomal pathway and is a cargo of multivesicular bodies

localizing to the lumen of these late endosomes for delivery to the vacuole (Beck *et al.*,

2012c; Spallek *et al.*, 2013). This pathway could be responsible for the flg22-induced FLS2

degradation, because chemicals affecting endosomal trafficking inhibit the degradation of this

receptor (Smith *et al.*, 2014). Our co-localization data strongly suggest that activated Cf-4 is

also targeted for vacuolar degradation through the late endosomal pathway, consistent with

the observation that Cf-4-GFP protein levels are reduced upon Avr4 elicitation (Supporting

Information Fig. S1a). Endosomal sorting for vacuolar degradation requires the transfer of

ubiquitin to the plasma membrane cargo and subsequent deubiquitination of the cargo at

multivesicular bodies (Beck *et al.*, 2012c). The ubiquitin E3 ligase CMPG1 and the

deubiquitinating enzyme UBP12 are known positive and negative regulators of Cf-4- and Cf-

9-mediated HR, respectively (Ewan *et al.*, 2011; Gilroy *et al.*, 2011), and given their

biochemical function could also be involved in the endocytosis and degradation of Cf

proteins.

Ligand-induced activation triggers the formation of a hetero-complex consisting of

Cf-4 and *Sl*SERK1/3. This differs from the situation with TLRs from mammals, which recruit

cytoplasmic kinases to initiate signaling, but in plants is reminiscent of the FLS2 pathway

that also depends on interaction with the co-receptor BAK1/SERK3 to signal. Endocytic

removal of activated PRRs, which is also known for TLR4 (Husebye *et al.*, 2006), might

provide a mechanism to regulate receptor presence at the plasma membrane and associated

events (Felix *et al.*, 1998; Beck *et al.*, 2012a; Smith *et al.*, 2014). Understanding how plant

cells regulate PRR subcellular localization is essential as pathogens are likely to target

components of the trafficking pathway to suppress plant defenses, as was recently found for

the *Phytophthora infestans* Avr3a effector (Chaparro-Garcia *et al.*) and bacterial HopM1

(Nomura *et al.*, 2011). This latter effector of *Pseudomonas syringae* targets a host ADP

ribosylation factor guanine nucleotide exchange factor, referred to as *At*MIN7/BEN1,

resulting in its breakdown. *At*MIN7/BEN1 is required for PAMP-triggered immunity and

regulates endosomal trafficking possibly at the TGN, where HopM1 can also be found

(Nomura *et al.*, 2006; Nomura *et al.*, 2011). It remains to be seen whether *C. fulvum* also

secretes effectors that enter inside plant cells and are able to alter host subcellular trafficking

to promote infection success.

Acknowledgements

We thank the members of the Robatzek laboratory for fruitful discussions, Brande Wulff for

reading the manuscript and Cyril Zipfel for providing *At*BAK1 constructs. Zeinu Mussa is

acknowledged for help with *N. benthamiana* transformations.

References

Bar M, Sharfman M, Ron M, Avni A. 2010. BAK1 is required for the attenuation of ethylene-inducing xylanase (Eix)-induced defense responses by the decoy receptor LeEix1. *The Plant Journal* **63**(5): 791-800.

Beck M, Heard W, Mbengue M, Robatzek S. 2012a. The INs and OUTs of pattern recognition receptors at the cell surface. *Current Opinion in Plant Biology* **15**(4): 367-374.

Beck M, Zhou J, Faulkner C, Mac D, Robatzek S. 2012c. Spatio-temporal cellular dynamics of the *Arabidopsis* flagellin receptor reveal activation status-dependent endosomal sorting. *The Plant Cell* **24**(10): 4205-4219.

Bendahmane A, Farnham G, Moffett P, Baulcombe DC. 2002. Constitutive gain-of-function mutants in a nucleotide binding site-leucine rich repeat protein encoded at the *Rx* locus of potato. *The Plant Journal* **32**(2): 195-204.

Benghezal M, Wasteneys GO, Jones DA. 2000. The C-terminal dilysine motif confers endoplasmic reticulum localization to type I membrane proteins in plants. *The Plant Cell* **12**(7): 1179-1201.

Boller T, Felix G. 2009. A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annual review of plant Biology* **60**: 379-407.

Broz P, Monack DM. 2013. Newly described pattern recognition receptors team up against intracellular pathogens. *Nature Reviews Immunology* **13**(8): 551-565.

Bücherl CA, van Esse GW, Kruis A, Luchtenberg J, Westphal AH, Aker J, van Hoek A, Albrecht C, Borst JW, de Vries SC. 2013. Visualization of BRI1 and BAK1(SERK3) membrane receptor heterooligomers during Brassinosteroid signaling. *Plant Physiology* **162**(4): 1911-1925.

Burr CA, Leslie ME, Orlowski SK, Chen I, Wright CE, Daniels MJ, Liljegren SJ. 2011. Cast away, a membrane-associated receptor-like kinase, inhibits organ abscission in Arabidopsis. *Plant Physiology* **156**(4): 1837-1850.

Chaparro-Garcia A, Schwizer S, Sklenar J, Yoshida K, Bos JIB, Schornack S, Jones AME, Bozkurt TO, Kamoun S. Phytophthora infestans RXLR-WY effector AVR3a associates with a Dynamin-Related Protein involved in endocytosis of a plant pattern recognition receptor. *biorXiv* **doi: 10.1101/012963**.

Chaparro-Garcia A, Wilkinson RC, Gimenez-Ibanez S, Findlay K, Coffey MD, Zipfel C,

Rathjen JP, Kamoun S, Schornack S. 2011. The receptor-like kinase Serk3/Bak1 is
required for basal resistance against the late blight pathogen *Phytophthora infestans* in
Nicotiana benthamiana. *PLoS ONE* **6**(1).

Chinchilla D, Zipfel C, Robatzek S, Kemmerling B, Nürnberger T, Jones JDG, Felix G,
Boller T. 2007. A flagellin-induced complex of the receptor FLS2 and BAK1 initiates
plant defence. *Nature* **448**(7152): 497-500.

Choi SW, Tamaki T, Ebine K, Uemura T, Ueda T, Nakano A. 2013. RABA members act in distinct
steps of subcellular trafficking of the FLAGELLIN SENSING2 receptor. *The Plant*
Cell **25**(3): 1174-1187.

Dettmer J, Hong-Hermesdorf A, Stierhof YD, Schumacher K. 2006. Vacuolar H+-ATPase
activity is required for endocytic and secretory trafficking in Arabidopsis. *The Plant*
Cell **18**(3): 715-730.

Ewan R, Pangestuti R, Thornber S, Craig A, Carr C, O'Donnell L, Zhang C,
Sadanandom A. 2011. Deubiquitinating enzymes AtUBP12 and AtUBP13 and their
tobacco homologue NtUBP12 are negative regulators of plant immunity. *New*
Phytologist **191**(1): 92-106.

Faulkner C, Robatzek S. 2012. Plants and pathogens: Putting infection strategies and
defence mechanisms on the map. *Current Opinion in Plant Biology* **15**(6): 699-707.

Felix G, Baureithel K, Boller T. 1998. Desensitization of the perception system for chitin
fragments in tomato cells. *Plant Physiology* **117**(2): 643-650.

Fradin EF, Abd-El-Haliem A, Masini L, van den Berg GCM, Joosten MHAJ, Thomma
BPHJ. 2011. Interfamily transfer of tomato *Ve1* mediates *Verticillium* resistance in
Arabidopsis. *Plant Physiology* **156**(4): 2255-2265.

Fradin EF, Zhang Z, Ayala JCJ, Castroverde CDM, Nazar RN, Robb J, Liu CM,
Thomma BPHJ. 2009. Genetic dissection of *Verticillium* wilt resistance mediated by
tomato Ve1. *Plant Physiology* **150**(1): 320-332.

Frei dit Frey NF, Mbengue M, Kwaaitaal M, Nitsch L, Altenbach D, Häweker H,
Lozano-Duran R, Njo MF, Beeckman T, Huettel B, et al. 2012. Plasma membrane
calcium ATPases are important components of receptor-mediated signaling in plant
immune responses and development. *Plant Physiology* **159**(2): 798-809.

Gabriëls SHEJ, Takken FLW, Vossen JH, De Jong CF, Liu Q, Turk SCHJ, Wachowski
LK, Peters J, Witsenboer HMA, de Wit PJGM, et al. 2006. cDNA-AFLP
combined with functional analysis reveals novel genes involved in the hypersensitive
response. *Molecular Plant-Microbe Interactions* **19**(6): 567-576.

Gabriëls SHEJ, Vossen JH, Ekengren SK, van Ooijen G, Abd-El-Haliem AM, van den
Berg GCM, Rainey DY, Martin GB, Takken FLW, de Wit PJGM, et al. 2007. An
NB-LRR protein required for HR signalling mediated by both extra- and intracellular
resistance proteins. *The Plant Journal* **50**(1): 14-28.

Gao M, Wang X, Wang D, Xu F, Ding X, Zhang Z, Bi D, Cheng YT, Chen S, Li X, et al.
2009. Regulation of cell death and innate immunity by two receptor-like kinases in
Arabidopsis. *Cell Host & Microbe* **6**(1): 34-44.

Geldner N, Dénervaud-Tendon V, Hyman DL, Mayer U, Stierhof YD, Chory J. 2009.
Rapid, combinatorial analysis of membrane compartments in intact plants with a
multicolor marker set. *The Plant Journal* **59**(1): 169-178.

Gilroy EM, Taylor RM, Hein I, Boevink P, Sadanandom A, Birch PRJ. 2011. CMPG1-
dependent cell death follows perception of diverse pathogen elicitors at the host
plasma membrane and is suppressed by Phytophthora infestans RXLR effector
AVR3a. *New Phytologist* **190**(3): 653-666.

Grefen C, Donald N, Hashimoto K, Kudla J, Schumacher K, Blatt MR. 2010. A
ubiquitin-10 promoter-based vector set for fluorescent protein tagging facilitates
temporal stability and native protein distribution in transient and stable expression
studies. *Plant J* **64**(2): 355-365.

Heese A, Hann DR, Gimenez-Ibanez S, Jones AME, He K, Li J, Schroeder JI, Peck SC,
Rathjen JP. 2007. The receptor-like kinase SERK3/BAK1 is a central regulator of
innate immunity in plants. *Proceedings of the National Academy of Sciences* **104**(29):
12217-12222.

Horsch RB, Fry JE, Hoffmann NL, Eichholtz D, Rogers SG, Fraley RT. 1985. A simple
and general method for transferring genes into plants. *Science* **227**(4691): 1229-1230.

Husebye H, Halaas Ø, Stenmark H, Tunheim G, Sandanger Ø, Bogen B, Brech A, Latz
E, Espevik T. 2006. Endocytic pathways regulate Toll-like receptor 4 signaling and
link innate and adaptive immunity. *EMBO Journal* **25**(4): 683-692.

Jehle AK, Fürst U, Lipschis M, Albert M, Felix G. 2013. Perception of the novel MAMP
eMax from different *Xanthomonas* species requires the *Arabidopsis* receptor-like
protein ReMAX and the receptor kinase SOBIR. *Plant Signaling & Behavior* **8**(12):
e27408.

Jones DA, Thomas CM, Hammond-Kosack KE, Balint-Kurti PJ, Jones JDG. 1994.
Isolation of the tomato *Cf-9* gene for resistance to *Cladosporium fulvum* by
transposon tagging. *Science* **266**(5186): 789-793.

Joosten MHAJ. 2012. Isolation of apoplastic fluid from leaf tissue by the vacuum infiltration-centrifugation technique. *Methods in molecular biology* **835**: 603-610.

Joosten MHAJ, Cozijnsen TJ, De Wit PJGM. 1994. Host resistance to a fungal tomato pathogen lost by a single base-pair change in an avirulence gene. *Nature* **367**(6461): 384-386.

Joosten MHAJ, de Wit PJGM. 1999. The tomato-*Cladosporium fulvum* interaction: A versatile experimental system to study plant-pathogen interactions. *Annual Review of Phytopathology* **37**: 335-367.

Kemmerling B, Schwedt A, Rodriguez P, Mazzotta S, Frank M, Qamar SA, Mengiste T, Betsuyaku S, Parker JE, Müssig C, et al. 2007. The BRI1-associated kinase 1, BAK1, has a brassinolide-independent role in plant cell-death control. *Current Biology* **17**(13): 1116-1122.

Lacomme C, Santa Cruz S. 1999. Bax-induced cell death in tobacco is similar to the hypersensitive response. *Proceedings of the National Academy of Sciences of the United States of America* **96**(14): 7956-7961.

Lemmon MA, Schlessinger J. 2010. Cell signaling by receptor tyrosine kinases. *Cell* **141**(7): 1117-1134.

Leslie ME, Lewis MW, Youn JY, Daniels MJ, Liljegren SJ. 2010. The EVERSHED receptor-like kinase modulates floral organ shedding in Arabidopsis. *Development* **137**(3): 467-476.

Liebrand TWH, Smit P, Abd-El-Haliem A, De Jonge R, Cordewener JHG, America AHP, Sklenar J, Jones AME, Robatzek S, Thomma BPHJ, et al. 2012. Endoplasmic reticulum-quality control chaperones facilitate the biogenesis of Cf receptor-like proteins involved in pathogen resistance of tomato. *Plant Physiology* **159**(4): 1819-1833.

Liebrand TWH, van den Berg GCM, Zhang Z, Smit P, Cordewener JHG, America AHP, Sklenar J, Jones AME, Tameling WIL, Robatzek S, et al. 2013. The receptor-like kinase SOBIR1/EVR interacts with receptor-like proteins in plant immunity against fungal infection. *Proceedings of the National Academy of Sciences of the United States of America* **110**(24): 10010-10015.

Liebrand TWH, van den Burg HA, Joosten MHAJ. 2014. Two for all: receptor-associated kinases SOBIR1 and BAK1. *Trends in Plant Science* **19**(2): 123-132.

Liu Z, Wu Y, Yang F, Zhang Y, Chen S, Xie Q, Tian X, Zhou JM. 2013. BIK1 interacts with PEPRs to mediate ethylene-induced immunity. *Proceedings of the National*

Academy of Sciences of the United States of America **110**(15): 6205-6210.

Lu D, Lin W, Gao X, Wu S, Cheng C, Avila J, Heese A, Devarenne TP, He P, Shan L.

2011. Direct ubiquitination of pattern recognition receptor FLS2 attenuates plant

innate immunity. *Science* **332**(6036): 1439-1442.

Monaghan J, Zipfel C. 2012. Plant pattern recognition receptor complexes at the plasma

membrane. *Current Opinion in Plant Biology* **15**: 349-357.

Moresco EMY, LaVine D, Beutler B. 2011. Toll-like receptors. *Current Biology* **21**(13):

R488-R493.

Nakagawa T, Kurose T, Hino T, Tanaka K, Kawamukai M, Niwa Y, Toyooka K,

Matsuoka K, Jinbo T, Kimura T. 2007. Development of series of gateway binary

vectors, pGWBs, for realizing efficient construction of fusion genes for plant

transformation. *Journal of Bioscience and Bioengineering* **104**(1): 34-41.

Navarro L, Zipfel C, Rowland O, Keller I, Robatzek S, Boller T, Jones JDG. 2004. The

transcriptional innate immune response to flg22. Interplay and overlap with Avr gene-

dependent defense responses and bacterial pathogenesis. *Plant Physiology* **135**(2):

1113-1128.

Nomura K, Debroy S, Lee YH, Pumplin N, Jones J, He SY. 2006. A bacterial virulence

protein suppresses host innate immunity to cause plant disease. *Science* **313**(5784):

220-223.

Nomura K, Mecey C, Lee YN, Imboden LA, Chang JH, He SY. 2011. Effector-triggered

immunity blocks pathogen degradation of an immunity-associated vesicle traffic

regulator in Arabidopsis. *Proceedings of the National Academy of Sciences of the*

United States of America **108**(26): 10774-10779.

Ntoukakis V, Schwessinger B, Segonzac C, Zipfel C. 2011. Cautionary Notes on the Use of

C-Terminal BAK1 Fusion Proteins for Functional Studies. *The Plant Cell* **23**(11):

3871-3878.

Piedras P, Rivas S, Dröge S, Hillmer S, Jones JDG. 2000. Functional, c-myc-tagged *Cf-9*

resistance gene products are plasma-membrane localized and glycosylated. *The Plant*

Journal **21**(6): 529-536.

Rivas S, Thomas CM. 2005. Molecular interactions between tomato and the leaf mold

pathogen *Cladosporium fulvum*. *Annual Review of Phytopathology* **43**: 395-436.

Robatzek S, Chinchilla D, Boller T. 2006. Ligand-induced endocytosis of the pattern

recognition receptor FLS2 in *Arabidopsis*. *Genes and Development* **20**(5): 537-542.

Roux M, Schwessinger B, Albrecht C, Chinchilla D, Jones A, Holton N, Malinovsky FG,

Tör M, de Vries S, Zipfel C. 2011. The *Arabidopsis* leucine-rich repeat receptor-like
kinases BAK1/SERK3 and BKK1/SERK4 are required for innate immunity to
hemibiotrophic and biotrophic pathogens. *The Plant Cell* **23**(6): 2440-2455.

Russinova E, Borst JW, Kwaaitaal M, Caño-Delgado A, Yin Y, Chory J, de Vries SC.
2004. Heterodimerization and endocytosis of Arabidopsis brassinosteroid receptors
BRI1 and AtSERK3 (BAK1). *The Plant Cell* **16**(12): 3216-3229.

Santiago J, Henzler C, Hothorn M. 2013. Molecular mechanism for plant steroid receptor
activation by somatic embryogenesis co-receptor kinases. *Science* **341**(6148): 889-
892.

Schwessinger B, Roux M, Kadota Y, Ntoukakis V, Sklenar J, Jones A, Zipfel C. 2011.
Phosphorylation-dependent differential regulation of plant growth, cell death, and
innate immunity by the regulatory receptor-like kinase BAK1. *PLoS Genetics* **7**(4):
e1002046.

Smith JM, Salamango DJ, Leslie ME, Collins CA, Heese A. 2014. Sensitivity to Flg22 Is
modulated by ligand-induced degradation and de novo synthesis of the endogenous
flagellin-receptor FLAGELLIN-SENSING2. *Plant Physiology* **164**(1): 440-454.

Spallek T, Beck M, Ben Khaled S, Salomon S, Bourdais G, Schellmann S, Robatzek S.
2013. ESCRT-I mediates FLS2 endosomal sorting and plant immunity. *PLoS Genetics*
9(12): e1004035.

Stergiopoulos I, de Wit PJGM. 2009. Fungal effector proteins. *Annual Review of*
Phytopathology **47**: 233-263.

Stergiopoulos I, van den Burg HA, Ökmen B, Beenen HG, van Liere S, Kema GHJ, de
Wit PJGM. 2010. Tomato Cf resistance proteins mediate recognition of cognate
homologous effectors from fungi pathogenic on dicots and monocots. *Proceedings of*
the National Academy of Sciences of the United States of America **107**(16): 7610-
7615.

Sun W, Cao Y, Jansen KL, Bittel P, Boller T, Bent AF. 2012. Probing the *Arabidopsis*
flagellin receptor: FLS2-FLS2 association and the contributions of specific domains
to signaling function. *The Plant Cell* **24**(3): 1096-1113.

Sun Y, Li L, Macho AP, Han Z, Hu Z, Zipfel C, Zhou J-m, Chai J. 2013. Structural basis
for flg22-induced activation of the *Arabidopsis* FLS2-BAK1 immune complex.
Science **342**(6158): 624-628.

Tameling WIL, Baulcombe DC. 2007. Physical association of the NB-LRR resistance
protein Rx with a Ran GTPase-activating protein is required for extreme resistance to

Potato virus X. The Plant Cell **19**(5): 1682-1694.

Thomma BPHJ, Nürnberger T, Joosten MHAJ. 2011. Of PAMPs and effectors: the

blurred PTI-ETI dichotomy. *The Plant Cell* **23**(1): 4-15.

Thomma BPHJ, Van Esse HP, Crous PW, De Wit PJGM. 2005. *Cladosporium fulvum*

(syn. *Passalora fulva*), a highly specialized plant pathogen as a model for functional

studies on plant pathogenic *Mycosphaerellaceae*. *Molecular Plant Pathology* **6**(4):

379-393.

Tintor N, Ross A, Kanehara K, Yamada Y, Fan L, Kemmerling B, Nürnberger T, Tsuda

K, Saijo Y. 2013. Layered pattern receptor signaling via ethylene and endogenous

elicitor peptides during *Arabidopsis* immunity to bacterial infection. *Proceedings of*

the National Academy of Sciences of the United States of America **110**(15): 6211-6216.

van den Burg HA, Harrison SJ, Joosten MHAJ, Vervoort J, de Wit PJGM. 2006.

Cladosporium fulvum Avr4 protects fungal cell walls against hydrolysis by plant

chitinases accumulating during infection. *Molecular Plant-Microbe Interactions*

19(12): 1420-1430.

van den Burg HA, Westerink N, Francoijs KJ, Roth R, Woestenenk E, Boeren S, de Wit

PJGM, Joosten MHAJ, Vervoort J. 2003. Natural disulfide bond-disrupted mutants

of AVR4 of the tomato pathogen *Cladosporium fulvum* are sensitive to proteolysis,

circumvent Cf-4-mediated resistance, but retain their chitin binding ability. *Journal of*

Biological Chemistry **278**(30): 27340-27346.

van der Hoorn RAL, van der Ploeg A, de Wit PJGM, Joosten MHAJ. 2001. The C-

terminal dilysine motif for targeting to the endoplasmic reticulum is not required for

Cf-9 function. *Molecular Plant-Microbe Interactions* **14**(3): 412-415.

Zhang L, Kars I, Essenstam B, Liebrand TWH, Wagemakers L, Elberse J, Tagkalaki P,

Tjoitang D, van den Ackerveken G, van Kan JAL. 2014. Fungal

endopolygalacturonases are recognized as microbe-associated molecular patterns by

the *Arabidopsis* receptor-like protein RESPONSIVENESS TO BOTRYTIS

POLYGALACTURONASES1. *Plant Physiology* **164**(1): 352-364.

Zhang W, Fraiture M, Kolb D, Löffelhardt B, Desaki Y, Boutrot F, Tör M, Zipfel C,

Gust AA, Brunner F. 2013. The *Arabidopsis thaliana* receptor-like protein RLP30

and receptor-like kinase SOBIR1/EVR mediate innate immunity toward necrotrophic

fungi. *The Plant Cell* **25**(10): 4227-4241.

Figure legends

Fig. 1 Cf-4 and *Sl*SOBIR1 are present at the plasma membrane. Confocal micrographs show *N. benthamiana* leaf epidermal cells transiently co-expressing the indicated Cf-4 and *Sl*SOBIR1 fusion proteins and plasma membrane-localized ACA8-mCherry. The first panels show GFP/YFP fluorescence, the second panels show mCherry fluorescence, the third panels depict the overlay images of the two fluorescence signals shown in the first and second panels. Overlay images indicate co-localization of the proteins fused to GFP or YFP and mCherry, as a yellow colour is produced. Dashed squares in these images are shown as detail pictures (magnified in the fourth panels). White lines in the detail pictures indicate the ROIs that correspond to intensity profiles in the last panels. Intensity profiles display grey value of pixels across the ROI in the green and red channels, on a scale of 1-300. Images were taken at three days post infiltration (dpi) for Cf-4-GFP and *Sl*SOBIR1-GFP and at two dpi for BiFC of Cf-4, C-terminally fused to the C-terminal half of YFP (YFPc), with *Sl*SOBIR1 C-terminally fused to the N-terminal half of YFP (YFPn); scale bars = 10 μm.

Fig. 2 *Sl*SOBIR1 constitutively localizes to endosomes. Confocal micrographs show *N. benthamiana* leaf epidermal cells transiently expressing *Sl*SOBIR1-GFP or Cf-4-GFP (left panels), either without or with co-expression of the indicated organelle markers fused to mCherry/RFP (middle, left panels). Co-localization in the overlay images is depicted by the development of a yellow colour (middle, right panels). Dashed squares in these images are shown as detail pictures (magnified in the right panels). Arrowheads in the detail pictures point at mobile vesicles in the upper right panel and indicate *Sl*SOBIR1 localization at endosomes in the two lower right panels. Images were taken at three dpi; scale bars = 10 μm.

Fig. 3 Cf-4 localizes to endosomes in a ligand-dependent manner and together with *Sl*SOBIR1. (**a**) Confocal micrographs show *N. benthamiana* leaf epidermal cells transiently expressing Cf-4-GFP without and with co-expression of ARA6-RFP and *Sl*SOBIR1-mCherry, treated with Avr4 or Avr2 (both at 100μM), as indicated. The left panels show GFP fluorescence, middle left panels show autofluorescence and RFP/mCherry fluorescence, middle right panels depict the overlay images of the two fluorescence signals shown in the left and middle left panels, and the right panels show detail pictures of the dashed squares. Arrowheads point at mobile vesicles positive for Cf-4-GFP and ARA6-RFP and indicate co-localization of Cf-4-GFP with ARA6-RFP and *Sl*SOBIR1-mCherry at endosomes. Images were taken at three dpi and 90 min after elicitation; scale bars = 10 μm. (**b**) Confocal

micrographs show *N. benthamiana* leaf epidermal cells transiently expressing Cf-4, C-terminally fused to the C-terminal half of YFP (YFPc) and *Sl*SOBIR1, C-terminally fused to the N-terminal half of YFP (YFPn) (left panels), without or with co-expression of ARA6-RFP (middle left panels) and treated with Avr4 or Avr2 (both at 100µM), as indicated. Co-localization between reconstituted YFP and RFP in the overlay images is depicted by the development of a yellow colour (middle, right panels). Dashed squares in these panels are shown as detail pictures (right panels). Arrowheads point at mobile vesicles in the upper right panel and indicate co-localization at endosomes in the middle right panel. Images were taken at three dpi and 90 min after elicitation; scale bars = 10 µm.

Fig. 4 Cf-4 endocytosis requires BAK1/SERK3. Leaves of Cf-4-*N. benthamiana* stable and *N. benthamiana* plants were TRV-silenced for *NbSERK3a/b* and *GUS* as a control for three weeks and subsequently used to transiently express *At*BAK1, *At*BAK1-KD, and ARA6/RabF1-RFP as indicated for three days. Confocal micrographs show Cf-4-GFP localisation upon treatment with Avr4 (100µM, left panels) and ARA6/RabF1-RFP-labelled endosomes (left panels), and detail pictures from dashed squares (middle panels). Arrowheads point at Cf-4-GFP- and ARA6/RabF1-RFP-positive vesicles. Quantification of Cf-4-GFP- and ARA6/RabF1-RFP-positive vesicles was done with EndoQuant (right panels, bars depict means ± 2 SE; n = 6; $p < 0.05$; statistical significant differences are indicated by asterisks). Note, much fewer Cf-4-GFP-positive vesicles are observed in *NbSERK3a/b*-silenced leaves compared to the *GUS*-silenced control. The amount of Cf-4-GFP-positive vesicles increased upon transient co-expression of *At*BAK1 but not when its kinase-inactive variant *At*BAK1-KD was co-expressed. No difference in the amount of ARA6/RabF1-RFP-positive endosomes was observed when *NbSERK3a/b* was silenced. Images were taken 90 min after Avr4 elicitation; scale bars = 10 µm. See Supporting information Fig. S10b,c for transcript abundance of silenced genes and protein levels of Cf-4.

Fig. 5 *Sl*SERK3a interacts with Cf-4. Co-immunoprecipitation from GFP-trap bead pull-downs on *N. benthamiana* co-expressing Cf-4-GFP, *Sl*SOBIR1-HA and *Sl*SERK3a-Myc. Leaf samples were taken at two dpi, after 60 min treatments without or with 10µM flg22 or 100µM Avr4, as indicated. Total proteins (Input) and immunoprecipitated proteins (IP) were subjected to SDS/PAGE and blotted. Blots were incubated with αGFP, αHA or αMyc antibodies for the detection of immunoprecipitated Cf-4-GFP and co-purifying *Sl*SOBIR1-HA and *Sl*SERK3a-Myc, respectively.

Fig. 6 *Sl*SERK3a is required for Avr4-triggered HR and immunity against *C. fulvum*. (a) Images show leaves of transgenic *N. benthamiana* plants stably expressing Cf-4-GFP that had been *TRV*-silenced for either *Cf-4*, *NbSOBIR1*, *NbSERK3a/b* or for *GUS* as control, and were subsequently agro-infiltrated with Avr4 (O.D.600 = 0.03), RxD460V (O.D.600 = 0.1) and BAX (O.D.600 = 0.5) as indicated. Images were taken three days after agro-infiltration. HR is observed as brownish cell death. The numbers below the panels indicate the occurrence of full HR, intermediate or no symptoms out of 36 Avr4 agro-infiltrations that were performed. (**b**) Images show leaves two weeks after inoculation with an Avr4-secreting, *GUS*-transgenic strain of *C. fulvum* of MM-Cf-0 tomato as a control, and Cf-4 tomato that had been inoculated with recombinant *TRV* constructs targeting *Cf-4*, *SlSERK3, SlSERK1* or *GUS* three weeks earlier. To visualize *C. fulvum* colonization, leaves were stained for GUS activity. The amount of successful colonization attempts (blue spots) versus the total amount of Cf-4 leaves that were sampled is indicated between parentheses.

Fig. 7 Model of the Cf-4, SOBIR1 and SERK3 subcellular trafficking pathways. Following its folding and maturation in the endoplasmic reticulum (ER), Cf-4 is predominantly trafficked to the plasma membrane, where it constitutively interacts with SOBIR1, which itself is constitutively endocytosed. Upon colonization of Cf-4 tomato leaves, *C. fulvum* secretes Avr4 into the apoplast, which is recognized by Cf-4 and induces interaction of Cf-4 with BAK1/SERK3. This complex is required for Cf-4-mediated immunity (not shown) and endocytosis into ARA7- and ARA6-positive compartments of the late endosomal pathway, destined for vacuolar degradation. Subcellular localization of markers (ACA8, MEMB12, VHA-a1, ARA6 and ARA7) used in this study is depicted in brown lettering. SV, secretory vesicle; V, vesicle; TGN, *trans*-Golgi network; EE, early endosome; LE, late endosome; MVB, multivesicular body.

Supporting Information

Fig. S1 GFP-tagged Cf-4 migrates with the predicted molecular weight and is functional. (**a**) Western blot analysis of Cf-4-GFP transiently and stably expressed in *N. benthamiana*, without and with Avr4 (100µM) treatment, as indicated. For transient expression, leaf samples were harvested at three days post infiltration (dpi) and Avr4 elicitation was done for 90 min. Blots were incubated with anti-GFP antibodies for the detection of Cf-4-GFP. CBB,

Coomassie Brilliant Blue staining. (**b**) Images of *N. benthamiana* wild type (WT) leaves (top

panel) and leaves transiently expressing Cf-4-GFP (bottom panel), treated with either Avr2 or

Avr4 (300µM), as indicated. Images were taken at five dpi and at four days after treatment

with the Avrs.

Fig. S2 Subcellular localization of Cf-4, *Sl*SOBIR1-like and *At*SOBIR1. Confocal

micrographs show *N. benthamiana* leaf epidermal cells stably expressing Cf-4-GFP and

transiently expressing *Sl*SOBIR1-like-GFP and *At*SOBIR1-GFP as indicated (left panels), and

co-expressing plasma membrane-localized ACA8-mCherry (middle panels). Overlay images

indicate co-localization of the proteins fused to GFP and mCherry, as a yellow colour is

produced (right panels). Images were taken at three dpi; scale bars = 10 µm.

Fig. S3 *Sl*SOBIR1-like-GFP and *At*SOBIR1-GFP co-localize with endosomal markers.

Confocal micrographs show *N. benthamiana* leaf epidermal cells transiently expressing

*Sl*SOBIR1-like-GFP or *At*SOBIR1-GFP (left panels), and co-expressed mCherry/RFP-tagged

organelle markers (middle, left panels). Overlay images indicate co-localization through

generation of a yellow colour (middle, right panels). Dashed boxes indicated in the middle

right panels are depicted as detail pictures on the right (right panels). Arrowheads point at co-

localizing endosomes. Images were taken at three dpi; scale bars = 10 µm. (**a**) Co-expression

of *Sl*SOBIR1-like-GFP/*At*SOBIR1-GFP with Golgi marker MEMB12-mCherry. (**b**) Co-

expression of *Sl*SOBIR1-like-GFP/*At*SOBIR1-GFP with *trans*-Golgi network marker VHA-

a1-RFP. (**c**) Co-expression of *Sl*SOBIR1-like-GFP/*At*SOBIR1-GFP with endosome marker

RFP-ARA7/RabF2b. (**d**) Co-expression of *Sl*SOBIR1-like-GFP/*At*SOBIR1-GFP with late

endosome marker ARA6/RabF1-RFP.

Fig. S4. Line intensity profiles of Cf-4, SOBIR1 and membrane markers. Confocal

micrographs show transient co-expression of Cf-4-GFP, *Sl*SOBIR1-GFP or Cf-4-

YFPc/*Sl*SOBIR1-YFPn with indicated membrane markers and effector treatments. Left

panels correspond to confocal micrographs displayed in (**a**) Fig. 1, (**b**) Fig. 2a, (**c**) Fig. 2b.

Dashed squares in these panels are shown as detail pictures (magnified in middle panels).

White lines in detail pictures indicate the ROIs that correspond to intensity profiles in the last

panels. Intensity profiles display grey value of pixels across the ROI in the green and red

channels on a scale of 1-300.

Fig. S5 *Sl*SOBIR1 does not interact with FLS2 in BiFC experiments. (**a**) Confocal
micrographs show *N. benthamiana* leaf epidermal cells transiently co-expressing FLS2 and
*Sl*SOBIR1 C-terminally fused to the C- or N-terminal halves of YFP (YFPc and YFPn,
respectively), as indicated. Left panels show absence or presence of YFP fluorescence, the
middle panels show autofluorescence, and the right panels show the overlay of the
fluorescence signals. Images were taken at two dpi, scale bars = 100 μm. (**b**) Confocal
micrographs of *N. benthamiana* leaf epidermal cells transiently co-expressing FLS2-YFPc
and FLS2-YFPn, either without or with flg22 treatment (10μM) for 90 min, as indicated. Left
panels show absence or presence of YFP fluorescence, the middle panels show
autofluorescence, and the right panels show the overlay of the fluorescence signals. Images
were taken at two dpi, scale bars = 10 μm.

Fig. S6 Cf-4-GFP endocytosis in stable transgenic plants. Confocal micrographs of Cf-4-
GFP-transgenic *N. benthamiana* plants show leaf epidermal cells expressing Cf-4-GFP, either
treated with Avr4 or Avr2 (both at 100 μM), as indicated. Left panels show GFP fluorescence,
middle left panels show autofluorescence, and middle right panels show the overlay of the
fluorescence signals shown in the left and middle left panels. Arrowheads in the detail
pictures (right panels) point at mobile vesicles (top right panel). Images were taken at 90
minutes after elicitation; scale bars = 10 μm.

Fig. S7 Cf-4 endocytosis requires SOBIR1 kinase activity. Leaves of Cf-4-GFP transgenic
N. benthamiana plants that had been TRV-silenced for *NbSOBIR1/-like* were used for
transient expression of *AtSOBIR1-Myc* and its kinase-inactive variant *AtSOBIR1-KD-Myc*. (**a**)
Confocal micrographs show Cf-4-GFP localisation upon treatment with Avr4 (100μM, left
panels) and detail pictures from dashed squares (middle panels). Arrowheads point at Cf-4-
GFP-positive vesicles. Images were taken at three weeks after inoculation with *TRV*, at three
dpi for transient expression, and at 90 min after elicitation; scale bars = 10 μm. (**b**)
Quantification of Cf-4-GFP-positive vesicles was done with EndoQuant (bars depict means ±
2 SE; n = 6; $p < 0.05$; statistical significant differences are indicated by asterisks). Transient
co-expression of *At*BAK1, but not its kinase-inactive variant *At*BAK1-KD, increased Cf-4-
GFP-positive vesicles in *NbSOBIR1/-like*-silenced leaves. (**c**) Immunoblots from extracted
total proteins of Cf-4-GFP-transgenic *N. benthamiana* plants that were TRV-silenced for
NbSOBIR1/-like and subsequently transiently transformed with *AtSOBIR1-Myc* or *AtSOBIR1-*
KD-Myc. Cf-4-GFP, *At*SOBIR1-Myc and *At*SOBIR1-KD-Myc were revealed with αGFP and

αMyc antibodies as indicated. Ponceau staining is shown for equal loading.

Fig. S8 Both *Sl*SERK1 and *Sl*SERK3a interact with the Cf-4. Co-immunoprecipitation

from GFP-trap bead pull-downs on *N. benthamiana* co-expressing Cf-4-GFP and *Sl*SOBIR1-

HA, and either *Sl*SERK3a-Myc or *Sl*SERK1-Myc. Leaf samples were taken at two dpi, after

60 min treatment with Avr2 or Avr4 (100μM), as indicated. Total proteins (Input) and

immunoprecipitated proteins (IP) were subjected to SDS/PAGE and blotted. Blots were

incubated with αGFP, αHA or αMyc antibodies for the detection of immunoprecipitated Cf-4-

GFP and co-purifying *Sl*SOBIR1-HA and *Sl*SERK3a-Myc or *Sl*SERK1-Myc, respectively.

Fig. S9 Avr9 induces interaction of Cf-9 with both *Sl*SERK1 and *Sl*SERK3a. Co-

immunoprecipitation from GFP-trap bead pull-downs on *N. benthamiana* co-expressing Cf-9-

GFP and *Sl*SOBIR1-HA, and either *Sl*SERK1-Myc or *Sl*SERK3a-Myc. Leaf samples were

taken at two dpi, after 60 min treatment with Avr9 or Avr4 (100μM), as indicated. Total

proteins (Input) and immunoprecipitated proteins (IP) were subjected to SDS/PAGE and

blotted. Blots were incubated with αGFP, αHA or αMyc antibodies for the detection of

immunoprecipitated Cf-9-GFP and co-purifying *Sl*SOBIR1-HA and *Sl*SERK1-Myc or

*Sl*SERK3a-Myc, respectively.

Fig. S10 *Sl*SERK3 is required for Avr4-induced HR. (**a**) Images show *N. benthamiana*

leaves from *TRV::NbSERK3a/b*-inoculated plants and *TRV*-inoculated controls, transiently

expressing Cf-4-GFP and treated with Avr4 protein (300μM). Cf-4-GFP was agro-infiltrated

at 3 weeks post inoculation with the recombinant *TRV* constructs and one day later the Avr4

protein was infiltrated. Images were taken at six days after Avr4 infiltration. HR is observed

as brownish cell death. (**b**) qRT-PCR analysis showing *NbSERK1*, *NbSERK3a/b* and

NbSOBIR1 expression in *N. benthamiana* inoculated with the indicated *TRV* VIGS constructs.

Expression of query genes was normalized to endogenous *NbActin* expression levels. Data

presented have been combined from twelve (*NbSOBIR1* expression) or 24 (*NbSERK1* and

NbSERK3 expression) individual qRT-PCRs, based on three independent biological replicates,

for each gene of which the expression was studied. Statistical differences (α=0.05) between

groups were calculated by a one-way ANOVA and different groupings are indicated. Error

bars represent the standard deviation. (**c**) Immunoblots from total protein extracts of Cf-4-

GFP-transgenic *N. benthamiana* plants that were TRV-silenced for *NbSERK3a/b* and

subsequently transiently transformed with *AtBAK1* and *AtBAK1-KD*. Cf-4-GFP was revealed

with αGFP. Ponceau staining is shown for equal loading. (**d**) RT-PCR on cDNA obtained

from total RNA extracts of Cf-4-GFP-transgenic *N. benthamiana* plants that were TRV-

silenced for *NbSERK3a/b* and subsequently transiently transformed with *AtBAK1* and

AtBAK1-KD.

Video S1. Mobile vesicles in transient Cf-4-GFP expression. Time series of confocal

micrographs of *N. benthamiana* leaf epidermal cells transiently expressing Cf-4-GFP and

treated with Avr4 (100 µM). Images were taken at 3 days post infiltration (dpi) and 1.5 hours

after Avr4 treatment. Frame counter is displayed in the top left. Acquisition speed is ~4

seconds per frame. Scale bar = 10 µm.

Video S2. Mobile vesicles in stable Cf-4-GFP expression. Time series of confocal

micrographs of *N. benthamiana* leaf epidermal cells stably expressing Cf-4-GFP and treated

with Avr4 (100 µM). Images were taken 1.5 hours after Avr4 treatment. Frame counter is

displayed in the top left. Acquisition speed is ~4 seconds per frame. Scale bar = 10 µm.

SlSOBIR1-GFP			overlay	detail
	autofluorescence			
	MEMB12-mCherry			
	VHA-a1-RFP			
	RFP-ARA7			
	ARA6-RFP			
Cf-4-GFP	autofluorescence			

(a)

| Cf-4-GFP | | overlay | detail |

+ Avr4 — autofluorescence

+ Avr4 — ARA6-RFP

+ Avr2 — ARA6-RFP

+ Avr4 — SlSOBIR1-mCherry

(b)

Cf-4-YFPc
SlSOBIR1-YFPn | | overlay | detail |

+ Avr4 — autofluorescence

+ Avr4 — ARA6-RFP

+ Avr2 — autofluorescence

(a)

TRV::Cf-4	TRV::NbSOBIR1	TRV::NbSERK3a/b	TRV control
full: 6/36	full: 5/36	full: 2/36	full: 27/36
intermediate: 7/36	intermediate: 4/36	intermediate: 8/36	intermediate: 9/36
none: 23/36	none: 27/36	none: 26/36	none: 0/36

Avr4, RxLR460V, Avr4, BAX labels on leaves

(b)

TRV::

SlSERK3a	SlSERK3a SlSERK1	Cf-4	GUS	no TRV
(364/36)	(98/31)	(>500/32)	(12/44)	

Cf-4   MM-Cf-0