Phylodynamics of H5N1 Highly Pathogenic Avian Influenza in Europe, 2005–2010: Potential for molecular surveillance of new outbreaks

Mohammad A. Alkhamis[1,3*], Brian R. Moore[2], Andres M. Perez[3]

[1]Environmental and Life Sciences Research Center, Kuwait Institute For Scientific Research, Kuwait City, Safat 13109, Kuwait; E-Mail: mkhamis@kisr.edu.kw (M.A.)
[2]Department of Evolution and Ecology, Center for Population Biology, University of California Davis, Davis, California, 95616, USA; E-Mail: brianmoore@ucdavis.edu (B.R.M.)
[3]Department of Veterinary Population Medicine, College of Veterinary Medicine, University of Minnesota, St. Paul, Minnesota, 55108, USA; E-Mail: aperez@umn.edu (A.P.)

*Author to whom correspondence should be addressed; mkhamis@kisr.edu.kw (M.A.); Tel: +96599144485; +15302195295

Abstract: Previous Bayesian phylogeographic studies of H5N1 highly pathogenic avian influenza viruses (HPAIVs) explored the origin and spread of the epidemic from China into Russia, indicating that HPAIV circulated in Russia prior to its detection there in 2005. In this study, we extend this research to explore the evolution and spread of HPAIV within Europe during the 2005–2010 epidemic, using all available sequences of the HA and NA gene regions that were collected in Europe and Russia during the outbreak. We use discrete-trait phylodynamic models within a Bayesian statistical framework to explore the evolution of HPAIV. Our results indicate that the genetic diversity and effective population size of HPAIV peaked between mid-2005 and early 2006, followed by drastic decline in 2007, which coincides with the end of the epidemic in Europe. Our results also suggest that domestic birds were the most likely source of the spread of the virus from Russia into Europe. Additionally, estimates of viral dispersal routes indicate that Russia, Romania, and Germany were key epicenters of these outbreaks. Our study quantifies the dynamics of a major European HPAIV pandemic and substantiates the ability of phylodynamic models to improve molecular surveillance of novel AIVs.

Keywords: H5N1; Highly Pathogenic Avian Influenza; phylodynamic models; Bayesian inference; phylogeography; Europe; Russia; Surveillance

1. Introduction

The epidemiology of the H5N1 highly pathogenic avian influenza viruses (HPAIVs) is characterized both by the far-reaching economic impact for commercial poultry production, and also by the risk of spread into mammalian species, including humans. Transmission into humans has mainly been attributed to direct contact of susceptible individuals with infected poultry, including activities associated with the slaughter and dressing of poultry for human consumption [1]. The maintenance and spread of avian influenza (AI) infection in domesticated poultry are mainly associated with the establishment of the virus in free-range (or backyard) poultry and its subsequent introduction to commercial animals via live-bird markets. Furthermore, wild birds are the natural reservoirs of the virus, and the movement of wild birds infected with HPAIV is known to play an important role in the spread, circulation, and maintenance of the virus through either direct or indirect contact with poultry [2–4].

The ancestral strain of H5N1 HPAIV, referred to as A/goose/Guangdong/1/96, emerged in commercial domesticated geese in the Guangdong province of China in 1996 [5]. Since then, the strain has undergone numerous genetic changes that have given rise to several distinct lineages, also referred to as clades [6]. To date, ten H5N1 HPAIV clades have been identified based on comparisons of 859 hemagglutinin (HA) gene sequences [7]. These viruses are now widespread in countries throughout Asia, Europe, and Africa. Between 2004 and 2005, a new H5N1 HPAIV strain (sub-clade 2.2) emerged that caused severe outbreaks in wild birds and poultry in northwestern China [6, 8]. In Europe, HPAI H5N1 was first detected at the end of 2005 in both poultry (in Russia, Romania, and Turkey) and wild birds (in Croatia) [2, 9–11]. In January 2006, the first human infection outside Southeast Asia was reported in Turkey. In February 2006, the first of occurrence of HPAI H5N1 in Northern Europe was identified from a wild mute swan (*Cygnus olor*) in Germany in the southwestern part of the Baltic Sea [11,12]. Between 2006 and 2010, the epidemic spread through 13 European Union (EU) member states located eastward of a line extending from south-eastern Sweden to south-western Italy. Both poultry and wild-bird populations were infected in Austria, Bulgaria, Czech Republic, Denmark, France, Greece, Hungary, Poland, Slovak Republic, Slovenia, Switzerland, and the UK [10, 13]. Phylogenetic analysis of HPAIVs isolated from European outbreaks distinguished at least two different sub-lineages of the 2005 Qinghai strain, which confirmed that the virus spread from southeastern Asia into Europe [10-13].

Since 2005, surveillance of avian influenza viruses in poultry and wild birds has become compulsory in the EU. Member states have conducted extensive molecular surveillance for AI, with most of the studies focused on genetic analysis of partially or fully sequenced AI viral genomes [14–18]. Molecular epidemiological studies of H5N1 HPAIV sequence data isolated between 2005–2010 in EU member states, for example, were focused on the molecular characterization, genetic classification, identification of genetic shifts and drifts, and the degree of relatedness to previously isolated viruses. Phylogenies of H5N1 HPAIV isolated in the EU were estimated using traditional parsimony, neighbor-joining, and maximum-likelihood methods. However, those methods typically ignore various sources of uncertainty, including uncertainty associated with estimates of the phylogenetic relationships, divergence times, and history of geographic dispersal [19]. Furthermore, epidemiological studies that explored risk factors and the spatial and temporal evolution of H5N1 HPAIV in the EU between 2005–2010 were mainly conducted separately from phylogenetic studies [20-24]. Phylodynamics is an emerging field that aims to characterize the joint evolutionary and epidemical behavior of rapidly

evolving infectious diseases using tools borrowed from the field of phylogenetics [25]. The probabilis-
tic models used in the phylodynamic approach are mostly pursued in a Bayesian statistical framework
[26]. This approach treats parameters of the phylodynamic model as random variables, such that each
parameter is described by a specified prior probability distribution (and a corresponding inferred poste-
rior probability distribution). Accordingly, the Bayesian approach provides a natural way to estimate
(and accommodate) uncertainty in the phylodynamic model parameters, including the virus phylogeny,
divergence times, and history of geographic spread [27].

Recent evidence suggests that Bayesian methods may be more accurate in assessing HPAIV evolu-
tion than traditional phylogenetic methods. For example, Lemey *et al.* (2009) [27] adopted a Bayesian
phylodymanic approach to re-analyze a H5N1 HPAIV dataset initially studied using conventional
methods [28, 29]. Results of the Bayesian analysis did not support the earlier conclusion regarding the
epidemiological link between Guangdong and Indonesia, but instead suggested that the ancestral
strains of the Indonesian outbreaks originated from the Hunan province in south central China [27].
Furthermore, these studies demonstrated the potential of Bayesian phylogeographic methods to esti-
mate the posterior probability of each geographic location at any point in the phylogenetic tree, and the
ability for phylogenetic parameters to be estimated from different genomic segments of the H5N1
HPAIV genome without assuming that the sequences shared an identical phylogenetic history [27].

Here, we extend the landmark study of Lemey *et al.* (2009) to estimate the history of HPAIV dis-
persal into and within Europe after its initial detection in Russia in 2005. Our data comprise 277 H5N1
HPAIV hemagglutinin (HA) and neuraminidase (NA) gene sequences (referred to as isolates) collected
between 2005 and 2010 in Europe and Russia. We adopt a discrete-trait phylodynamic model to esti-
mate both the history of viral migration between geographic areas, and also to infer the movement of
the virus among host species (wild versus domesticated birds). We also estimate temporal changes in
viral population size, and discuss the potential of phylodynamic methods to improve AI surveillance.
Our study provides quantitative estimates of the mechanisms that led to the rapid spread of HPAIV
throughout Europe in 2005–2010 and further illustrates the potential of this approach to improve the
prevention and control of novel emerging AIVs.

2. Results and Discussion

Our analyses of the viral population dynamics for both HA and NA genes reveal a strong increase in
genetic diversity (and, therefore, effective population size) in mid-2005 followed by a second increase
in early 2006 (Figure 1B and C). This coincides with the highest frequency of reported cases of H5N1
infections during the epidemic, when several independent introductions of H5N1 viruses belonging to
sub-clade 2.2 have been confirmed (Figure 1A)[15, 16, 30, 31]. The relatively high viral genetic diver-
sity during the early stage of the epidemic is consistent with the introduction of the virus into a novel
region, with a naïve population, where it was exposed to selection pressure imposed by environmental
and demographic factors [10, 23, 31]. We also inferred a third (and relatively minor) change in viral
genetic diversity during mid-2007, which coincides with the entry of the virus into new EU countries.
From 2005 to 2007 conditions seem to have been relatively favorable for the virus to perpetuate in the
susceptible population and, given the genetic diversity, reassortment may have occurred during this
time frame. The viral genetic diversity is then inferred to decline toward 2007 and to stabilize until

2010. The inferred decrease of viral genetic diversity coincides with the decline of the epidemic, sug-
gesting that prevailing epidemiological factors, such as host density or weather conditions, in 2007–
2010 were not sufficient to favor the establishment of the newly emerging H5N1 strain in the popula-
tion. In contrast to typical AIV epidemics (which exhibit seasonal variation in viral genetic diversity)
the genetic diversity of the H5N1 virus did not exhibit seasonal variation in genetic diversity during
the 2005–2010 period of the epidemic in Europe and Russia, which suggests that transmission of the
virus was influenced by factors different from those typically observed in AIV epidemics.

121
122

Figure 1. (A) Temporal distribution of H5N1 HPAI outbreaks (per year) in domesticated
poultry and wild birds in Europe and Russia from between 2005 and 2010. **(B)** and **(C)**
Bayesian Skyline plots (BSP) illustrating temporal changes in the relative genetic diversity
of H5N1 HPAIV isolates from outbreaks in Europe and the Russia between May, 2005 and
June, 2010 estimated from the HA and NA gene sequences. Line plots summarize esti-
mates of the effective population size (N_eT), a measure of genetic diversity, for HA (above)
and NA (below) gene regions; the shaded regions correspond to the 95% HPD.

Our analyses indicate that domestic birds were the most probable ancestral host type for both genes.
Domestic birds are inferred to be the most probable host type along many branches of the HA and NA
phylogenies, suggesting that the evolution of the H5N1 HPAIV virus largely occurred in association
with domestic-bird populations. By contrast, the large proportion of wild-bird isolates (Figure 2) and
the inferred high transition rates between host types (Table 1) suggest that wild birds also played an
important role in viral dispersal between host populations and geographic areas. This finding is con-
sistent with a number of alternative hypotheses, including scenarios where (1) domestic birds were the
source of the spread of H5N1 HPAIV into Europe, or (2) where the HPAI strains were first transmitted
from non-EU wild birds to non-EU domestic birds (with little or no genetic change), and later transmit-
ted from non-EU domestic birds to EU domestic birds (Figure 2). The first scenario may be explained
by the illegal introduction of domestic birds into Europe, which has been implicated in the spread of
other diseases within some European countries [34]. The second scenario assumes very low selective
pressure on the virus within—and rapid transmission of the virus among—wild-bird populations,
which would explain the high mortality in wild-bird populations and the reported prevalence of the
virus along migratory routes [33]. These two seemingly contradictory scenarios may be reconciled as
follows: H5N1 HPAIV was initially transmitted from non-EU wild birds to domestic poultry as a less
virulent and non-reassortant strain of the virus. Subsequently, the virulent strain of H5N1 HPAIV that
caused the EU epidemic emerged as a reassortant strain from poultry, as suggested elsewhere [32]. The
high density of poultry populations may have resulted in elevated contact and transmission rates, ulti-
mately enhancing opportunities for viral evolution in poultry compared to wild birds. Additionally, the
three sequences isolated from non-avian species were obtained from hosts that prey on birds (two from
cats and one from a marten), which is both consistent with a wild-bird origin of infection (Table 1) and
with the results of previous studies [11, 35].
Our divergence-time estimates (Figure 2) suggest that H5N1 HPAIV originated and experienced re-
assortment among hosts between 1995–1996 (Figure S1). These results are consistent with the frequent
reports of highly reassorted H5N1 HPAIVs introductions into eastern and southeastern Asia since the
emergence of the ancestral strain in 1996 (A/goose/ Guangdong /1/96) [5, 36].

Table 1. Bayes factor (BF) tests for non-zero transition rates between host types for each
gene region (HA, NA). BF values > 6 indicate significant rates of exchange between host
types.

Gene	BF	From	To
HA	11049.7	Wild	Other
HA	11049.7	Other	Wild
HA	171.4	Domestic	Wild
NA	11049.7	Wild	Other
NA	11049.7	Other	Wild
'NA	183.0	Domestic	Wild

Figure 2. Maximum clade credibility (MCC) trees for H5N1 HPAIV hemagglutinin (HA) and neauraminidase (NA) gene regions (A and B, respectively). Branch lengths are rendered proportional to absolute time (see timescales), the branches are colored according to the most probable host type (wild birds, domestic birds, or other), and the branch thickness is proportional to the posterior probability of the inferred host state. Branches where the host state is uncertain (where the posterior probability of any hosts <0.5) are colored black. The posterior probabilities for the ancestral host states are shown in the upper left panel for each tree.

Results of our discrete phylodynamic analyses indicate that Russia, Romania, and Germany were
important epicenters of the H5N1 HPAIV epidemic in Europe. Specifically, our results suggest that the
virus first originated and accumulated in Russia, with subsequent significant dispersal rates out of Rus-
sia into Romania (Figures 3–5, S2). Later, significant dispersal rates occurred between Romania and
Germany. In fact, exchange between Russia-Romania and Romania-Germany represent the most sig-
nificant dispersal rates of the virus during the course of the epidemic (Figure 5, Table S3). Finally, in
late 2005 and early 2006 the virus spread independently from Romania and Germany, respectively into
other areas of the EU (Figure S3); a scenario that is consistent with the results of previous studies [1,
14-16, 30].

Figure 3. Maximum clade credibility (MCC) tree for H5N1 HPAIV hemagglutinin (HA) and neauraminidase (NA) gene regions. Branch lengths are rendered proportional to absolute time (see timescales), the branches are colored according to the most probable geographic location, and the branch thickness is proportional to the posterior probability of the inferred area (see inset legends). Color gradients along branches reflect the inferred changes in the posterior probability of the corresponding areas.

Figure 4. Posterior probabilities of ancestral areas of H5N1 HPAIV hemagglutinin (HA; left) and neauraminidase (NA; right) gene regions collected in Europe and the Russia.

Figure 5. Bayes factor (BF) test for significant non-zero dispersal rates in H5N1 HPAIV. Only rates supported by a BF greater than six are indicated. The color gradient of lines correspond to the probability of the inferred dispersal routes; blue lines and red lines indicate relatively weak and strong support, respectively. The maps are based on satellite pictures available in Google Earth (http://earth.google.com). A: HA gene. B: NA gene. The maps are based on satellite images sourced from the NASA World Wind Java SDK (http://worldwind.arc.nasa.gov/java/).

The inferred geographic expansion of the virus is consistent with our estimates of the viral effective population size; the geographic expanse and the genetic diversity of the virus increased between 2005 and 2006, and subsequently declined between 2007 and 2010 (Figure 1 and S3). The geographic expansion of the virus entailed two major waves. The first wave began in early 2005 from Russia, intensified in Romania, and resulted in the spread of the virus into Germany, Ukraine, and Turkey. The se-

cond wave began in early 2006, also commencing in Russia, intensified in both Romania and Germa-
ny, and resulted in the spread of the virus into Switzerland, Czech Republic, Denmark, and Sweden
[14, 15, 17, 30, 37]. Additionally, a minor wave in 2007 emerged and intensified within Germany, and
resulted in the spread of the virus into France, Hungary, and Poland [15].

The results of our phylodynamic analyses generally appear to be robust (Table 2). The significance
(p-value <0.001) of the observed association-index (AI) values and their 95% credible intervals for
both genes indicate a strong relationship between the inferred phylogeny of the H5N1 virus and the
discrete traits (geographic area and host type), and also indicate that both host type and geography
played important roles in the transmission of the virus. By contrast, the Kullback-Leibler (KL) values
for the discrete host-type models were fairly small for both genes (suggesting relatively poor model
fit), whereas the KL values for the geographic models were large, indicating good fit the between
model and the geographic data [27].

Table 2. Kullback-Leibler divergence and Association Index statistics assessing the fit of
the viral data to the discrete phylodynamic models.

Tree	Kullback-Leibler	Association Index
Host Type		
HA	0.71	6.01 (4.97,7.06)*
NA	0.72	7.22 (5.97,8.44)*
Location (by Countries)		
HA	3.24	9.1(7.87,10.3)*
NA	1.68	10.45(9.29,11.58)*

217   * Statistically significant (*p-value* < 0.001).

Although generally robust, the results of this study are based on analyses that necessarily entailed
several compromises, including: (1) imprecise geographic information on the isolation of H5N1 se-
quences; and; (2) incomplete and possibly biased sampling of H5N1 isolates. First, precise lati-
tude/longitude coordinates were unavailable for 65.8% of the sequences used in this study; instead
only the country of isolation was recorded. In these cases, we used the centroid of the country as a
proxy for the location, which is apt to depart substantially from the true location in most cases, which
may bias our inferences of the phylogeographic history of the H5N1 virus. This sampling issue pre-
cludes our use of the continuous phylodynamic model described by Lemey et al [38], as these models
require more precise information on the locations of the viral samples. Unfortunately, this information
in typically not recorded for publically available viral sequence data in GenBank or disease databases.
The greatest obstacle for applying continuous phylodynamic models to the H5N1 system is therefore
mainly an issue of data confidentiality. The impact of incomplete spatial metadata on the performance
of phylogeographic models has been discussed elsewhere [39, 40]. Inferences under the phylodynamic
models—like all statistical inference methods—assume that we have either a complete or random
sample of sequence data. In the present case, this requires that the H5N1 sequences were collected
randomly with respect to time (between ~1990 and 2010), geographic location, and host type. Myriad
practical considerations almost certainly result in very strongly biased samples, and the impact of these
departures from random sampling on the estimates are difficult to quantify. Although certainly incom-
plete, and almost certainly non-random, our study is based on all available sequence data for the HA

and NA gene regions associated with the H5N1 epidemic in the EU during 2005–2010, and therefore
reflects our best understanding based of the available data.

Phylodynamic analysis has not yet been widely embraced as a resource by public health agencies to
support the design of disease-surveillance strategies. For example, most of the published literature on
avian influenza—including disease summary reports and peer-reviewed scientific studies—upon which
global disease agencies base their decision-making process primarily include numerical summaries of
reported cases in each country, risk maps based purely on spatial and temporal distribution of the dis-
ease, and conventional phylogenetic analysis of sequenced isolates from these outbreaks [13–18, 22–
24, 42]. The 2009 AI global pandemic demonstrates the ability of phylodynamic analyses to provide
novel insights for decisions regarding animal and public health [42]. Our phylodynamic analyses of
an H5N1 HPAIV sequence dataset and associated metadata provides insights on the origin of the out-
break in Europe (*e.g.*, the ancestral location or host type) and the temporal and the spatial progression
of this epidemic. These inferences could be used to inform prevention and control measures to block
the virus at the source (*e.g.*, high-risk geographical areas or bird species), which can limit the spread of
the virus both to the domestic poultry population or to uninfected geographic areas. Furthermore, the
ability to infer and predict routes of viral transmission has clear implications for informing the selec-
tion of appropriate strains for vaccine production to more effectively control new reassortant strains of
AIV in future epidemics. Accordingly, incorporating phylodynamic analyses as a standard tool for the
molecular surveillance of AI might support the development of more effective (and cost effective) pol-
icy decisions for the control of this virus in high-risk regions.

3. Conclusions

The purpose of this study was to illustrate the potential of a Bayesian phylodynamic approach to in-
form AI surveillance and control programs and associated decision-making processes. Our results indi-
cate that the peak in viral genetic diversity between 2005–2006 coincided with its geographic spread
from Russia into Romania and Germany, and throughout the rest of the EU. This expansion was fol-
lowed by a drastic decline in viral genetic diversity, which corresponds with the epidemic decline in
most EU member states. Our findings corroborate previous conclusions that wild birds were the most
important vectors for the spread of the disease. However, our analyses indicate that domestic (rather
than wild) birds were the source of the epidemic, and that viral reassortment most likely occurred in
domestic species. Finally, our results provide further demonstration of the potential of the Bayesian
phylodynamic approach to effectively study the dynamics of disease origin and spread, which promis-
es to maximize the utility of molecular sequence data for the surveillance and control of emerging dis-
eases. These results will contribute to the design of surveillance and control strategies for H5N1
HPAIV in Europe and provide a methodological framework for molecular epidemiological modeling
of emerging novel HPAIVs at regional, national, and international scales. This work can also assist
global and European disease agencies in their effort to improve the efficiency of their surveillance sys-
tems by sharing and improving their genomic databases in the course of emergence of new AIVs.

4. Material and Methods

4.1 Sequence data

Our dataset comprised complete (or nearly complete) H5N1 HPAIV haemagglutinin (HA) and neuraminidase (NA) gene sequences isolated in Europe and the Russia between May, 2005 and June, 2010, including information on the date, location, and host from which the sequences were isolated. These data were obtained from the Global Initiative on Sharing All influenza data (GISAID) public database (http://platform.gisaid.org), with the exception of sequences from Romania, which have been described elsewhere [46]. Laboratory recombinants or highly cultured sequences were excluded from the dataset. The length of the HA and NA gene regions ranged from 1621–1803 and 1211–1458 bp, respectively. Our dataset comprised a total of 347 sequences, of which 277 (79.8%) were from Europe and Russia, and 70 (20.2%) were from Asian and African countries that were included to root the phylogenetic tree (Table S1). Furthermore, we cross-matched each retrieved sequence with the NCBI Influenza Virus Resource (IVR) (http://www.ncbi.nlm.nih.gov/genomes/FLU/FLU.html) database to obtain information on the date, location, and host of origin of the sample. Because both IVR and GISAID databases do not report the exact longitude and latitude (and occasionally the actual date of collection), we consulted the metadata field for each sequence isolate to approximate the location and date of collection, as well as to specify the host species. We identified metadata for each sequence by comparing strain names between databases (*e.g.*, A/swan/Germany/6/2007), and, when available, we cross-matched information with the corresponding publication [11, 47-52]. We specified the location for each sample (latitude-longitude) using the centroid of the country from which samples were isolated. We converted the collection date for each sequence into fractional years (decimal days) in order to estimate divergence times. If only the year of isolation was available (1.2% of the sequences), we specified the ages as the mid-point of the corresponding year. Similarly, if only the month of isolation was available (10.4% of the sequences), we recorded the date as the mid-point of the corresponding month. Finally, we classified host species as either *wild birds*, *domestic birds*, or *other* (non-avian hosts). Table S2 summarizes the profile of 277 H5N1 HPAIV sequences isolated from reported outbreaks in Europe and the Russia between May 2005, and June 2010. Finally, an epidemic curve was plotted using approximately 1130 outbreak reports retrieved from the Food and Agriculture Organization of the United Nation (FAO) Global Animal Disease Information System EMPRES-i (http://empres-i.fao.org/). The outbreak data included dates and locations of detected cases in domesticated poultry and wild birds in Europe and Russia between 2005 and 2010.

4.2 Preliminary phylogenetic analysis

We aligned the H5N1 HPAIV HA and NA gene regions using MUSCLE version 3.8 [43]. The resulting alignments were adjusted manually to ensure that these protein-coding gene regions remained in frame (assessed by amino-acid translation using Mesquite version 3.01) [44]. We removed all sequences (25%) with 100% nucleotide identity and both gene regions were found to be free of homologous recombination using Recombination Detection Program version 3 (RDP3) [45]. For each gene region, we selected a mixed substitution model (assignment of data partitions to substitution models) by first defining six data subsets (the codon positions of each gene region), and then selecting among

mixed-substitution model using the Bayesian Information Criterion (BIC) implemented in Partition-
Finder v 1.1 [46]. Finally, we assessed the degree of topological (in)congruence between the two gene
regions by performing maximum-likelihood estimates of the phylogeny for the individual gene regions
under the selected mixed-substitution model; these analyses entailed 100 non-parametric bootstrap
replicate searches of each gene region using RAxML version 8 [47].

4.3 Divergence-time estimation

We estimated divergence times for the H5N1 HPAIV sequence dataset using the (relaxed) molecu-
lar-clock models implemented in BEAST v 1.8 [48]. We used the mixed-substitution model for each
gene region identified in the preliminary analyses detailed above. We assessed the fit of the sequence
data to three branch-rate prior models: (1) a strict molecular clock model (which assumes that substitu-
tion rates are stochastically constant across branches of the tree); (2) the uncorrelated exponential re-
laxed clock (UCED) model, which assumes that substitution rates on adjacent branches are sampled
from a shared exponential distribution, and; (3) the uncorrelated lognormal relaxed clock (UCLN)
model, which assumes that substitution rates on adjacent branches are drawn from a shared lognormal
distribution. We assessed the fit of the sequence data to these three branch-rate models by comparing
their corresponding marginal likelihoods, which were estimated using Tracer version 1.6 [48, 49].
Bayes factor (BF) comparisons indicated that the UCED branch-rate model provided the best fit for
both HA and NA gene regions (Bayes factor >25 for the log marginal likelihood). Estimation of diver-
gence times also requires a node-age model; we assumed a Bayesian skyline coalescent tree prior,
which allowed us to estimate changes in the effective population size through time [50]. Isolation dates
of the sequences were used to calibrate divergence-time estimates.
We approximated the joint posterior probability density of these model parameters using the Mar-
kov Chain Monte Carlo (MCMC) algorithms implemented in BEAST v 1.8. Each MCMC simulation
was run for 1×10^8 cycles, and thinned by sampling every 10,000 cycle. We performed four replicate
MCMC simulations to aid in assessing performance of the simulations. We assessed MCMC conver-
gence by comparing the parameter estimates from independent analyses for each parameter, ensuring
that the estimates of the marginal posterior probability densities were effectively identical for the four
independent chains. We also assessed convergence by the estimating effective sample sizes (ESS) for
each parameter using Tracer, and assessed mixing based on the acceptance ratios. Those evaluations
suggested that the MCMC algorithms provided a reliable approximation of the posterior probability
density, and suggested that the first 10% of the samples (the "burn-in") should be discarded. We sum-
marized posterior results as a maximum clade credibility (MCC) tree using TreeAnnotator. We then
generated a Bayesian skyride plot to infer the population demographics of H5N1 HPAIV HA and NA
gene segments in Europe and the Russia. We plotted the inferred effective population size of the virus
between 2005–2010 in terms of relative genetic diversity (N_eT), where N_e is the effective population
size and T the generation time [51].

4.4 Estimation of geographic history under the discrete phylodynamic model

We incorporated information on the geographic locations where sequences were isolated to describe
the spatial dynamics of viral epidemics following a procedure described by Lemey *et al.* [27]. Briefly,

the approach jointly estimates phylogeny and history of discrete traits in a Bayesian statistical frame-
work, which enables inference of the timing of viral dispersal patterns while accommodating phyloge-
netic uncertainty. Here, the discrete states are geographic areas, and the goal is to estimate the history
of viral migration (transitions) between these areas through time. This is achieved using a model aver-
aging approach (based on Bayesian stochastic search variable selection; BSSVS) to describe the spatial
evolution of H5N1 HPAIV epidemic. We modeled the geographic transition of the H5N1 HPAIV be-
tween areas as discrete states under a continuous-time Markov model comprising a number of non-
zero transition rates that were identified by means of BSSVS. We assessed the fit of the H5N1 HPAIV
data to a number of candidate discrete phylogeographic models (Table S4), including both symmetric
and asymmetric discrete-trait models with irreversible and reversible transitions, respectively (using a
mean-one exponential prior for the rate parameters). Accordingly, our analyses were based on a com-
posite phylogenetic model that comprised: (1) the previously selected mixed-substitution model; (2)
the strict, UCLN, UCED branch-rate models to describe variation in substitution rate across branches;
(3) the coalescent Gaussian Markov Random field (GMRF) Bayesian Skyride model as a prior on the
node times in the tree, and; (4) the symmetric and asymmetric discrete-state phylodynamic models to
describe the history of viral migration between discrete geographic areas. Accordingly, we explored a
total of six candidate composite phylodynamic models (the symmetric and asymmetric variants of the
geographic model and the three branch-rate models).

We assessed the relative fit of the six candidate phylodynamic models to the HA and NA sequence
datasets using Bayes factors. To this end, we first estimated the marginal likelihood for each of the six
candidate phylodynamic models from the resulting posterior samples using the posterior simulation-
based analogue of Akaike's information criterion (AICm) [52]. Bael et al. [53] have recently shown
the AICm to provide more reliable estimates of the marginal likelihood than the harmonic-mean esti-
mator. We then used the marginal-likelihood estimates to compute Bayes factors to select among the
candidate models. The UCED branch-rate model was again preferred for both HA and NA gene seg-
ments (BF > 25). Bayes factor (BF) comparisons indicated that the symmetric UCED branch-rate
model with irreversible transitions provided the best fit for both HA and NA gene regions (Bayes fac-
tor >25 for the log marginal likelihood). The posterior probability distribution of trees sampled under
the preferred model was summarized as an MCC consensus tree with posterior probabilities of geo-
graphic areas plotted at interior nodes using FigTree version 1.4 [54]. We used SPREAD version 1.0.6
[55] to assess the strength of transition rates between discrete geographic areas, using a BF cutoff = 6
to identify non-zero transition rates. Finally, we generated a keyhole markup language (KML) file to
visualize the geographic migration of the virus.

4.5 Exploring the evolution of H5N1 HPAIV host infection

We modeled the evolutionary movement of H5N1 HPAIV within and between host types (domestic
birds, wild birds, other). To this end, we again adopted a discrete-trait model, where the discrete states
in this case correspond to the host type, where the objective is to infer the history of H5N1 HPAIV
migration between hosts through time. The number of non-zero transition rates in the model was again
estimated using BSSVS. The relative strength of transition rates (*e.g.*, wild → domestic) was estimated
using Bayes factors. Following [64], we estimated the ancestral states (host type) at internal nodes of

the tree under a composite phylogenetic model that included: (1) the previously selected mixed-substitution model to describe evolution of nucleotides over the tree with branch lengths; (2) the preferred uncorrelated lognormal model to describe the variation of substitution rates across branches of the tree [56]; (3) a constant population-size coalescent model to describe the temporal distribution of node heights in the phylogeny, and; (4) the asymmetric discrete-state phylodynamic model to describe the history of viral migration between host types as a continuous-time Markov process (with a number of non-zero transition rates determined by the BSSVS procedure, using a uniform prior with mean = 1 for the rate parameters). We summarized the posterior probability distribution of parameter estimates as an MCC consensus tree with the posterior probability of host state mapped to interior nodes of the tree using FigTree. Additionally, we assessed the strength of transition rates between states (hosts) by means of Bayes factors using SPREAD with a cutoff of six to identify non-zero transition rates. Use of the asymmetric discrete trait model allowed us to assess the strength of directionality between states (*e.g.*, wild → domestic and domestic → wild bird).

4.6 Assessing uncertainty in discrete-trait mappings and association statistics

We used the Kullback-Leibler (KL) divergence statistic [57] to accommodate phylogenetic uncertainty in the discrete-trait estimates (for host type and geographic location). The KL divergence measures the departure between prior probability distribution and the corresponding posterior probability distribution for a given parameter. The premise is simple: the posterior probability distribution inferred for a given parameter is the updated version of prior probability distribution—it is updated by the information in the data via the likelihood function. Accordingly, if the data contain little information regarding the value of a parameter, its posterior probability distribution will closely resemble the corresponding prior probability distribution, and the KL divergence between these two probability distributions will therefore be small [27]. We calculated the KL divergence for each selected posterior distribution of trees using a function written by Razavi [58] implemented in Matlab v 2013a [59]. The Association Index (AI) was calculated using Bayesian Tip-Significance Testing (BaTS) software version 1.0 to test the hypothesis that a taxon with a given trait (host type or geographic location) are more likely to share traits with adjoining taxa than that expected by chance [60].

Acknowledgments

We are grateful to Matthew L. Scotch for assistance with Matlab. This research was supported by National Science Foundation (NSF) grants DEB-0842181, DEB-0919529, DBI-1356737, and DEB-1457835 awarded to B.R.M..

Author Contributions

M.A. and B.R.M. conceived and designed the study. M.A. and A.P. analyzed the data. B.R.M. provided advise regarding the analyses. M.A., B.R.M. and A.P. wrote the paper.

Conflicts of Interest

The authors declare no conflict of interest.

References and Notes

1. Kilpatrick, A. M.; Chmura, A. A.; Gibbons, D. W.; Fleischer, R. C.; Marra, P. P.; Daszak, P., Predicting the global spread of H5N1 avian influenza. *Proc Natl Acad Sci U S A* **2006,** 103, (51), 19368-73.

2. Brown, I. H., Summary of avian influenza activity in Europe, Asia, and Africa, 2006-2009. *Avian diseases* **2010,** 54, (1 Suppl), 187-93.

3. Sims, L. D.; Domenech, J.; Benigno, C.; Kahn, S.; Kamata, A.; Lubroth, J.; Martin, V.; Roeder, P., Origin and evolution of highly pathogenic H5N1 avian influenza in Asia. *The Veterinary record* **2005,** 157, (6), 159-64.

4. Bonn, D., Wild birds, poultry, and avian influenza. *Lancet Infect Dis* **2006,** 6, (5), 262.

5. Xu, X.; Subbarao; Cox, N. J.; Guo, Y., Genetic characterization of the pathogenic influenza A/Goose/Guangdong/1/96 (H5N1) virus: similarity of its hemagglutinin gene to those of H5N1 viruses from the 1997 outbreaks in Hong Kong. *Virology* **1999,** 261, (1), 15-9.

6. Mukhtar, M. M.; Rasool, S. T.; Song, D.; Zhu, C.; Hao, Q.; Zhu, Y.; Wu, J., Origin of highly pathogenic H5N1 avian influenza virus in China and genetic characterization of donor and recipient viruses. *The Journal of general virology* **2007,** 88, (Pt 11), 3094-9.

7. WHO/OIE/FAO H5N1 Evolution Working Group. Toward a unified nomenclature system for highly pathogenic avian influenza virus (H5N1). *Emerging infectious diseases* **2008,** 14, (7), e1.

8. Chen, H.; Li, Y.; Li, Z.; Shi, J.; Shinya, K.; Deng, G.; Qi, Q.; Tian, G.; Fan, S.; Zhao, H.; Sun, Y.; Kawaoka, Y., Properties and dissemination of H5N1 viruses isolated during an influenza outbreak in migratory waterfowl in western China. *J Virol* **2006,** 80, (12), 5976-83.

9. European Commission (EC), A report on surveys for avian influenza in poultry in member states during 2005, available at: http://ec.europa.eu/food/animal/diseases/controlmeasures/avian/eu_resp_surveillance_en.htm. (Accessed on 1st January 2014)

10. Globig, A.; Staubach, C.; Beer, M.; Koppen, U.; Fiedler, W.; Nieburg, M.; Wilking, H.; Starick, E.; Teifke, J. P.; Werner, O.; Unger, F.; Grund, C.; Wolf, C.; Roost, H.; Feldhusen, F.; Conraths, F. J.; Mettenleiter, T. C.; Harder, T. C., Epidemiological and ornithological aspects of outbreaks of highly pathogenic avian influenza virus H5N1 of Asian lineage in wild birds in Germany, 2006 and 2007. *Transbound Emerg Dis* **2009,** 56, (3), 57-72.

11. Weber, S.; Harder, T.; Starick, E.; Beer, M.; Werner, O.; Hoffmann, B.; Mettenleiter, T. C.; Mundt, E., Molecular analysis of highly pathogenic avian influenza virus of subtype H5N1 isolated from wild birds and mammals in northern Germany. *J Gen Virol* **2007,** 88, (Pt 2), 554-8.

12. Starick, E.; Beer, M.; Hoffmann, B.; Staubach, C.; Werner, O.; Globig, A.; Strebelow, G.; Grund, C.; Durban, M.; Conraths, F. J.; Mettenleiter, T.; Harder, T., Phylogenetic analyses of highly pathogenic avian influenza virus isolates from Germany in 2006 and 2007 suggest at least three separate introductions of H5N1 virus. *Vet Microbiol* **2008,** 128, (3-4), 243-52.

13. Breed, A. C.; Harris, K.; Hesterberg, U.; Gould, G.; Londt, B. Z.; Brown, I. H.; Cook, A. J., Surveillance for avian influenza in wild birds in the European Union in 2007. *Avian Dis* **2010,** 54, (1 Suppl), 399-404.

14. Oprisan, G.; Coste, H.; Lupulescu, E.; Oprisoreanu, A. M.; Szmal, C.; Onita, I.; Popovici, N.; Ionescu, L. E.; Bicheru, S.; Enache, N.; Ceianu, C.; Czobor, F.; Olaru, E.; Alexandrescu, V.; Radu, D. L.; Onu, A.; Popa, M. I., Molecular analysis of the first avian influenza H5N1 isolates from fowl in Romania. *Roum Arch Microbiol Immunol* **2006,** 65, (3-4), 79-82.

15. Hofmann, M. A.; Renzullo, S.; Baumer, A., Phylogenetic characterization of H5N1 highly pathogenic avian influenza viruses isolated in Switzerland in 2006. *Virus Genes* **2008,** 37, (3), 407-13.

16. Kiss, I.; Gyarmati, P.; Zohari, S.; Ramsay, K. W.; Metreveli, G.; Weiss, E.; Brytting, M.; Stivers, M.; Lindstrom, S.; Lundkvist, A.; Nemirov, K.; Thoren, P.; Berg, M.; Czifra, G.; Belak, S., Molecular characterization of highly pathogenic H5N1 avian influenza viruses isolated in Sweden in 2006. *Virol J* **2008,** 5, 113.

17. Zohari, S.; Gyarmati, P.; Thoren, P.; Czifra, G.; Brojer, C.; Belak, S.; Berg, M., Genetic characterization of the NS gene indicates co-circulation of two sub-lineages of highly pathogenic avian influenza virus of H5N1 subtype in Northern Europe in 2006. *Virus Genes* **2008,** 36, (1), 117-25.

18. Nagy, A.; Vostinakova, V.; Pindova, Z.; Hornickova, J.; Cernikova, L.; Sedlak, K.; Mojzis, M.; Dirbakova, Z.; Machova, J., Molecular and phylogenetic analysis of the H5N1 avian influenza virus caused the first highly pathogenic avian influenza outbreak in poultry in the Czech Republic in 2007. *Veterinary microbiology* **2009,** 133, (3), 257-63.

19. Ronquist, F., Bayesian inference of character evolution. *Trends Ecol Evol* **2004,** 19, (9), 475-81.

20. Jewell, C. P.; Kypraios, T.; Christley, R. M.; Roberts, G. O., A novel approach to real-time risk prediction for emerging infectious diseases: a case study in Avian Influenza H5N1. *Prev Vet Med* **2009,** 91, (1), 19-28.

21. Martinez, M.; Munoz, M. J.; De La Torre, A.; Iglesias, I.; Peris, S.; Infante, O.; Sanchez-Vizcaino, J. M., Risk of introduction of H5N1 HPAI from Europe to Spain by wild water birds in autumn. *Transbound Emerg Dis* **2009,** 56, (3), 86-98.

22. Fink, M.; Fernandez, S. R.; Schobesberger, H.; Koefer, J., Geographical spread of highly pathogenic avian influenza virus H5N1 during the 2006 outbreak in Austria. *J Virol* **2010,** 84, (11), 5815-23.

23. Reperant, L. A.; Fuckar, N. S.; Osterhaus, A. D.; Dobson, A. P.; Kuiken, T., Spatial and temporal association of outbreaks of H5N1 influenza virus infection in wild birds with the 0 degrees C isotherm. *PLoS Pathog* **2010,** 6, (4), e1000854.

24. Alkhamis, M.; Willeberg, P.; Carlsson, U.; Carpenter, T.; Perez, A., Alternative scan-based approaches to identify space-time clusters of highly pathogenic avian influenza virus H5N1 in wild birds in Denmark and Sweden in 2006. *Avian Dis* **2012,** 56, (4 Suppl), 1040-8.

25. Grenfell, B. T.; Pybus, O. G.; Gog, J. R.; Wood, J. L.; Daly, J. M.; Mumford, J. A.; Holmes, E. C., Unifying the epidemiological and evolutionary dynamics of pathogens. *Science* **2004,** 303, (5656), 327-32.

26. Minin, V. N.; Suchard, M. A., Fast, accurate and simulation-free stochastic mapping. *Philos Trans R Soc Lond B Biol Sci* **2008,** 363, (1512), 3985-95.

27. Lemey, P.; Rambaut, A.; Drummond, A. J.; Suchard, M. A., Bayesian phylogeography finds its roots. *PLoS Comput Biol* **2009,** 5, (9), e1000520.

28. Wallace, R. G.; Fitch, W. M., Influenza A H5N1 immigration is filtered out at some international borders. *PLoS One* **2008,** 3, (2), e1697.

29. Wallace, R. G.; Hodac, H.; Lathrop, R. H.; Fitch, W. M., A statistical phylogeography of influenza A H5N1. *Proc Natl Acad Sci U S A* **2007,** 104, (11), 4473-8.

30. Bragstad, K.; Jorgensen, P. H.; Handberg, K.; Hammer, A. S.; Kabell, S.; Fomsgaard, A., First introduction of highly pathogenic H5N1 avian influenza A viruses in wild and domestic birds in Denmark, Northern Europe. *Virol J* **2007,** 4, 43.

31. Rao, D. M.; Chernyakhovsky, A.; Rao, V., Modeling and analysis of global epidemiology of avian influenza. *Environmental Modelling & Software* **2009,** 24, (1), 124-134.

32. Vijaykrishna, D.; Bahl, J.; Riley, S.; Duan, L.; Zhang, J. X.; Chen, H.; Peiris, J. S.; Smith, G. J.; Guan, Y., Evolutionary dynamics and emergence of panzootic H5N1 influenza viruses. *PLoS Pathogens* **2008,** 4, (9), e1000161.

33. Olsen, B.; Munster, V. J.; Wallensten, A.; Waldenstrom, J.; Osterhaus, A. D.; Fouchier, R. A., Global patterns of influenza a virus in wild birds. *Science* **2006,** 312, (5772), 384-8.

34. Ward, M. P.; Maftei, D.; Apostu, C.; Suru, A., Geostatistical visualisation and spatial statistics for evaluation of the dispersion of epidemic highly pathogenic avian influenza subtype H5N1. *Vet Res* **2008,** 39, (3), 22.

35. Mushtaq, M. H.; Juan, H.; Jiang, P.; Li, Y.; Li, T.; Du, Y.; Mukhtar, M. M., Complete genome analysis of a highly pathogenic H5N1 influenza A virus isolated from a tiger in China. *Archives of virology* **2008,** 153, (8), 1569-74.

36. Ellis, T. M.; Bousfield, R. B.; Bissett, L. A.; Dyrting, K. C.; Luk, G. S.; Tsim, S. T.; Sturm-Ramirez, K.; Webster, R. G.; Guan, Y.; Malik Peiris, J. S., Investigation of outbreaks of highly pathogenic H5N1 avian influenza in waterfowl and wild birds in Hong Kong in late 2002. *Avian pathology : journal of the W.V.P.A* **2004,** 33, (5), 492-505.

37. Nagy, A.; Machova, J.; Hornickova, J.; Tomci, M.; Nagl, I.; Horyna, B.; Holko, I., Highly pathogenic avian influenza virus subtype H5N1 in Mute swans in the Czech Republic. *Vet Microbiol* **2007,** 120, (1-2), 9-16.

38. Lemey, P.; Rambaut, A.; Welch, J. J.; Suchard, M. A., Phylogeography takes a relaxed random walk in continuous space and time. *Molecular biology and evolution* **2010,** 27, (8), 1877-85.

39. Scotch, M.; Sarkar, I. N.; Mei, C.; Leaman, R.; Cheung, K. H.; Ortiz, P.; Singraur, A.; Gonzalez, G., Enhancing phylogeography by improving geographical information from GenBank. *J Biomed Inform* **2011,** 44 (Suppl 1), S44-7.

40. Tahsin, T.; Beard, R.; Rivera, R.; Lauder, R.; Wallstrom, G.; Scotch, M.; Gonzalez, G., Natural language processing methods for enhancing geographic metadata for phylogeography of zoonotic viruses. *AMIA Jt Summits Transl Sci Proc* **2014,** (2014), 102-11.

41. Szeleczky, Z.; Dan, A.; Ursu, K.; Ivanics, E.; Kiss, I.; Erdelyi, K.; Belak, S.; Muller, C. P.; Brown, I. H.; Balint, A., Four different sublineages of highly pathogenic avian influenza H5N1 introduced in Hungary in 2006-2007. *Veterinary microbiology* **2009,** 139, (1-2), 24-33.

42. Pybus, O. G.; Fraser, C.; Rambaut, A., Evolutionary epidemiology: preparing for an age of
genomic plenty. *Philos Trans R Soc Lond B Biol Sci* **2013,** 368, (1614), 20120193.=

43. Edgar, R. C., MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res **2004,** 32, (5), 1792-7.

44. Maddison, W. P.; Maddison, D. R. Mesquite : a modular system for evolutionary analysis.
Version 2.75. http://mesquiteproject.org

45. Martin, D. P.; Lemey, P.; Lott, M.; Moulton, V.; Posada, D.; Lefeuvre, P., RDP3: a flexible and
fast computer program for analyzing recombination. *Bioinformatics* **2010,** 26, (19), 2462-3.

46. Lanfear, R.; Calcott, B.; Ho, S. Y.; Guindon, S., Partitionfinder: combined selection of
partitioning schemes and substitution models for phylogenetic analyses. *Molecular biology and*
evolution **2012,** 29, (6), 1695-701.

47. Stamatakis, A., RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **2014**.

48. Drummond, A. J.; Rambaut, A., BEAST: Bayesian evolutionary analysis by sampling trees.
BMC evolutionary biology **2007,** 7, 214.

49. Suchard, M. A.; Weiss, R. E.; Sinsheimer, J. S., Bayesian selection of continuous-time Markov
chain evolutionary models. *Molecular biology and evolution* **2001,** 18, (6), 1001-13.

50. Drummond, A. J.; Rambaut, A.; Shapiro, B.; Pybus, O. G., Bayesian coalescent inference of
past population dynamics from molecular sequences. *Molecular biology and evolution* **2005,**
22, (5), 1185-92.

51. Minin, V. N.; Bloomquist, E. W.; Suchard, M. A., Smooth skyride through a rough skyline:
Bayesian coalescent-based inference of population dynamics. *Molecular biology and evolution*
2008, 25, (7), 1459-71.

52. Raftery, A.; Newton, M.; Satagopan, J.; Krivitsky, P., Estimating the integrated likelihood via
posterior simulation using the harmonic mean identity. Oxford University Press: 2007.

53. Baele, G.; Lemey, P.; Bedford, T.; Rambaut, A.; Suchard, M. A.; Alekseyenko, A. V.,
Improving the accuracy of demographic and molecular clock model comparison while
accommodating phylogenetic uncertainty. Molecular biology and evolution 2012, 29, (9),
2157-67.

54. Shapiro, B.; Rambaut, A.; Drummond, A. J., Choosing appropriate substitution models for the
phylogenetic analysis of protein-coding sequences. *Molecular biology and evolution* **2006,** 23,
(1), 7-9.

55. Bielejec, F.; Rambaut, A.; Suchard, M. A.; Lemey, P., SPREAD: spatial phylogenetic
reconstruction of evolutionary dynamics. *Bioinformatics* **2011,** 27, (20), 2910-2.

56. Lu, L.; Lycett, S. J.; Leigh Brown, A. J., Reassortment patterns of avian influenza virus internal
segments among different subtypes. *BMC evolutionary biology* **2014,** 14, 16.

57. Kullback, S.; Leibler, R., On information and sufficiency. *Annals of Mathematical Statistics*
1951, 22, 79–86.

58. Razavi, N. Kullback–Leibler Divergence, **2008**.

59. MathWorks *Matlab*, 2013a; **2012**.

60. Parker, J.; Rambaut, A.; Pybus, O. G., Correlating viral phenotypes with phylogeny:
accounting for phylogenetic uncertainty. *Infect Genet Evol* **2008,** 8, (3), 239-46.

Supplementary Materials

Table S1 Bayes factor test for non-zero rates for location (country) state model for both
HA and NA genes. Large BF values indicate most likely routes of dispersal between coun-
try's location states.

Gene	BF	Between	
HA	77823.6	Russia	Romania
HA	77823.6	Germany	Austria
HA	77823.6	Germany	Czech Republic
HA	8639.4	Turkey	Romania
HA	317.0	Romania	Germany
HA	274.4	Poland	Germany
HA	90.6	Romania	Hungary
HA	87.4	Romania	Italy
HA	48.5	Ukraine	Russia
HA	46.4	Switzerland	Germany
HA	43.4	Germany	France
HA	16.0	Romania	Bosnia
HA	13.9	Sweden	Romania
HA	11.9	Romania	Bulgaria
HA	11.1	Romania	Croatia
HA	9.8	Slovenia	Austria
HA	7.7	Sweden	Denmark
HA	7.0	Germany	Denmark
NA	77823.6	Romania	Germany
NA	2984.9	Russia	Romania
NA	2502.1	Turkey	Romania
NA	441.2	Germany	Czech Republic
NA	409.8	Switzerland	Germany
NA	240.0	Sweden	Denmark
NA	190.9	Romania	Hungary
NA	149.2	Poland	Germany
NA	108.2	Germany	France
NA	87.0	Ukraine	Russia
NA	57.9	Germany	Austria
NA	27.5	Romania	Italy
NA	16.3	Romania	Bulgaria
NA	15.4	Slovenia	Austria
NA	15.3	Sweden	Germany
NA	11.1	Romania	Bosnia
NA	7.6	United Kingdom	Germany
NA	6.9	Romania	Croatia

599 Bayes factors (BF) > 6 indicate significant non-zero dispersal
600

601   **Table S2** GISAID database Accession numbers and isolate names matched with other pub-
602   lic databases used in this study for both HA and NA gene segments, with the exception of
603   sequences isolated from Romania.

Epi_Flu #	Isolate_Name
EPI_ISL_7012	A/chicken/Afghanistan/1573_7/2006
EPI_ISL_66178	A/coot/Voecklabruck/1585/2006
EPI_ISL_66194	A/duck/Leibnitz/243/2006
EPI_ISL_66195	A/duck/Mellach/335/2006
EPI_ISL_66196	A/duck/Schaerding/1398/2006
EPI_ISL_66199	A/duck/Wels/2025/2006
EPI_ISL_66200	A/duck/Wien/1836/2006
EPI_ISL_66201	A/egret/Wien/1977/2006
EPI_ISL_66202	A/goose/Wien/1966/2006
EPI_ISL_66219	A/swan/Krems/2354/2006
EPI_ISL_66220	A/swan/Krems/2547/2006
EPI_ISL_66221	A/swan/Krems/2675/2006
EPI_ISL_66222	A/swan/Laakirchen/2703/2006
EPI_ISL_66223	A/swan/Mellach/215/2006
EPI_ISL_66225	A/swan/Perg/1358/2006
EPI_ISL_66226	A/swan/Schaerding/1499/2006
EPI_ISL_66227	A/swan/Schaerding/1806/2006
EPI_ISL_66228	A/swan/Schwechat/2538/2006
EPI_ISL_66229	A/swan/Voecklabruck/1484/2006
EPI_ISL_66230	A/swan/Wien/1410/2006
EPI_ISL_66231	A/swan/Wien/2323/2006
EPI_ISL_74836	A/Anas platyrhynchos/Belgium/09-762-P1/2008
EPI_ISL_64083	A/Cygnus olor/BIH/1/2006
EPI_ISL_105998	A/common buzzard/Bulgaria/38WB/2010
EPI_ISL_1254	A/Goose/Guangdong/1/96
EPI_ISL_1909	A/goose/Guangdong/3/1997
EPI_ISL_10006	A/chicken/Hebei/718/2001
EPI_ISL_15275	A/duck/Guangxi/27/2003
EPI_ISL_4533	A/duck/Guangdong/173/2004
EPI_ISL_64893	A/Mallard/Huadong/hn/2005
EPI_ISL_78004	A/chicken/Xinjiang/27/2006
EPI_ISL_75197	A/great_crested_grebe/Qinghai/1/2009
EPI_ISL_6386	A/cygnus olor/Croatia/1/2005
EPI_ISL_15685	A/Cygnus olor/Czech Republic/6111/06
EPI_ISL_15689	A/Cygnus olor/Czech Republic/10814/06
EPI_ISL_15690	A/turkey/Czech Republic/10309-3/07
EPI_ISL_15691	A/chicken/Czech Republic/11242-38/07
EPI_ISL_15692	A/Cygnus olor/Czech Republic/10732/07
EPI_ISL_15693	A/Cygnus olor/Czech Republic/10662/06
EPI_ISL_69720	A/buzzard/Denmark/6370/06
EPI_ISL_69730	A/tufted duck/Denmark/6431/06
EPI_ISL_69724	A/tufted duck/Denmark/6540/06

EPI_ISL_69725	A/peregrine/Denmark/6632/06
EPI_ISL_69731	A/grey lag goose/Denmark/6692/06
EPI_ISL_69732	A/whooper swan/Denmark/7224/06
EPI_ISL_69722	A/whooper swan/Denmark/7275/06
EPI_ISL_69721	A/great crested grebe/Denmark/7498/06
EPI_ISL_64348	A/duck/Egypt/2253_3/2006
EPI_ISL_120302	A/chicken/Egypt/9403_NAMRU3/2007
EPI_ISL_33865	A/chicken/Egypt/083/2008
EPI_ISL_89195	A/chicken/Egypt/CL6/2007
EPI_ISL_2988	A/mute swan/France/06299/2006
EPI_ISL_31666	A/mute swan/France/070203/2007
EPI_ISL_74112	A/teal/Germany/Wv632/2005
EPI_ISL_10142	A/swan/Germany/R65/2006
EPI_ISL_2985	A/stone marten/Germany/R747/2006
EPI_ISL_10142	A/swan/Germany/R65/2006
EPI_ISL_10333	A/cat/Germany/606/2006
EPI_ISL_68227	A/swan/Bavaria/25/2006
EPI_ISL_68226	A/swan/Bavaria/24/2006
EPI_ISL_68225	A/swan/Bavaria/23/2006
EPI_ISL_68228	A/tufted duck/Bavaria/26/2006
EPI_ISL_11353	A/falcon/Bavaria/15/2006
EPI_ISL_11359	A/swan/Bavaria/21/2006
EPI_ISL_11355	A/swan/Bavaria/17/2006
EPI_ISL_11354	A/swan/Bavaria/16/2006
EPI_ISL_11352	A/swan/Bavaria/14/2006
EPI_ISL_11350	A/mute swan/Bavaria/12/2006
EPI_ISL_11360	A/great crested grebe/Bavaria/22/2006
EPI_ISL_11358	A/goosander/Bavaria/20/2006
EPI_ISL_11356	A/goosander/Bavaria/18/2006
EPI_ISL_11351	A/buzzard/Bavaria/13/2006
EPI_ISL_11348	A/eagle owl/Bavaria/10/2006
EPI_ISL_11357	A/goldeneye duck/Bavaria/19/2006
EPI_ISL_11349	A/common buzzard/Bavaria/11/2006
EPI_ISL_68235	A/swan/Bavaria/33/2006
EPI_ISL_68234	A/swan/Bavaria/32/2006
EPI_ISL_68231	A/swan/Bavaria/29/2006
EPI_ISL_68230	A/swan/Bavaria/28/2006
EPI_ISL_68233	A/great crested grebe/Bavaria/31/2006
EPI_ISL_68236	A/goosander/Bavaria/34/2006
EPI_ISL_68232	A/common pochard/Bavaria/30/2006
EPI_ISL_68229	A/tufted duck/Bavaria/27/2006
EPI_ISL_3011	A/mute_swan/Germany/R1359/07
EPI_ISL_24700	A/mallard/Bavaria/10/2007
EPI_ISL_27795	A/domestic_goose/Germany/R1400/2007
EPI_ISL_27769	A/domestic_duck/Germany/R2048/2007
EPI_ISL_27768	A/domestic_duck/Germany/R1779/2007
EPI_ISL_27797	A/great crested grebe/Germany/R1406/07

EPI_ISL_68254	A/Canada goose/Bavaria/4/2007
EPI_ISL_68255	A/swan/Bavaria/5/2007
EPI_ISL_68252	A/swan/Bavaria/2/2007
EPI_ISL_68253	A/mute swan/Bavaria/3/2007
EPI_ISL_63430	A/black_nested_ grebe/Germany/R1393/2007
EPI_ISL_68260	A/greylag goose /Bavaria/11/2007
EPI_ISL_68259	A/swan/Bavaria/9/2007
EPI_ISL_68258	A/swan/Bavaria/8/2007
EPI_ISL_68257	A/swan/Bavaria/7/2007
EPI_ISL_68256	A/swan/Bavaria/6/2007
EPI_ISL_68261	A/wild duck/Bavaria/12/2007
EPI_ISL_27767	A/domestic duck/Germany/R1772/2007
EPI_ISL_27773	A/chicken/Germany/R3234/2007
EPI_ISL_27771	A/chicken/Germany/R3272/2007
EPI_ISL_27772	A/chicken/Germany/R3294/2007
EPI_ISL_2963	A/whooper swan/Germany/R88/06
EPI_ISL_10140	A/mallard/Bavaria/1/2006
EPI_ISL_25455	A/domestic duck/Germany/R874/2008
EPI_ISL_32413	A/duck/Germany/R854/2008
EPI_ISL_25455	A/domestic duck/Germany/R874/2008
EPI_ISL_3091	A/Duck/Hong Kong/380.5/2001
EPI_ISL_67724	A/chicken/Hong_Kong/409.1/2002
EPI_ISL_25699	A/grey_heron/Hong_Kong/1046/2008
EPI_ISL_31503	A/mute swan/Hungary/3472/2006
EPI_ISL_31502	A/mute swan/Hungary/4571/2006
EPI_ISL_31501	A/goose/Hungary/14756/2006
EPI_ISL_11425	A/goose/Hungary/3413/2007
EPI_ISL_63515	A/goose/Hungary/2823/2/2007
EPI_ISL_65649	A/quail/Yogjakarta/UT1075/2004
EPI_ISL_30650	A/chicken/East_Java/UT6031/2007
EPI_ISL_10369	A/turkey/Israel/345/2006
EPI_ISL_6383	A/mallard/Italy/835/2006
EPI_ISL_21034	A/cygnus olor/Italy/742/2006
EPI_ISL_6891	A/cygnus olor/Italy/808/2006
EPI_ISL_6930	A/turkey/Ivory Coast/4372_2/2006
EPI_ISL_63599	A/falcon/Kuwait/0286/2007
EPI_ISL_6248	A/chicken/Nigeria/957_20/2006
EPI_ISL_6412	A/duck/Niger/914/2006
EPI_ISL_29083	A/chicken/Sihala/NARC3303.4/2006
EPI_ISL_27762	A/chicken/Poland/R3248/2007
EPI_ISL_27763	A/turkey/Poland/R3249/2007
EPI_ISL_20922	A/swan/Poland/1242-139V08/2006
EPI_ISL_20921	A/hawk/Poland/937-138V08/2006
EPI_ISL_20920	A/swan/Poland/467-136V08/2006
EPI_ISL_20919	A/goosander/Poland/502-137V08/2006
EPI_ISL_20918	A/buzzard/Poland/MB266B-141V08/2007
EPI_ISL_20917	A/chicken/Poland/38-140V08/2007

EPI_ISL_20916	A/swan/Poland/305-135V08/2006
EPI_ISL_11388	A/goose/Krasnoozerskoe/627/2005
EPI_ISL_11387	A/goose/Suzdalka/10/2005
EPI_ISL_10106	A/Cygnus olor/Astrakhan/Ast05-2-9/2005
EPI_ISL_10103	A/Cygnus olor/Astrakhan/Ast05-2-1/2005
EPI_ISL_10101	A/Cygnus olor/Astrakhan/Ast05-2-8/2005
EPI_ISL_10051	A/Cygnus olor/Astrakhan/Ast05-2-7/2005
EPI_ISL_10050	A/Cygnus olor/Astrakhan/Ast05-2-4/2005
EPI_ISL_11389	A/turkey/Suzdalka/12/05
EPI_ISL_9768	A/grebe/Novosibirsk/29/2005
EPI_ISL_11391	A/chicken/Tula/4/05
EPI_ISL_11386	A/chicken/Omsk/14/05
EPI_ISL_11385	A/chicken/Suzdalka/06/05
EPI_ISL_10137	A/chicken/Kurgan/05/2005
EPI_ISL_10138	A/duck/Kurgan/08/2005
EPI_ISL_9769	A/duck/Novosibirsk/56/2005
EPI_ISL_10121	A/Cygnus olor/Astrakhan/Ast05_2_10/2005
EPI_ISL_10053	A/Cygnus olor/Astrakhan/Ast05_2_5/2005
EPI_ISL_10052	A/Cygnus olor/Astrakhan/Ast05_2_6/2005
EPI_ISL_10008	A/Cygnus olor/Astrakhan/Ast05_2_2/2005
EPI_ISL_10048	A/Cygnus olor/Astrakhan/Ast05_2_3/2005
EPI_ISL_75725	A/Cygnus olor/Caspian Sea/2006
EPI_ISL_11390	A/chicken/Krasnodar/123/06
EPI_ISL_64332	A/chicken/Reshoty/02/2006
EPI_ISL_10543	A/grebe/Tyva/Tyv06_1/2006
EPI_ISL_10440	A/Grebe/Tyva/Tyv06_8/2006
EPI_ISL_10435	A/grebe/Tyva/Tyv06_2/06
EPI_ISL_64356	A/duck/Omsk/1822/2006
EPI_ISL_14423	A/chicken/Rostov/22_1/2007
EPI_ISL_63440	A/chicken/Domodedovo/MK/2007
EPI_ISL_11823	A/chicken/Moscow/2/2007
EPI_ISL_13577	A/chicken/Krasnodar/300/07
EPI_ISL_13848	A/Cygnus_cygnus/Krasnodar/329/07
EPI_ISL_20040	A/rook/Rostov_on_Don/26/2007
EPI_ISL_15359	A/pigeon/Rostov_on_Don/6/2007
EPI_ISL_14419	A/chicken/Rostov_on_Don/35/2007
EPI_ISL_15357	A/starling/Rostov_on_Don/39/2007
EPI_ISL_15358	A/muscovy duck/Rostov_on_Don/51/2007
EPI_ISL_63010	A/chicken/Primorje/1/2008
EPI_ISL_32668	A/bean_goose/Tyva/10/2009
EPI_ISL_32667	A/grebe/Tyva/3/2009
EPI_ISL_80265	A/great_crested_grebe/Tyva/22/2010
EPI_ISL_84604	A/grebe/Tyva/2/2010
EPI_ISL_31936	A/black-headed gull/Tyva/115/2009
EPI_ISL_28437	A/chicken/Primorsky/85/2008
EPI_ISL_9967	A/chicken/Kurgan/3/2005
EPI_ISL_64866	A/duck/Novosibirsk/02/05

EPI_ISL_10436	A/duck/Tuva/01/2006
EPI_ISL_10361	A/chicken/Krasnodar/01/2006
EPI_ISL_10360	A/chicken/Mahachkala/05/2006
EPI_ISL_64343	A/common gull/Chany/P/2006
EPI_ISL_31937	A/great crested grebe/Tyva/120/2009
EPI_ISL_25228	A/falcon/Saudi_Arabia/D1936/2007
EPI_ISL_15688	A/Mergus albellus/Slovakia/Vh212/2006
EPI_ISL_27786	A/Peregrine falcon/Slovakia/Vh242/2006
EPI_ISL_6414	A/swan/Slovenia/760/2006
EPI_ISL_27791	A/cygnus olor/Slovenia/156/06
EPI_ISL_27792	A/Ardea cinerea/Slovenia/185/06
EPI_ISL_27793	A/Anas platyrhynchos/Slovenia/359/2006
EPI_ISL_27794	A/Anas acuta/Slovenia/470/06
EPI_ISL_6249	A/chicken/Sudan/1784_7/2006
EPI_ISL_7014	A/chicken/Sudan/2115_10/2006
EPI_ISL_13230	A/tufted duck/Sweden/V526/2006
EPI_ISL_13231	A/goosander/Sweden/V539/2006
EPI_ISL_13233	A/tufted duck/Sweden/V599/2006
EPI_ISL_24705	A/eagle owl/Sweden/V618/2006
EPI_ISL_13240	A/smew/Sweden/V820/2006
EPI_ISL_24704	A/eagle owl/Sweden/V1218/2006
EPI_ISL_13249	A/herring gull/Sweden/V1116/2006
EPI_ISL_13247	A/tufted duck/Sweden/V1027/2006
EPI_ISL_13246	A/tufted duck/Sweden/V998/2006
EPI_ISL_13245	A/canada goose/Sweden/V978/2006
EPI_ISL_13241	A/mute swan/Sweden/V827/2006
EPI_ISL_69212	A/common pochard/Switzerland/WV4080110/2008
EPI_ISL_11231	A/common coot/Switzerland/V544/2006
EPI_ISL_11917	A/tufted duck/Switzerland/V504/2006
EPI_ISL_12822	A/mute swan/Switzerland/V68/2006
EPI_ISL_12823	A/little grebe/Switzerland/V330/2006
EPI_ISL_12824	A/duck/Switzerland/V389/2006
EPI_ISL_12825	A/duck/Switzerland/V426/2006
EPI_ISL_12826	A/duck/Switzerland/V487/2006
EPI_ISL_12827	A/common pochard/Switzerland/V505/2006
EPI_ISL_12828	A/mallard/Switzerland/V537/2006
EPI_ISL_12829	A/common pochard/Switzerland/V558/2006
EPI_ISL_12830	A/common pochard/Switzerland/V592/2006
EPI_ISL_12831	A/common pochard/Switzerland/V762/2006
EPI_ISL_10107	A/turkey/Turkey/1/2005
EPI_ISL_83693	A/Pigeon/Turkey-Agri/09rs2841-8/2005
EPI_ISL_83692	A/Pigeon/Turkey-Kars/09rs2841-7/2005
EPI_ISL_83700	A/Chicken/Turkey-Edirne/09rs2843-5/2008
EPI_ISL_83699	A/Chicken/Turkey-Batman/09rs2842-92/2007
EPI_ISL_83698	A/Chicken/Turkey-Bitlis/09rs2841-93/2006
EPI_ISL_83697	A/Chicken/Turkey-Batman/09rs2841-71/2006
EPI_ISL_83696	A/Chicken/Turkey-Mus/09rs2841-51/2006

EPI_ISL_83695	A/Chicken/Turkey-Mus/09rs2841-50/2006
EPI_ISL_83694	A/Chicken/Turkey-Isparta/09rs2841-37/2006
EPI_ISL_10340	A/chicken/Crimea/08/2005
EPI_ISL_103335	A/swan/England/AV26-70/2008
EPI_ISL_103350	A/swan/England/AV380-2326/2008
EPI_ISL_99850	A/Common Buzzard/Berlin/1/2006
EPI_ISL_6924	A/chicken/Egypt/2253_1/2006
EPI_ISL_6925	A/turkey/Egypt/2253_2/2006
EPI_ISL_6384	A/duck/Ivory Coast/1787_18/2006
EPI_ISL_2932	A/chicken/Nigeria/SO494/2006
EPI_ISL_66863	A/chicken/Nigeria/08RS848-117/2007
EPI_ISL_66864	A/chicken/Nigeria/08RS848-118/2007
EPI_ISL_66867	A/chicken/Nigeria/08RS848-121/2007
EPI_ISL_292	A/whooper swan/Mongolia/3/05
EPI_ISL_293	A/whooper swan/Mongolia/4/05
EPI_ISL_372	A/whooper swan/Mongolia/2/06
EPI_ISL_548	A/common goldeneye/Mongolia/12/2006
EPI_ISL_9940	A/migratory duck/Jiangxi/2300/2005
EPI_ISL_9939	A/migratory duck/Jiangxi/2295/2005
EPI_ISL_9938	A/migratory duck/Jiangxi/2136/2005
EPI_ISL_9937	A/migratory duck/Jiangxi/1701/2005
EPI_ISL_9936	A/migratory duck/Jiangxi/1657/2005
EPI_ISL_9935	A/migratory duck/Jiangxi/1653/2005
EPI_ISL_3903	A/duck/Zhejiang/11/2000
EPI_ISL_3900	A/duck/Shanghai/35/2002
EPI_ISL_3891	A/duck/Fujian/19/2000
EPI_ISL_68660	A/chicken/Hong Kong/786-2/1997
EPI_ISL_966	A/Chicken/Hong Kong/728/97
EPI_ISL_62720	A/bar-headed goose/Mongolia/X54/2009
EPI_ISL_62738	A/whooper swan/Mongolia/8/2009
EPI_ISL_76273	A/whooper swan/Mongolia/1/2010
EPI_ISL_76635	A/whooper swan/Mongolia/21/2010
EPI_ISL_93807	A/duck/Hokkaido/WZ101/2010
EPI_ISL_93808	A/duck/Hokkaido/WZ83/2010
EPI_ISL_131214	A/chicken/Shimane/1/2010
EPI_ISL_131215	A/mute swan/Toyama/1/2010
EPI_ISL_123277	A/chicken/Pakistan-Khi/NARC-520/2008
EPI_ISL_123318	A/crow/Pakistan-Khi/NARC-11672/2008

605   **Table S3** Summary profile of H5N1 HPAIV sequences isolated from reported outbreaks in
606   Europe and the Russia between May 2005 and June 2010 ($N = 277$).

Country	N.seq	Collection date	Host type
Austria	20	February 2006-April 2006	Wild
Belgium	1	November 2008	Wild
Bosnia	1	February 2006	Wild
Bulgaria	1	March 2010	Wild
Croatia	1	October 2005	Wild
Czech Republic	6	March 2006-July 2007	Wild & domestic
Denmark	8	March 2006-April 2006	Wild
France	2	February 2006-June 2007	Wild
Germany	57	November 2005-October 2008	Wild, domestic, & Other
Hungary	5	February 2006-January 2007	Wild
Italy	3	February 2006	Wild
Poland	9	March 2006-December 2007	Wild & domestic
Romania	78	October 2005-March 2010	Wild, domestic, & Other
Russia	51	July 2005-January 2010	Wild & domestic
Slovakia	2	February 2006	Wild
Slovenia	5	February 2006-March 2006	Wild
Sweden	11	March 2006-April 2006	Wild
Switzerland	13	February 2006-February 2008	Wild
Ukraine	1	May 2005	Domestic
United Kingdom	2	January 2008	Wild

609   **Table S4** Candidate phylogeographic models for HA and NA gene segments explored for relative fit.
610

Model	Sites	Discrete-traits	Branch rates	Trees
1			Strict clock	
2		Symmetric reversible	UCLN	
3			UCED	Coalescent: GMRF
4	GTR+GAMMA		Strict clock	Bayesian Skyride
5		Asymmetric irreversible	UCLN	
6			UCED	

611
612

Figure S1. Maximum clade credibility (MCC) trees for H5N1 HPAIV hemagglutinin (HA) and neauraminidase (NA) gene regions (A and B, respectively). Branch lengths are rendered proportional to absolute time (see timescales), and the branch thickness is proportional to the posterior probability of the inferred host state. Nodes correspond to median ages and the blue horizontal bars at nodes represent the corresponding 95% HPDs for divergence-time estimates.

2003 2004 2005 2006 2007 2008 2009 2010   2003 2004 2005 2006 2007 2008 2009 2010

Figure S2. Maximum clade credibility (MCC) trees for H5N1 HPAIV hemagglutinin (HA)
and neauraminidase (NA) gene regions (A and B, respectively). Branch lengths are ren-
dered proportional to absolute time (see timescales), and the branch thickness is propor-
tional to the posterior probability of the inferred location state. Nodes correspond to medi-
an ages and the blue horizontal bars at nodes represent the corresponding 95% HPDs for
divergence-time estimates.

Figure S3. Temporal dynamics of the spatial diffusion of H5N1 HPAIV. The figure consists of snap shots of the dispersal pattern of avian influenza between May 2005 and June 2010. Lines between locations represent branches in the MCC tree along which spatial transmission occurs. The diameters of circles are proportional to square root of the number of MCC branches maintaining a particular location state at each time-point. The blue and red color gradients reflect the relative age of the transitions for each gene (older-recent, respectively). A: HA gene. B: NA gene. The maps are based on satellite pictures available in NASA World Wind (http://worldwind.arc.nasa.gov/java/).

File S1 Temporal dynamics of the spatial diffusion of H5N1 HPAIV based on the HA gene. The
KML file demonstrates the dispersal pattern of avian influenza between May 2005 and June 2010.
Lines between locations represent branches in the MCC tree along which spatial transmission occurs.
The diameters of circles are proportional to square root of the number of MCC branches maintaining a
particular location state at each time-point. The blue and red color gradients reflect the relative age of
the transitions for HA gene (older-recent, respectively).