Dis-integrating the fly: A mutational perspective on phenotypic integration and covariation.

Running title: Mutational effects on environmental covariation

Authors: Annat Haber[1,2] & Ian Dworkin[1,3,4]

Author affiliations:

1)   BEACON Center for the study of Evolution in Action. Michigan State University.

2)   Department of Zoology, Tel Aviv University

3)   Department of Integrative Biology, Michigan State University.

4)   Department of Biology, McMaster University.

Correspondence: annat22@gmail.com, dworkin@mcmaster.ca

Key words: Drosophila, integration, canalization, morphometrics, covariation, mutation

Number of words: 5230

Number of tables: 3

Number of figures: 7

Data will be archived in DRYAD

<u>Abstract</u>

The structure of environmentally induced phenotypic covariation can influence the

effective strength and magnitude of natural selection. Yet our understanding of the factors that

contribute to and influence the evolutionary lability of such covariation is poor. Most studies

have examined either environmental variation, without accounting for covariation, or examined

phenotypic and genetic covariation, without distinguishing the environmental component. In this

study we examined the effect of mutational perturbations on different properties of

environmental covariation, as well as mean shape. We use strains of *Drosophila melanogaster*

bearing well-characterized mutations known to influence wing shape, as well as naturally-derived

strains, all reared under carefully-controlled conditions and with the same genetic background.

We find that mean shape changes more freely than the covariance structure, and that different

properties of the covariance matrix change independently from each other. The perturbations

affect matrix orientation more than they affect matrix size or eccentricity. Yet, mutational effects

on matrix orientation do not cluster according to the developmental pathway that they target.

These results suggest that it might be useful to consider a more general concept of

'decanalization', involving all aspects of variation and covariation.

Introduction

The evolutionary response to natural selection requires phenotypic variation. As such, the

mechanisms generating phenotypic variation and covariation (**P**) are of fundamental importance.

Most studies focus on the genetic component in **P**, summarized as the genetic covariance matrix

G, due to its role in the response to natural selection as summarized in the multivariate version

of the breeder's equation (Lande 1979; Lande and Arnold 1983). Thus, the generation of

heritable components of variation, governed by mutation and recombination, is broadly studied.

However non-heritable sources of phenotypic variation, often summarized through **E**, remain

important for our understanding of both the magnitude and direction of response to natural

selection through its contributions to **P**. The basic measure of the strength of selection - the

selection differential - depends on the phenotypic variance of a trait via the covariance of the

phenotype and fitness (the Robertson-Price identity; Robertson 1966; Price 1970). More

generally, the magnitude of the selection differential is a function of both the shape of the

underlying fitness profile (W(z)) and the current population mean and variance for the trait (V_z)

associated with fitness. In addition to its influence on the magnitude of selection, **E** (via **P**)

influences the response to selection as summarized in Lande's (1979) equation. When

considering a multivariate set of phenotypes, the influence of covariation in **E** becomes

increasingly relevant, influencing the orientation of the response to selection. Although

environmental covariation may not facilitate adaptive evolution, it can certainly impede it, and

possibly bias its trajectory towards a secondary peak (Burger 1986). Yet, our understanding of

how this environmental covariation is generated during development is incomplete, as is our

understanding of how it evolves.

Mutational and environmental perturbations have often been used to examine how the

environmental component of phenotypic variation changes, particularly in the context of

canalization (Waddington 1952; Dworkin 2005a,b; Levy and Siegal 2008; Hallgrímsson et al. 2009;

Paaby and Rockman 2014; and references therein). Numerous studies have found an increase in

phenotypic variance in response to perturbations, often attributed to decanalization, and due in

part to a release of cryptic genetic variation. However, both increases and decreases of

environmental variance have also been observed without the release of genetic variation. Most

studies (e.g., Dworkin 2005a; Levy and Siegal 2008), however, have examined the effect of

mutational perturbations on the variance of a trait, or the total variance across several traits,

whereas in multivariate systems other aspects of covariation need to be considered.

Covariance matrices have a number of important properties, namely matrix size, shape

(eccentricity), and orientation (Jones et al. 2003; Hohenlohe and Arnold 2008), as illustrated in

Figure 1. The size of a matrix is measured as the total variance (sum of the individual trait

variances) and quantifies variation overall, irrespective of how it is distributed among different

directions (Fig. 1A). The orientation of the matrix refers to the direction in which the primary axes

of covariation are pointing (Fig. 1B), and is determined by the relative contribution of each trait

to each eigenvector. Thus, a change in matrix orientation reflects a change in the direction in

which most of the variation is concentrated. The eccentricity of the matrix describes its shape –

how much it deviates from a hypersphere – and it quantifies how evenly the variance is

distributed among the different directions (Fig. 1C), irrespective of the orientation, and,

depending on how it is measured, irrespective of matrix size. A more eccentric matrix (i.e. more

cigar-shaped) means that there is considerably more variation along the first axis relative to

other axes.

Previous studies provide mixed expectations as to how the different properties of the

covariance matrix would change relative to each other and relative to mean shape. Hallgrímsson

et al. (2009) postulated that when the variance increases due to perturbations, matrix

eccentricity can either increase or decrease, depending on how the perturbed developmental

process contributes to the original patterns of covariation. Eccentricity would increase with total

variance if the targeted pathway plays a major role in integrating the measured traits, thus

adding variance to the major axis of covariation. When the perturbed pathway does not

contribute much to the major axis of covariation, changes in variance are distributed among the

smaller subsequent eigenvalues, thus reducing eccentricity.

The association between changes to matrix orientation and eccentricity has mostly been

studied so far for **P** and **G** rather than **E**, making expectations for **E** far less clear. Simulations by

Jones et al. (2003, 2004, 2007, 2012) and Revell (2007) suggest that the orientation of **G** is more

likely to change than eccentricity under most conditions, including effects due to population size,

magnitude, and orientation of mutational correlation and stabilizing selection, and different

modes of directional selection. In addition, higher eccentricity enhances the stability of the

orientation under most conditions. The magnitude and stability of eccentricity and matrix size, on

the other hand, are mostly influenced by population size rather than mutational correlations and

selection. Empirical studies have found evidence that both eccentricity and orientation can be

either stable or labile across populations and species at different phylogenetic scales (Arnold et

al. 2008; de Oliveira et al. 2009; Porto et al. 2009; Haber 2015a,b). Unfortunately it is not clear

whether the expectations from these studies (examining **G** and **P**) are relevant to how **E** changes.

Changes in **G** reflect a combination of influences from mutation and other evolutionary forces.

The structure of **E**, on the other hand, reflects the interaction between the environment and

development. Thus, while strong directional selection can deplete genetic covariation, altering

matrix eccentricity and size of **G**, it is not clear *a priori* that this should alter the matrix properties

of **E** as well.

Empirical studies have found mutational effects on matrix orientation of **E** that could not

be readily associated with developmental pathways or mutational effect size (Hallgrímsson et al.

2006; Debat et al. 2009; Jamniczky and Hallgrímsson 2009). In addition, studies have found both

increases and decreases of eccentricity (Hallgrímsson et al. 2009). Most studies, however, did not

include a direct comparison of the different properties of **E** and shape variation across a wide

range of mutations. Thus, we are left with little understanding of the lability of **E**.

Drosophila wing shape is an ideal system to address these questions, and in particular

how mutational perturbations influence the different aspects of the covariance structure of **E**.

The development of the wing of *D. melanogaster* has been a model system for over 70 years, and

it is one of the best genetically characterized model systems. The extensive set of genetic tools,

mutant lines, and ability to generate many genetically identical individuals reared under common

environments enables the study of mutational effects on **E**. Variation in both size and shape of

the *Drosophila* wing has a long history of research, and sophisticated high dimensional

representations of shape can be generated in a high throughput manner (Houle et al. 2003;

Pitchers et al. 2013). Studies of de-canalization for *Drosophila* wing shape, and the influence of

mutations on aspects of **E** have had mixed results. While mutant perturbation can have profound

effects on mean shape (Weber 2005, Breuker et al. 2006; Debat et al. 2006; Dworkin and Gibson

2006; Debat et al. 2009, Debat et al. 2011), the influence on variance has been more mixed, with

some studies seeing a general increase (Debat et al. 2009, Debat et al. 2011) and others seeing

both increase and decrease (Debat et al. 2006; Dworkin and Gibson 2006). Some evidence for the

effect of mutational and environmental perturbations on matrix orientation has been shown in

these studies, but other aspects of the covariance structure have not been examined, and

changes to covariance and shape have not been compared directly.

To better understand the evolutionary lability of the **E** matrix, we used a set of induced

mutations measured against a common co-isogenic wild type (Dworkin and Gibson 2006; Debat

et al. 2009), as well as a panel of strains derived from natural populations, and examined the

relationship between mean shape and different aspects of environmental covariance. Thus, all

strains were reared under the same carefully-controlled conditions, and all mutant lines had the

same genetic background, minimizing unknown sources of variation. We observe that mean

shape varies more freely than the covariance structure, and that matrix orientation varies more

than – and independently from – its size and eccentricity.

<u>Material and Methods</u>

Drosophila strains

Insertional mutations (caused by the insertion of P-elements, marked with w^+) in genes in

the TGF-β and EGFR signaling pathways were provided from the Bloomington Stock Center (Table

1). These represent a subset of the alleles that were first described in Dworkin and Gibson (2006).

All insertions were initially introgressed into the wild-type lab strain Samarkand (Sam) for at least

10 generations. The Samarkand wild type was marked with w^- (white eyes) so flies with insertions

could be distinguished by a rescue of the eye color phenotype. Introgressions were performed by

repeated backcrossing of females bearing the insertion to males of Sam. Prior to generating flies

for the experiments described in this study, the alleles were backcrossed to Sam for an additional

four generations to remove any *de novo* mutations that had accumulated in these lines (relative

to Sam) since the original introgression procedure. Selection was based entirely on the presence

of the eye colour marker, precluding unwitting selection for wing phenotypes. All crosses were

performed using standard cornmeal-molasses media, in a 24° C Percival incubator on a 12/12-hr

light/dark cycle with 60% relative humidity (Dworkin and Gibson 2006). The mutant strains used

here are part of a larger unpublished study (Dworkin), and were chosen in part based on sample

size (>50/genotype).

Experimental setup

After the additional four generations of backcrossing back to Samarkand, crosses between

each mutation and Samarkand were set up in vials, allowing females to lay for 2-3 days, so that

egg density was low (generally less than 60 flies/vial). The temperature of the incubator was

maintained at 24° C, and monitored carefully for fluctuations, and vial position was randomized

within the incubator on a daily basis to reduce any possible edge effects. As larvae crawled out of

the media, a piece of paper towel was added to each vial to provide additional pupation space.

After eclosion and sclerotization flies were separated, on the basis of eye color, into wild-type

individuals without the P-element-induced mutations (w-) and heterozygotes for the P-element,

and then stored in 70% ethanol.

Strains derived from natural populations from North Carolina and Maine.

Iso-female lines of D. melanogaster were established separately in the summer of 2004 at

a Peach orchard in West End, North Carolina (NC2) and in a blueberry field in Cherryfield Maine

(courtesy of Marty Kreitman). To generate inbred lines, full-sib mating was performed for 15-20

generations (Goering et al 2009; Reed et al 2010). Flies were reared at 25C in a 12/12 light/dark

cycle at 50% humidity. Seven Maine and five NC2 lines, each with at least 51 males were used

here.

Data collection

A single wing from each fly was dissected and mounted in 70% glycerol (>20 wings per

genotype/replicate on average; males only). Wings were imaged at 40X magnification on an

Olympus DP30BW camera mounted on an Olympus BX51 microscope using 'DP controller' V3.1.1

software. All images were saved as greyscale tiff files. To extract landmark and semi-landmark

data, we followed a modified protocol for the use of the "WINGMACHINE" software (Houle et al.

2003) as detailed in Pitchers et al. (2013). First, we used "tpsDig2" (Rohlf 2010) software to

record the coordinates of the two starting landmarks needed by WINGMACHINE. We then used

WINGMACHINE to automatically fit nine B-splines to the veins and margins of the wing (Fig. 2A).

We reviewed each splined image and manually adjusted the control points as necessary. The x

and y coordinates of 14 landmarks and 34 semi-landmarks (Fig. 2A) were extracted using CPR

(Marquez and Houle 2014). The data were checked for visual outliers on scatter plots, and

putative outliers were examined, and either fixed in WINGMACHINE or deleted. The resulting

dataset was then passed on to **R** for further analysis.

Preparation of data

All specimens from all genotypes were superimposed together using Generalized

Procrustes Analysis with function *gpagen* in the **R** package *geomorph 2.1.1* (Adams and Otarola-

Castillo 2013). The semi-landmarks were optimized using minimum Procrustes distance. The

whole dataset was then projected into the tangent space. Replicate effect was corrected by

subtracting the replicate mean from each specimen and adding the grand mean of that genotype

(Sokal and Rohlf 1995; Marroig and Cheverud 2004). The natural strains were similarly corrected

for locality (Maine *vs.* North Carolina). Following these corrections, data were regressed on

centroid size within each genotype, and added to the predicted value for that genotype mean

size.

The size-regressed dataset was then passed on to LORY for calculating the interpolated

Jacobian-based data (Marquez et al. 2012). LORY uses spatial interpolation to evaluate shape

deformation at points throughout the wing. It starts by creating a tessellated grid, with the

landmarks and semi-landmarks as the vertices (Fig. 3A). The centroids of the resulting triangles

are considered the evaluation points. Given an interpolation function, it then calculates the

Jacobian matrix that describes the shape deformation at each point (e.g., Fig. 3B). The log-

transformed determinant of the Jacobian matrix quantifies the amount of expansion or

contraction at each evaluation point relative to the mean configuration. Thus, the interpolation

takes into account information from the whole configuration of landmarks, as well as from the

rest of the sample through the mean configuration, while quantifying local shape changes. The

resulting dataset is multivariate, rather than multidimensional, in the sense that each variable is

meaningful in its own right and the whole dataset can be analyzed using conventional

multivariate analyses (Marquez et al. 2012).

The interpolation function necessarily assumes a model for the distribution of shape

changes across the configuration. The most common interpolation function used in geometric

morphometrics studies is the Thin Plate Spline (TPS), which assumes the object is rigid and

therefore penalizes local deformations more than global ones. In this study we compared TPS

with another interpolation function implemented in LORY – the Elastic Body Spline (EBS) – that

assumes the object is elastic and penalizes local deformations less than global ones (Marquez et

al. 2012). Therefore, like the Procrustes superimposition, TPS tends to spread the variation more

globally than EBS. For this study system, we consider EBS to be a more suitable model *a priori*,

because the development of the fruit fly wing has been shown to proceed in a relatively

compartmentalized manner in which variation tends to be locally contained (Zecca and Struhl

2002; Barrio and Celis 2004; Cook et al. 2004; Martín and Morata 2006; Ziv et al. 2012). At the

same time, results based on Procrustes data were highly correlated with the Jacobian-based

data, using both TPS and EBS (Figs. S1-S2). Therefore, we chose to focus here on results based on

EBS. Results for TPS and Procrustes data are provided in the supplemental material. Each dataset

was reduced to its first 30 dimensions using Principal Component Analysis, covering 99% of the

variation in the data (Fig. S3). All subsequent analyses are based on these PCA scores.

Quantification of variation and covariation

Variance-covariance (VCV) matrices were quantified based on their size, orientation, and

shape (see Fig. 1). Matrix size was calculated as the trace of the matrix (i.e., total variance, or

sum of its eigenvalues). Matrix shape was characterized based on its eccentricity (Jones et al.

2003), and calculated as the ratio of the largest eigenvalue to the total variance, which is the

inverse of Kirkpatrick (2009)'s effective number of dimensions (Kirkpatrick 2009):

$$\text{Eccentricity} = \frac{\lambda_1}{\sum_{i=1}^{P} \lambda_i}$$

231   In addition, we calculated the relative standard deviation of the eigenvalues (rSDE), also

232 known as integration level (Pavlicev et al. 2009; Haber 2011). However, rSDE yielded very similar

233 results to the above measure of eccentricity (Fig. S4), and will not be considered further.

234   Similarity in matrix orientation was quantified using Random Skewers (Cheverud et al.

235 1983; Cheverud and Marroig 2007; Marroig et al. 2011), which measures the average similarity

236 between two matrices in their response to random unit vectors (representing selection). We

237 used 5000 random vectors drawn from a normal distribution with mean 0 and variance 1,

238 normed to unit length (Marroig et al. 2011). A covariance distance metric was then calculated as

239 $\sqrt{(1-r^2)}$, where r is the Random Skewers value for that pair of matrices, and normalized using

240 Fisher's z-transformation (Jamniczky and Hallgrímsson 2009). Two other distance metrics were

241 calculated as well: a modification of Krzanowski's (1979) metric following Zelditch et al. (2006),

242 and the relative eigenvalues metric of Mitteroecker and Bookstein (2009). However, these two

243 methods proved unstable for our dataset under resampling, yielding unreliable CI's. At the same

244 time, their estimated values followed the same pattern as Random Skewers (see online

245 supplementary material). Moreover, Random Skewers is the only method of the three that is

246 related to evolutionary theory (Hansen and Houle 2008). Therefore, we present below only

247 results based on Random Skewers.

248   Shape distances were calculated as the Euclidean distance for both the interpolated data

249 and the Procrustes-based data, after projecting the Procrustes data to tangent space.

250   Confidence intervals for shape distances, matrix size, and eccentricity, were estimated

using a non-parametric bootstrap procedure with a BCa correction (DiCiccio and Efron 1996;

Carpenter and Bithell 2000) using 999 iterations. The BCa correction was necessary because the

pseudovalue distribution is expected to be biased upward when the statistic is bounded by zero,

and to depend on its mean when bounded by both zero and one. These confidence intervals

were used for evaluating significant differences as well.

Confidence intervals for covariance distance were estimated using a Jackknife procedure.

Each Jackknife pseudovalue was calculated by leaving out one of the specimens in one of the two

samples compared. The confidence interval was calculated as the 95% of the Jackknife

distribution, without applying a bias correction. The Jackknife was preferred in this case because

the bootstrap (both parametric and non-parametric) consistently resulted in distributions that

were highly biased upward, often excluding the observed value, invalidating the BCa correction.

This is a commonly found phenomenon in similar studies that has not yet been properly

addressed in the literature to the best of our knowledge. Here we found that Jackknife without a

bias correction provided fairy symmetric distributions around the observed values with

reasonably wide confidence intervals, whereas the bias-corrected jackknife yielded extremely

narrow intervals that often excluded their observed value (i.e., they were "over-corrected").

In order to further compare covariance changes with shape changes we used Principal

Coordinates Analysis to generate a shape space and a covariance space, based on all pairwise

distances. Thus, in each of these spaces, genotypes are located based on how different they are

from each other in either shape or matrix orientation. The covariance space was then

superimposed onto the shape space using a symmetric Procrustes superimposition including

scaling (function *protest* in **R** package *vegan 2.0-10*; Oksansen et al. 2013). The sum of squared

deviations between the two spaces, after superimposition, was used as a measure of correlation

between them. In addition, we calculated the disparity of mutants and naturally-derived strains -

for shape and covariance - as the sum of variances of their respective scores in the joint space.

We used MANOVA (Pillai's Λ) to test the effect of pathway on the mutants' PCoA scores in both

shape and covariance spaces.

All analyses were carried out in **R** 3.0.3. All scripts and data will be made available on

DRYAD and on the Dworkin lab github repository.

Results

Both mean shape and matrix orientation vary considerably among mutant and

naturally-derived strains.

All mutants and naturally-derived strains differ significantly in their mean shape from the

Sam wild type (Table 2 and Fig. 4), as determined by the lack of overlap between their confidence

intervals and the benchmark of zero distance from Sam (horizontal dashed line in figure 4). Most

mutant strains are more similar to the Sam wild type than any of the naturally-derived strains.

Yet, two of the mutants (*Omb*^{md653} and Bs^{k07909}) show magnitudes of shape change as extreme as

any of the naturally derived strains, indicating that the range of mutations included here provide

a reasonable representation of what we might observe in nature.

The orientation of the covariance matrix of all strains – mutants as well as naturally

derived – also differ significantly from the Sam wild type (Table 2 and Fig. 5): none of the

covariance distances include zero in their confidence interval. Unlike mean shape, however, most

of the mutant strains differ from Sam just as much as the naturally-derived strains do.

Several mutant strains have lower environmental variance than their wild type

Surprisingly, most mutations resulted in a lower total variance than their co-isogenic Sam

wild-type (Table 2 and Fig. 4). However, only 4 of the 12 mutant strains fall below the confidence

interval of Sam (Table 2). One of the significant strains ($tkv^{kg01923}$) is part of the TGF-β pathway,

and the other three (S^{k09530}, $rho^{kg07115}$, spi^{s3547}) contribute to EGFR signaling (Table 1). The only

strain that has substantially and significantly higher total variance is Omb^{md653}, which also differs

greatly in its mean shape. The variance within each of the naturally-derived strains is mostly

higher than that of Sam, but only 3 of the 12 naturally-derived strains are above the confidence

interval of Sam. In addition, there is a positive linear relationship between the total variance

within each mutant strain and its shape distance from Sam across the mutant strains (Fig. 4). This

is consistent with the common expectation that greater decanalization is associated with larger

changes to mean shape.

A positive non-linear association between changes in mean shape and matrix

orientation.

There is also a weak, but positive association between covariance distance (orientation)

and shape distance (Fig. 5), relative to the Sam wild type. This relationship, however, is not linear,

suggesting that the covariance distance is somewhat bounded at the upper range. These findings

are further supported by comparing the shape and covariance PCoA spaces (Fig. 6). These spaces

are based on all pairwise comparisons, rather than comparing to Sam only, and were

superimposed for comparability. The Procrustes superimposition scales the two spaces to a

common scale, centers, and rotates them so that they are comparable in terms of both

magnitude and direction of change. The grey solid lines indicate the deviation between the two

spaces. The Procrustes sum of squares 0.669, and the coefficient of determination between the

two spaces is small yet significantly different from zero (R^2 = 0.33; p≤0.001 with 999

permutations). Thus, the two spaces are substantially different from each other, but not

unrelated. The first PCo1 axis mostly separates the naturally-derived strains from the

mutants+Sam, for both shape and covariance. The disparity of shape is more than twice as large

as that of the covariance structure when strain Omb^{md653} is included, and twice as large when it is

excluded (Table 3; calculations include all 24 PCoA dimensions that resulted in eigenvalues larger

than zero). The disparity of both shape and covariance is larger for the naturally-derived strains

than for the mutants, both with and without Omb^{md653}. The mutants do not cluster by pathway

(Fig. 6B), for either shape (Pillai's Λ=0.44, p=0.24) or covariance distance (Pillai's Λ=0.34, p=0.52).

Most mutations do not influence matrix eccentricity, despite effects on mean shape,

matrix size, and matrix orientation.

Most of the mutations do not have a substantial effect on the eccentricity of the matrix

(i.e., matrix shape). Most values fall well within the confidence interval of their co-isogenic Sam

wild type (Table 2). The only mutation that made a substantial (and significant) increase in

eccentricity is aos^{W11}. Interestingly, aos^{W11} does not differ much from Sam for total variance (Fig.

7A). Omb^{md653}, on the other hand, differs significantly and substantially from Sam in its total

variance and shape (see Table 2), and yet is very similar in its eccentricity to Sam. Most of the

naturally-derived strains have a higher eccentricity than Sam (Fig. 7A), showing a greater

difference from Sam than the mutants do, as well as a slight tendency toward a positive

association with the total variance. However, this association depends mostly on the two most

deviating strains. When eccentricity is calculated using the relative standard deviation of the

eigenvalues (rSDE), a few additional strains differ significantly from Sam (Table 2), but the

relationship between eccentricity and total variance is still weak (Fig. S12). Similarly, there is no

clear association between eccentricity and shape distance from Sam (figure 7B), and no

relationship between eccentricity and covariance distance from Sam (Fig. 7C). The same picture

emerges using other metrics of covariance distance, based on both Procrustes and TPS data (Figs.

S10-S11).

Discussion

The pattern and magnitude of phenotypic covariation remains central to selection theory

(Robertson 1966; Price 1970; Lande and Arnold 1983). Although it cannot facilitate adaptation,

the environmental component of phenotypic covariation, **E**, can impede and divert the

population response to selection (Burger 1986). Despite the importance of this for our

understanding the response to selection, factors that influence the lability of **E** are poorly

understood. For univariate measures, environmental and genetic stressors have been shown to

increase variance in **E** in numerous situations (Dworkin 2005; Hallgrímsson et al. 2009; Paaby and

Rockman 2014; and references therein), but little is known about how such perturbations

influence other properties of integration and covariation.

Using the wing of *Drosophila melanogaster* as a model system, we found that mean shape

and matrix orientation vary among the co-isogenic mutants, as well as the naturally-derived

strains (Table 1 and Fig. 5). By superimposing the covariance space onto shape space we were

able to compare their distribution in terms of both magnitude and direction of change and show

that shape has changed to a greater extent than matrix orientation (Fig. 6). In contrast, mutations

had little influence on the total variance and eccentricity of the matrix. Whereas total variance is

positively associated with mean shape, it is not associated with eccentricity, nor is eccentricity

associated with matrix orientation. All together, these findings suggest that the potential of the

covariance structure to change is more limited than that of mean shape, and that different

properties of the covariance structure can change independently from each other. It is worth

noting that some of the environmental effects (such as rearing temperature and density) were

highly controlled in this study, both within and among strains. Thus, our experimental design may

represent a lower bound for the environmental contributions to covariation, and the patterns

observed are likely an underestimate of the possible effect sizes.

Contrary to the common expectation (Dworkin 2005; Hallgrímsson et al. 2009; and

references therein), the variance of most mutants decreases relative to their co-isogenic wild

type, Samarkand (Sam) (albeit significant for only 4/12 genotypes). Equally puzzling is the

observation that the natural strains all have higher variance than Sam rather than distributed

around Sam. Since the individuals within each strain are genetically identical, the decrease in

total variance within the mutants, and the increase within the natural strains, likely reflects a

higher, and lower, canalization of wing shape, respectively (Hall et al 2007). Such mixed results

have been observed also by Dworkin and Gibson (2006) for mutant strains, using a smaller set of

landmarks. A related study (Debat et al 2009) showed a more consistent increase in total

variance, even for some of the same mutations. However, the genetic background of the wild

type strains in Debat et. al (2009) was Oregon-R (rather than Samarkand), which has been shown

to be more sensitive to mutational perturbations under some conditions (Chari & Dworkin 2013).

Increases in the environmental component of phenotypic variance have been explained in

previous studies by a nonlinear relationship between the trait mean and its underlying

developmental parameters (Klingenberg and Nijhout 1999; Hallgrímsson et al. 2009). A steeper

slope for relationship between underlying developmental parameters and phenotypic variation

implies greater variation is expressed around the mean phenotype, and therefore a less canalized

phenotype. It often assumed that the wild type is at its most canalized state, and therefore when

development is perturbed the mean is expected to shift to a less canalized section on the curve

and express greater variation. It is possible, however, that the mean would shift into more

canalized regions, rather than less canalized, resulting in a decrease of variance rather than an

increase. It is also possible to have system-wide canalizing factors (such as chaperon proteins;

Rutherford and Lindquist 1998) operate simultaneously, masking the effects of local

perturbations and stabilizing the variance (Hallgrímsson et al. 2006). Or, the mutations could

increase redundancy and complexity thus increasing stability (Hallgrímsson et al. 2006; Levy and

Siegal 2008). All of these possible explanations would be consistent with the relationship we

found between changes of mean shape and total variance (relative to the Sam co-isogenic wild

type).

A possible explanation for the increase in variance within the natural strains could be

related to the time and amount of inbreeding among the strains (Whitlock & Fowler 1999;

Whitlock et al. 2002). Inbreeding in *Drosophila* does represent a form of "genetic stress" as

deleterious recessive alleles are made homozygous (and their effects are phenotypically

expressed). We would therefore expect the natural strains to be de-canalized as well. However,

the Sam wild-type has also undergone very extensive inbreeding, and is near isogenic (Chandler

et al. 2014). Thus, a satisfactory explanation for this particular observation eludes us.

In accord with previous studies (Dworkin and Gibson 2006; Debat et al. 2009; Jamniczky

and Hallgrímsson 2009), covariance matrices of the mutant strains do not cluster according to the

developmental pathway that they presumably affect. This could points to the complexity of the

developmental system, further supporting the palimpsest model suggested by Hallgrímsson et al.

(2009). It could also mean that these pathways do not contribute to the structure of **E** as

effectively and consistently as previously postulated. A detailed analysis of the covariance

pattern is needed to further investigate the extent to which the modularity structure of the fruit

fly wing follows *a priori* developmental models, and how it is disrupted by developmental

perturbations.

It is difficult to compare differences in eccentricity to differences in orientation directly

because of their different dimensionality and somewhat different method for calculating CI's.

However, mutant eccentricity varies from that of Sam within a smaller portion of its possible

range (0.2-0.43; 23%) compared to how much their orientation varies from Sam (0.37-0.77; 40%

without the Fisher's *z*-transformation). These findings are in accord with simulations by Jones et

al. (2003, 2004, 2012) for **G**, suggesting that eccentricity is likely to vary less than orientation for

a given population size under most combinations of genetic architecture, selection regimes, and

phylogenetic scale. It is also consistent with empirical evidence for **P** from Haber (2015a,b),

showing that matrix orientation has varied greatly among closely-related ruminant species,

whereas eccentricity has remained largely the same throughout most of bovids history, and only

varied within 33% of its possible range among other ruminants.

Previous studies have shown an increase in variance to be clearly associated with

decanalization. In this study, however, we find that mutants altered the orientation of the

covariance matrix more often than its total variance. Mutations do not only affect trait means

and variances, but aspects of covariances as well, though this has not been part of the general

formulation so far. Thus, our study suggests that it might be useful to consider a more general

concept of decanalization.

Acknowledgements: We thank Ellen Larsen, Jeff Conner and Will Pitchers for comments

that improved clarity of this manuscript. This material is based in part on work supported by the

National Science Foundation under Cooperative Agreement No. DBI-0939454. Any opinions,

findings and conclusions or recommendations expressed in this material are those of the

author(s) and do not necessarily reflect the views of the National Science Foundation. This work

was supported by NIH grant 1R01GM094423-01 to ID. Funding for AH was provided by the

Council of Higher Education, Israel, and BEACON Center for the Study of Evolution in Action at

Michigan State University.

Literature cited

Adams, D. C., and E. Otarola-Castillo. 2013. geomorph: an R package for the collection and

analysis of geometric morphometric shape data. Methods in Ecology and Evolution 4:393-

399.

Arnold, S. J., R. Burger, P. A. Hohenlohe, B. C. Ajie, and A. G. Jones. 2008. Understanding the

evolution and stability of the G-matrix. Evolution 62:2451-2461.

Arnold, S. J., R. Bürger, P. A. Hohenlohe, B. C. Ajie, and A. G. Jones. 2008. Understanding the

evolution and stability of the G-matrix. Evolution 62:2451-2461.

Barrio, R., and C. de, Jose F. 2004. Regulation of *spalt* expression in the *Drosophila* wing blade in

response to the Decapentaplegic signaling pathway. Proceedings of the National Academy

of Sciences of the United States of America 101:6021-6026.

Breuker, C. J., J. S. Patterson, and C. P. Klingenberg. 2006. A Single Basis for Developmental

Buffering of *Drosophila* Wing Shape. PLoS ONE 1:e7.

Burger, R. 1986. Constraints for the evolution of functionally coupled characters: A nonlinear

analysis of a phenotypic model. Evolution 40:182-193.

Camara, M., C. Ancell, and M. Pigliucci. 2000. Induced mutations: a novel tool to study

phenotypic integration and evolutionary constraints in *Arabidopsis thaliana*. Evolutionary

Ecology Research 2:1009-1029.

Carpenter, J., and J. Bithell. 2000. Bootstrap confidence intervals: when, which, what? A

practical guide for medical statisticians. Statistics in Medicine 19:1141-1164.

Chari, S., and I. Dworkin. 2013. The Conditional Nature of Genetic Interactions: The

Consequences of Wild-Type Backgrounds on Mutational Interactions in a Genome-Wide

Modifier Screen. PLoS Genetics 9:e1003661.

Chandler, C. H., S. Chari, D. Tack, and I. Dworkin. 2014. Causes and Consequences of Genetic

Background Effects Illuminated by Integrative Genomic Analysis. Genetics 196:1321-1336.

Cheverud, J. M., J. J. Rutledge, and W. R. Atchley. 1983. Quantitative Genetics of Development:

Genetic Correlations Among Age-Specific Trait Values and the Evolution of Ontogeny.

Evolution 37:895-905.

Cook, O., B. Biehs, and E. Bier. 2004. *brinker* and *optomotor-blind* act coordinately to initiate

development of the L5 wing vein primordium in *Drosophila*. Development 131:2113-2124.

de Oliveira, F. B., A. Porto, and G. Marroig. 2009. Covariance structure in the skull of Catarrhini:

a case of pattern stasis and magnitude evolution. Journal of Human Evolution 56:417-430.

Debat, V., S. Bloyer, F. Faradji, N. Gidaszewski, N. Navarro, P. Orozco-terWengel, V. Ribeiro, C.

Schlötterer, J. S. Deutsch, and F. Peronnet. 2011. Developmental Stability: A Major Role

for *Cyclin G* in *Drosophila melanogaster*. PLoS Genetics 7:e1002314.

Debat, V., A. Debelle, and I. Dworkin. 2009. Plasticity, canalization, and developmental stability

of the *Drosophila* wing: joint effects of mutations and developmental temperature.

Evolution 63:2864-2876.

Debat, V., C. C. Milton, S. Rutherford, C. P. Klingenberg, and A. A. Hoffmann. 2006. Hsp90 and

the quantitative variation of wing shape in *Drosophila melanogaster*. Evolution 60:2529-

2538.

DiCiccio, T. J., and B. Efron. 1996. Bootstrap Confidence Intervals. Statistical Science 11:189-212.

Dworkin, I. 2005a. A study of canalization and developmental stability in the sternopleural

bristle system of *Drosophila melanogaster*. Evolution 59:1500-1509.

Dworkin, I. 2005b. Canalization, cryptic variation, and developmental buffering: a critical

examination and analytical perspective. Pp. 131-158 *in* B. Hallgrímsson and B. K. Hall, eds.

Variation, a central concept in biology

Dworkin, I., and G. Gibson. 2006. The EGF-R and TGF-ß signal transduction pathways contribute

to variation for wing shape in *Drosophila melanogaster*. Genetics 173:1417–1431.

Goering, L. M., P. K. Hunt, C. Heighington, C. Busick, P. Pennings, J. Hermisson, S. Kumar, and G.

Gibson. 2009. Association of orthodenticle with natural variation for early embryonic

patterning in *Drosophila melanogaster*. Journal of Experimental Zoology Part B: Molecular

and Developmental Evolution 312B:841-854.

Haber, A. 2011. A Comparative Analysis of Integration Indices. Evolutionary Biology 38:476-488.

Haber, A. 2015a. The Evolution of Morphological Integration in the Ruminant Skull. Evolutionary

Biology 42:99-114.

Haber, A. 2015b. Phenotypic Covariation And Morphological Diversification In The Ruminant

Skull. bioRxiv doi: http://dx.doi.org/10.1101/017533.

Hall, M. C., I. Dworkin, M. C. Ungerer, and M. Purugganan. 2007. Genetics of

microenvironmental canalization in *Arabidopsis thaliana*. Proceedings of the National

Academy of Sciences 104:13717.

Hallgrímsson, B., H. A. Jamniczky, N. Young, C. Rolian, T. Parsons, J. Boughner, and R. Marcucio.

2009. Deciphering the Palimpsest: Studying the Relationship Between Morphological

Integration and Phenotypic Covariation. Evolutionary Biology 36:355-376.

Hallsson, L. R., and M. Björklund. 2012. Selection in a fluctuating environment leads to

decreased genetic variation and facilitates the evolution of phenotypic plasticity. Journal

of Evolutionary Biology 25:1275-1290.

Hansen, T. F., and D. Houle. 2008. Measuring and comparing evolvability and constraint in

multivariate characters. Journal of Evolutionary Biology 21:1201-1219.

Hohenlohe, P. A., and S. J. Arnold. 2008. MIPoD: A Hypothesis-Testing Framework for

Microevolutionary Inference from Patterns of Divergence. The American Naturalist

171:366-385.

Houle, D., J. Mezey, P. Galpern, and A. Carter. 2003. Automated measurement of Drosophila

wings. BMC Evolutionary Biology 3:25.

Jamniczky, H. A., and B. Hallgrímsson. 2009. A Comparison of Covariance Structure in Wild and

Laboratory Muroid Crania. Evolution 63:1540-1556.

Jones, A. G., S. J. Arnold, and R. Bürger. 2003. Stability of the G-Matrix in a Population

Experiencing Pleiotropic Mutation, Stabilizing Selection, and Genetic Drift. Evolution

57:1747-1760.

Jones, A. G., S. J. Arnold, and R. Bürger. 2004. Evolution and Stability of the G-Matrix on a

Landscape with a Moving Optimum. Evolution 58:1639-1654.

Jones, A. G., S. J. Arnold, R. Bürger, and C. Goodnight. 2007. The mutation matrix and the

evolution of evolvability. Evolution 61:727-745.

Jones, A. G., R. Bürger, S. J. Arnold, P. A. Hohenlohe, and J. C. Uyeda. 2012. The effects of

stochastic and episodic movement of the optimum on the evolution of the G-matrix and

the response of the trait mean to selection. Journal of Evolutionary Biology 25:2210-2231.

Kirkpatrick, M. 2009. Patterns of quantitative genetic variation in multiple dimensions. Genetica

136:271-284.

Klingenberg, C. P., and H. F. Nijhout. 1999. Genetics of Fluctuating Asymmetry: A Developmental

Model of Developmental Instability. Evolution 53:358-375.

Lande, R. 1979. Quantitative Genetic Analysis of Multivariate Evolution, Applied to Brain: Body

Size Allometry. Evolution 33:402-416.

Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters.

Evolution 37:1210-1226.

Le Rouzic, A., J. Álvarez-Castro, and T. F. Hansen. 2013. The Evolution of Canalization and

Evolvability in Stable and Fluctuating Environments. Evol Biol Evolutionary Biology 40:317-

340.

Ledón-Rettig, C. C., D. W. Pfennig, A. J. Chunco, and I. Dworkin. 2014. Cryptic Genetic Variation

in Natural Populations: A Predictive Framework. Integrative and Comparative Biology

54:783-793.

Levy, S. F., and M. L. Siegal. 2008. Network Hubs Buffer Environmental Variation in

Saccharomyces cerevisiae. PLoS Biology 6:e264.

Marquez, E. J., R. Cabeen, R. P. Woods, and D. Houle. 2012. The Measurement of Local Variation

in Shape. Evolutionary Biology 39:419-439.

Marroig, G., and J. M. Cheverud. 2004. Cranial evolution in sakis (*Pithecia*, Platyrrhini) I:

Interspecific differentiation and allometric patterns. American Journal of Physical

Anthropology 125:266-278.

Marroig, G., D. Melo, A. Porto, H. Sebastião, and G. Garcia. 2011. Selection Response

Decomposition (SRD): A New Tool for Dissecting Differences and Similarities Between

Matrices. Evolutionary Biology 38:225-241.

Martín, F. A., and G. Morata. 2006. Compartments and the control of growth in the Drosophila

wing imaginal disc. Development 133:4421-4426.

Mitteroecker, P., and F. Bookstein. 2009. The ontogenetic trajectory of the phenotypic

covariance matrix, with examples from craniofacial shape in rats and humans. Evolution

63:727-737.

Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. G. O'Hara, G. L. Simpson, P.

Solymos, H. M. H. Stevens, and H. Wagner. 2013. *vegan*: Community Ecology Package.

Paaby, A. B., and M. V. Rockman. 2014. Cryptic genetic variation: evolution's hidden substrate.

Nature Reviews Genetics 15:247-258.

Pavlicev, M., J. Cheverud, and G. Wagner. 2009. Measuring Morphological Integration Using

Eigenvalue Variance. Evolutionary Biology 36:157-170.

Pitchers, W, Pool, JE, Dworkin, I: Altitudinal clinal variation in wing size and shape in African

Drosophila melanogaster: one cline or many? Evolution 2013, 67:438–452

Porto, A., F. de Oliveira, L. Shirai, V. De Conto, and G. Marroig. 2009. The Evolution of

Modularity in the Mammalian Skull I: Morphological Integration Patterns and Magnitudes.

Evolutionary Biology 36:118-135.

Price, G. R. 1970. Selection and covariance. Nature 227:520-521.

Reed, L. K., S. Williams, M. Springston, J. Brown, K. Freeman, C. E. DesRoches, M. B. Sokolowski,

and G. Gibson. 2010. Genotype-by-Diet Interactions Drive Metabolic Phenotype

Variation in *Drosophila melanogaster*. Genetics 185:1009-1019.

Revell, L. J. 2007. The G matrix under fluctuating correlational mutation and selection. Evolution

61:1857-1872.

Robertson, A. 1966. A mathematical model of the culling process in dairy cattle. Animal Science

8:95-108.

Rohlf, F. 2010. tpsDig, digitize landmarks and outlines. 2.16. Department of ecology and

evolution. Stony Brook: State University

Rutherford, S. L., and S. Lindquist. 1998. Hsp90 as a capacitor for morphological evolution.

Nature 396:336-342.

Sokal, R. R., and F. J. Rohlf. 1995. Biometry. W. H. Freeman and company, New York.

Vermeij, G. J. 1973. Adaptation, Versatility, and Evolution. Systematic Zoology 22:466-477.

Waddington, C. H. 1952. Selection of the Genetic Basis for an Acquired Character. Nature

169:625-626.

Weber, K., N. Johnson, D. Champlin, and A. Patty. 2005. Many P-Element Insertions Affect Wing

Shape in *Drosophila melanogaster*. Genetics 169:1461-1475.

Willmore, K. E., C. P. Klingenberg, and B. Hallgrímsson. 2005. The relationship between

fluctuating asymmetry and environmental variance in rhesus macaque skulls. Evolution

59:898-909.

Willmore, K. E., N. M. Young, and J. T. Richtsmeier. 2007. Phenotypic Variability: Its

Components, Measurement and Underlying Developmental Processes. Evolutionary

Biology Evolutionary Biology 34:99-120.

Whitlock, M. C., P. C. Phillips, and K. Fowler. 2002. Persistence of changes in the genetic

covariance matrix after a bottleneck. Evolution 56:1968-1975.

Whitlock, M. C., and K. Fowler. 1999. The Changes in Genetic and Environmental Variance With

Inbreeding in *Drosophila melanogaster*. Genetics 152:345-353.

Zecca, M., and G. Struhl. 2002. Subdivision of the *Drosophila* wing imaginal disc by EGFR-

mediated signaling. Development 129:1357-1368.

Zelditch, M. L., J. Mezey, H. D. Sheets, B. L. Lundrigan, and G. J. Theodore. 2006. Developmental

regulation of skull morphology II: ontogenetic dynamics of covariance. Evolution &

Development 8:46-60.

Ziv, O., R. Finkelstein, Y. Suissa, T. Dinur, G. Deshpande, and O. Gerlitz. 2012. Inverse regulation

of target genes at the brink of the BMP morphogen activity gradient. Journal of Cell

Science 125:5811-5818.

603
<u>Figures</u>

Figure 1: Illustration of the different matrix properties. Ellipses represent 95% confidence interval

of a normal bivariate distribution, and arrows (ellipse axes) represent orthogonal axes of

covariaton. Matrices are presented as bivariate for illustration, but the principles are the same

for multivariate. Ellipses A, B, and C represent a change in only one property each, relative to the

ellipse in the middle. **A**, smaller matrix size (lower total variance); **B**, perpendicular orientation; **C,**

higher eccentricity.

Figure 2: **A,** A representative wing image with fitted splines and the location of landmarks and

semilandmarks along them. **B,** Individual variation in the sample (grey) relative to Sam (black),

mutant means (green), and means for naturally-derived lines (red).

Figure 3: Illustrating the LORY method for interpolating shape changes from Procrustes data of

genotype Bs^{k07909}. **A**, the tessellation scheme for choosing evaluation points. The landmarks and

semi-landmarks (dots) are used as vertices. The centroids of the resulting triangles (crosses) are

the final evaluation points. **B**, shape differences between the genotype mean and Sam inferred

by calculating the Jacobians around each evaluation point, scaled by 100 (crosses and crossed

circles). Bars represent the direction and change of differences between Sam and the sample

mean at the landmarks and semi-landmarks based on the Procrustes data, scaled by 10X.

Figure 4: Mean-shape distance from Sam plotted against the total variance within each strain.

Bars are 95% confidence intervals. Dashed lines represent values for the Sam wild type.

Figure 5: Covariance distance (i.e orientation differences), estimated by Random Skewers,

plotted against mean-shape distance, relative to the Sam wild type.

Figure 6: Covariance space superimposed onto shape space. Both spaces are based on all

pairwise distances. Solid grey lines indicate the deviation between the two spaces for a given

genotype. **A,** All genotypes; **B,** Mutants only, including the Sam wild type. Mutants that affect the

TGF-β pathway are marked by *t*. All other mutants affect the EGFR pathway

Figure 7: The association between eccentricity and total variance within each genotype (**A**), their

mean-shape distance from Sam (**B**) and their covariance distance from Sam (**C**)

638
639 Table 1: Information about the genotypes included in this study
640

Genotype	N	Wild Type	Allele	Gene		Pathway
Sam	175	Sam	Wild type	Wild type		-
10385	81	Sam	k05115	*Epidermal growth factor Receptor*	$Egfr^{k05115}$	Egfr
10413	88	Sam	k07909	*Blistered*	Bs^{k07909}	Egfr
10418	84	Sam	k09530	*Star*	S^{k09530}	Egfr
10462	89	Sam	s3547	*spitz*	spi^{s3547}	Egfr
10514	83	Sam	k02507	*Patched*	Ptc^{k02507}	Hh
12772	90	Sam	BG01610	*scribbler*	$sbb^{BG01610}$	TGF-β
14321	89	Sam	kg07115	*rhomboid*	$rho^{kg07115}$	Egfr
14403	76	Sam	kg01923	*thickveins*	$tkv^{kg01923}$	TGF-β
2513	86	Sam	W11	*argos*	aos^{W11}	Egfr
3045	92	Sam	md653	*Optometer blind/Bifid*	Omb^{md653}	TGF-β
6342	92	Sam	225-3	*crossveinless-2*	$cv\text{-}2^{225\text{-}3}$	TGF-β
6372	82	Sam	mip-w[+]	*GTPase activating protein1*	$Gap1^{mip\text{-}w[+]}$	Egfr
MA10	63	MA	WT	WT	-	-
MA12	62	MA	WT	WT	-	-
MA44	52	MA	WT	WT	-	-
MA53	55	MA	WT	WT	-	-
MA61	56	MA	WT	WT	-	-
MA70	57	MA	WT	WT	-	-
MA71	55	MA	WT	WT	-	-
NC2	54	NC	WT	WT	-	-
NC30	51	NC	WT	WT	-	-
NC53	58	NC	WT	WT	-	-
NC9	51	NC	WT	WT	-	-
NC94	61	NC	WT	WT	-	-

641
642
643

643 Table 2: Matrix properties of all genotypes, and their distance from Sam in terms of matrix orientation
644 (cov.dis) and mean shape (shape.dis). All measures are based on Jacobians, using EBS, and are therefore
645 proportional to the mean configuration of that genotype.

Genotype	Total variance	rSDE	Eccentricity[a]	cov.dis[b]	shape.dis
Sam (wild type)	0.13 (0.12,0.15)	0.28 (0.26,0.29)	0.23 (0.19,0.26)	0	0
aos^{W11}	0.12 (0.11,0.15)	*0.35 (0.29,0.41)	*0.34 (0.25,0.42)	*0.54 (0.53,0.55)	*0.25 (0.22,0.29)
Omb^{md653}	*0.18 (0.16,0.21)	0.29 (0.26,0.29)	0.21 (0.18,0.23)	*0.68 (0.67,0.68)	*1.23 (1.19,1.27)
$cv\text{-}2^{225\text{-}3}$	0.12 (0.1,0.14)	0.29 (0.28,0.3)	0.23 (0.19,0.26)	*0.44 (0.44,0.45)	*0.22 (0.18,0.25)
$Gap1^{mip\text{-}w[+]}$	0.12 (0.11,0.15)	*0.31 (0.28,0.32)	0.24 (0.19,0.3)	*0.65 (0.64,0.66)	*0.43 (0.38,0.46)
$Egfr^{k05115}$	0.12 (0.11,0.14)	*0.31 (0.28,0.31)	0.25 (0.21,0.28)	*0.52 (0.51,0.53)	*0.23 (0.2,0.26)
Bs^{k07909}	0.13 (0.12,0.15)	*0.30 (0.27,0.31)	0.26 (0.2,0.3)	*0.56 (0.55,0.57)	*0.63 (0.6,0.65)
S^{k09530}	*0.10 (0.09,0.12)	0.29 (0.27,0.3)	0.23 (0.19,0.27)	*0.42 (0.41,0.43)	*0.13 (0.1,0.14)
spi^{s3547}	*0.11 (0.10,0.12)	0.29 (0.27,0.29)	0.24 (0.19,0.28)	*0.50 (0.5,0.51)	*0.38 (0.34,0.41)
Ptc^{k02507}	0.12 (0.11,0.14)	0.29 (0.27,0.29)	0.20 (0.18,0.21)	*0.55 (0.54,0.56)	*0.53 (0.5,0.56)
$sbb^{BG01610}$	0.13 (0.11,0.15)	*0.30 (0.28,0.31)	0.25 (0.19,0.3)	*0.52 (0.52,0.53)	*0.37 (0.34,0.4)
$rho^{kg07115}$	*0.10 (0.1,0.12)	0.29 (0.27,0.29)	0.22 (0.18,0.25)	*0.39 (0.39,0.4)	*0.17 (0.14,0.19)
$tkv^{kg01923}$	*0.10 (0.09,0.12)	*0.30 (0.28,0.3)	0.23 (0.19,0.25)	*0.60 (0.59,0.61)	*0.3 (0.27,0.33)
MA10	0.15 (0.13,0.18)	*0.34 (0.3,0.34)	*0.28 (0.23,0.33)	*0.63 (0.62,0.64)	*0.98 (0.94,1.02)
MA12	0.13 (0.12,0.15)	*0.31 (0.3,0.32)	0.26 (0.21,0.28)	*0.55 (0.54,0.56)	*0.92 (0.89,0.96)
MA44	0.15 (0.13,0.18)	*0.34 (0.3,0.36)	*0.28 (0.21,0.37)	*0.58 (0.57,0.59)	*0.82 (0.79,0.85)
MA53	*0.17 (0.15,0.21)	*0.32 (0.3,0.33)	*0.27 (0.21,0.33)	*0.60 (0.59,0.61)	*1.01 (0.96,1.05)
MA61	0.13 (0.12,0.16)	*0.35 (0.32,0.36)	*0.28 (0.23,0.35)	*0.67 (0.66,0.68)	*1.09 (1.05,1.12)
MA70	0.12 (0.1,0.14)	*0.30 (0.29,0.30)	0.21 (0.17,0.22)	*0.71 (0.7,0.73)	*1.02 (0.99,1.06)
MA71	*0.18 (0.15,0.22)	*0.30 (0.29,0.30)	0.21 (0.19,0.22)	*0.66 (0.65,0.67)	*1.33 (1.28,1.38)
NC2	0.12 (0.11,0.14)	0.29 (0.28,0.28)	0.20 (0.16,0.23)	*0.83 (0.82,0.84)	*1.2 (1.16,1.24)
NC30	0.13 (0.11,0.16)	*0.30 (0.30,0.30)	0.20 (0.19,0.20)	*0.68 (0.67,0.69)	*0.96 (0.93,0.99)
NC53	0.13 (0.11,0.15)	*0.31 (0.29,0.31)	0.25 (0.19,0.29)	*0.61 (0.6,0.62)	*1.12 (1.08,1.15)
NC9	0.15 (0.13,0.19)	*0.40 (0.33,0.47)	*0.38 (0.28,0.49)	*0.70 (0.7,0.72)	*0.58 (0.53,0.63)
NC94	*0.21 (0.19,0.24)	*0.45 (0.40,0.49)	*0.43 (0.34,0.49)	*1.02 (1.01,1.03)	*1.1 (1.04,1.15)

646 * Falls outside the confidence interval of Sam
647 [a] Measured as the ratio between the largest eigenvalue and the total variance.
648 [b] Difference in orientation based on Random Skewers.

649
650
651
652 Table 3: Disparity of shape and covariance based on their joint PCoA space (see Fig. 6). Covariance space
653 is superimposed onto shape space for comparability. Calculations include all 24 PCoA dimensions that
654 resulted in eigenvalues larger than zero. Numbers in parentheses indicate the equivalent calculations
655 without strain 3045.

	Total	Mutants	Naturally-derived
Shape	0.042 (0.038)	0.021 (0.011)	0.029
Covariance	0.019 (0.019)	0.012 (0.011)	0.023

Figure1: Illustration of the different matrix properties. Ellipses represent 95% confidence interval of a normal bivariate distribution, and arrows (ellipse axes) represent orthogonal axes of covariaton. Matrices are presented as bivariate for illustration, but the principles are the same for multivariate. Ellipses A, B, and C represent a change in only one property each, relative to the ellipse in the middle. A, smaller matrix size (lower total variance); B, perpendicular orientation; C, higher eccentricity.

658
659

A

B

Figure 2: **A.** A representative wing image with fitted splines and the location of landmarks and semilandmarks along them. **B.** Individual variation in the sample (grey) relative to Sam (black), mutant means (orange), and means for naturally-derived lines (purple).

660
661

661
662

Figure 3: Illustrating the LORY method for interpolating shape changes from Procrustes data of genotype *Bs*^*k07909*. **A**, the tessellation scheme for choosing evaluation points. The landmarks and semi-landmarks (dots) are used as vertices. The centroids of the resulting triangles (crosses) are the final evaluation points. **B**, shape differences between the genotype mean and Sam inferred by calculating the Jacobians around each evaluation point, scaled by 100 (crosses and crossed circles). Bars represent the direction and change of differences between Sam and the sample mean at the landmarks and semi-landmarks based on the Procrustes data, scaled by 10X.

663
664

Figure 4: Mean-shape distance from Sam plotted against the total variance within each strain. Bars are 95% confidence intervals. Dashed lines represent values for the Sam wild type.

665
666

Figure 5: Covariance distance (i.e, difference in orientation), estimated by Random Skewers, plotted against mean-shape distance, relative to the Sam wild type. Bars are 95% confidence intervals.

Figure 6: Covariance space superimposed onto shape space. Both spaces are based on all pairwise distances. Solid grey lines indicate the deviation between the two spaces for a given genotype. **A,** All genotypes; **B,** Mutants only, including the Sam wild type. Mutants that affect the TGF-β pathway are marked by *t*. All other mutants affect the EGFR pathway

A

B

C

Figure 7: The association between eccentricity and total variance within each genotype (**A**), their mean-shape distance from Sam (**B**) and their covariance distance from Sam (**C**)