Estimating the effect of competition on trait evolution using maximum likelihood inference

Jonathan Drury[1]*, Julien Clavel[1], Marc Manceau[1], & Hélène Morlon[1]

[1]*Institut de Biologie de l'Ecole Normale Supérieure (IBENS), CNRS, Inserm, Ecole Normale*

Supérieure, PSL Research University, F-75005 Paris, France

*Correspondence: drury@biologie.ens.fr, +33 (7) 83 11 43 21, *IBENS, 46 rue d'Ulm, F-75005*

Paris, France

Running head: INTERSPECIFIC COMPETITION & TRAIT EVOLUTION

Abstract

Many classical ecological and evolutionary theoretical frameworks posit that competition

between species is an important selective force. For example, in adaptive radiations, resource

competition between evolving lineages plays a role in driving phenotypic diversification and

exploration of novel ecological space. Nevertheless, current models of trait evolution fit to

phylogenies and comparative datasets are not designed for incorporating the effect of

competition. The most advanced models in this direction are diversity-dependent models where

evolutionary rates depend on lineage diversity. However, these models still treat changes in traits

in one branch as independent of the value of traits on other branches, thus ignoring the effect of

species similarity on trait evolution. Here, we consider a model where the evolutionary dynamics

of traits involved in interspecific interactions are influenced by species similarity in trait values

and where we can specify which lineages are in sympatry. We develop a maximum-likelihood

based approach to fit this model to combined phylogenetic and phenotypic data. Using

simulations, we demonstrate that the approach accurately estimates the simulated parameter

values across a broad range of parameter space. Additionally, we develop tools for specifying the

biogeographic context in which trait evolution occurs. In order to compare models, we also apply

these biogeographic methods to specify which lineages interact sympatrically for two diversity-

dependent models. Finally, we fit these various models to morphological data from a classical

adaptive radiation (Greater Antillean *Anolis* lizards). We show that models that account for

competition and geography perform better than other models. The matching competition model

is an important new tool for studying the influence of interspecific interactions, in particular

competition, on phenotypic evolution. More generally, it constitutes a step toward a better

integration of interspecific interactions in many ecological and evolutionary processes.

Keywords: interspecific competition, trait evolution, phylogenetic comparative methods,

adaptive radiation, community phylogenetics, *Anolis*, maximum likelihood

Interactions between species can be strong selective forces. Indeed, many classical

evolutionary theories assume that interspecific competition has large impacts on fitness.

Character displacement theory (Brown and Wilson 1956; Grant 1972; Pfennig and Pfennig

2009), for example, posits that interactions between species, whether in ecological or social

contexts, drive adaptive changes in phenotypes. Similarly, adaptive radiation theory (Schluter

2000) has been a popular focus of investigators interested in explaining the rapid evolution of

phenotypic disparity (Grant and Grant 2002; Losos 2009; Mahler et al. 2013; Weir and Mursleen

2013), and competitive interactions between species in a diversifying clade are a fundamental

component of adaptive radiations (Schluter 2000; Losos and Ricklefs 2009; Grant and Grant

2011).

Additionally, social interactions between species, whether in reproductive (Gröning and

Hochkirch 2008; Pfennig and Pfennig 2009) or agonistic (Grether et al. 2009, 2013) contexts, are

important drivers of changes in signal traits used in social interactions. Several evolutionary

hypotheses predict that geographical overlap with closely related taxa should drive divergence in

traits used to distinguish between conspecifics and heterospecifics (e.g., traits involved in mate

recognition; Wallace 1889; Fisher 1930; Dobzhansky 1940; Mayr 1963; Gröning and Hochkirch

2008; Ord and Stamps 2009; Ord et al. 2011). Moreover, biologists interested in speciation have

often argued that interspecific competitive interactions are important drivers of divergence

between lineages that ultimately leads to reproductive isolation. Reinforcement, or selection

against hybridization (Dobzhansky 1937, 1940), for example, is often thought to be an important

phase of speciation (Grant 1999; Coyne and Orr 2004; Rundle and Nosil 2005; Pfennig and

Pfennig 2009).

In addition to the importance of interspecific competition in driving phenotypic

divergence between species, competitive interactions are also central to many theories of

community assembly, which posit that species with similar ecologies exclude each other from

the community (Elton 1946). In spite of the importance of interspecific competition to these key

ecological and evolutionary theories, the role of competition in driving adaptive divergence and

species exclusion from ecological communities has been historically difficult to measure (Losos

2009), because both trait divergence and species exclusion resulting from competition between

lineages during their evolutionary history has the effect of eliminating competition between those

lineages at the present. Community phylogeneticists have aimed to solve part of this conundrum

by analyzing the phylogenetic structure of local communities: assuming that phylogenetic

similarity between two species is a good proxy for their ecological similarity, competitive

interactions are considered to have been more important in shaping communities comprised of

phylogenetically (and therefore ecologically) distant species (Webb et al. 2002; Cavender-Bares

et al. 2009). However, there is an intrinsic contradiction in this reasoning, because using

phylogenetic similarity as a proxy for ecological similarity implicitly (or explicitly) assumes that

traits evolved under a Brownian model of trait evolution, meaning that species interactions had

no effect on trait divergence (Kraft et al. 2007; Cavender-Bares et al. 2009; Mouquet et al. 2012;

Pennell and Harmon 2013).

More generally, and despite the preponderance of classical evolutionary processes that

assume that interspecific interactions have important fitness consequences, existing phylogenetic

models treat trait evolution within a lineage as independent from traits in other lineages. For

example, in the commonly used Brownian motion and Ornstein-Uhlenbeck models of trait

evolution (Cavalli-Sforza & Edwards 1967, Felsenstein 1988, Hansen and Martins 1996), once

an ancestor splits into two daughter lineages, the trait values in those daughter lineages do not

depend on the trait values of sister taxa. Some investigators have indirectly incorporated the

influence of interspecific interactions by fitting models where evolutionary rates at a given time

depend on the diversity of lineages at that time (e.g., the "diversity-dependent" models of Mahler

et al. 2010, Weir and Mursleen 2013). While these models capture some parts of the interspecific

processes of central importance to evolutionary theory, such as the influence of ecological

opportunity, they do not explicitly account for trait-driven interactions between lineages, as trait

values in one lineage do not vary directly as a function of trait values in other evolving lineages.

Recently, Nuismer and Harmon (2015) proposed a model where the evolution of a

species' trait depends on other species' traits. In particular, they consider a model, which they

refer to as the model of phenotype matching, where the probability that an encounter between

two individuals has fitness consequences declines as the phenotypes of the individuals become

more dissimilar. The consequence of the encounter on fitness can be either negative if the

interaction is competitive, resulting in character divergence (matching competition, e.g. resource

competition), or positive if the interaction is mutualistic, resulting in character convergence

(matching mutualism, e.g. Müllerian mimicry). Applying Lande's formula (Lande 1976) and

given a number of simplifying assumptions—importantly that all lineages evolve in sympatry

and that competitive interactions are approximately equivalent across sympatric taxa— this

model yields a simple prediction for the evolution of a population's mean phenotype.

Here, we develop inference tools for fitting a simple version of the matching competition

model (i.e., the phenotype matching model of Nuismer and Harmon incorporating competitive

interactions between lineages) to combined phylogenetic and trait data. We begin by showing

how to compute likelihoods associated with this model. Next, we use simulations to explore the

statistical properties of maximum likelihood estimation of the matching competition model

(parameter estimation as well as model identifiability). While the inclusion of interactions

between lineages is an important contribution to quantitative models of trait evolution, applying

the matching competition model to an entire clade relies on the assumption that all lineages in

the clade are sympatric. However, this assumption will be violated in most empirical cases, so

we also developed a method for incorporating data on the biogeographical overlap between

species for this model and for the linear and exponential diversity-dependent trait models of Weir

& Mursleen (2013), wherein the evolutionary rate at a given time in a tree varies as a function of

the number of lineages in the reconstructed phylogeny at that time (see also Mahler et al. 2010).

We then fit the model to data from a classical adaptive radiation: Greater Antillean *Anolis*

lizards (Harmon et al. 2003; Losos 2009). Many lines of evidence support the hypothesis that

resource competition is responsible for generating divergence between species in both habitat use

(e.g., Pacala and Roughgarden 1982) and morphology (Schoener 1970; Williams 1972; see

review in Losos 1994). Thus, we can make an *a priori* prediction that model comparison will

uncover a signature of competition in morphological traits that vary with habitat and resource

use. Given the well-resolved molecular phylogeny (Mahler et al. 2010, 2013) and the relatively

simple geographical relationships between species (i.e., many species are restricted to single

islands, Rabosky and Glor 2010; Mahler and Ingram 2014), the Greater Antillean *Anolis* lizards

provide a good test system for exploring the effect of competition on trait evolution using the

matching competition model.

METHODS

Likelihood Estimation of the Matching Competition Model

We consider the evolution of a quantitative trait under the matching competition model of

Nuismer & Harmon (2015) wherein trait divergence between lineages will be favored by

selection. We make the assumption that the outcome of competitive interactions is similar

between all members of an evolving clade rather than sensitive to pairwise phenotypic similarity

(i.e., that α in Eq. 1 of Nuismer and Harmon 2015 is small). This assumption is crucial, as it

ensures that the evolution of a population's mean phenotype is given by a linear model (Eq. S38

in Nuismer and Harmon 2015). Importantly, this implies that the expected distribution of trait

values on a given phylogeny follows a multivariate normal distribution (Manceau *et al.*, in prep),

as is the case for classical models of quantitative trait evolution (Hansen and Martin 1996,

Harmon et al. 2010, Weir and Mursleen 2013). In our current treatment of the model, we remove

stabilizing selection to focus on the effect of competition (see Discussion). Under these two

simplifying assumptions, the mean trait value for lineage *i* after an infinitesimally small time step

dt is given by (Eq. S38 in Nuismer and Harmon 2015 with $\psi = 0$):

$z_i(t + dt) = z_i(t) + S\big(\mu(t) - z_i(t)\big)dt + \sigma dB_i$ (Eq. 1)

where $z_i(t)$ is the mean trait value for lineage *i* at time t, $\mu(t)$ is the mean trait value for the

entire clade at time t, *S* measures the strength of interaction (more intense competitive

interactions are represented by larger negative values), and drift is incorporated as Brownian

motion σdB_i with mean = 0 and variance = $\sigma^2 dt$, Note that when $S = 0$ or $n = 1$ (i.e., when a

species is alone), this model reduces to Brownian motion. Under the model specified by Eq. 1, if

a species trait value is greater (or smaller) than the trait value average across species in the clade,

the species' trait will evolve towards even larger (or smaller) trait values. We discuss the

strengths and limitations of this formulation of the matching competition in the Discussion.

Given that the expected distribution of trait values on a phylogeny under the matching

competition model specified in Eq. 1 follows a multivariate normal distribution, it is entirely

described with its expected mean vector (made of terms each equal to the character value at the

root of the tree) and variance-covariance matrix. Nuismer & Harmon (2015) provide the system

of ordinary differential equations describing the evolution of the variance and covariance terms

through time (their Eqs.10b and 10c). These differential equations can be integrated numerically

from the root to the tips of phylogenies to compute expected variance-covariance matrices for a

given set of parameter values and the associated likelihood values given by the multivariate

normal distribution.

Additionally, to relax the assumption that all of the lineages in a clade coexist

sympatrically, we included a term to specify which lineages co-occur at any given time-point in

the phylogeny, which can be inferred, e.g., by biogeographical reconstruction. We define

piecewise constant coexistence matrices $\mathbf{A}$, where $\mathbf{A}_{i,j}$ equals 1 at time t if i and j are sympatric at

that time, and 0 otherwise (Fig. 1). The evolution of the trait value for lineage i is then given by:

$z_i(t + dt) = z_i(t) + S\left(\left(\frac{1}{n_i}\sum_{l=1}^{n}\mathbf{A}_{i,l}\, z_l(t)\right) - z_i(t)\right)dt + \sigma dB_i$ (Eq. 2)

where $n_i = \sum_{j=1}^{n} \mathbf{A}_{ij}$ is the number of lineages interacting with lineage i at time t (equal to the

number n of lineages in the reconstructed phylogeny at time t if all species are sympatric) such

that trait evolution is only influenced by sympatric taxa.

We show (Appendix S1) that the corresponding system of ordinary differential equations

describing the evolution of the variance and covariance terms through time is:

$\dfrac{dv_{i,i}}{dt} = -\dfrac{2S(n_i-1)}{n_i} v_{i,i} + \dfrac{2S}{n_i}\left(\sum_{\substack{l=1 \\ (l\neq i)}}^{n} \mathbf{A}_{i,l}v_{l,i} \right) + \sigma^2$ (Eq. 3a)

$\dfrac{dv_{i,j}}{dt} = -S\left(\dfrac{n_i-1}{n_i} + \dfrac{n_j-1}{n_j} \right) v_{i,j} + \dfrac{S}{n_i}\sum_{\substack{k=1 \\ k\neq i}}^{n} \mathbf{A}_{i,k}v_{k,j} + \dfrac{S}{n_j}\sum_{\substack{l=1 \\ l\neq j}}^{n} \mathbf{A}_{j,l}v_{l,i}$ (Eq. 3b)

where $v_{i,i}$ is the variance for each species i at time t and $v_{i,j}$ is the covariance for each species

pair i,j at time t. Using numerical integration, we solve this system of ordinary differential

equations from the root of the tree to the tips in order to calculate the values of the variance-

covariance matrix expected under the model for a given phylogeny and set of parameter values.

Specifically, Eq. 3a and 3b dictate the evolution of the variance and covariance values along the

branches of the tree; at a given branching event, the variance and covariance values associated to

the two daughter species are simply inherited from those of the ancestral species. With the

expected variance-covariance matrix at present, we calculate the likelihood for the model using

the likelihood function for a multivariate-normal distribution (e.g. Harmon et al. 2010). Then,

using standard optimization algorithms, we identify the maximum likelihood values for the

model parameters. The matching competition model has three free parameters: σ^2, S and the

ancestral state z_0 at the root. As with other models of trait evolution, the maximum likelihood

estimate for the ancestral state is computed through GLS using the estimated variance-covariance

matrix (Grafen 1989; Martins and Hansen 1997).

We used the ode function in the R package deSolve (Soetaert et al. 2010) to perform the

numerical integration of the differential equations using the "lsoda" solver, and the Nelder-Mead

algorithm implemented in the optim function to perform the maximum likelihood optimization.

Codes for these analyses are freely available on github (https://github.com/hmorlon/PANDA)

and included the R package RPANDA (Morlon et al. MS).

Figure 1. Illustration of geography matrices (defined between each node and after each dispersal event inferred, e.g., by stochastic mapping) delineating which lineages interact in sympatry in an imagined phylogeny. These matrices were used to identify potentially interacting lineages for the matching competition and both diversity-dependent models of character evolution (see Eqs. 3-5 in the main text). *Anolis* outline courtesy of Sarah Werning, licensed under Creative Commons.

Incorporating Geography into Diversity-Dependent Models

Using the same geography matrix **A** described above for the matching competition model

(Fig. 1), we modified the diversity-dependent linear and exponential models of Weir & Mursleen

(2013) to incorporate biological realism into the models, because ecological opportunity is only

relevant within rather than between biogeographical regions. The resulting variance-covariance

matrices, **V**, of these models have the elements:

$$\mathbf{V}_{ij} = \sum_{m=2}^{M}(\sigma_0^2 + bn_i)(\max(s_{ij} - t_{m-1}, 0) - \max(s_{ij} - t_m, 0)) \qquad \text{(Eq. 4)}$$

for the diversity-dependent linear model, and

$$\mathbf{V}_{ij} = \sum_{m=2}^{M}(\sigma_0^2 \times e^{rn_i})(\max(s_{ij} - t_{m-1}, 0) - \max(s_{ij} - t_m, 0)) \qquad \text{(Eq. 5)}$$

for the diversity-dependent exponential model, where σ_0^2 is the rate parameter at the root of the

tree, *b* and *r* are the slopes in the linear and exponential models, respectively, s_{ij} is the shared

path length of lineages *i* and *j* from the root of the phylogeny to their common ancestor, n_i is the

number of sympatric lineages (as above) between times t_{m-1} and t_m (where t_1 is 0, the time at the

root, and t_M is the total length of the tree) (Weir & Mursleen 2013). When *b* or *r* = 0, these

models reduce to Brownian motion. For the linear version of the model, we constrained the

maximum likelihood search such that the term $(\sigma_0^2 + bn_i)$ in Eq. 3 ≥ 0 to prevent the model

from having negative evolutionary rates at any t_m.

Simulation-based Analysis of Statistical Properties of the Matching Competition Model

To verify that the matching competition model can be reliably fit to empirical data, we

simulated trait datasets to estimate its statistical properties (i.e., parameter estimation and

identifiability using AICc). For all simulations, we began by first generating 100 pure-birth trees

using TreeSim (Stadler 2014). To determine the influence of the number of tips in a tree, we ran

simulations on trees of size $n = 20$, 50, 100, and 150. We then simulated continuous trait datasets

by applying the matching competition model recursively from the root to the tip of each tree

(Paradis 2012), following Eq. 1, assuming that all lineages evolved in sympatry. For these

simulations, we set $\sigma^2 = 0.05$ and systematically varied S (-1.5, -1, -0.5, -0.1, or 0). Finally, we

fit the matching competition model to these datasets using the ML optimization described above.

To determine the ability of the approach to accurately estimate simulated parameter

values, we first compared estimated parameters to the known parameters used to simulate

datasets under the matching competition model (S and σ^2). We also quantified the robustness of

these estimates in the presence of extinction by estimating parameters for datasets simulated on

birth-death trees; in addition, we compared the robustness of the matching competition model to

extinction to that of the diversity-dependent models. These two latter sets of analyses are

described in detail in the Supplementary Appendix 2.

To assess the ability to correctly identify the matching competition model when it is the

generating model, we compared the fit (measured by AICc, Burnham and Anderson 2002) of this

model to other commonly used trait models on the same data (i.e. data simulated under the

matching competition model). Specifically, we compared the matching competition model to (1)

Brownian motion (BM), (2) Ornstein-Uhlenbeck/single-stationary peak model (OU, Hansen &

Martin 1996), (3) exponential time-dependent (TD_{exp}, i.e., the early burst model, or the ACDC

model with the rate parameter set to be negative, Blomberg et al. 2003; Harmon et al. 2010), (4)

linear time-dependent evolutionary rate (TD$_{lin}$, Weir and Mursleen 2013), (5) linear rate

diversity-dependent (DD$_{lin}$, Mahler et al. 2010; Weir and Mursleen 2013), and (6) exponential

rate diversity-dependent (DD$_{exp}$, Weir and Mursleen 2013). These models were fitted using

geiger (Harmon et al. 2008) when available there (BM, OU, TD$_{exp}$, TD$_{lin}$), or using our own

codes, available in RPANDA (Morlon et al. MS) when they were not available in geiger (DD$_{lin}$,

DD$_{exp}$). With the exception of TD$_{exp}$, which we restricted to have decreasing rates through time

since it has recently been shown that the accelerating rates version of the model is unidentifiable

from OU (Uyeda et al. 2015), we did not restrict the ML search for the parameters in TD$_{lin}$ or DD

models.

We assessed the identifiability of other trait models against the matching competition

model by calculating the fit of this model to datasets simulated under the same trait models

mentioned above. For BM and OU models, we generated datasets from simulations using

parameter values from the appendix of Harmon et al. 2010 scaled to a tree of length 400 (BM, σ^2

= 0.03; OU, σ^2 = 0.3, α = 0.06). For both the linear and exponential versions of the time- and

diversity-dependent models, we simulated datasets with starting rates of σ^2 = 0.6 and ending

rates of σ^2 = 0.01, declining with a slope determined by the model and tree (e.g., for time-

dependent models, the slope is a function of the total height of the tree; for the TD$_{exp}$ model,

these parameters result in a total of 5.9 half-lives elapsing from the root to the tip of the tree,

Slater and Pennell 2014). In another set of simulations, we fixed the tree size at 100 tips and

varied parameter values to determine the effect of parameter values on identifiability (see

Results). As above, we calculated the AICc for all models for each simulated dataset.

Finally, to understand how removing stabilizing selection from the likelihood of the

matching competition model affects our inference in the presence of stabilizing selection, we

simulated datasets with both matching competition and stabilizing selection on 100 tip trees,

across a range of parameter space (S = -1, -0.5, and 0, α = 0.05, 0.5, and 5, holding σ^2 at 0.05).

We fit BM, OU, and matching competition models to these simulated datasets. All simulations

were performed using our own codes, available in RPANDA (Morlon 2014).

Fitting the Matching Competition Model of Trait Evolution to Caribbean Anolis *Lizards*

To determine whether the matching competition model is favored over models that ignore

interspecific interactions in an empirical system where competition likely influenced character

evolution, we fit the matching competition model to a morphological dataset of adult males from

100 species of Greater Antillean *Anolis* lizards and the time calibrated, maximum clade

credibility tree calculated from a Bayesian sample of molecular phylogenies (Mahler et al. 2010,

2013; Mahler and Ingram 2014). We included the first four size-corrected phylogenetic principal

components from a set of 11 morphological measurements, collectively accounting for 93% of

the cumulative variance explained (see details in Mahler et al. 2013). Each of these axes is

readily interpretable as a suite of specific morphological characters (see Discussion), and

together, the shape axes quantified by these principal components describe the morphological

variation associated with differences between classical ecomorphs in Caribbean anoles (Williams

1972). In addition to the matching competition model, we fit the six previously mentioned

models (BM, OU, TD_{exp}, TD_{lin}, DD_{exp}, and DD_{lin}) separately to each phylogenetic PC axis in the

Anolis dataset.

For the matching competition model and diversity-dependent models, to determine the

influence of uncertainty in designating clades as sympatric and allopatric, we fit the model for

each trait using 101 sets of geography matrices (i.e., **A** in Eq. 1b, 2, & 3, see Fig. 1): one where

all lineages were set as sympatric, and the remaining 100 with biogeographical reconstructions

from the output of the make.simmap function in phytools (Revell 2012). To simplify the ML

optimization, we restricted S to take negative values while fitting the matching competition

model including the biogeographical relationships among taxa.

Figure 2. Parameter estimation under the matching competition model. As tree size increases and/or the magnitude of competition increases (i.e., the S parameter in the matching competition model becomes more negative), so does the accuracy of ML parameter estimates of (A) S ($n = 100$ for each tree size and S value combination; red horizontal lines indicate the simulated S value) and (B) σ^2 ($n = 500$ for each tree size; red horizontal lines indicate the simulated value). In a small number of cases (7/2000), the ML estimate for σ^2 was unusually large (> 0.25), and we removed these rare cases for plotting. The numbers below the violin plots in (B) show the number of outliers removed for each tree size.

RESULTS

Statistical Properties of the Matching Competition Model

Across a range of *S* values, maximum likelihood optimization returns reliable estimates

of parameter values for the matching competition model (Fig. 2). As the number of tips

increases, so does the reliability of maximum likelihood parameter values (Fig. 2). Parameter

estimates remain reliable in the presence of extinction, unless the extinction fraction is very large

(i.e., ≥ 0.6; Supplementary Appendix 2). When datasets are simulated under the matching

competition model, model selection using AICc generally picks the matching competition model

as the best model (Figs. 3, S1); the strength of this discrimination depends on both the *S* value

used to simulate the data and the size of the tree (Figs. 3, S1).

Figure 3. AICc support for datasets simulated under the matching competition (MC) model increases with tree size and with increasing levels of competition (i.e., increasingly negative *S* values). The dotted line denotes 10%.

Simulating datasets under BM, OU, DD_{exp}, and DD_{lin} generating models, we found that in

most scenarios, and in most parameter space, these models are distinguishable from the matching

competition model (Fig. 4a,b,e,f, Fig. S2). As with the matching competition model, the ability

to distinguish between models using AICc generally increases with increasing tree sizes (Fig. 4)

and with increasing magnitude of parameter values (Fig. S2). When character data were

simulated under a TD_{lin} model of evolution, the matching competition and/or the diversity-

dependent models tended to have lower AICc values than the TD_{lin} model, especially among

smaller trees (Figure 4d). For data generated under a TD_{exp} model, model selection always

favored the matching competition model over the TD_{exp} model (Fig. 4c).

As the strength of stabilizing selection increases relative to the strength of competition

(i.e., α as increases relative to S) AICc model selection shifts from favoring the matching

competition model (under large S, small α scenarios) to favoring the OU model (under small S,

large α scenarios) (Fig. S3). Likewise, maximum likelihood increasingly underestimates the

value of S as the value of α increases (Fig. S4).

Competition in Greater Antillean Anolis *Lizards*

For the first four phylogenetic principal components describing variation in *Anolis*

morphology, we found that models that incorporate species interactions fit the data better than

models that ignore them (Table 1). PC1, which describes variation in hindlimb/hindtoe length

(Mahler et al. 2013), is fit best by the matching competition model. PC2, which describes

variation in body size (snout vent length) is fit best by the linear diversity-dependent model.

PC3, which describes variation in forelimb/foretoe length, and PC4, which describes variation in

lamellae number are fit with mixed support across the models included, but with models

incorporating species interactions providing the best overall fits.

Figure 4. Identifiability simulation results for the matching competition (MC) model. When the generating model is either (A) BM, (B) OU, (E) DD$_{exp}$ (for larger trees) or (F) DD$_{lin}$, the generating model is largely favored by model selection. However, both (C) TD$_{exp}$ and (D) TD$_{lin}$ (for smaller trees) are erroneously rejected as the generating model. The dotted lines denote 10%.

337   Additionally, for every PC axis, the best-fit models were ones that incorporated the geographic

338   relationships among species in the tree, and these conclusions were robust to uncertainty in

339   ancestral reconstructions of sympatry (Table 1).

Table 1. Comparison of model fits for the first four phylogenetic principal components of a morphological dataset of Greater Antillean anoles. Models run incorporating geography matrices are indicated by "+ GEO", and models with the lowest AICc for each trait are shaded and written in bold text. Parameter values presented follow the nomenclature of Eqs. 2-4 in the main text, and k represents the number of parameters estimated for each model. Note that TD_{exp} is the ACDC model (or the early-burst model when $r < 0$). OU model weights were excluded because the ML estimates of α equaled 0 for all PC axes, and thus the OU model was equivalent to BM. Median (standard error) of parameter estimates, ΔAICc values, and Akaike weights are presented for fits across 100 sampled stochastic maps of *Anolis* biogeography (standard errors are omitted for Akaike weights < 0.05).

341
342
343

Trait	Model	k	σ^2	b	r	S	ln($\mathcal{L}$)	ΔAICc	Akaike weights
pPC1	BM	2	0.0033	—	—	—	-13.68	21.36 (0.19)	<0.01
	OU	3	0.0033	—	—	—	-13.68	--	--
	TDexp	3	0.0324	—	-0.068	—	-5.20	6.51 (0.19)	0.03
	TDlinear	3	0.0113	-0.019	—	—	-4.88	5.89 (0.19)	0.04
	DDexp	3	0.0184	—	-0.028	—	-4.37	4.87 (0.19)	0.06 (0.004)
	DDexp + GEO	3	0.0087 (1.48E-5)	—	-0.043 (7.29E-5)	—	-8.00 (0.050)	12.05 (0.19)	<0.01
	DDlin	3	0.0089	-0.00008	—	—	-4.89	5.91 (0.19)	0.04
	DDlin + GEO	3	0.0060 (6.25E-6)	-0.00011 (1.24E-7)	—	—	-8.23 (0.034)	12.49 (0.18)	<0.01
	MC$_{sym}$	3	0.0010	—	—	-0.037	-3.67	2.96 (0.19)	0.16 (0.01)
	MC + GEO	**3**	**0.0010 (6.75E-6)**	**—**	**—**	**-0.038 (0.00017)**	**-1.94 (0.10)**	**0 (0.031)**	**0.68 (0.02)**
pPC2	BM	2	0.0027	—	—	—	-4.69	9.64 (0.036)	0.01
	OU	3	0.0027	—	—	—	-4.69	--	--
	TDexp	3	0.0046	—	-0.014	—	-4.30	10.99 (0.036)	<0.01
	TDlinear	3	0.0047	-0.011	—	—	-4.23	10.85 (0.036)	<0.01
	DDexp	3	0.0041	—	-0.006	—	-4.27	10.94 (0.036)	<0.01
	DDexp + GEO	3	0.0068 (1.36E-5)	—	-0.039 (9.72E-5)	—	0.51 (0.024)	1.37 (0.015)	0.33 (0.002)
	DDlin	3	0.0042	-0.00002	—	—	-4.21	10.82 (0.036)	<0.01
	DDlin + GEO	**3**	**0.0054 (4.24E-6)**	**-0.00010 (9.17E-8)**	**—**	**—**	**1.20 (0.017)**	**0 (0)**	**0.64 (0.001)**
	MC$_{sym}$	3	0.0021	—	—	-9.9e-3	-3.95	9.79 (0.036)	<0.01
	MC + GEO	3	0.0018 (2.44E-6)	—	—	-0.015 (4.67E-5)	-2.94 (0.010)	8.30 (0.047)	0.01
pPC3	BM	2	0.0010	—	—	—	45.57	2.56 (0.021)	0.09 (0.0003)
	OU	3	0.0010	—	—	—	45.57	--	--
	TDexp	3	0.0020	—	-0.019	—	46.30	3.22 (0.021)	0.06 (0.0002)
	TDlinear	3	0.0019	-0.013	—	—	46.41	3.02 (0.021)	0.07 (0.0003)
	DDexp	3	0.0017	—	-0.008	—	46.40	3.02 (0.021)	0.07 (0.0003)
	DDexp + GEO	3	0.0015 (1.48E-6)	—	-0.017 (3.72E-5)	—	46.79 (0.006)	2.24 (0.024)	0.10 (0.0006)
	DDlin	3	0.0017	-0.000009	—	—	46.46	2.90 (0.021)	0.08 (0.0003)
	DDlin + GEO	3	0.0014 (8.76E-7)	-0.000016 (2.96E-8)	—	—	46.68 (0.005)	2.44 (0.023)	0.09 (0.0005)
	MC$_{sym}$	3	0.0007	—	—	-0.012	46.75	2.33 (0.021)	0.10 (0.0004)
	MC + GEO	**3**	**0.0006 (6.15E-7)**	**—**	**—**	**-0.017 (3.38E-5)**	**47.91 (0.011)**	**0 (0)**	**0.32 (0.002)**
pPC4	BM	2	0.0006	—	—	—	69.07	2.50 (0.016)	0.06 (0.0002)
	OU	3	0.0006	—	—	—	69.07	--	--
	TDexp	3	0.0015	—	-0.025	—	70.55	1.66 (0.016)	0.09 (0.0003)
	TDlinear	3	0.0012	-0.013	—	—	70.45	1.86 (0.016)	0.08 (0.0003)
	DDexp	3	0.0012	—	-0.010	—	70.52	1.73 (0.016)	0.08 (0.0003)
	DDexp + GEO	3	0.0010 (1.18E-6)	—	-0.020 (4.38E-5)	—	71.28 (0.011)	0.13 (0.020)	0.18 (0.001)
	DDlin	3	0.0011	-0.000006	—	—	70.39	1.99 (0.016)	0.07 (0.0002)
	DDlin + GEO	3	0.0009 (5.77E-7)	-0.000009 (1.98E-8)	—	—	70.78 (0.008)	1.12 (0.016)	0.11 (0.0006)
	MC	3	0.0004	—	—	-0.015	71.1	0.57 (0.016)	0.15 (0.0005)
	MC + GEO	**3**	**0.0004 (4.21E-7)**	**—**	**—**	**-0.016 (3.56E-5)**	**71.34 (0.009)**	**0 (0.012)**	**0.19 (0.001)**

DISCUSSION

The inference methods we present here represent an important new addition to the

comparative trait analysis toolkit. Whereas previous models had not accounted for the influence

of trait values in other lineages on character evolution, the matching competition model takes

these into account. Furthermore, extending both the matching competition model and two

diversity-dependent trait evolution models to incorporate geographic networks of sympatry

further extends the utility and biological realism of these models.

We found that the matching competition model has increasing AICc support and

accuracy of parameter estimation with increasing tree sizes and competition strength. We also

found that, for most of the generating models we tested, AICc-based model selection does not

tend to erroneously select the matching competition model (i.e., these models are identifiable

from the matching competition model). As with all other models, the statistical properties of the

matching competition model will depend on the size and shape of a particular phylogeny as well

as specific model parameter values. Future investigators can employ other approaches, such as

phylogenetic Monte Carlo and posterior predictive simulations directly on their empirical trees

(Boettiger et al. 2012, Slater & Pennell 2014), to assess the confidence they can have in their

results.

We did, however, find that data generated under time-dependent models were often fit

better by models that incorporate interspecific interactions (i.e., density-dependent and matching

competition models) (Fig. 4c,d). This was especially true for the TD_{exp} model, often referred to

as the early-burst model—the matching competition model nearly always fit data generated

under the TD_{exp} model better than the TD_{exp} model (Fig. 4c). We do not view this as a major

limitation of the model for two reasons. First, the TD_{exp} model is known to be statistically

difficult to estimate on neontological data alone (Harmon et al. 2010; Slater et al. 2012a; Slater

and Pennell 2014). Secondly, and more importantly, time-dependent models are not process-

based models, but rather incorporate time since the root of a tree as a proxy for ecological

opportunity or available niche space (Harmon et al. 2010; Mahler et al. 2010; Slater 2015). The

matching competition and density-dependent models explicitly account for the interspecific

competitive interactions that time-dependent models purport to model, thus we argue that these

process-based models are more biologically meaningful than time-dependent models (Moen and

Morlon 2014).

We did not incorporate stabilizing selection in our model. Preliminary analyses suggested

that S and α are not identifiable, as competition and stabilizing selection operate in opposite

directions. As a result, when trait data are simulated with simultaneous stabilizing selection and

matching competition, the strength of competition is underestimated. In addition, which model is

chosen by model selection depends on the ratio of the strength of attraction toward an optimum

to the strength of competition, with Brownian model being selected at equal strengths (Figs. S3,

S4). Given that many traits involved in competitive interactions are also likely to have been

subject to stabilizing selection (i.e., extreme trait values eventually become targeted by negative

selection), statistical inference under the matching competition model without stabilizing

selection is likely to underestimate the true effect of competition on trait evolution. Future work

aimed at directly incorporating stabilizing selection in the inference tool could provide a more

accurate quantification of the effect of competition, although dealing with the non-identifiability

issue may require incorporating additional data such as fossils.

Because the matching competition model depends on the mean trait values in an evolving

clade, maximum likelihood estimation is robust to extinction, whereas the diversity-dependent

models are less so (Appendix S2, Figs. S5-S8). Nevertheless, given the failure of maximum

likelihood to recover accurate parameter estimates of the matching competition model at high

levels of extinction ($\mu : \lambda \geq 0.6$), we suggest that these models should not be used in clades where

the extinction rate is known to be particularly high. In such cases, it would be preferable to

modify the inference framework presented here to include data from fossil lineages (Slater et al.

2012a) by adapting the ordinary differential equations described in Eq. 3a and 3b for non-

ultrametric trees.

For all of the traits we analyzed, we found that models incorporating both the influence of

other lineages and the specific geographical relationships among lineages were the most strongly

supported models (though less strikingly for PC3 and PC4). Incorporating uncertainty in

biogeographical reconstruction, which we encourage future investigators to do in general,

demonstrated that these conclusions were robust to variation in the designation of allopatry and

sympatry throughout the clade. The matching competition model is favored in the phylogenetic

principal component axis describing variation in relative hindlimb size. Previous research

demonstrates that limb morphology explains between-ecomorph variation in locomotive

capabilities and perch characteristics (Losos 1990, 2009; Irschick et al. 1997), and our results

suggest that the evolutionary dynamics of these traits have been influenced by the evolution of

limb morphology in other sympatric lineages. These results support the assumption that

interspecific interactions resulting from similarity in trait values are important components of

adaptive radiations (Losos 1994, Schluter 2000), a prediction that has been historically difficult

to test (Losos 2009, but see Mahler et al. 2010). In combination with previous research

demonstrating a set of convergent adaptive peaks in morphospace to which lineages are attracted

(Mahler et al. 2013), our results suggest that competition likely played an important role in

driving lineages toward these distinct peaks. Because we expect the presence of selection toward

optima to lead to underestimation of the S parameter in the matching competition model (Figs.

S3, S4), we would have likely detected an even stronger effect of competition in *Anolis* dataset if

we had included stabilizing selection. Recently, Uyeda and colleagues (2015) demonstrated that

the use of principal components can bias inferences of trait evolution. We used BM-based

phylogenetic PC axes here, which should reduce this potential bias (Revell 2009). We recognize

that there is some circularity in assuming BM in order to compute phylogenetic PC axes before

fitting other trait models to these axes; a general solution to address this circularity problem

remains to be found (Uyeda et al. 2015). Uyeda & colleagues suggested that using phylogenetic

PC axes sorts the traits according to specific models. In the Greater Antillean *Anolis* lizards, the

first axes are easily interpretable as specific suite of traits relevant to competitive interactions,

and our results suggest that competition played an important role in shaping the evolution of

these traits.

The linear version of Nuismer & Harmon's (2015) model (Eq. 1) results from making the

simplifying assumption that competitive interactions are approximately equivalent across all

sympatric taxa. We used this version here, since currently available likelihood tools for trait

evolution rely on the multivariate normal distribution, which is to be expected only for this linear

form of the model. The current formulation (Eq. 1) corresponds to a scenario in which the rate of

phenotypic evolution in a lineage gets higher as the lineage deviates from the mean phenotype,

although character displacement theory, for example, posits that selection for divergence should

be the strongest when species are most ecologically similar (Brown and Wilson 1956).

Nevertheless, the developments presented here provide an important new set of tools for

investigating the impact of interspecific interactions on trait evolution, and researchers can

perform posterior simulations to assess the realism of the resulting inference. Future

development of likelihood-free methods, such as Approximate Bayesian Computation (Slater et

al. 2012b; Kutsukake and Innan 2013), may be possible for fitting the version of the model in

which the outcome of competitive interactions depend on distance in trait space.

We imagine that the matching competition model and biogeographical implementations

of diversity-dependent models will play a substantial role in the study of interspecific

competition. For example, by comparing the fits of the matching competition model with other

models that do not include competitive interactions between lineages, biologists can directly test

hypotheses that make predictions about the role of interspecific interactions in driving trait

evolution. In other words, while the effect of competition has been historically difficult to detect

(Losos 2009), it may be detectable in the contemporary distribution of trait values and their

covariance structure (Hansen and Martins 1996; Nuismer and Harmon 2015). The ability to

consider trait distributions among species that arise from a model explicitly accounting for the

effect of species interactions on trait divergence is also an important step toward a more coherent

integration of macroevolutionary models of phenotypic evolution in community ecology.

There are many possible extensions of the tools developed in this paper. In the future,

empirical applications of the model can be implemented with more complex geography matrices

that are more realistic for mainland taxa (e.g., using ancestral biogeographical reconstruction,

Ronquist and Sanmartín 2011; Landis et al. 2013), and can also specify degrees of sympatric

overlap (i.e., syntopy). Additionally, the current version of the model is rather computationally

expensive with larger trees (with 100 or more tips). Further work developing an analytical

solution to the model may greatly speed up the likelihood calculation and permit the inclusion of

stabilizing selection.

The current form of the model assumes that the degree of competition is equal for all

interacting lineages. Future modifications of the model, such as applications of stepwise AICc

algorithms (Alfaro et al. 2009; Thomas and Freckleton 2012; Mahler et al. 2013) or reversible-

jump Markov Chain Monte Carlo (Pagel and Meade 2006; Eastman et al. 2011; Rabosky 2014;

Uyeda and Harmon 2014), may be useful to either identify more intensely competing lineages or

test specific hypotheses about the strength of competition between specific taxa. Improvements

could also be made on the formulation itself of the evolution of a species' trait as a response to

the phenotypic landscape in which the species occurs. Moreover, a great array of extensions will

come from modeling species interactions not only within clades, but also among interacting

clades, as in the case of coevolution in bipartite mutualistic or antagonistic networks, such as

plant-pollinator or plant-herbivore systems.

Acknowledgements

We thank J. Weir for providing R code for diversity-dependent models and E. Lewitus,

O. Missa, F. Anderson, L. Harmon, and two anonymous reviewers for helpful comments on the

manuscript. This research was funded by the Agence Nationale de la Recherche (grant CHEX-

ECOEVOBIO) and the European Research Council (grant 616419-PANDA) to HM.

References

References

Alfaro M.E., Santini F., Brock C., Alamillo H., Dornburg A., Rabosky D.L., Carnevale G.,
Harmon L.J. 2009. Nine exceptional radiations plus high turnover explain species diversity
in jawed vertebrates. Proc. Natl. Acad. Sci. 106:13410–13414.

Blomberg S.P., Garland T., Ives A.R. 2003. Testing for phylogenetic signal in comparative data:
behavioral traits are more labile. Evolution. 57:717–745.

Boettiger C., Coop G., Ralph P. 2012. Is your phylogeny informative? Measuring the power of
comparative methods. Evolution. 66:2240–2251.

Brown W.L., Wilson E.O. 1956. Character displacement. Syst. Zool. 5:49.

Cavender-Bares J., Kozak K.H., Fine P.V.A., Kembel S.W. 2009. The merging of community
ecology and phylogenetic biology. Ecol. Lett. 12:693–715.

Coyne J.A., Orr H.A. 2004. Speciation. Sunderland, MA: Sinauer Associates.

Dobzhansky T. 1937. Genetics and the Origin of Species. New York, NY: Columbia University
Press.

Dobzhansky T. 1940. Speciation as a stage in evolutionary divergence. Am. Nat. 74:312–321.

Eastman J.M., Alfaro M.E., Joyce P., Hipp A.L., Harmon L.J. 2011. A novel comparative
method for identifying shifts in the rate of character evolution on trees. Evolution. 65:3578–
3589.

Elton C. 1946. Competition and the structure of ecological communities. J. Anim. Ecol.:54–68.

Fisher R.A. 1930. The Genetical Theory of Natural Selection. Oxford, UK: Oxford University
Press.

Grafen A. 1989. The phylogenetic regression. Philos. Trans. R. Soc. Lond. B. Biol. Sci.
326:119–157.

Grant P.R., Grant B.R. 2002. Adaptive radiation of Darwin's finches: Recent data help explain
how this famous group of Galápagos birds evolved, although gaps in our understanding
remain. Am. Sci. 90:130–139.

Grant P.R., Grant B.R. 2011. How and Why Species Multiply: the Radiation of Darwin's
Finches. Princeton, NJ: Princeton University Press.

Grant P.R. 1972. Convergent and divergent character displacement. Biol. J. Linn. Soc. 4:39–68.

Grant P.R. 1999. Ecology and Evolution of Darwin's finches. Princeton, NJ: Princeton
University Press.

507 Grether G.F., Anderson C.N., Drury J.P., Kirschel A.N.G., Losin N., Okamoto K., Peiman K.S.
508   2013. The evolutionary consequences of interspecific aggression. Ann. N. Y. Acad. Sci.
509   1289:48–68.

510 Grether G.F., Losin N., Anderson C.N., Okamoto K. 2009. The role of interspecific interference
511   competition in character displacement and the evolution of competitor recognition. Biol.
512   Rev. 84:617–635.

513 Gröning J., Hochkirch A. 2008. Reproductive interference between animal species. Q. Rev. Biol.
514   83:257–282.

515 Hansen T.F., Martins E.P. 1996. Translating between microevolutionary process and
516   macroevolutionary patterns: The correlation structure of interspecific data. Evolution.
517   50:1404–1417.

518 Harmon L.J., Losos J.B., Jonathan Davies T., Gillespie R.G., Gittleman J.L., Bryan Jennings W.,
519   Kozak K.H., McPeek M.A., Moreno-Roark F., Near T.J., Purvis A., Ricklefs R.E., Schluter
520   D., Schulte J.A., Seehausen O., Sidlauskas B.L., Torres-Carvajal O., Weir J.T., Mooers
521   A.T. 2010. Early bursts of body size and shape evolution are rare in comparative data.
522   Evolution. 64:2385–2396.

523 Harmon L.J., Schulte J.A., Larson A., Losos J.B. 2003. Tempo and mode of evolutionary
524   radiation in iguanian lizards. Science 301:961–964.

525 Harmon L.J., Weir J.T., Brock C.D., Glor R.E., Challenger W. 2008. GEIGER: investigating
526   evolutionary radiations. Bioinformatics. 24:129–131.

527 Irschick D.J., Vitt L.J., Zani P.A., Losos J.B. 1997. A comparison of evolutionary radiations in
528   mainland and Caribbean Anolis lizards. Ecology. 78:2191–2203.

529 Kraft N.J.B., Cornwell W.K., Webb C.O., Ackerly D.D. 2007. Trait evolution, community
530   assembly, and the phylogenetic structure of ecological communities. Am. Nat. 170:271–
531   283.

532 Kutsukake N., Innan H. 2013. Simulation-based likelihood approach for evolutionary models of
533   phenotypic traits on phylogeny. Evolution. 67:355–367.

534 Landis M.J., Matzke N.J., Moore B.R., Huelsenbeck J.P. 2013. Bayesian analysis of
535   biogeography when the number of areas is large. Syst. Biol. 62:789–804.

536 Losos J.B., Ricklefs R.E. 2009. Adaptation and diversification on islands. Nature. 457:830–836.

537 Losos J.B. 1990. The evolution of form and function: morphology and locomotor performance in
538   West Indian *Anolis* lizards. 44:1189–1203.

539 Losos J.B. 1994. Integrative approaches to evolutionary ecology: *Anolis* lizards as model

systems. Annu. Rev. Ecol. Syst. 25:467–493.

Losos J.B. 2009. Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles.
Los Angeles, CA: University of California Press.

Mahler D.L., Ingram T., Revell L.J., Losos J.B. 2013. Exceptional convergence on the
macroevolutionary landscape in island lizard radiations. Science 341:292–5.

Mahler D.L., Ingram T. 2014. Phylogenetic comparative methods for studying clade-wide
convergence. In: Garamszegi L., editor. Modern Phylogenetic Comparative Methods and
Their Application in Evolutionary Biology. New York, NY: Springer. p. 425–450.

Mahler D.L., Revell L.J., Glor R.E., Losos J.B. 2010. Ecological opportunity and the rate of
morphological evolution in the diversification of greater Antillean anoles. Evolution.
64:2731–2745.

Martins E.P., Hansen T.F. 1997. Phylogenies and the comparative method: a general approach to
incorporating phylogenetic information into the analysis of interspecific data. Am. Nat.
149:646–667.

Mayr E. 1963. Animal Species and Evolution. Cambridge, MA: Harvard University Press.

Moen D., Morlon H. 2014. Why does diversification slow down? Trends Ecol. Evol. 29:190–
197.

Morlon H, Lewitus E, Condamine FL, Manceau M., Clavel, J., Drury, J. *in revision*. RPANDA:
an R package for macroevolutionary analyses on phylogenetic trees. R package version 1.0.
http://CRAN.R-project.org/package=RPANDA. (in revision at Methods in Ecology &
Evolution)

Mouquet N., Devictor V., Meynard C.N., Munoz F., Bersier L.-F., Chave J., Couteron P.,
Dalecky A., Fontaine C., Gravel D., Hardy O.J., Jabot F., Lavergne S., Leibold M.,
Mouillot D., Münkemüller T., Pavoine S., Prinzing A., Rodrigues A.S.L., Rohr R.P.,
Thébault E., Thuiller W. 2012. Ecophylogenetics: advances and perspectives. Biol. Rev.
87:769–785.

Nuismer S.L., Harmon L.J. 2015. Predicting rates of interspecific interaction from phylogenetic
trees. Ecol. Lett. 18:17–27.

Ord T.J., King L., Young A.R. 2011. Contrasting theory with the empirical data of species
recognition. Evolution. 65:2572–2591.

Ord T.J., Stamps J.A. 2009. Species identity cues in animal communication. Am. Nat. 174:585–
593.

Pacala S., Roughgarden J. 1982. Resource partitioning and interspecific competition in two two-

573  species insular *Anolis* lizard communities. Science 217:444–446.

574 Pagel M., Meade A. 2006. Bayesian analysis of correlated evolution of discrete characters by
575  reversible-jump Markov chain Monte Carlo. Am. Nat. 167:808–825.

576 Paradis E. 2011. Analysis of Phylogenetics and Evolution with R. New York, NY: Springer.

577 Pennell M.W., Harmon L.J. 2013. An integrative view of phylogenetic comparative methods:
578  connections to population genetics, community ecology, and paleobiology. Ann. N. Y.
579  Acad. Sci. 1289:90–105.

580 Pfennig K.S., Pfennig D.W. 2009. Character displacement: Ecological and reproductive
581  responses to a common evolutionary problem. Q. Rev. Biol. 84:253–276.

582 Rabosky D.L., Glor R.E. 2010. Equilibrium speciation dynamics in a model adaptive radiation of
583  island lizards. Proc. Natl. Acad. Sci. 107:22178–22183.

584 Rabosky D.L. 2014. Automatic detection of key innovations, rate shifts, and diversity-
585  dependence on phylogenetic trees. PLoS One. 9:e89543.

586 Revell L.J. 2009. Size-correction and principal components for interspecific comparative studies.
587  Evolution. 63:3258–3268.

588 Revell L.J. 2012. phytools: An R package for phylogenetic comparative biology (and other
589  things). Methods Ecol. Evol. 3:217–223.

590 Ronquist F., Sanmartín I. 2011. Phylogenetic methods in biogeography. Annu. Rev. Ecol. Evol.
591  Syst. 42:441–464.

592 Rundle H.D., Nosil P. 2005. Ecological speciation. Ecol. Lett. 8:336–352.

593 Schluter D. 2000. The Ecology of Adaptive Radiation. Oxford, UK: Oxford University Press.

594 Schoener T.W. 1970. Size patterns in West Indian *Anolis* lizards. II. Correlations with the sizes
595  of particular sympatric species-displacement and convergence. Am. Nat. 104:155–174.

596 Slater G.J., Harmon L.J., Alfaro M.E. 2012a. Integrating fossils with molecular phylogenies
597  improves inference of trait evolution. 66:3931–3944.

598 Slater G.J., Harmon L.J., Wegmann D., Joyce P., Revell L.J., Alfaro M.E. 2012b. Fitting models
599  of continuous trait evolution to incompletely sampled comparative data using approximate
600  Bayesian computation. Evolution. 66:752–762.

601 Slater G.J., Pennell M.W. 2014. Robust regression and posterior predictive simulation increase
602  power to detect early bursts of trait evolution. Syst. Biol. 63:293–308.

603 Soetaert K., Petzoldt T., Setzer R.W. 2010. Solving differential equations in R: package deSolve.
604  J. Stat. Softw. 33:1–25.

Stadler T. 2014. TreeSim: Simulating trees under the birth-death model. R package version

2.1. http://CRAN.R-project.org/package=TreeSim.

Thomas G.H., Freckleton R.P. 2012. MOTMOT: models of trait macroevolution on trees.
Methods Ecol. Evol. 3:145–151.

Uyeda J.C., Caetano D.S., Pennell M.W. 2015. Comparative analysis of principal components
can be misleading. Syst. Biol. 64:677–689.

Uyeda J.C., Harmon L.J. 2014. A novel Bayesian method for inferring and interpreting the
dynamics of adaptive landscapes from phylogenetic comparative data. Syst. Biol. 63:902–
918.

Wallace A.R. 1889. Darwinism. 2007 facs., Cosimo, Inc.

Webb C.O., Ackerly D.D., McPeek M.A., Donoghue M.J. 2002. Phylogenies and community
ecology. Annu. Rev. Ecol. Syst.:475–505.

Weir J.T., Mursleen S. 2013. Diversity-dependent cladogenesis and trait evolution in the
adaptive radiation of the auks (Aves: Alcidae). Evolution. 67:403–416.

Williams E.E. 1972. The origin of faunas. Evolution of lizard congeners in a complex island
fauna: A trial analysis. Evol. Biol. 6:47–89.

622
623

624   **Supplementary Material**

626   Appendix S1 & S2

628   Table S1

630   Figure S1-S8

1 Supplementary Appendix 1

Considering that n lineages are interacting at time t, each trait i evolves following the equation :

$$dz_i(t) = z_i(t + dt) - z_i(t) = S\left(\left(\frac{1}{n_i}\sum_{l=1}^{n} A_{i,l}z_l(t)\right) - z_i(t)\right)dt + \sigma dB_i(t)$$

Where $A_{i,l}$ is equal to 1 if lineages i and l are sympatric, and to 0 otherwise, $n_i = \sum_{l=1}^{n} A_{i,l}$ is the total number of lineages in sympatry with lineage i, and $B_i(t)$ represents standard Brownian motion.

Here, we present the derivation of Equations 3a and 3b from the main text. To make the derivation easier to follow, we drop the dependence on time t, replacing $z_i(t)$ with z_i and $B_i(t)$ with B_i.

First, applying the Itô formula to these stochastic processes gives us :

$$
\begin{aligned}
d\left(z_i z_j\right) = & \ z_i dz_j + z_j dz_i + d < z_i, z_j > \\
= & \ S\left(\left(\frac{1}{n_j}\sum_{l=1}^{n} A_{j,l}z_l z_i\right) - z_j z_i\right)dt + \sigma z_i dB_j \\
+ & \ S\left(\left(\frac{1}{n_i}\sum_{k=1}^{n} A_{i,k}z_k z_j\right) - z_j z_i\right)dt + \sigma z_j dB_i \\
+ & \ \sigma^2 1_{i=j}dt
\end{aligned}
$$

where $1_{i=j}$ equals one if $i = j$ and zero otherwise.

Taking this expectation, it follows that :

$$
\begin{aligned}
\frac{d}{dt}\mathbb{E}\left(z_i z_j\right) = & \ S\left(\left(\frac{1}{n_j}\sum_{l=1}^{n} A_{j,l}\mathbb{E}(z_l z_i)\right) - \mathbb{E}(z_j z_i)\right)dt \\
+ & \ S\left(\left(\frac{1}{n_i}\sum_{k=1}^{n} A_{i,k}\mathbb{E}(z_k z_j)\right) - \mathbb{E}(z_j z_i)\right)dt \\
+ & \ \sigma^2 1_{i=j}dt
\end{aligned}
$$

Moreover, we get :

$$\frac{d}{dt}\mathbb{E}(z_i) = S\left(\left(\frac{1}{n_i}\sum_{l=1}^{n} A_{i,l}\mathbb{E}(z_l)\right) - \mathbb{E}(z_i)\right)dt$$

which leads to :

$$
\begin{aligned}
\frac{d}{dt}\left(\mathbb{E}(z_i)\mathbb{E}(z_j)\right) = & \ \mathbb{E}(z_j)\frac{d}{dt}\mathbb{E}(z_i) + \mathbb{E}(z_i)\frac{d}{dt}\mathbb{E}(z_j) \\
= & \ S\left(\left(\frac{1}{n_i}\sum_{k=1}^{n} A_{i,k}\mathbb{E}(z_k)\mathbb{E}(z_j)\right) - \mathbb{E}(z_i)\mathbb{E}(z_j)\right)dt \\
+ & \ S\left(\left(\frac{1}{n_j}\sum_{l=1}^{n} A_{j,l}\mathbb{E}(z_l)\mathbb{E}(z_i)\right) - \mathbb{E}(z_j)\mathbb{E}(z_i)\right)dt
\end{aligned}
$$

Taking together these different parts gives us the ODE satisfied by all covariances (denoted $v_{i,j} = \text{Cov}(z_i, z_j)$) :

$$
\begin{aligned}
\frac{d}{dt}v_{i,j} = & \ \frac{d}{dt}\left(\mathbb{E}\left(z_i z_j\right) - \mathbb{E}(z_i)\mathbb{E}(z_j)\right) \\
= & \ -2Sv_{i,j} + \frac{S}{n_i}\sum_{k=1}^{n} A_{i,k}v_{k,j} + \frac{S}{n_j}\sum_{l=1}^{n} A_{j,l}v_{l,i} + \sigma^2 1_{i=j}
\end{aligned} \tag{1}
$$

Using these derivations, the variance terms $(i = j)$ are calculated using:

$$\frac{d}{dt}v_{i,i} = -\frac{2S(n_i - 1)}{n_i}v_{i,i} + \frac{2S}{n_i}\sum_{\substack{l=1 \\ l \neq i}}^{n} A_{i,l}v_{l,i} + \sigma^2 \tag{2}$$

The covariance terms $(i \neq j)$ are calculated using:

$$\frac{d}{dt}v_{i,j} = -S\left(\frac{n_i - 1}{n_i} + \frac{n_j - 1}{n_j}\right)v_{i,j} + \frac{S}{n_i}\sum_{\substack{k=1 \\ k \neq i}}^{n} A_{i,k}v_{k,j} + \frac{S}{n_j}\sum_{\substack{l=1 \\ l \neq j}}^{n} A_{j,l}v_{l,i} \tag{3}$$

In the case where lineages i and j are in sympatry, this formula simplifies to:

$$\frac{d}{dt}v_{i,j} = -\frac{2S(n_i - 1)}{n_i}v_{i,j} + \frac{S}{n_i}\left(\sum_{\substack{k=1 \\ k \neq i}}^{n} A_{i,k}v_{k,j} + \sum_{\substack{l=1 \\ l \neq j}}^{n} A_{j,l}v_{l,i}\right) \tag{4}$$

To solve the ODEs for the variance and covariance terms from the root to the tip, we begin by fixing the variance v_0 for the process at the root to 0. At each speciation event, the starting value for both the variance of each of the new lineages and the covariance between the two new lineages is the variance of the immediate ancestor at the time of the speciation event, and the starting value for the covariance between the new lineages and any other persisting lineage is set to the value of the covariance between the persisting lineage and the ancestor of the new lineages at the time of speciation.

Supplementary Appendix 2: **Estimating the effect of extinction on parameter estimation for**

the matching competition and density-dependent models.

Given that the matching competition and diversity-dependent models take into account

the number of interacting lineages, extinction may affect our ability to recover true parameter

values. To estimate the impact of extinction, we simulated 100 trees with 100 extant species,

varying the extinction fraction (μ:λ = 0.2, 0.4, 0.6, and 0.8). As above, we recursively simulated

traits using the matching competition model with $\sigma^2 = 0.05$ and $S = $ -1.5, -1, -0.5, -0.1, or 0, and

the linear and exponential diversity-dependent models with starting rates of $\sigma^2 = 0.6$ and ending

rates of $\sigma^2 = 0.01$. We then estimated the maximum likelihood parameter estimates for the

generating models by fitting the models to the trait values for extant species and the tree with

extinct lineages removed. In the case of the matching competition model, because many

simulated birth-death trees with high extinction rates have substantially older root ages, the

simulated trait datasets for some trees had very large variances. For these biologically unrealistic

trait datasets (i.e., variance in trait values $\geq 1 \times 10^8$), ML does not yield reliable parameter

estimates, so we removed them from further analyses (the sample size of included simulations is

reported in Fig. S5, S6).

Parameter estimates are quite robust to extinction under the matching competition model

(Fig. S5, S6), and much more so than under both diversity-dependent models (Fig. S7, S8).

Under the matching competition model, the maximum likelihood optimization returns reliable

estimates of S and σ^2 values used to simulate datasets on trees with extinct lineages (Fig. S5,

S6), although the estimates become much less reliable with larger extinction fractions, likely

because simulations under the matching competition model were unbounded, resulting in trait

datasets with biologically unrealistic variances. Under both diversity-dependent models, the

magnitude of both the slope and σ^2 parameter values are increasingly underestimated with

increasing extinction fractions (Fig. S7, S8).

662
663
664

665 *Supplementary Figure 1.* As tree size and/or the degree of competition (*S*) increases, model

666 selection becomes more reliable. Comparison of Akaike weights (median & 90% CIs) for NH,

667 BM, OU, and EB models when simulated under various levels of competition (*S* = -1.5, -1, -0.5,

668 -0.1, and 0) for trees with 20, 50, 100, and 150 tips.

670
671
672

Supplementary Figure 2. Identifiability simulation results for the matching competition model as

a function of varying parameter values of the generating models. Parameter values are (A) σ^2 for

BM, (B) α for OU (σ^2 was fixed at 0.3), and the σ^2 value at the root for (C) TD_{exp}, (D) TD_{lin}, (E)

DD_{exp}, and (F) DD_{lin} (for C-F, σ^2 at the present was fixed at 0.01).

Supplementary Figure 3. The effect of incorporating stabilizing selection into trait evolution on

model selection. For datasets generated under the matching competition model with stabilizing

selection included, as the ratio of the strength competition (*S*) to the strength of selection toward

an optimum (*α*) varies, so does the model preferred by model selection.

Supplementary Figure 4. The effect of incorporating stabilizing selection into trait evolution on

parameter estimation. As the strength of stabilizing selection increases (i.e., as α increases),

maximum likelihood under the matching competition model underestimates the true *S* value used

to simulate datasets. Positive *S* values represent selection toward, rather than away, from the

clade mean and are thus expected when the ratio of α to *S* is large. The horizontal red line

represents the simulated *S* value, and the dashed horizontal line represents *S* = 0.

Supplementary Figure 5. Simulation results showing the effect of varying the extinction fraction

on estimation of the *S* parameter for the matching competition model. Red horizontal lines

indicate the simulated *S* values, and numbers above sets of simulations indicate the sample size

of included simulations under those scenarios (see main text for more details).

696
697

698   *Supplementary Figure 6.* Simulation results showing the effect of varying the extinction fraction

699   on estimation of the σ^2 parameter for the matching competition model. Red horizontal lines

700   indicate the simulated σ^2 value (0.05), the numbers below sets of simulations indicate the sample

701   size of included simulations under those scenarios (see main text for more details), and the

702   number in parentheses indicate sample size after σ^2 values > 0.25 were removed.

704
705

Supplementary Figure 7. Simulation results showing the effect of varying the extinction fraction

on slope (top) and σ^2 (bottom) parameters for the exponential diversity-dependent model.

Increasing extinction levels result in increasingly underestimated slope values and σ^2 parameters.

Red horizontal lines indicate the simulated parameter values.

Supplementary Figure 8. Simulation results showing the effect of varying the extinction fraction

on slope (top) and sigma-squared (bottom) parameters for the linear diversity-dependent model.

Increasing extinction levels result in increasingly underestimated slope values and σ^2 parameters.

Red horizontal lines indicate the simulated parameter values.