1 **Estimating the effect of competition on trait evolution using maximum likelihood inference**

2 Jonathan Drury[1]*, Julien Clavel[1], & Hélène Morlon[1]

3 [1]*Département de Biologie, École Normale Supérieure, 46 rue d'Ulm, 75005 Paris, France*

4 *Correspondence: drury@biologie.ens.fr, +33 (7) 83 11 43 21

5 Running head: INTERSPECIFIC COMPETITION & TRAIT EVOLUTION

7 **Abstract**

8   Many classical ecological and evolutionary theoretical frameworks posit that competition

9 between species is an important selective force. For example, in adaptive radiations, resource

10 competition between evolving lineages plays a role in driving phenotypic diversification and

11 exploration of novel ecological space. Nevertheless, current models of trait evolution fit to

12 phylogenies and comparative datasets are not well designed for incorporating the effect of

13 competition. The most advanced models in this direction are diversity-dependent models where

14 evolutionary rates depend on lineage diversity. However, these models still treat changes in traits

15 in one branch as independent of the value of traits on other branches, thus ignoring the effect of

16 species similarity on trait evolution. Here, we consider a model where the evolutionary dynamics

17 of traits involved in interspecific interactions are influenced by species similarity in trait values

18 and where we can specify which lineages are in sympatry. We develop a maximum-likelihood

19 based approach to fit this model to combined phylogenetic and phenotypic data. Using

20 simulations, we demonstrate that the approach accurately estimates the simulated parameter

21 values across a broad range of parameter space. Additionally, we develop tools for specifying the

22 biogeographic context in which trait evolution occurs. In order to compare models, we also apply

23 these biogeographic methods to specify which lineages interact sympatrically for two diversity-

24 dependent models. Finally, we fit these various models to morphological data from a classical

25 adaptive radiation (Greater Antillean *Anolis* lizards). We show that models that account for

26 competition and geography perform better than other models. Given the importance of

27 interspecific interactions, in particular competition, in many ecological and evolutionary

28 processes, this model is an important new tool for using comparative datasets to study

29 phenotypic evolution.

Keywords: interspecific competition, phylogenetic comparative methods, adaptive radiation, trait

evolution, *Anolis*, maximum likelihood

Interactions between species can be strong selective forces. Indeed, many classical

evolutionary theories assume that interspecific competition has large impacts on fitness.

Character displacement theory (Brown and Wilson 1956; Grant 1972; Pfennig and Pfennig

2009), for example, posits that interactions between species, whether in ecological or social

contexts, drive adaptive changes in phenotypes. Similarly, adaptive radiation theory has been a

popular focus of investigators interested in explaining the rapid evolution of phenotypic disparity

(Grant and Grant 2002; Losos 2009; Mahler et al. 2013; Weir and Mursleen 2013), and

competitive interactions between species in a diversifying clade are a fundamental component of

adaptive radiations (Schluter 2000; Losos and Ricklefs 2009; Grant and Grant 2011).

Additionally, social interactions between species, whether in reproductive (Gröning and

Hochkirch 2008; Pfennig and Pfennig 2009) or agonistic (Grether et al. 2009, 2013) contexts, are

important drivers of changes in signal traits used in social interactions. Several evolutionary

hypotheses predict that geographical overlap with closely related taxa should drive divergence in

traits used to distinguish between conspecifics and heterospecifics (Wallace 1889; Fisher 1930;

Dobzhansky 1940; Mayr 1963; Gröning and Hochkirch 2008; Ord and Stamps 2009; Ord et al.

2011). Moreover, biologists interested in speciation have often argued that interspecific

competitive interactions are important drivers of divergence between lineages that ultimately

leads to reproductive isolation. Reinforcement, or selection against hybridization (Dobzhansky

1937, 1940), for example, is often thought to be an important phase of speciation (Grant 1999;

Coyne and Orr 2004; Rundle and Nosil 2005; Pfennig and Pfennig 2009).

Despite the preponderance of classical evolutionary processes that assume that

interspecific interactions have important fitness consequences, existing phylogenetic models of

trait evolution treat trait evolution within a lineage as independent from traits in other lineages.

For example, in the commonly-used Brownian motion and Ornstein-Uhlenbeck models of trait

evolution (Cavalli-Sforza & Edwards 1967, Felsenstein 1988), once an ancestor splits into two

daughter lineages, the trait values in those daughter lineages do not depend on the trait values of

sister taxa. Some investigators have indirectly incorporated the influence of interspecific

interactions by fitting models where evolutionary rates at a given time depend on the diversity of

lineages at that time (e.g., the "diversity-dependent" models of Mahler et al. 2010, Weir and

Mursleen 2013). While these models capture some parts of the interspecific processes of central

importance to evolutionary theory, such as the influence of ecological opportunity, they do not

explicitly account for trait-driven interactions between lineages, as trait values in one lineage do

not vary directly as a function of trait values in other evolving lineages.

Recently, Nuismer and Harmon (2015) proposed a model where the evolution of a

species' trait depends on other species' traits. In particular, they consider a model, which they

refer to as the model of phenotype matching, where the probability that an encounter between

two individuals has fitness consequences declines as the phenotypes of the individuals become

more dissimilar. The consequence of the encounter on fitness can be either negative if the

interaction is competitive, resulting in character divergence (matching competition, e.g. resource

competition), or positive if the interaction is mutualistic, resulting in character convergence

(matching mutualism, e.g. Müllerian mimicry). Applying Lande's formula (Lande 1976) and

given a number of simplifying assumptions—importantly that all lineages evolve in sympatry—

this model yields a simple prediction for the evolution of a population's mean phenotype.

Here, we develop inference tools for fitting the matching competition model (i.e., the

phenotype matching model of Nuismer and Harmon incorporating competitive interactions

between lineages) to combined phylogenetic and trait data. We begin by showing how to

compute likelihoods associated with this model. Next, we use simulations to explore the

statistical properties of maximum likelihood estimation of the matching competition model

(parameter estimation as well as the power and type I error rates). While the inclusion of

interactions between lineages is an important contribution to quantitative models of trait

evolution, applying the matching competition model to an entire clade relies on the assumption

that all lineages in the clade are sympatric. However, this assumption will be violated in most

empirical cases, so we also developed a method for incorporating data on the biogeographical

overlap between species for this model and for the linear and exponential diversity-dependent

trait models of Weir & Mursleen (2013), wherein the evolutionary rate at a given time in a tree

varies as a function of the number of lineages in the reconstructed phylogeny at that time (see

also Mahler et al. 2010).

We then fit the model to data from a classical adaptive radiation: Greater Antillean *Anolis*

lizards (Harmon et al. 2003; Losos 2009). Many lines of evidence support the hypothesis that

resource competition is responsible for generating divergence between species in both habitat use

(e.g., Pacala and Roughgarden 1982) and morphology (Schoener 1970; Williams 1972; see

review in Losos 1994). Thus, we can make an *a priori* prediction that model comparison will

uncover a signature of competition in morphological traits that vary with habitat and resource

use. However, in spite of the importance of interspecific competition to adaptive radiation

theory, the role of competition in driving adaptive divergence has been historically difficult to

measure (Losos 2009) because trait divergence resulting from competition between lineages

during their evolutionary history should have the effect of eliminating competition between those

lineages at the present. Nevertheless, the signature of such competition may be detectable in the

contemporary distribution of trait values and their covariance structure (Hansen and Martins

1996; Nuismer and Harmon 2015), and as such, the matching competition model should provide

a good fit to morphological data collected on anoles. Given the well-resolved molecular

phylogeny (Mahler et al. 2010, 2013) and the relatively simple geographical relationships

between species (i.e., species are restricted to single islands, Rabosky and Glor 2010; Mahler and

Ingram 2014), the Greater Antillean *Anolis* lizards provide a good test system for exploring the

effect of competition on trait evolution using the matching competition model.

METHODS

Likelihood Estimation of the Matching Competition Model

We consider the evolution of a quantitative trait under the matching competition model of

Nuismer & Harmon (2015) wherein trait divergence between lineages will be favored by

selection. In our version of the model, we remove stabilizing selection to focus on the effect of

competition. The evolution of a population's mean phenotype is thus given by (Eq. S38 in

Nuismer & Harmon 2015 with $\psi = 0$):

$\bar{z}_i' = \bar{z}_i + S(\mu - \bar{z}_i) + \delta$ (Eq. 1)

where drift is incorporated as a Gaussian variable δ with mean $= 0$ and variance $= \sigma^2$ (the

Brownian motion rate parameter) and S measures the strength of interaction (more intense

competitive interactions are represented by larger negative values), $\bar{z}_i'$ is the mean trait value for

lineage i after a infinitesimally small time step, and μ is the mean trait value for the entire clade

at the beginning of that time step. Hence, if a species trait value is greater (respectively smaller)

than the trait value average across species in the clade, the species' trait will evolve towards even

larger (respectively smaller) trait values.

Like several other models of quantitative trait evolution (Harmon et al. 2010, Weir &

Mursleen 2013), the expected distribution of trait values on a given phylogeny under the

matching competition model follows a multivariate normal distribution with mean vector made

of terms each equal to the character value at the root of the tree and variance-covariance matrix

determined by the parameter values and phylogeny. Nuismer & Harmon (2015) provide the

system of ordinary differential equations describing the evolution of the variance and covariance

terms through time (their Eqs.10b and 10c). These differential equations can be integrated

numerically from the root to the tips of phylogenies to compute expected variance-covariance

matrices and the associated likelihood values given by the multivariate normal distribution.

Additionally, to relax the assumption that all of the lineages in a clade coexist

sympatrically, we included a term to specify which lineages co-occur at any given time-point in

the phylogeny, which can be inferred, e.g., by biogeographical reconstruction. The resulting

system of ordinary differential equations describing the evolution of the variance and covariance

terms through time are:

$$\frac{dv_i^2}{dt} = -\left(\frac{2(n_i-1)}{n_i}S\right)v_i^2 + \frac{2S}{n_i}\left(\sum_{\substack{j=1 \\ (j\neq i)}}^{n}\mathbf{A}_{i,j}v_{i,j}\right) + \sigma^2 \qquad\text{(Eq. 2a)}$$

$$\frac{dv_{i,j}}{dt} = -\left(\frac{2(n_i-1)}{n_i}S\right)v_{i,j} + \frac{S}{n_i}\left(\sum_{\substack{k=1 \\ k\neq j}}^{n}\mathbf{A}_{i,k}v_{i,k} + \sum_{\substack{l=1 \\ l\neq i}}^{n}\mathbf{A}_{j,l}v_{j,l}\right) \qquad\text{(Eq. 2b)}$$

where v_i is the variance for each species i at time t and $v_{i,j}$ is the covariance for each species pair

i,j at time t. Eq. 2b describes the evolution of the covariance at time t if species i and j are in

sympatry at time t . If they are not, $\frac{dv_{i,j}}{dt} = 0$ such that the covariance between species i and j

remains fixed, The terms σ^2 and S are as specified above, $\mathbf{A}_{i,j}$ equals 1 at time t if i and j are

sympatric at that time, and 0 otherwise (Fig. 1), and $n_i = \sum_{j=1}^{n} A_{ij}$ is the number of lineages

interacting with lineage i at time t (equal to the number n of lineages in the reconstructed

phylogeny at time t if all species are sympatric). Here, we consider the matrix $\mathbf{A}$ to be a block

diagonal matrix, such that sets of interacting species are non-overlapping, as in the case of *Anolis*

lizards (Fig. 1). This implies that if species i and j are sympatric, then $n_i = n_j$ and $v_{i,j} = v_{j,i}$.

Note that when $S = 0$ or $n = 1$ (i.e., when a species is alone), this model reduces to Brownian

motion. This model has three free parameters: σ^2, S and the ancestral state z_0 at the root. As with

other models of trait evolution, the maximum likelihood estimate for the ancestral state is

computed through GLS using the estimated variance-covariance matrix (Grafen 1989; Martins

and Hansen 1997).

We used the ode function in the R package deSolve (Soetaert et al. 2010) to perform the

numerical integration of the differential equations using the "lsoda" solver, and the Nelder-Mead

algorithm implemented in the optim function to perform the maximum likelihood optimization.

Codes for these analyses are freely available on github (https://github.com/hmorlon/PANDA)

and included the R package RPANDA (Morlon 2014).

Incorporating Geography into Diversity-Dependent Models

Using the same geography matrix **A** described above for the matching competition model

(Fig. 1), we modified the diversity-dependent linear and exponential models of Weir & Mursleen

(2013) to incorporate biological realism into the models, because ecological opportunity is only

relevant within rather than between biogeographical regions. The resulting variance-covariance

matrices, **V**, of these models have the elements:

$\mathbf{V}_{ij} = \sum_{m=2}^{M}(\sigma_0^2 + bn_i)(\max(s_{ij} - t_{m-1}, 0) - \max(s_{ij} - t_m, 0))$ (Eq. 3)

for the diversity-dependent linear model, and

$\mathbf{V}_{ij} = \sum_{m=2}^{M}(\sigma_0^2 \times e^{rn_i})(\max(s_{ij} - t_{m-1}, 0) - \max(s_{ij} - t_m, 0))$ (Eq. 4)

for the diversity-dependent exponential model, where σ_0^2 is the rate parameter at the root of the

tree, *b* and *r* are the slopes in the linear and exponential models, respectively, s_{ij} is the shared

path length of lineages *i* and *j* from the root of the phylogeny to their common ancestor, n_i is the

number of sympatric lineages (as above) between nodes at times t_{m-1} and t_m (where t_1 is 0, the

time at the root, and t_M is the total length of the tree) (Weir & Mursleen 2013). When *b* or *r* = 0,

these models reduce to Brownian motion.

Simulation-based Analysis of Statistical Properties of the Matching Competition Model

To verify that the matching competition model can be reliably fit to empirical data, we

simulated trait datasets to estimate its statistical properties (i.e., parameter estimation, power, and

type I error rates). For all simulations, we began by first generating 100 pure-birth trees using

TreeSim (Stadler 2014). To determine the influence of the number of tips in a tree, we ran

simulations on trees of size $n = 20$, 50, 100, and 150. We then simulated continuous trait datasets

by applying the matching competition model recursively from the root to the tip of each tree

(Paradis 2011), following Eq. 1. For these simulations, we set $\sigma^2 = 0.05$ and systematically

varied S (-1.5, -1, -0.5, -0.1, or 0). Finally, we fit the matching competition model to these

datasets using the ML optimization described above.

To determine the ability of the approach to accurately estimate simulation parameter

values, we first compared estimated parameters to the known parameters used to simulate

datasets under the matching competition model (S and σ^2). We also quantified the robustness of

these estimates in the presence of extinction by estimating parameters for datasets simulated on

birth-death trees; in addition, we compared the robustness of the matching competition model to

extinction to that of the diversity-dependent models. These two latter sets of analyses are

described in detail in the Supplementary Appendix 1.

To assess the power of the matching competition model, we compared the fit of this

model to other commonly used trait models on the same data (i.e. data simulated under the

matching competition model). Specifically, we compared the matching competition model to (1)

Brownian motion (BM), (2) Ornstein-Uhlenbeck/single-stationary peak model (OU, Hansen &

Martin 1996), (3) exponential time-dependent (TD$_{exp}$, i.e., the early burst model, or the ACDC

model with the rate parameter set to be negative, Blomberg et al. 2003; Harmon et al. 2010), (4)

linear time-dependent evolutionary rate (TD$_{lin}$, Weir and Mursleen 2012), (5) linear rate

diversity-dependent (DD$_{lin}$, Mahler et al. 2010; Weir and Mursleen 2013), and (6) exponential

rate diversity-dependent (DD$_{exp}$, Weir and Mursleen 2013). These models were fitted using

geiger (Harmon et al. 2008) when available there (BM, OU, TD$_{exp}$, TD$_{lin}$), or using our own

codes, available in RPANDA (Morlon 2014) when they were not available in geiger (DD$_{lin}$,

DD$_{exp}$). Support for each model was determined using Akaike weights (Burnham and Anderson

2002).

We assessed the type I error rate of the matching competition model by calculating the fit

of this model to datasets simulated under the same trait models mentioned above. For BM and

OU models, we generated datasets from simulations using parameter values from the appendix of

Harmon et al. 2010 scaled to a tree of length 400 (BM, $\sigma^2 = 0.03$; OU, $\sigma^2 = 0.3$, $\alpha = 0.06$). For

both the linear and exponential versions of the time- and diversity-dependent models, we

simulated datasets with starting rates of $\sigma^2 = 0.6$ and ending rates of $\sigma^2 = 0.01$, declining with a

slope determined by the model and tree (e.g., for time-dependent models, the slope is a function

of the total height of the tree; for the TD$_{exp}$ model, these parameters result in a total of 5.9 half-

lives elapsing from the root to the tip of the tree, Slater and Pennell 2014). These simulations

were performed using our own codes, available in RPANDA (Morlon 2014). As above, we

calculated the Akaike weights for all models for each simulated dataset.

Fitting the Matching Competition Model of Trait Evolution to Caribbean Anolis *Lizards*

To determine whether the matching competition model is favored over models that ignore

interspecific interactions in an empirical system where competition is known to have influenced

character evolution, we fit the matching competition model to a morphological dataset on adult

males of 100 species of Greater Antillean *Anolis* lizards and the time calibrated, maximum clade

credibility tree calculated from a Bayesian sample of molecular phylogenies (Mahler et al. 2010,

2013; Mahler and Ingram 2014). We included the first four size-corrected phylogenetic principal

components from a set of 11 morphological measurements, collectively accounting for 93% of

the cumulative variance explained (see details in Mahler et al. 2013). Together, the shape axes

quantified by these principal components describe the morphological variation associated with

differences between classical ecomorphs in Caribbean anoles (Williams 1972). In addition to the

matching competition model, we fit the six previously mentioned models (BM, OU, TD_{exp}, TD_{lin},

DD_{exp}, and DD_{lin}) to the *Anolis* dataset.

For the matching competition model and diversity-dependent models, to determine the

influence of designating clades as sympatric and allopatric, we fit the model for each trait using

two sets of geography matrices (i.e., **A** in Eq. 1b, 2, & 3, see Fig. 1): one where all lineages were

set as sympatric, and another following the biogeographical reconstruction presented in Fig 18.3

of Mahler and Ingram (2014). To simplify the ML optimization, we restricted *S* to take negative

values while fitting the matching competition model including the biogeographical relationships

among taxa.

RESULTS

Statistical Properties of the Matching Competition Model

Across a range of *S* values, maximum likelihood optimization returns reliable estimates

of parameter values (Fig. 2). As the number of tips increases, so does the reliability of maximum

likelihood parameter values (Fig. 2). Parameter estimates remain reliable in the presence of

extinction, unless the extinction fraction is very large (Supplementary Appendix 1). When

datasets are simulated under the matching competition model, model selection generally picks

the matching competition model as the best model (Figs. 3, S1); the strength of this

discrimination depends on both the S value used to simulate the data and the size of the tree

(Figs. 3, S1).

We found that the matching competition model behaves similarly to a model of declining

evolutionary rates through time, as the rate parameter values of the TD_{exp} model estimated on

data generated under the matching competition model became increasingly negative under

increasing levels of competition (Fig. S2). The dynamics of this declining rate were better

described by an exponential decline in rates than by a linear one (Table S1).

Simulating datasets under BM, OU, DD_{exp}, and DD_{lin} generating models, we found that

there is a reasonably low type I error rate for the matching competition model (Fig. 4a,b,e,f).

When character data were simulated under a TD_{lin} model of evolution, the matching competition

and/or the diversity-dependent models tended to have lower AICc values than the TD_{lin} model,

especially among smaller trees (Figure 4d). For data generated under a TD_{exp} model, model

selection always favored the matching competition model over the TD_{exp} model (Fig. 4c).

Competition in Greater Antillean Anolis *Lizards*

For the first four phylogenetic principal components describing variation in *Anolis*

morphology, we found that models that incorporate species interactions fit the data better than

models that ignore them (Table 1). PC1 and PC3, which describe variation in hindlimb/hindtoe

and forelimb/foretoe length (Mahler et al. 2013), respectively, are fit best by the matching

competition model. PC2 and PC4, which describe variation in body size (snout vent length) and

lamellae number, respectively, are fit best by the exponential diversity-dependent model.

Additionally, in every case, models that incorporated the geographic relationships among species

in the tree outperformed models that assumed that lineages interact in sympatry at the clade level

(Table 1).

DISCUSSION

The inference methods we present here represent an important new addition to the

comparative trait analysis toolkit. Whereas previous models had not accounted for the influence

of trait values in other lineages on character evolution, the matching competition model takes

these into account. Furthermore, extending both the matching competition model and two

diversity-dependent trait evolution models to incorporate geographic networks of sympatry

further extends the utility and biological realism of these models.

We found that the matching competition property has increasing power and accuracy of

parameter estimation with increasing tree sizes and competition strength. We also found that, for

most of the generating models we tested (but see below), the matching competition model is not

erroneously favored using model selection (i.e., there is a reasonably low type-I error rate). As

with all other models, the statistical properties of the matching competition model will depend on

the size and shape of a particular phylogeny as well as specific model parameter values. Future

investigators can employ other approaches, such as phylogenetic Monte Carlo and posterior

predictive simulations directly on their empirical trees (Boettiger et al. 2012, Slater & Pennell

2014), to assess the confidence they can have in their results

We found that data generated under time-dependent models were often fit better by

models that incorporate interspecific interactions (i.e., density-dependent and matching

competition models) (Fig. 4c,d). This was especially true for the TD_{exp} model, often referred to

as the early-burst model—the matching competition model nearly always fit data generated

under the TD_{exp} model better than the TD_{exp} model (Fig. 4c). We do not view this as a major

limitation of the model for two reasons. First, the TD_{exp} model is known to be statistically

difficult to estimate on neontological data alone (Harmon et al. 2010; Slater et al. 2012; Slater

and Pennell 2014). Accordingly, since the matching competition model also describes declining

evolutionary rates from the root to the tip of the tree (Fig. S2, Table S1), it is perhaps not

surprising that the matching competition model fits data generated under the TD_{exp} model well.

Secondly, and more importantly, time-dependent models are not process-based models, but

rather incorporate time since the root of a tree as a proxy for ecological opportunity or available

niche space (Harmon et al. 2010; Mahler et al. 2010; Slater 2015). The matching competition and

density-dependent models explicitly account for the interspecific competitive interactions that

time-dependent models purport to model, thus we argue that these process-based models are

more biologically meaningful than time-dependent models (Moen and Morlon 2014).

Because the matching competition model depends on the mean trait values in an evolving

clade, maximum likelihood estimation is robust to extinction, whereas the diversity-dependent

models are less so (Appendix S1, Figs. S3-S6). Nevertheless, given the failure of maximum

likelihood to recover accurate parameter estimations at high levels of extinction ($\mu : \lambda \geq 0.6$), we

suggest that these models should not be used in clades where the extinction rate is known to be

particularly high. In such cases, it would be preferable to adapt the inference framework

presented here to include data from fossil lineages (Slater et al. 2012), which could be easily

done by integrating the ordinary differential equations described in Eq. 2a and 2b on non-

ultrametric trees.

For all of the traits we analyzed, we found that models incorporating both the influence of

other lineages and the specific geographical relationships among lineages were the most strongly

supported models (though less strikingly for PC3 and PC4). The matching competition model is

favored in the PCs describing variation in relative limb size. Previous research demonstrates that

limb morphology explains between-ecomorph variation in locomotive capabilities and perch

characteristics (Losos 1990, 2009; Irschick et al. 1997), and our results suggest that the

evolutionary dynamics of these traits have been influenced by the evolution of limb morphology

in other sympatric lineages. These results support the assumption that interspecific interactions

resulting from similarity in trait values are important components of adaptive radiations (Losos

1994, Schluter 2000), a prediction that has been historically difficult to test (Losos 2009, but see

Mahler et al. 2010). In combination with previous research demonstrating a set of convergent

adaptive peaks to which lineages are attracted (Mahler et al. 2013), our results suggest that

competition likely played an important role in driving lineages toward these distinct peaks.

We imagine that the matching competition model and biogeographical implementations

of diversity-dependent models will play a substantial role in comparative studies of interspecific

competition. There are many possible extensions of the tools developed in this paper. In the

current implementation of interspecific geographic overlap, the matrices of sympatry/allopatry

are assumed to be block diagonal matrices. In the future, the model can be extended to more

complex geography matrices that are more realistic for mainland taxa (e.g., using ancestral

biogeographical reconstruction, Ronquist and Sanmartín 2011; Landis et al. 2013), and can also

specify degrees of sympatric overlap (i.e., syntopy). Additionally, although the current version of

the model is rather computationally expensive with larger trees (with 100 or more tips), future

work to speed up the calculation of the variance-covariance matrix under the matching

competition model would allow the inclusion of uncertainty in biogeographical reconstruction

(e.g., from samples of stochastic maps). The current form of the model assumes that the degree

of competition is equal for all interacting lineages. Future modifications of the model, such as

applications of stepwise AICc algorithms (Alfaro et al. 2009; Thomas and Freckleton 2012;

Mahler et al. 2013) or reversible-jump Markov Chain Monte Carlo (Pagel and Meade 2006;

Eastman et al. 2011; Rabosky 2014; Uyeda and Harmon 2014), may be useful to either identify

more intensely competing lineages or test specific hypotheses about the strength of competition

between specific taxa. Improvements could also be made on the formulation itself of the

evolution of a species' trait as a response to the phenotypic landscape in which the species

occurs. The current formulation (Eq. 1) corresponds to a scenario in which the rate of phenotypic

evolution on a lineage gets higher as the lineage deviates from the mean phenotype, although

character displacement theory, for example, posits that selection for divergence should be the

strongest when species are most ecologically similar (Brown & Wilson 1956). Finally, a great

array of extensions will come from modeling species interactions not only within clades, but also

among interacting clades, as in the case of coevolution in bipartite mutualistic or antagonistic

networks, such as plant-pollinator or plant-herbivore systems (Bartosek et al. 2012).

In addition to uncovering a signature of competitive interactions in traits used in

exploitative competition for resources (e.g., trait-axes along which adaptive radiations proceed,

Schluter 2000), the tools presented here can be used to test several hypotheses that have

historically been difficult to test using phylogenetic comparative methods. For example, various

hypotheses about the dynamics of traits involved in social interactions that posit that interspecific

interactions have played an important role in shaping contemporary trait values (e.g., Pfennig

and Pfennig 2012; Grether et al. 2013) are beginning to be tested at comparative scales (Seddon

et al. 2013; Tobias et al. 2013; Martin et al. 2015). However, the tools available for such analyses

have been limited (e.g., sister-taxa comparisons) relative to those available to test

macroevolutionary hypotheses about within-taxa processes. By comparing the fits of the

matching competition model with other models that do not include competitive interactions

between lineages, biologists can directly test hypotheses that make predictions about the role of

interspecific interactions in driving trait evolution.

Acknowledgements

We thank M. Manceau for his help troubleshooting the inference approach, J. Weir for

providing R code for diversity-dependent models, and E. Lewitus and O. Missa for helpful

comments on the manuscript. This research was funded by the Agence Nationale de la Recherche

(grant CHEX-ECOEVOBIO); and the European Research Council (grant 616419-PANDA to

HM).

References

Alfaro M.E., Santini F., Brock C., Alamillo H., Dornburg A., Rabosky D.L., Carnevale G.,

Harmon L.J. 2009. Nine exceptional radiations plus high turnover explain species diversity

in jawed vertebrates. Proc. Natl. Acad. Sci. 106:13410–13414.

Bartosek K., Pienaar J., Mostad P., Andersson S., Hansen T.F. 2012. A phylogenetic

comparative method for studying multivariate adaptation. J. of Theor. Biol. 314:204-215.

Blomberg S.P., Garland T., Ives A.R. 2003. Testing for phylogenetic signal in comparative data:

behavioral traits are more labile. Evolution. 57:717–745.

Boettiger C., Coop G., Ralph P. 2012. Is Your Phylogeny Informative ? Measuring the Power of

Comparative Methods. Evolution 66:2240–2251.

Brown W.L., Wilson E.O. 1956. Character Displacement. Syst. Zool. 5:49-62.

Burnham K.P., Anderson D.R. 2002. Model selection and multimodel inference: a practical

information-theoretic approach. New York, NY: Springer.

Cavalli-Sforza, L.L., Edwards, A.W.F. 1967. Phylogenetic analysis: Models and estimation

procedures. Amer. J. of Hum. Gen. 19:233-257.

Coyne J.A., Orr H.A. 2004. Speciation. Sunderland, MA: Sinauer Associates.

Dobzhansky T. 1937. Genetics and the Origin of Species. New York, NY: Columbia University

Press.

Dobzhansky T. 1940. Speciation as a stage in evolutionary divergence. Am. Nat. 74:312–321.

Eastman J.M., Alfaro M.E., Joyce P., Hipp A.L., Harmon L.J. 2011. A novel comparative
method for identifying shifts in the rate of character evolution on trees. Evolution 65:3578–
3589.

Felsenstein, J. 1988. Phylogenies and quantitative characters. Ann. Rev. of Ecol., Evol., &
Systematics. 19:445-471.

Fisher R.A. 1930. The Genetical Theory of Natural Selection. Oxford, UK.: Oxford University
Press.

Grafen A. 1989. The phylogenetic regression. Philos. Trans. R. Soc. Lond. B. Biol. Sci.
326:119–157.

Grant P.R., Grant B.R. 2002. Adaptive radiation of Darwin's finches: Recent data help explain
how this famous group of Galápagos birds evolved, although gaps in our understanding
remain. Am. Sci. 90:130–139.

Grant P.R., Grant B.R. 2011. How and Why Species Multiply: the Radiation of Darwin's
Finches. Princeton, NJ: Princeton University Press.

Grant P.R. 1972. Convergent and divergent character displacement. Biol. J. Linn. Soc. 4:39–68.

Grant P.R. 1999. Ecology and Evolution of Darwin's finches. Princeton, NJ: Princeton
University Press.

Grether G.F., Anderson C.N., Drury J.P., Kirschel A.N.G., Losin N., Okamoto K., Peiman K.S.

2013. The evolutionary consequences of interspecific aggression. Ann. N. Y. Acad. Sci.

1289:48–68.

Grether G.F., Losin N., Anderson C.N., Okamoto K. 2009. The role of interspecific interference

competition in character displacement and the evolution of competitor recognition. Biol.

Rev. 84:617–635.

Gröning J., Hochkirch A. 2008. Reproductive interference between animal species. Q. Rev. Biol.

83:257–282.

Hansen T.F., Martins E.P. 1996. Translating between microevolutionary process and

macroevolutionary patterns: The correlation structure of interspecific data. Evolution

50:1404–1417.

Harmon L.J., Losos J.B., Jonathan Davies T., Gillespie R.G., Gittleman J.L., Bryan Jennings W.,

Kozak K.H., McPeek M.A., Moreno-Roark F., Near T.J., Purvis A., Ricklefs R.E., Schluter

D., Schulte J.A., Seehausen O., Sidlauskas B.L., Torres-Carvajal O., Weir J.T., Mooers

A.T. 2010. Early bursts of body size and shape evolution are rare in comparative data.

Evolution 64:2385–2396.

Harmon L.J., Schulte J.A., Larson A., Losos J.B. 2003. Tempo and mode of evolutionary

radiation in iguanian lizards. Science 301:961–964.

Harmon L.J., Weir J.T., Brock C.D., Glor R.E., Challenger W. 2008. GEIGER: investigating

evolutionary radiations. Bioinformatics. 24:129–131.

Irschick D.J., Vitt L.J., Zani P.A., Losos J.B. 1997. A comparison of evolutionary radiations in

mainland and Caribbean Anolis lizards. Ecology. 78:2191–2203.

Landis M.J., Matzke N.J., Moore B.R., Huelsenbeck J.P. 2013. Bayesian analysis of

biogeography when the number of areas is large. Syst. Biol. 62:789–804.

Losos J.B., Ricklefs R.E. 2009. Adaptation and diversification on islands. Nature. 457:830–836.

Losos J.B. 1990. The evolution of form and function: morphology and locomotor performance in

West Indian Anolis lizards. 44:1189–1203.

Losos J.B. 1994. Integrative Approaches to Evolutionary Ecology: Anolis Lizards as Model

Systems. Annu. Rev. Ecol. Syst. 25:467–493.

Losos J.B. 2009. Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles.

Los Angeles, CA: University of California Press.

Mahler D.L., Ingram T., Revell L.J., Losos J.B. 2013. Exceptional convergence on the

macroevolutionary landscape in island lizard radiations. Science 341:292–5.

Mahler D.L., Ingram T. 2014. Phylogenetic Comparative Methods for Studying Clade-Wide

Convergence. In: Garamszegi L., editor. Modern Phylogenetic Comparative Methods and

Their Application in Evolutionary Biology. New York, NY: Springer. p. 425–450.

Mahler D.L., Revell L.J., Glor R.E., Losos J.B. 2010. Ecological opportunity and the rate of

morphological evolution in the diversification of greater Antillean anoles. Evolution

64:2731–2745.

Martin P.R., Montgomerie R., Lougheed S.C. 2015. Color patterns of closely related bird species

are more divergent at intermediate levels of breeding-range sympatry. Am. Nat. 185:443–

451.

Martins E.P., Hansen T.F. 1997. Phylogenies and the comparative method: a general approach to

incorporating phylogenetic information into the analysis of interspecific data. Am. Nat.

149:646–667.

Mayr E. 1963. Animal Species and Evolution. Cambridge, MA: Harvard University Press.

Moen D., Morlon H. 2014. Why does diversification slow down? Trends Ecol. Evol. 29:190–

197.

Morlon H. 2014. RPANDA: Phylogenetic ANalyses of DiversificAtion. R package version 1.0.

https://github.com/hmorlon/PANDA

Nuismer S.L., Harmon L.J. 2015. Predicting rates of interspecific interaction from phylogenetic

trees. Ecol. Lett. 18:17–27.

Ord T.J., King L., Young A.R. 2011. Contrasting theory with the empirical data of species

recognition. Evolution 65:2572–2591.

Ord T.J., Stamps J.A. 2009. Species identity cues in animal communication. Am. Nat. 174:585–

593.

Pacala S., Roughgarden J. 1982. Resource partitioning and interspecific competition in two two-

species insular Anolis lizard communities. Science 217:444–446.

Pagel M., Meade A. 2006. Bayesian analysis of correlated evolution of discrete characters by

reversible-jump Markov chain Monte Carlo. Am. Nat. 167:808–825.

Paradis E. 2011. Analysis of Phylogenetics and Evolution with R. New York, NY: Springer.

Pfennig D.W., Pfennig K.S. 2012. Evolution's Wedge: Competition and the Origins of Diversity.

Los Angeles, CA: Univ of California Press.

Pfennig K.S., Pfennig D.W. 2009. Character displacement: ecological and reproductive

responses to a common evolutionary problem. Q. Rev. Biol. 84:253–276.

Rabosky D.L., Glor R.E. 2010. Equilibrium speciation dynamics in a model adaptive radiation of

island lizards. Proc. Natl. Acad. Sci. U. S. A. 107:22178–22183.

Rabosky D.L. 2014. Automatic detection of key innovations, rate shifts, and diversity-

dependence on phylogenetic trees. PLoS One. 9:e89543.

Ronquist F., Sanmartín I. 2011. Phylogenetic methods in biogeography. Annu. Rev. Ecol. Evol.

Syst. 42:441–464.

Rundle H.D., Nosil P. 2005. Ecological speciation. Ecol. Lett. 8:336–352.

Schluter D. 2000. The Ecology of Adaptive Radiation. Oxford, UK: Oxford University Press.

Schoener T.W. 1970. Size Patterns in West Indian Anolis Lizards. II. Correlations with the Sizes

of Particular Sympatric Species-Displacement and Convergence. Am. Nat. 104:155–174.

Seddon N., Botero C.A., Tobias J.A., Dunn P.O., Macgregor H.E., Rubenstein D.R., Uy J.A.,

Weir J.T., Whittingham L.A., Safran R.J. 2013. Sexual selection accelerates signal

evolution during speciation in birds. Proc. R. Soc. B Biol. Sci. 280:20131065.

Slater G.J., Harmon L.J., Alfaro M.E. 2012. Integrating fossils with molecular phylogenies

improves inference of trait evolution. 66:3931–3944.

Slater G.J., Pennell M.W. 2014. Robust regression and posterior predictive simulation increase

power to detect early bursts of trait evolution. Syst. Biol. 63:293–308.

Soetaert K., Petzoldt T., Setzer R.W. 2010. Solving Differential Equations in R: Package

deSolve. J. Stat. Softw. 33:1–25.

Stadler T. 2014. TreeSim: Simulating trees under the birth-death model. R package version 2.1.

http://CRAN.R-project.org/package=TreeSim.

Thomas G.H., Freckleton R.P. 2012. MOTMOT: models of trait macroevolution on trees.

Methods Ecol. Evol. 3:145–151.

Tobias J.A., Cornwallis C.K., Derryberry E.P., Claramunt S., Brumfield R.T., Seddon N. 2013.

Species coexistence and the dynamics of phenotypic evolution in adaptive radiation. Nature.

506:359–363.

Uyeda J.C., Harmon L.J. 2014. A novel Bayesian method for inferring and interpreting the

dynamics of adaptive landscapes from phylogenetic comparative data. Syst. Biol. 63:902–

918.

Wallace A.R. 1889. Darwinism. 2007 facsimile, Cosimo, Inc.

Weir J.T., Mursleen S. 2013. Diversity-dependent cladogenesis and trait evolution in the

adaptive radiation of the auks (Aves: Alcidae). Evolution 67:403–416.

Williams E.E. 1972. The Origin of Faunas. Evolution of Lizard Congeners in a Complex Island

Fauna: A Trial Analysis. Evol. Biol. 6:47–89.

  Table & Figure legends

Table 1. Comparison of model fits for the first four phylogenetic principal components of a

morphological dataset of Greater Antillean anoles. Models run incorporating geography matrices

are indicated by "+ GEO", and models with the lowest AICc for each trait are shaded and written

in bold text. Parameter values presented follow the nomenclature of Eqs. 2-4 in the main text,

and k represents the number of parameters estimated for each model. Note that TD_{exp} is the

ACDC model (or the early-burst model when $r < 0$). OU model weights were excluded because

α was estimated to be 0 for each trait, and thus the OU model was equivalent to BM.

Figure 1. Illustration of geography matrices (defined between each node) delineating which

lineages interact in sympatry in an imagined phylogeny. These matrices were used to identify

potentially interacting lineages for the matching competition and both diversity-dependent

models of character evolution (see Eqs. 2-4 in the main text). *Anolis* outline courtesy of Sarah

Werning, licensed under Creative Commons (http://creativecommons.org/licenses/by/3.0/).

Figure 2. Parameter estimation under the matching competition model. As tree size increases

and/or the magnitude of competition increases (i.e., the S parameter in the matching competition

model becomes more negative), so does the accuracy of ML parameter estimates of (A) S ($n =$

100 for each tree size and S value combination; red horizontal lines indicate the simulated S

value) and (B) the accuracy of ML parameter estimates of σ^2 ($n = 500$ for each tree size; red

horizontal lines indicate the simulated σ^2 value). The numbers below the violin plots in (B) show

the number of outliers removed for plotting.

Figure 3. The statistical power of the matching competition increases with tree size and with

increasing levels of competition (i.e., increasingly negative S values).

Figure 4. Type-I error simulation results for the matching competition model. When the

generating model is either (A) BM, (B) OU, (E) DD_{exp} (for larger trees) or (F) DD_{lin}, the

generating model is largely favored by model selection. However, both (C) TD_{exp} and (D) TD_{lin}

(for smaller trees) are erroneously rejected as the generating model.

552
553

554 **Table 1.**

Trait	Model	k	σ^2	b	r	S	$\ln(\mathcal{L})$	ΔAICc	Akaike weights
pPC1	BM	2	0.0033	—	—	—	-13.68	22.40	<0.01
	OU	3	0.0033	—	—	—	-13.68	24.52	—
	TDexp	3	0.0324	—	-0.068	—	-5.20	7.55	0.02
	TDlinear	3	0.0113	-0.019	—	—	-4.88	6.92	0.03
	LDexp	3	0.0184	—	-0.028	—	-4.37	5.90	0.04
	LDexp + GEO	3	0.0086	—	-0.043	—	-7.28	11.72	<0.01
	LDlin	3	0.0089	-0.00008	—	—	-4.89	6.94	0.02
	LDlin + GEO	3	0.0042	-0.000042	—	—	-10.34	17.84	<0.01
	MC$_{sym}$	3	0.0010	—	—	-0.037	-3.67	4.50	0.08
	MC + GEO	**3**	**0.0010**	**—**	**—**	**-0.037**	**-1.42**	**0**	**0.80**
pPC2	BM	2	0.0027	—	—	—	-4.69	8.55	0.01
	OU	3	0.0027	—	—	—	-4.69	10.67	—
	TDexp	3	0.0046	—	-0.014	—	-4.30	9.89	<0.01
	TDlinear	3	0.0047	-0.011	—	—	-4.23	9.75	<0.01
	LDexp	3	0.0041	—	-0.006	—	-4.27	9.84	<0.01
	LDexp + GEO	**3**	**0.0067**	**—**	**-0.039**	**—**	**0.65**	**0**	**0.84**
	LDlin	3	0.0041	-0.00002	—	—	-4.21	9.72	<0.01
	LDlin + GEO	3	0.0035	-0.000035	—	—	-1.57	4.44	0.09
	MC$_{sym}$	3	0.0021	—	—	-9.9e-3	-3.95	9.19	<0.01
	MC + GEO	3	0.0019	—	—	-0.014	-2.90	7.10	0.02
pPC3	BM	2	0.0010	—	—	—	45.57	1.4	0.11
	OU	3	0.0010	—	—	—	45.57	3.5	—
	TDexp	3	0.0020	—	-0.019	—	46.30	2.06	0.08
	TDlinear	3	0.0019	-0.013	—	—	46.41	1.85	0.09
	LDexp	3	0.0017	—	-0.008	—	46.40	1.86	0.09
	LDexp + GEO	3	0.0015	—	-0.017	—	46.67	1.33	0.11
	LDlin	3	0.0017	-0.000009	—	—	46.46	1.74	0.09
	LDlin + GEO	3	0.0013	-0.000014	—	—	46.58	1.5	0.10
	MC$_{sym}$	3	0.0007	—	—	-0.012	46.75	1.17	0.12
	MC + GEO	**3**	**0.0007**	**—**	**—**	**-0.015**	**47.33**	**0**	**0.22**
pPC4	BM	2	0.0006	—	—	—	69.07	1.96	0.07
	OU	3	0.0006	—	—	—	69.07	4.09	—
	TDexp	3	0.0015	—	-0.015	—	70.55	1.12	0.10
	TDlinear	3	0.0012	-0.011	—	—	70.45	1.32	0.09
	LDexp	3	0.0012	—	-0.0061	—	70.52	1.18	0.10
	LDexp + GEO	**3**	**0.0010**	**—**	**-0.019**	**—**	**71.11**	**0**	**0.17**
	LDlin	3	0.0011	-0.00002	—	—	70.39	1.45	0.08
	LDlin + GEO	3	0.0008	-0.000035	—	—	70.66	0.92	0.11
	MC	3	0.0004	—	—	-0.015	71.1	0.02	0.17
	MC + GEO	3	0.0004	—	—	-0.013	70.6	1.03	0.10

558   Figure 1.
559
560

562   Figure 2.

563
564

565   Figure 3

567
568   Figure 4.
569

570
571
572
573

574 **Supplementary Material**

576 Appendix S1

578 Table S1

580 Figure S1-S6

Supplementary Appendix 1: **Estimating the effect of extinction on parameter estimation for**

the matching competition and density-dependent models.

Given that the matching competition and diversity-dependent models take into account

the number of interacting lineages, extinction may affect our ability to recover true parameter

values. To estimate the impact of extinction, we simulated 100 trees with 100 extant species,

varying the extinction fraction (μ: $\lambda = 0.2$, 0.4, 0.6, and 0.8). As above, we recursively simulated

traits using the matching competition model with $\sigma^2 = 0.05$ and $S = -1.5$, -1, -0.5, -0.1, or 0, and

the linear and exponential diversity-dependent models with starting rates of $\sigma^2 = 0.6$ and ending

rates of $\sigma^2 = 0.01$. We then estimated the maximum likelihood parameter estimates for the

generating models by fitting the models to the trait values for extant species and the tree with

extinct lineages removed. In the case of the matching competition model, because many

simulated birth-death trees with high extinction rates have substantially older root ages, the

simulated trait datasets for some trees had very large variances. For these biologically unrealistic

trait datasets (i.e., variance in trait values $\geq 1\times10^8$), ML does not yield reliable parameter

estimates, so we removed them from further analyses (the sample size of included simulations is

reported in Fig. S3, S4).

Parameter estimates are quite robust to extinction under the matching competition model

(Fig. S3, S4), and much more so than under both diversity-dependent models (Fig. S5,S6). Under

the matching competition model, the maximum likelihood optimization returns reliable estimates

of S and σ^2 values used to simulate datasets on trees with extinct lineages (Fig. S3, S4), although

the estimates become much less reliable with larger extinction fractions, likely because

simulations under the matching competition model were unbounded, resulting in trait datasets

with biologically unrealistic variances. Under both diversity-dependent models, the magnitude of

both the slope and σ^2 parameter values are increasingly underestimated with increasing

extinction fractions (Fig. S5, S6).

612
613

614 **Table S1.** Data generated under the matching competition model are better fit by the TD_{exp}

615 model than by the TD_{lin} model. $\Delta AICc$ [90% CI] presented for the matching competition (MC)

616 and time dependent models.

tips	S value	MC	TD_{exp}	TD_{lin}
20	-1.5	0 [0, 0.68]	6.04 [3.20, 10.53]	12.17 [1.27, 67.92]
	-1	0 [0, 2.17]	3.96 [1.87, 7.47]	4.18 [0.78, 25.03]
	-0.5	0 [0, 2.76]	2.74 [1.70, 4.06]	2.42 [0.97, 6.36]
	-0.1	2.37 [0, 2.79]	2.79 [1.83, 5.04]	2.1 [0.85, 3.11]
	0	2.47 [0, 2.98]	2.79 [2.17, 6.0]	2.06 [0.88, 3.62]
50	-1.5	0 [0,0]	8.46 [5.57, 12.06]	63.24 [11.75, 216.11]
	-1	0 [0,0]	5.51 [3.45, 8.49]	17.26 [3.50, 67.67]
	-0.5	0 [0, 1.57]	2.53 [1.36, 5.01]	1.87 [0.51, 8.29]
	-0.1	1.83 [0, 2.37]	2.27 [1.44, 4.09]	1.86 [0.51, 3.2]
	0	1.98 [0.09, 2.45]	2.27 [1.37, 4.62]	1.61 [0.05, 3.39]
100	-1.5	0 [0,0]	10.56 [7.11, 13.92]	208.76 [61.32, 442.7]
	-1	0 [0,0]	7.02 [4.93, 8.84]	45.69 [6.44, 149.18]
	-0.5	0 [0, 1.43]	3.34 [1.61, 4.62]	3.67 [0.63, 15.55]
	-0.1	1.60 [0, 2.12]	2.04 [0.99, 2.45]	1.95 [0.61, 4.36]
	0	1.83 [0, 2.11]	2.13 [0.98, 4.41]	1.52 [0, 4.77]
150	-1.5	0 [0,0]	11.77 [8.74, 14.23]	392.29 [177.79, 835.42]
	-1	0 [0,0]	7.63 [5.80, 10.14]	80.58 [27.51, 232.06]
	-0.5	0 [0, 1.0]	3.50 [2.35, 6.11]	5.71 [0.42, 29.24]
	-0.1	1.45 [0, 2.17]	1.83 [0.81, 2.61]	1.6 [0.19, 6.49]
	0	1.81 [0, 2.18]	2.08 [1.02, 4.45]	1.67 [0, 5.98]

Supplementary Figure 1. As tree size and/or the degree of competition (*S*) increases, model

selection becomes more reliable. Comparison of Akaike weights (median & 90% CIs) for NH,

BM, OU, and EB models when simulated under various levels of competition (*S* = -1.5, -1, -0.5,

-0.1, and 0) for trees with 20, 50, 100, and 150 tips.

Supplementary Figure 2. Maximum likelihood estimates of ACDC rate parameters (i.e., TDexp

where rate parameters can take both positive and negative values) fit to data generated under the

matching competition model. As the level of competition increases (i.e., as *S* becomes

increasingly negative), the estimated rate parameter of the ACDC model becomes increasingly

negative.

Supplementary Figure 3. Simulation results showing the effect of varying the extinction fraction

on estimation of the *S* parameter for the matching competition model. Red horizontal lines

indicate the simulated *S* values, and numbers above sets of simulations indicate the sample size

of included simulations under those scenarios (see main text for more details).

639
640

641   *Supplementary Figure 4.* Simulation results showing the effect of varying the extinction fraction

642   on estimation of the σ^2 parameter for the matching competition model. Red horizontal lines

643   indicate the simulated σ^2 value (0.05), the numbers below sets of simulations indicate the sample

644   size of included simulations under those scenarios (see main text for more details), and the

645   number in parentheses indicate sample size after σ^2 values > 0.25 re removed.

647
648

Supplementary Figure 5. Simulation results showing the effect of varying the extinction fraction

on slope (top) and σ^2 (bottom) parameters for the exponential diversity-dependent model.

Increasing extinction levels result in increasingly underestimated slope values and σ^2 parameters.

Red horizontal lines indicate the simulated parameter values.

Supplementary Figure 6. Simulation results showing the effect of varying the extinction fraction

on slope (top) and sigma-squared (bottom) parameters for the linear diversity-dependent model.

Increasing extinction levels result in increasingly underestimated slope values and σ^2 parameters.

Red horizontal lines indicate the simulated parameter values.