1 **Brain Transcriptional Profiles of Male Alternative Reproductive Tactics in**
2 **Bluegill Sunfish**
3

4 Charlyn G. Partridge[*§], Matthew D. MacManes[†], Rosemary Knapp[¶], Bryan D. Neff[*]
5
[*]Department of Biology, University of Western Ontario, London, ON, N6C 1B1
Canada
[†]Department of Molecular, Cellular, and Biomedical Sciences, University of New
Hampshire, Durham, NH 03824, USA
[¶]Department of Biology, University of Oklahoma, Norman, OK 73019, USA
[§]Current address: Annis Water Resources Institute, Grand Valley State University,
Muskegon, MI 49441, USA

Abstract

Bluegill sunfish are one of the classic systems for studying male alternative reproductive tactics (ARTs) in teleost fishes. In this species, there are two distinct life histories: parental and cuckolder, encompassing three reproductive tactics, parental, satellite, and sneaker. The parental tactic is fixed, whereas individuals who enter the cuckolder life history transition from the sneaker to the satellite tactic as they grow. For this study, we used RNAseq to characterize the brain transcriptome of the three male tactics during spawning to identify gene categories associated with each tactic and identify potential candidate genes influencing their different spawning behaviors. We found that sneaker males had higher levels of gene differentiation compared to the other two tactics, suggesting that life history does not exclusively drive differential gene expression. Sneaker males had high expression in ionotropic glutamate receptor genes, specifically AMPA receptors, which may be important for increased working spatial memory while attempting to cuckold nests in bluegill colonies. We also found significant expression differences in several candidate genes involved in ARTs that were previously identified in other species and suggest a previously undescribed role for cytosolic 5'-nucleotidase II (*nt5c2*) in influencing parental male behavior during spawning.

 Introduction

Understanding the genetic mechanisms influencing variation in behavior can

provide insight into how different behavioral phenotypes within populations evolve

and are maintained. One important area of research on behavioral phenotypes

focuses on alternative reproductive tactics (ARTs), which are found in a wide array

of taxa (Gross 1996; Mank & Avise 2006; Taborsky *et al.* 2008; Taborsky &

Brockman 2010). ARTs typically consist of larger males practicing a "territorial"

tactic that maintain and protect breeding territories and smaller "sneaking" males

that sneak fertilizations rather than compete with territorial males (Taborsky

1998). The mechanisms underlying the expression of ARTs can differ significantly

across species. In some cases, tactics are fixed for life (fixed tactics) (Taborsky

1998) and often represent distinct life histories. Fixed tactics can occur through

either inherited genetic polymorphisms (Lank *et al.* 1995; Shuster & Sassaman

1997), condition-dependent switches that are triggered prior to sexual maturation

(Taborsky 1996; Gross 1996; Gross & Repka 1998), or a combination of genetic and

environmental factors (Piché *et al.* 2008, Neff & Svensson 2013). In other cases,

individuals can exhibit different tactics throughout their reproductive life, either as

they grow or in response to changing social context (plastic tactics or status-

dependent tactics) (Gross 1996; Taborsky 1996; Taborsky *et al*. 2008). Recent

advances in genome sequencing, such as RNA sequencing (RNAseq), now allow

behavioral ecologists to explore what genetic pathways contribute to behavioral

variation among mating tactics and examine if the pathways differ among species or

between individuals that exhibit fixed versus plastic tactics.

Over the past few years there have been numerous studies examining how

differences in gene expression correlate with behaviors adopted by male ARTs

(Aubin-Horth *et al*. 2005; Renn *et al.* 2008; Fraser *et al.* 2014; Schunter *et al.* 2014;

Stiver *et al.* 2015). Most of these studies have found a large number of genes within

the brain that vary in expression among tactics during mating. For example, a recent

study examining gene expression differences in the ocellated wrasse (*Symphodus*

ocellatus) found 1,048 differentially expressed genes when comparing sneakers to

two other male tactics (nesting and satellite) and to females (Stiver *et al.* 2015). In

the black-faced blenny (*Tripterygion delaisi*), RNAseq identified approximately 600

transcripts differentially expressed within the brains of sneaker versus territorial

males (Schunter *et al.* 2014). In a third study, approximately 2,000 transcripts were

differentially expressed between intermediately-sized sailfin molly (*Poecilia*

latipinna) males that primarily perform courtship behaviors compared to small

males that only perform sneaking behaviors (Fraser *et al.* 2014). Changes in social

context also led to a larger response (i.e. changes in gene expression) in

intermediate-sized males that show higher levels of tactic plasticity when compared

to small sneaker males (Fraser *et al*. 2014), suggesting that genes driving neural

response during mating may differ between plastic and fixed tactics.

With the increase in genomic studies examining differential gene expression

among male ARTs, there are a growing number of candidate genes suggested to

drive the behavioral differences among tactics. Schunter *et al*. (2014) proposed a list

of potential candidate genes based on a number of studies that included

gonadotropin releasing hormone (*gnrh*), arginine vasotocin (*avt*), cytochrome P450

family 19 subfamily A polypeptide 1 (*cyp19a1*), ependymin (*epd*), galanin (*gal*),

stomatostatin (*sstr1* and *sstr3*), and early growth response 1 (*egr1*). Many of these

genes are involved in hormone regulation and mating behavior, and differences in

expression levels have been observed among mating tactics in different fish species

(Table 1). For example, the product of the *cyp19a1b* gene is aromatase B, a key

enzyme responsible for the conversion of androgens to estrogens within radial glial

cells of adult fish (Forlano and Bass 2005; Le Page *et al.* 2010). *Cyp19a1* plays an

important role in sex determination and sex change in fish (Nakamura & Kobayashi

2005; Black *et al.* 2005; Marsh *et al.* 2006) and higher levels of gene expression have

been observed in territorial males compared to sneaker males in peacock blennies

(*Salaria pavo*) (Gonçalves *et al.* 2008), black-faced blennies (*Tripterygion delaisi*)

(Schunter *et al*. 2015), and an African cichlid (*Astatotilapia burtoni*) (Renn *et al.*

2008). As more genomic data become available, the number of candidate genes in

this list will likely increase and evaluating gene responses across taxa will aid in

determining whether similar genetic pathways drive ART behaviors across different

species.

One of the best-studied fish species with male ARTs is the bluegill sunfish

(*Lepomis macrochirus*). In this species, males have two distinct life histories:

parental and cuckolder. In Lake Opinicon (Ontario, Canada), parental males mature

at around seven years old and construct nests, court females, and provide care to

young (Gross 1982). Cuckolder males mature at a significantly younger age, around

two years old (Gross 1982). Rather than competing with parental males for access

to females, cuckolders initially use a "sneaking" tactic to dart in and out of nests

while parental males and females are spawning. As they grow, typically around an

age of 4 years, cuckolder males transition into "satellite" males by taking on female-

like coloration and behaviors (Dominey 1980; Gross 1982). Satellite males use this

female mimicry to deceive a parental male that he has two true females in his nest

(Neff & Gross 2001). The parental and cuckolder life histories are fixed – once a

male adopts the parental or cuckolder life history, he remains in that life history for

life (Gross & Charnov 1980). However, within the cuckolder life history, mating

tactics are ontologically plastic, with males apparently transitioning from the

sneaker tactic to the satellite tactic as they age (Gross & Charnov 1980).

While the spawning behavior, reproductive success, and hormone profiles of

bluegill have been studied extensively (Gross & Charnov 1980; Kindler *et al.* 1980;

Kindler *et al*. 1991; Neff 2001; Neff 2004; Knapp & Neff 2007), the genetic factors

influencing behavioral differences during spawning are less clear (Partridge *et al*

2015). Thus, for this study, we used RNAseq to characterize the brain transcriptome

of the three spawning male tactics (parental, sneaker, and satellite), in addition to

non-spawning parental males, to examine how differences in gene expression may

relate to behavioral variation among the tactics. Specifically, we (1) assessed

whether or not there is a greater difference in gene expression profiles between

fixed tactics (parental versus the two cuckolder tactics) than between tactics within

a plastic life history (sneaker versus satellite), (2) identified specific gene categories

that are expressed for each tactic, and (3) examined the expression of potential

candidate genes associated with ARTs from other fish species to determine if they

also differentiate the ARTs in bluegill.

Methods and Materials

Bluegill Sampling

In June 2013, bluegill sunfish were collected from Lake Opinion near Queen's

University Biological Station (QUBS), Elgin, Ontario, Canada. A total of 12 parental

males, 12 sneaker males, 13 satellite males, and 12 females were collected in the act

of spawning directly from the bluegill colony. An additional 12 non-nesting parental

males were collected four days prior to spawning (as determined once spawning at

these colonies began). Individuals were euthanized using clove oil, total body length

was measured, and brains were immediately dissected out and stored in RNAlater

(Life Technologies, Carlsbad, CA). Brains remained in RNAlater at 4°C for 24 hours

and were then transferred to fresh cryovials, flash frozen, and kept in liquid

nitrogen until they were transported on dry ice to the University of Western

Ontario. Samples were then stored at -80°C until RNA extraction.

Total RNA Extraction

Total RNA was extracted using a standard Trizol (Life Technologies,

Carlsbad, CA) extraction. RNA was submitted to the London Genomics Center at the

University of Western Ontario and quality was assessed using a 2100 Bioanalyzer

(Agilent Technologies, Palo Alto, CA). Four individuals from each group (spawning

parental males, non-spawning parental males, sneaker males, satellite males, and

females), for a total of 20 individuals, were submitted to the Michigan State

University Research Technology Support Facility - Genomics Center for cDNA

Library Construction and Sequencing. Individuals used for this study had RIN (RNA

Integrity Number) values ranging from 9.2-9.9.

cDNA Library Construction and Sequencing

The cDNA libraries were constructed for each individual using Illumina

TrueSeq Stranded mRNA Library Preparation Kits LT (Illumina, San Diego, CA), with

each individual receiving a uniquely identifiable index tag. The quality of each

library was evaluated and the 20 individuals were multiplexed into a single sample

that was subsequently run on two lanes of an Illumina HiSeq2500 Rapid Run flow

cell (v1). Sequencing was performed on paired end 2 x 150 bp format reads and

bases were called using Illumina Real Time Analysis software (v1.17.21.3). Reads

from each individual were identified based on their unique index tag, separated, and

converted to fastq files using Illumina Bcl2fastq v1.8.4. Sequencing produced an

average of 14.5 million reads per individual, with over 90% of the reads having a Q-

score >30.

De novo Transcriptome Assembly and Reference Transcriptome

Prior to assembly, read quality was assessed using FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc). Nucleotides whose

quality score was below PHRED=2 were trimmed using Trimmomatic version 0.32

(Bolger *et al.* 2014), following recommendations from MacManes (2014). The

reference transcriptome was assembled *de novo* using Trinity version 2.04 (Haas *et*

al. 2013, Grabherr *et al.* 2011). One representative of each of the five groups

(spawning parental male, non-spawning parental male, sneaker male, satellite male,

and female) was used to construct a combined reference transcriptome. The five

representatives selected for the reference were the individuals with the highest

number of reads within their group and, a total of 85 million paired-end reads were

assembled. The assembly was conducted with both normalized and non-normalized

reads and normalization was performed using Trinity's *in silico* normalization

program. To test the completeness of the transcriptome, reads from samples not

used in the assembly were mapped back to the transcriptome using Burrows-

Wheeler Aligner (bwa)-mem version 0.7.12 (Li 2013), and >90% of those reads

aligned, which is comparable to the rate of mapping for the individuals used in the

assembly (92%).

TransDecoder was used to identify protein-coding regions within the

assembled transcriptome. Transcripts that contained protein-coding regions or

transcripts that blasted to compete coding sequences (cds) and non-coding RNA

(ncRNA) from *Tetraodon nigroviridis, Lepisosteus oculatus, Xiphophorus*

maculatus, Oryzias latipes, Takifugu rubripes, Latimeria chalumnae, Astyanax

mexicanus, Danio rerio, or *Poecilia formosa* (downloaded from Ensembl) comprised

the reference transcriptome used for both read alignment and to estimate transcript

counts.

Read Alignment and Transcript Counts

Reads from each individual were separately aligned to the reference

transcriptome using bwa-mem 0.7.10 (Li 2013). At least 85% of all reads from each

individual mapped back to the reference, with the majority aligning 90% of reads or

higher. The sequence alignment/map (sam) files were then converted to a binary

format (bam) using Samtools 0.1.19 (Li *et al.* 2009). Transcript counts for each

individual were obtained using the program eXpress 1.5.1 (Roberts & Pachter

2013). Differential gene expression was determined using the R statistical package

edgeR 3.6.8 (Robinson *et al.* 2010). Low abundance transcripts were filtered out,

leaving 19,804 transcripts for differential analysis. Transcript counts were

normalized to account for differences in cDNA library size among individuals and

dispersion parameters were estimated using Tagwise dispersion estimates.

Differences in gene expression comparing paired treatments were calculated using

an Exact-test for binomial distribution. Genes with p-values lower than 0.05 after

false discovery rate (FDR) correction were determined to be statistically significant.

Multidimensional clustering analysis was used to cluster individuals together based

on the biological coefficient of variation.

Gene Annotation and Enrichment Analysis

Both the reference transcriptome and transcripts differentially expressed

among groups were blasted using Blastx against a custom-assembled fish protein

database. This database consisted of Ensembl protein databases of 13 different fish

species: Amazon molly (*Poecilia formosa*), zebrafish (*Danio rerio*), blind cave tetra

(*Astyanax mexicanus*), cod (*Gadus morhua*), coelancanth (*Latimeria chalumnae*),

Japanese pufferfish (*Takifugu rubripes*), sea lamprey (*Petromyzon marinus*), medaka

(*Oryzias latipes*), platyfish (*Xiphophorus maculatus*), spotted gar (*Lepisosteus*

oculatus), three-spined stickleback (*Gasterosteus aculeatus*), green-spotted

pufferfish (*Tetradon nigroviridis*), and Nile tilapia (*Oreochromis niloticus*). Blast hits

with e-values less than 1×10^{-10} were considered significant. Ensembl IDs from the

blast hits were then converted into GO term identifiers using Biology Database

Network (bioDBnet) (http://biodbnet.abcc.ncifcrf.gov/db/dbFind.php).

For purposes of gene annotation and enrichment analysis, we focused on

transcripts within the reference transcriptome that were not filtered out of the data

set due to low transcript expression (total of 19,804 transcripts). To examine which

GO terms were significantly enriched within this set, unique Ensembl IDs from

Blastx were converted to Ensemble IDs associated with stickleback homologs using

the R package biomaRt 2.20.0. Enrichment analysis was then conducted on the

homologs using the BioMart portal (http://central.biomart.org/enrichment).

For the transcripts that were differentially expressed among behavioral

groups, enrichment analysis was conducted using a Fisher Exact test to examine

whether the proportion of genes within each GO category was significantly higher

than what would be expected based upon the proportion of genes assigned to that

GO term within the reference transcriptome. To ensure adequate statistical power,

only GO terms with at least 10 transcripts within each category were included in the

statistical analysis. A FDR correction was applied to control for multiple testing and

GO terms with p-values < 0.05 were considered to be significant.

Results

<u>Reference Transcriptome</u>

This study presents the first reference transcriptome for the brain of bluegill

sunfish. The fully assembled transcriptome consisted of 272,189 transcripts. Of

these, 72,189 transcripts contained complete coding sequences or blasted to cds or

ncRNA from the customized Ensembl fish database. These 72,189 transcripts were

then used as the reference transcriptome for alignment and mapping. The mean

transcript length within the reference transcriptome was 2,024 bp, with N50 =

3,106 bp and N90 = 1,018 bp. The largest transcript consisted of 27,880 bp.

Approximately 82% of the transcripts had only one isoform, while 18% (12,951

transcripts) had two or more isoforms.

For GO enrichment analysis, we only examined the 19,804 transcripts within

the reference transcriptome that passed the filtering process. Of these, 18,108 had

significant Blastx hits with Ensembl gene IDs (Table S1, Supporting Information), of

which 12,224 transcripts had stickleback homologs that could be used to examine

GO term enrichment for bluegill sunfish compared to the stickleback genome. The

GO terms with significant enrichment included translation, catabolism, vesicle-

mediated transport, biosynthesis, small molecule metabolism, and generation of

precursor metabolites and energy (Fig. 1 A & B).

<u>Differential Gene Expression among Groups</u>

Based on the biological coefficient of variation, sneaker males grouped separately

from the other male tactics. However, non-spawning parental males, spawning

parental males, and satellite males showed a large amount of overlap in how

transcripts varied in their expression levels (Fig. 2). The largest number of

differentially expressed genes was observed when comparing spawning parental

males to sneaker males, followed by satellite males compared to sneaker males,

spawning parental males compared to satellite males, and then spawning parental

males compared to non-spawning parental males (Table 2). Analysis of sex

differences in differential expression will be presented in a companion paper

(Partridge et al. *in preparation*).

Differential Expression between Life Histories

<u>Spawning Parental Males versus Sneaker Males</u>. A total of 9,279 transcripts were

differentially expressed between parental males and sneaker males. Of these, 4,537

showed higher expression in parental males (Table S2, Supporting Information),

and 4,742 transcripts showed higher expression in sneaker males (Table S3,

Supporting Information). Enrichment analysis of GO terms associated with

differentially expressed genes showed that the biological processes most enriched

in parental males included translational initiation, translation elongation, and

oxidation-reduction processes (Table S4, Supporting Information). The molecular

function GO terms associated with translational processes included structural

constituents of the ribosome and translation initiation factor activity (Table S4,

Supporting Information). Biological processes enriched with genes displaying

higher expression in sneaker males included ion transport, hemophilic cell adhesion

(primarily due to protocadherin- and cadherin-related genes), the ionotropic

glutamate receptor signaling pathway, protein phosphorylation, and synaptic

transmission (Fig. 3A). Similarly, significantly enriched molecular function terms

included ionotropic glutamate receptor activity and protein binding (Table S4,

Supporting Information).

Spawning Parental Males versus Satellite Males. A total of 1,141 transcripts were

differentially expressed between parental males and satellite males. Of these, 676

displayed higher expression in parental males (Table S5, Supporting Information)

and 465 showed higher expression in satellite males (Table S6, Supporting

Information). Only one GO term related to biological processes was enriched in

parental males compared to satellite males and this was oxidation-reduction

processes. Significantly enriched molecular functions included iron ion binding,

oxidoreductase activity, and heme binding. Biological process terms enriched with

genes showing higher expression in satellite males included ion transport, and

enriched molecular function terms included nucleic acid binding, ion channel

activity, and GTP binding (Table S4, Supporting Information).

Differential Expression within Life Histories

Satellite Males verses Sneaker Males. There were 2,590 transcripts differentially

expressed between satellite males and sneaker males. Of these transcripts, 1,261

showed higher expression in satellite males (Table S7, Supporting Information), and

1,329 showed higher expression in sneaker males (Table S8, Supporting

Information). Biological processes enriched with genes displaying higher expression

in satellite males included translation, embryo development, and cell cycle

regulation. Molecular processes enriched in satellite males included structural

constituents of the ribosome, heme binding, and oxidoreductase activity (Table S4,

Supporting Information). Transcripts showing higher expression in sneaker males

were involved in biological processes related to ionotropic glutamate receptor

signaling pathways and mRNA processing (Fig. 3B). Molecular functions were

primarily related to ionotropic glutamate receptor activity and protein binding

(Table S4, Supporting Information).

Spawning Parental Males verses Non-Spawning Parental Males. A total of 137

transcripts were differentially expressed between spawning and non-spawning

parental males. The majority of these transcripts (132 transcripts) showed higher

expression in spawning males (Table S9, Supporting Information). Genes with the

highest expression in spawning males compared to non-spawning males were MHC

II antigen beta chain, cytosolic 5'-nucleotidase II (*nt5c2*), cAMP responsive element

modulator (*crem*), cysteine dioxygenase type 1 (*cdo1*), and an uncharacterized

protein. Only eight transcripts showed higher expression in non-spawning parental

males. These were nuclear receptor subfamily 1 group D member 4b (*nr1d4b*),

neuronal tyrosine-phosphoinositide-3-kinase adaptor 2 (*nyap2*), sphingosine-1-

phosphate receptor 4 (*s1pr4*), gamma-aminobutyric acid A receptor beta 3 (*gabrb3*),

and four uncharacterized proteins. Due to the limited number of transcripts

differentially expressed between these two groups, the number of transcripts

assigned to each GO term was too small to have adequate statistical power to

perform an enrichment analysis for this comparison.

Potential Candidate Genes Associated with ART Spawning Behavior

We observed differential expression in a number of transcripts previously

identified as potential candidate genes (described in Table 1) associated with

differences in ART spawning behaviors (Table 3). In our data set, the candidate

genes cytochrome P450 family 19 subfamily A polypeptide 1b (*cyp19a1b*),

ependymin (*epd*), and galanin (*gal*) showed higher expression in parental males

compared to sneaker males. *Epd* also had higher expression in satellite males

compared to sneakers. Early growth response 1 (*egr1*) showed higher expression in

both satellite and sneaker males relative to spawning parental males. Somatostatin

1 (*sstr1*) showed higher expression in sneaker males compared to satellite males,

but no differences in other comparisons between tactics. No differences in

expression related to gonadotropin releasing hormone (*gnrh*), arginine vasotocin

(*avt*), or somatostatin 3 (*sstr3*) were observed between any of our groups.

In addition to these previously identified candidate genes, transcripts that

displayed some of the highest differences in expression between parental males and

all other male phenotypes (including non-spawning males) were related to cytosolic

5'-nucleotidase II (*nt5c2*). Multiple isoforms were expressed, with log2 fold changes

ranging from 1.5 – 4.8 times higher in parental males compared to other male

groups (Fig. 4). Consistent with the finding for GO term enrichment, transcripts that

showed the highest levels of expression in sneaker males compared to other groups

were related to glutamate receptor genes, particularly AMPA ionotropic glutamate

receptors (Table S3, Supporting Information).

357   **Discussion**

Bluegill sunfish are a classic system for examining behavioral differences in

ARTs. In this study, we generated and assembled the first bluegill brain

transcriptome, and we identified candidate genes that contribute to differences in

male spawning tactics. The main differences in gene expression were found between

sneaker males when compared to the two other male tactics. Generally, sneaker

males showed higher expression in transcripts influencing neural activity, whereas

parental and satellite males exhibited higher expression in genes related to

translation and oxidoreductase activity.

One of our key findings is that a shared life history does not appear to be a

driving factor influencing similarity in gene expression in the brain of male tactics.

In bluegill, parental and cuckolder life histories are fixed, but within the cuckolder

life history, males transition from the sneaking to the satellite tactic as they age

(Gross 1982; Gross & Charnov 1991). Our data showed that, regardless of whether

comparisons were made across fixed (parental versus sneaker or parental versus

satellite) or plastic (sneaker versus satellite) tactics, sneaker males showed the

highest level of differentiation in gene transcription. Similar results have been

observed in the ocellated wrasse, *Symphodus ocellatus*, where sneaker males also

showed the greatest number of differentially expressed genes compared to nesting

and satellite males (Stiver *et al.* 2015). The expression differences in sneakers

compared to the other tactics in bluegill and the ocellated wrasse are likely partially

due to age because sneaker males are typically younger than satellite and parental

or territorial males. Indeed, a recent study in the short-lived fish *Nothobranchius*

furzeri found that genes related to translation elongation and ribosomal proteins are

up-regulated with age (Baumgart *et al*. 2014). Both translation elongation and

ribosomal proteins showed higher expression in parental and satellite males

compared to sneaker males in our data set. Additionally, the behaviors exhibited by

sneaker males during spawning differ in fundamental aspects from those of the

other male tactics. In the ocellated wrasse, for example, satellite and nesting males

cooperatively protect the nest from sneakers and other egg predators (Taborsky *et*

al. 1987); in bluegill, satellite and parental males associate closely with the female

throughout spawning, whereas sneakers dart in and out of the nest. These

differences in spawning tactics likely also contribute to the differences in gene

expression observed in the two studies. Thus, age and spawning tactic are

important contributors to gene expression patterns across ARTS, and life history is

not exclusively responsible for these differences.

Identifying distinct gene categories expressed by one ART type compared to

another provides information regarding the genes influencing behavioral

differences during spawning. Previous studies in sailfin mollies, *Poecilia latipinna,*

and Atlantic salmon, *Salmo salar*, indicate that sneaker males have increased

expression in genes related to neurotransmission and learning (Aubin-Horth *et al*.

2005; Fraser *et al*. 2014). We found that the GO terms consistently enriched in

bluegill sneaker males compared to both parental and satellite males were the

ionotropic glutamate signaling pathway and ionotropic gluatamate receptor activity.

Ionotropic gluatamate receptors are primarily excitatory neurotransmitter

receptors and play an important role in fast synaptic transmission (reviewed in

Lamprecht & LeDoux 2004). Two of these receptors, NMDA and AMPA, play

important roles in memory function and spatial learning (reviewed in Riedel *et al.*

2003). Blocking NMDA receptors impairs learning new spatial locations in goldfish

(Gómez *et al.* 2006). Futhermore, mice with impaired AMPA receptors, while

showing normal spatial learning, have impaired working spatial memory (i.e. their

ability to alter their spatial choice in response to changing environments is

impaired) (Reisel *et al.* 2002). We propose that increased expression of genes

related to spatial memory, particularly related to working spatial memory, could be

important for bluegill sneakers during spawning as they attempt to gain access to

nests while avoiding detection not only by the parental males, but also predators

that are common around the colony (Gross & MacMillian 1981). Bluegill sneakers

must also position themselves in close proximity to females so they can time sperm

release to coincide with female egg release (Stoltz & Neff 2006). Similarly, sailfin

molly sneakers, who also show enrichment in ionotropic glutamate related genes

(Fraser *et al.* 2014), probably benefit from increased working spatial memory

because they must successfully position themselves by the female for quick and

successful copulations. In this context, increased expression in gene pathways that

improve neural function related to working spatial memory are probably especially

beneficial to the sneaker tactic to increase their reproductive success.

There are a number of candidate genes that have been proposed to drive the

expression of alternative mating tactics (Schunter *et al*. 2015). In our study of

bluegill, we corroborate some of these candidates. For example, *cyp19a1b*, *epd*, and

gal had higher expression levels in parental males compared to sneaker males. The

expression patterns for all three genes are similar to what has been observed in

cichlids (Renn *et al*. 2008). In addition, expression of *epd* is lower in rainbow trout,

Oncorhynchus mykiss, males that use a sneaking tactic versus males that are

dominant and territorial (Sneedon *et al*. 2011), which is also consistent with our

findings. In contrast, the one candidate gene that responded opposite to

expectations was *egr1*. *Egr1* expression was lower in bluegill parental males

compared to sneaker or satellite males although previous work on cichlids found

that expression of this gene increases when subdominant males transition into

dominant males (Burmeister *et al*. 2005). However, *egr1* is an important

transcription factor involved in neural plasticity (Jones *et al*. 2001), so it may be

involved in regulating the switch from one tactic to another. Consequently, in

bluegill, this gene would be more important for individuals that alter their tactic

(sneaker to satellite) than for the fixed parental tactic. Taken together, our results

corroborate a role for *cyp19a1b*, *epd, gal, and egr1* as candidate genes contributing

to behavioral differences in ARTs across species.

We also found a transcript that may have a previously unrecognized function

in influencing male spawning behavior. Transcripts corresponding to splice variants

of cytosolic 5'-nucleotidase II (*nt5c2*) were significantly higher in parental males

when compared to all other male groups, including non-spawning males. The

protein product of *nt5c2* regulates purine metabolism (Bretonnet *et al.* 2005;

Walldén *et al.* 2007). Moreover, in the African cichlid, *A. burtoni*, uridine kinase

(*udk*) expression, a gene with similar function to *nt5c2*, is significantly higher in

dominant relative to subordinate males (Renn *et al.* 2008). While neither *nt5c2* nor

udk have been directly associated with spawning behavior in fishes, there is

evidence suggesting that altered expression levels of *nt5c2* in the brain can

significantly influence anxiety, mania, schizophrenia, and aggressive behaviors in

humans (Page *et al.* 2007), and altering levels of uridine in mice affects their level of

aggression toward an intruder (Kawasaki *et al.* 2013). Furthermore, in bluegill,

parental males display high levels of aggression to obtain nesting sites, circumvent

cuckoldry, and prevent egg predation by brood predators (Avila 1976; Colgan *et al.*

1979; Gross 1979; Gross & Macmillian 1981). The high levels of *nt5c2* expression in

spawning parental males suggests that this gene may have a role in influencing

parental male spawning behaviors in bluegill. Future work should examine how

nt5c2 influences mating behaviors in the ARTs of this and other species.

In summary, our work describes differences in gene expression profiles in

the brains of bluegill male ARTs during spawning. The largest differences in

expression levels were observed when comparing sneakers to parental and satellite

males, suggesting that, in bluegill, tactic is more related to differences in gene

expression than is life history. Consistent with other studies, our work demonstrates

that sneaker males have greater expression of genes involved in neural function

relative to more territorial-type males, particularly in relation to working spatial

memory, as mediated by ionotropic glutamate receptors. We found support for the

previously identified candidate genes *cyp19a1b*, *epd, gal, and egr1* contributing to

behavioral differences in ARTs, but we also show evidence for a novel candidate

gene, *nt5c2,* implicated in these differences. We suggest that *nt5c2* may have a role

in mediating courtship or territorial behaviors within this species, and we

recommend that future work should characterize this gene further in other species.

474   **Acknowledgments**

We thank Scott Colborne for his help in collecting bluegill, Dave Bridges for

providing the R script to convert Ensemble IDs to stickleback homologs, and David

Winter and Jeramia Ory for providing Python script used in the bioinformatics

analyses. We also thank Shawn Garner, Tim Hain, Lauren Kordonowy, and Lindsay

Havens for helpful comments on the manuscript. Funding for this project was

supported through the Natural Sciences and Engineering Research Council of

Canada (NSERC) to B.D.N. and the University of Oklahoma Department of Biology to

R.K.

References

Avila VL (1976) A field study of nesting behavior of male bluegill sunfish (*Lepomis macrochirus* Rafinesque). *American Midland Naturalist*, **96**, 195-206.

Aubin-Horth N, Landry CR, Letcher BH, Hofmann HA (2005) Alternative life histories shape brain gene expression profiles in males of the same population. *Proceedings of the Royal Society B: Biological Sciences*, **272**, 1655-1662.

Baumgart M, Groth, M, Priebe S, Savino A, Testa G, Dix A, Ripa R, Spallotta F, Gaetano C, Ori M, Tozzini ET, Guthke R, Platzer M, Cellerino A (2014) RNA-seq of the aging brain in the short-lived fish *N. furzeri* – conserved pathways and novel genes associated with neurogenesis. *Aging Cell*, **13,** 965-974.

Black MP, Balthazart J, Baillien M, Grober MS (2005) Socially induced and rapid increases in aggression are inversely related to brain aromatase activity in a sex-changing fish, *Lythrypnus dalli*. *Proceedings of the Royal Society of London B: Biological Sciences*, **272**, 2435-2440

Bloch GJ, Butler PC, Kohlert JG, Bloch DA (1993) Microinjection of galanin into the medial preoptic nucleus facilitates copulatory behavior in the male rat. *Physiology and Behavior*, **54**, 615-624.

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics*, btf170.

Bretonnet AS, Jordheim LP, Dumontet C, Lancelin JM (2005) Regulation and activity of cytosolic 5' nucleotidase II: A bifunctional allosteric enzyme of the Haloacid Dehalogenase superfamily involved in cellular metabolism. *FEBS Letters*, **579**, 3363-3368.

Burmeister SS, Jarvis ED, Fernald RD (2005) Rapid behavioral and genomic responses to social opportunity. *PLOS Biology*. e363

Colgan PW, Nowell WA, Gross MR, Grant JW (1979) Aggressive habituation and rim circling in the social organization of bluegill sunfish (*Lepomis macrochirus*). *Environmental Biology of Fishes*, **4**, 29-36.

Dominey WJ (1980) Female mimicry in male bluegill sunfish – a genetic polymorphism? *Nature*, **284**, 546-548.

Forlano PM, Bass AH (2005) Seasonal plasticity of brain aromatase mRNA expression in glia: divergence across sex and vocal phenotypes. *Neurobiology*, **65**, 37-49.

532   Fraser BA, Janowitz I, Thairu M, Travis J, Hughes KA (2014) Phenotypic and genomic
533   plasticity of alternative male reproductive tactics in sailfin mollies.
534   *Proceedings of the Royal Society B: Biological Sciences*, **281**, 20132310.
535

536   Gonçalves D, Teles M, Alpedrinha J, Oliveira RF (2008) Brain and gonadal aromatase
537   activity and steroid hormone levels in female and polymorphic males of the
538   peacock blenny *Salaria pavo. Hormones and Behavior*, **54**, 717-725.
539

540   Gómez Y, Vargas JP, Portavella M, López JC (2006) Spatial learning and goldfish
541   telencephalon NMDA receptors. *Neurobiology of Learning and Memory*, **85**,
542   252-262.
543

544   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
545   Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, di
546   Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A
547   (2011) Full-length transcriptome assembly from RNA-Seq data without a
548   reference genome. *Nature Biotechnology*, **29**, 644-652.
549

550   Greenwood AK, Wark AR, Fernald RD, Hofmann HA (2008) Expression of arginine
551   vasotocin in distinct preoptic regions is associated with dominant and
552   subordinate behaviour in an African cichlid fish. *Proceedings of the Royal
553   Society B: Biological Sciences*, **275**, 2393-2402.
554

555   Grober MS, George AA, Watkins KK, Carneiro LA, Oliveira RF (2002) Forebrain AVT
556   and courtship in fish with male alternative reproductive tactics. *Brain
557   Research Bulletin*, **57**, 423-425.
558

559   Gross MR (1979) Cuckoldry in sunfishes (*Lepomis*: Centrarchidae). *Canadian Journal
560   of Zoology*, **57**, 1507-1509.
561

562   Gross MR (1982) Sneakers, satellites and parentals: Polymorphic mating strategies
563   in North American sunfishes. *Zeitschrift für Tierpsychologie*, **60**, 1-26.
564

565   Gross MR (1996) Alternative reproductive strategies and tactics: diversity within
566   sexes. *Trends in Ecology and Evolution*, **11**, 92-98.
567

568   Gross MR, Charnov EL (1980) Alternative male life histories in bluegill sunfish.
569   *Proceedings of the National Academy of Sciences USA*, **77**, 6937-6940.
570

571   Gross MR, MacMillian AM (1981) Predation and the evolution of colonial nesting in
572   bluegill sunfish (*Lepomis macrochirus*). *Behavioral Ecology and Sociobiology*,
573   **8**, 163-174.
574

575   Gross MR, Repka J (1998) Stability with inheritance in the conditional strategy.
576   *Journal of Theoretical Biology*, **192**, 445-453.
577

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB,
MacManes MD, Ott M, Orvis J, Pochet N, Strozzi F, Weeks N, Westerman R,
William T, Dewey CN, Henschel R, LeDuc RD, Friedman N, Regev A (2013) *De*
Novo transcript sequence reconstruction from RNA-Seq using the Trinity
platform for reference generation and analysis. *Nature Protocols*, **8,** 1494–
1512.

Jones MW, Errington ML, French PJ, Fine A, Bliss TVP, Garel S, Charnay P, Bozon B,
Laroche S, Davis S (2001) A requirement for the immediate early gene Zif268
in the expression of LTP and long-term memories. *Nature Neuroscience*, **4**,
289-296.

Kindler PM, Philipp DP, Gross MR, Bahr JM (1989) Serum 11-ketotestosterone and
testosterone concentrations associated with reproduction in male bluegill
(*Lepomis macrochirus*: Centrarchidae). *General and Comparative*
Endocrinology, **75**, 446–453.

Kindler PM, Bahr JM, Philipp DP (1991) The effects of exogenous 11-
ketotestosterone, testosterone, and cyproterone acetate on prespawning and
parental care behaviors of male bluegill. *Hormones and Behavior*, **25**, 410–
423.

Kawasaki A, Suzuki N, Endo K, Ashida N (2013) Novel effects of uridine on
behavioral changes due to social isolation stress in mice. *Food Science and*
Technology Research, **19**, 455-461.

Knapp R, Neff BD (2007) Steroid hormones in bluegill, a species with male
alternative reproductive tactics including female mimicry. *Biology Letters,* **3**,
628-631.

Lank DB, Smith CM, Hanotte O, Burke T, Cooke F (1995) Genetic polymorphism for
alternative mating behaviour in lekking male ruff *Philomachus pugnax.*
Nature, **378**, 59-62.

Lamprecht R, LeDoux J (2004) Structural plasticity and memory. *Nature Reviews*
Neuroscience, **5**, 45-54.

Le Page Y, Diotel N, Vaillant C, Pellegrini E, Anglade I, Mérot Y, Kah O (2010)
Aromatase, brain sexualization and plasticity: the fish paradigm. *European*
Journal of Neuroscience, **32**, 2105-2115.

Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. arXiv:1303.3997v2 [q-bio.GN]

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R, 1000 Genome Project Data Processing Subgroup (2009) The

Sequence alignment/map (SAM) format and SAMtools. *Bioinformatics*, **25**,
2078-2079.
626

627 MacManes MD (2014) On the optimal trimming of high-throughput mRNA sequence
628   data. *Frontiers in Genetics,* **5**, 13. doi:10.3389/fgene.2014.00013.
629

630 Mank JE, Avise JC (2006) The evolution of reproductive and genomic diversity in
631   ray-finned fishes: insights from phylogeny and comparative analysis. *Journal*
632   *of Fish Biology*, **69**, 1-27.
633

634 Marsh KE, Creutz LM, Hawkins MB, Godwin J (2006) Aromatase immunoreactivity in
635   the bluehead wrasse brain, *Thalassoma bifasciatum*: Immunolocalization and
636   co-regionalization with arginine vasotocin and tyrosine hydroxylase. *Brain*
637   *Research*, **1126**, 91-101.
638

639 Nakamura M, Kobayashi Y (2005) Sex change in coral reef fish. *Fish Physiology and*
640   *Biochemistry*, **31**, 117-122.
641

642 Neff BD (2001) Genetic paternity analysis and breeding success in bluegill sunfish
643   (*Lepomis macrochirus*). *Journal of Heredity*, **92**, 111-119.
644

645 Neff BD (2004) Increased performance of offspring sired by parasitic males in
646   bluegill sunfish. *Behavioral Ecology*, **15**, 327-331.
647

648 Neff BD, Gross MR (2001) Dynamic adjustment of parental care in response to
649   perceived paternity. *Proceedings of the Royal Society of London B*, **268**, 1559-
650   1565.
651

652 Neff BD, Svensson EI (2013) Polyandry and alternative mating tactics. *Philosophical*
653   *Transactions of the Royal Society of London B*, **368**, 20120045.
654

655 Page T, Yu A, Fontanesi J, Nyhan WL (1997) Developmental disorder associated with
656   increased cellular nucleotidase activity. *Proceedings of the National Academy*
657   *of Sciences USA*, **94**, 11601-11606.
658

659 Piché J, Hutchings JA, Blanchard W (2008) Genetic variation in threshold reaction
660   norms for alternative reproductive tactics in Atlantic salmon, *Salmo salar*.
661   *Proceedings of the Royal Society B: Biological Sciences*, **275**, 1571-1575.
662

663 Reisel D, Bannerman DM, Schmitt WB, Deacon RMJ, Flit J, Borchardt T, Seeburg PH,
664   Rawlins JNP (2002) Spatial memory dissociations in mice lacking GluR1.
665   *Nature Neuroscience*, **5**, 868-873.
666

Renn SC, Aubin-Horth N, Hofmann HA (2008) Fish and chips: functional genomics of
social plasticity in an African cichlid fish. *Journal of Experimental Biology*,
211, 3041-3056.

Riedel G, Platt B, Micheau J (2003) Glutamate receptor function in learning and
memory. *Behavioural Brain Research*, **140**, 1-47.

Roberts A, Pachter L (2013) Streaming fragment assignment for real-time analysis
of sequencing experiments. *Nature Methods*, **10**, 1-47.

Robinson MD, McCarthy DJ, Smyth GK (2010) EdgeR: a Bioconductor package for
differential expression analysis of digital gene expression data.
Bioinformatics, **26**, 139-140.

Schunter C, Vollmer SV, Macpherson E, Pascual M (2014) Transcriptome analysis
and differential gene expression in a non-model fish species with alternative
mating tactics. *BMC Genomics*, **15**, 167.

Schuster SM, Sassaman C (1997) Genetic interaction between male mating strategy
and sex ratio in a marine isopod. *Nature*, **388**, 373-377.

Sneddon LU, Schmidt R, Fang Y, Cossins AR (2011) Molecular correlates of social
dominance: A novel role for ependymin in aggression. *PLoS One*. **6**, e18181.

Stiver KA, Harris RM, Townsend JP, Hofmann HA, Alonzo SH (2015) Neural gene
expression profiles and androgen levels underlie alternative reproductive
tactics in the ocellated wrasse, *Symphodus ocellatus. Ethology*, **121**, 152-167.

Stoltz JA, Neff BD (2006) Male size and mating tactic influence proximity to females
during sperm competition in bluegill sunfish. *Behavioral Ecology and
Sociobiology*, **59**, 811-818.

Taborsky M (1998) Sperm competition in fish: 'Bourgeois' males and parasitic
spawning. *Trends in Ecology and Evolution*, **13**, 222-227.

Taborsky M, Brockmann HJ (2010) Alternative reproductive tactics and life history
phenotypes. In: *Animal Behavior: Evolution and Mechanisms* (Ed. P. Kappeler).
Springer, Berlin, 537-586.

Taborsky M, Hudde B, Wirtz P (1987) Reproductive behaviour and ecology of
Symphodus (Crenilabrus) *ocellatus*, a European wrasse with four types of
male behaviour. *Behaviour*, **102**, 82-117.

Taborsky M, Oliviera RF, Brockmann HJ (2008) The evolution of alternative
reproductive tactics: concepts and questions. In: *Alternative Reproductive
Tactics: An Integrative Approach* (Eds. Oliveira RF, Taborksy M, Brockmann
HJ) Cambridge University Press, 1-21.

Thomson JS, Watts PC, Pottinger TG, Sneddon, LU (2011) Physiological and genetic
correlates of boldness: Characterizing the mechanisms of behavioral
variation in rainbow trout, *Oncorhynchus mykiss*. *Hormones and Behavior*, **59**,
67-74.
719
720 Walldén K, Stenmark P, Nyman T, Flodin S, Gräslund S, Loppnau P, Norlund P
721   (2007). Crystal structure of human cytosolic 5'-nucleotidase II. Insights into
722   allosteric regulation and substrate recognition. *Journal of Biological*
723   *Chemistry*, **282**, 17828-17836.
724
725 Wu Z, Autry AE, Bergan JF, Watabe-Uchida M, Dulac CG (2014) Galanin neurons in
726   the medial preoptic area govern parental behavior. *Nature*, **509**, 325-330.
727
728

729   **Author Contribution**

730 C.G.P, R.K., and B.D.N. designed the experiment and collected samples. M.D.M.

731 assembled the transcriptome. C.G.P. performed bioinformatic and statistical

732 analyses and wrote the manuscript. All authors provided comments, contributed to

733 the manuscript, and approved the final manuscript.

735   **Data Accessibility**

736   All raw sequence files are available on the Sequence Read Archive (SRA)

737 through BioProject ID: PRJNA287763. Environmental data, RNA quality information,

738 assembled transcriptome, the transcript count matrix, and R code for differential

739 gene analysis are available on Dryad (http://dx.doi.org/10.5061/dryad.82fd8 and

740 http://dx.doi.org/10.5061/dryad.10hh7).

742   **Supporting Information**

743 Table S1: Annotated Reference Transcriptome
744
Table S2: Transcripts with significantly higher expression in bluegill parental males
compared to sneaker males.

Table S3: Transcripts with significantly higher expression in bluegill sneaker males
compared to parental males.

Table S4: Biological Process and Molecular Function GO terms that are significantly
enriched with genes differentially expressed between tactics

Table S5: Transcripts with significantly higher expression in bluegill parental males
compared to satellite males.

Table S6: Transcripts with significantly higher expression in bluegill satellite males
compared to parental males.

Table S7: Transcripts with significantly higher expression in bluegill satellite males
compared to sneaker males.

Table S8: Transcripts with significantly higher expression in bluegill sneaker males
compared to satellite males.

Table S9: Transcripts with significantly higher expression in spawning parental
males compared to non-spawning parental males.

Table 1: Proposed candidate genes (Schunter *et al*. 2014) influencing teleost alternative reproductive
tactics (ARTs). POA = Pre-optic area

Proposed Candidate Genes	Function	Relationship to ARTs
Arginine Vasotocin (*avt*)	Non-mammalian homolog of vasopressin. Activates some aspects of sexual behavior	↑ in posterior POA of territorial cichlid males, but ↑ anterior POA of non-territorial (Greenwood *et al*. 2008); ↓ density of *avt* mRNA in POA in parental blenny males (Grober *et al*. 2002)
Gonadotrophin Releasing Hormone (*gnrh*)	Regulates release of lutenizing hormone and follicle-stimulating hormone from the pituitary gland	↑ in territorial cichlid males (Renn *et al*. 2008)
Cytochrome P450 family 19, subfamily A, polypeptide 1 (c*yp19a1*)	Brain aromatase. Key enzyme in estrogen biosynthesis	↑ in territorial cichlid males (Renn *et al*. 2008); ↑ territorial blenny males (Gonçalves *et al*. 2008); ↑ territorial black-faced blenny males (Schunter *et al.* 2014); ↓ in the sonic motor nucleus of nesting type 1 (territorial) male plainfin midshipmen compared to type II (female mimic) males (Forlano *et al.* 2005)
Ependymin (*epd*)	Glycoprotein associated with neuroplasticity and neuronal regeneration. Also affects aggression levels in zebrafish (Sneddon *et al* 2011); associated with stress in trout (Thomson *et al.* 2011)	↑ in territorial cichlid males (Renn *et al.* 2008); ↓ in subordinate trout males (Sneddon *et al.* 2011)
Galanin/GMAP prepropeptide (*gal*)	Neuropeptide that influences neurotransmitters. Associated with sexual behaviors (Bloch *et al.* 1993), and parental care (Wu *et al.* 2014)	↑ in territorial cichlid males (Renn *et al.* 2008)
Somatostatin (*sst*)	Neuropeptide that regulates endocrine pathways. Also affects neurotransmitters	↑ in territorial blenny males (Schunter *et al.* 2014); ↑ in territorial cichlid males (Renn *et al*. 2008)
Early growth response 1 (*egr1*)	Transcripton factor that influences neural plasticity	↑ when subdominant cichlid males switch to dominant (Burmeister *et al.* 2005)

778   Table 2: Number of transcripts differentially expressed (DE) between each male bluegill tactic,
779   including the number of transcripts without Blastx hits, the number with unique Ensembl IDs, and
780   the number of transcripts assigned to specific GO terms

Comparisons	Num of DE Transcripts	Num of DE Transcripts without Blastx Hits	Num with Unique Ensembl Gene IDs	Num DE Gene IDs with GO Annotation
Parental vs Sneaker	9,279	516	5,396	2,430
Parental vs Satellite	1,141	82	879	317
Satellite vs Sneaker	2,590	184	1,852	351
Parental vs Non-Spawner	140	6	102	70

782 Table 3: Gene expression differences (Log2 fold change) among male tactics for proposed candidate
783 genes (see Table 1).

Proposed Candidate Genes	Isoform ID	Comparison between Male Tactics (Log2 Fold Change)			
		Parent vs Sneak	Parent vs Sat	Sat vs Sneak	Non-Spawn vs Spawn Parent
Arginine Vasotocin (*avt*)	c34708_g2_i1	0.45 [0.32]	-0.98 [0.09]	0.54 [0.33]	-0.74 [0.5]
Gonadotrophin Releasing Hormone *(gnrh)*	63124_g1_i1	0.76 [0.5]	0.32 [0.87]	0.44 [0.77]	-0.77 [0.88]
Cytochrome P450 19a 1b (c*yp19a1b*)	c48084_g2_i1	**0.93 [0.0002]**	0.64 [0.06]	0.28 [0.4]	-0.39 [0.58]
Ependymin *(epd)*	c44195_g1_i5	**1.54 [1.4 x 10⁻⁸]**	0.66 [0.07]	**0.89 [0.007]**	0.51 [0.45]
Galanin/GMAP prepropeptide (*gal*)	c41071_g5_i2	**1.12 [0.0001]**	0.53 [0.91]	-0.59 [0.1]	-0.09 [0.97]
Somatostatin 1 *(sstr1)*	c3001_g1_i1	0.53 [0.15]	-0.39 [0.49]	**0.93 [0.03]**	-0.27 [0.88]
Somatostatin 3 (*sstr3*)	c46547_g6_i1	0.001 [1]	-0.25 [0.54]	0.25 [0.48]	-0.15 [0.9]
Early growth response 1 (*egr1*)	c37907_g1_i1	**-0.74 [0.02]**	**-0.91 [0.03]**	0.16 [0.72]	0.63 [0.42]

784
785 Values in brackets represent p-values after false discovery rate correction. Values in bold are
786 significant at $p < 0.05$. Parent = parental male, Sneak = sneaker male, Sat = satellite male.
787
788

Reference Transcriptome

A) Enriched Biological Processes

B) Enriched Molecular Functions

789
790 Fig. 1: GO terms related (A) biological processes and (B) molecular function that
791 were significantly enriched in the bluegill reference transcriptome relative to the
792 stickleback genome. Boxes of similar color can be grouped into the same GO term
793 hierarchy. The size of each box reflects the –log10 p-value of the GO term.

813
814
815 Fig. 2: Multi-dimensional space (MDS) plot based on the biological coefficient of
816 variation (bcv) among bluegill male ARTs. Red triangles: sneaker males, green
817 pluses: satellite males, black circles: spawning parental males, blue x: non-spawning
818 parental males.

825
826
827 Figure 3: GO terms significantly enriched by genes with higher expression in
828 sneaker males compared to (A) parental males and (B) satellite males. Boxes of
829 similar color are grouped into the same GO term hierarchy. Box size reflects the –
830 log10 p-value of the GO term.

849
850 Figure 4: Log2 fold changes of sneaker, satellite, and non-spawning (NS) parental
851 males relative to spawning parental males for cytosolic 5'-nucleotidase II (*nt5c2*). *
852 indicates fold changes that are significantly different with p-values < 0.05 after FDR
853 correction.
854