1   **Original Article**

2   **Title:** Species range expansion constrains the ecological niches of

3   resident butterflies.

Running header: Range expansion and changes in species interactions.

Authors:

Hélène Audusseau, Department of Zoology, Stockholm University, Stockholm, Sweden.

Maryline Le Vaillant, Department of Zoology, Stockholm University, Stockholm, Sweden.

Niklas Janz, Department of Zoology, Stockholm University, Stockholm, Sweden.

Sören Nylin, Department of Zoology, Stockholm University, Stockholm, Sweden.

Bengt Karlsson, Department of Zoology, Stockholm University, Stockholm, Sweden.

Reto Schmucki, Muséum National D'Histoire Naturelle, 55 rue Buffon, CP 51 75005 PARIS,
Centre d'Ecologie et des Sciences de la Conservation, Sorbonne Universités, MNHN-CNRS-
UPMC, Paris, France & Centre de Synthèse et d'Analyse sur la Biodiversité, Aix-en-Provence,
France.

Correspondence:

Hélène Audusseau, Department of Zoology, Svante Arrhenuis väg 18B, Stockholm University, 106
91 Stockholm, Sweden.

Phone: +468 164058

E-mail: helene.audusseau@zoologi.su.se

Word count: abstract 295 words; main body of the text 6959 words.

ABSTRACT

Aim Changes in community composition resulting from environmental changes modify biotic

interactions and affect the distribution and density of local populations. Such changes are currently

occurring in nettle-feeding butterflies in Sweden where *Araschnia levana* has recently expanded

its range northward and is now likely to interact with the resident species (*Aglais urticae* and *Aglais*

io). Butterfly occurrence data collected over years and across regions enabled us to investigate

how a recent range expansion of *A. levana* may have affected the environmental niche of resident

species.

Location We focused on two regions of Sweden (Skåne and Norrström) where *A. levana* has and

has not established, and two time-periods (2001-2006, 2009-2012) during its establishment in

Skåne.

Methods We performed two distinct analyses in each region using the PCA-env and the framework

described in Broennimann *et al*. (2012). First, we described the main sources of variation in the

environment. Second, in each time-period and region, we characterized the realized niches of our

focal species across topographic and land use gradients. Third, we quantified overlaps and

differences in realized niches between and within species over time.

Results In Skåne, *A. levana* has stabilized its distribution over time while the distribution of the

native species has shifted. These shifts depicted a consistent pattern of avoiding overlap between

the native species and the environmental space occupied by *A. levana,* and it was stronger for *A.*

urticae than for *A. io.* In both regions, we also found evidence of niche partitioning between native

species.

Main conclusions Interspecific interactions are likely to affect local species distributions. It

appears that the ongoing establishment of *A. levana* has modified local biotic interactions, and

induced shifts in resident species' distributions. Among the mechanisms that can explain such

patterns of niche partitioning, parasitoid-driven apparent competition may play an important role

in this community.

KEYWORDS biotic interactions, citizen science, community composition, environmental niche

model, nettle-feeding butterflies, ordination technique, parasitoid-driven indirect competition,

realized niche.

INTRODUCTION

Changes in community composition resulting from environmental changes modify biotic

interactions that are likely to affect the distribution and the density of local populations. To better

predict the widely recorded species' geographical and environmental shifts (Parmesan, 2006) it is

crucial to first define and understand species' environmental niches and how they are shaped by

local biotic interactions (Davis *et al.*, 1998; Tylianakis *et al.*, 2008; Gilman *et al.*, 2010).

Investigations of species co-occurrence across the landscape can provide useful insights for better

understanding the effects of species interactions on their distributions, and how community

composition is maintained or changed locally in a context of global change.

Northern regions offer a highly suitable model system to investigate how changes in species

interactions can induce shifts in realized niches. At high latitudes, the impacts of climate change

are most pronounced (IPCC, 2014), making it more likely to detect a signature of species'

responses to recent warming. Moreover, as species' northern ranges vary greatly and their

poleward shifts are not synchronized (Pöyry et *al.*, 2009), changes in community composition are

expected to alter interspecific interactions (González-Megías et *al.*, 2008; Devictor et *al.*, 2012)

and potentially lead to new interactions.

A good way to examine niche shifts and niche partitioning is to measure and test the overlap and

the difference in the environmental space occupied (i.e. realized niche) over time and across

species (Warren et *al.*, 2008; Broennimann et *al.*, 2012). Recent developments in statistical

methods and computational techniques have enabled better estimations of species-environment

relationships and thereby contributed to identify the processes and factors shaping species' realized

niches. In particular, Environmental Niche Models (ENMs) have been used to model and predict

species distributions according to changes in climatic and environmental variables, considered to

be the main drivers of species distribution at large and small spatial-scales (Berry et *al.*, 2002;

Thuiller et *al.*, 2005). However, ENMs that are based on the relationship between species

distribution and abiotic factors have often shown some discrepancy between the potential and the

realized niches. This strongly suggests that other processes such as biotic interactions have

substantial effect on species distribution and should be accounted for (Leathwick, 1998; Pellissier

et *al.*, 2012; Tingley et *al.*, 2014). Better consideration of biotic interactions in distribution models

assessing geographic ranges has greatly refined our predictions at both small and large scales

(Heikkinen et *al.*, 2007; Araújo & Luoto, 2007). Yet, even though ENMs and ordination techniques

have been successfully used to further our understanding of the role of biotic interactions by

comparing the realized niches of species (Schweiger et *al.*, 2012; Mason et *al.*, 2014), they have

rarely been used to understand how changes in local biotic interactions may induce shifts in species

realized niches (Wisz *et al.*, 2013). Furthermore, the reliability and accuracy of statistical models

in predicting the importance of biotic interactions are often limited by the availability of extensive

occurrence data, which are also often biased towards particular groups of species.

In the last decades, the amount of data collected, organized, and made available through public

databases has increased substantially. The use of such databases comes nonetheless with important

challenges as they cumulate data collected with no standardized sampling design and by observers

with different levels of expertise. Therefore, data contained in such databases have the drawback

of being prone to show biases in the region and habitat covered, lacking independence between

replicates, and having no explicit measures of sampling effort. However, considering the rapid

expansion of programs collating data through volunteer contribution of citizen observers, the real

potential of these large datasets is growing and many of them remain largely unexploited. This has

led to increased efforts being made to develop robust approaches to estimate and compare the

realized niche from occurrence and spatial environmental data, independently of the spatial

resolution and sampling biases that are often inherent to species occurrence data. In this context,

Broennimann *et al.* (2012) developed and tested an analysis framework to quantify niche overlap

and test for niche equivalency and similarity (*cf.* Warren *et al.*, 2008), using Principal Component

Analysis to define the environmental space (PCA-env). In contrast to ENM methods, the PCA-env

has been shown to be more reliable for defining the environmental space when tested on both

simulated and real case data (Broennimann *et al.*, 2012).

Here, we investigate the realized niche of three nettle-feeding butterflies (*Aglais urticae, Aglais*

io, and *Araschnia levana*) and examine how they vary in space and time as an illustration of the

potential role of interspecific interactions in shaping species distributions in the context of ongoing

climate change. We use occurrence data available through the internet reporting system

Artportalen (www.artportalen.se), a public database of species records based on citizens'

contribution in Sweden, to explore niche partitioning of nettle-feeding butterflies at the local scale.

In Sweden, *A. urticae* and *A. io* are common native species, easy to identify, and well represented

in the database with a large number of records available across the country (over 15 000 records

per species for the period 2001-2012). On the other hand, *A. levana,* for which the first anecdotal

observation reported in Sweden is from 1982 (Eliasson *et al*., 2005), is known to be expanding its

range northward and is now well established in the southern part of Sweden (Betzholtz *et al.*,

2013). All three species are specialists with larvae feeding on stinging nettle, *Urtica dioica,* and

showing overlapping phenology (see Fig. S1 in Appendix S1 in Supporting Information). While

no obvious direct competition has been documented, the three species are known to share common

parasitoids (Hinz & Horstmann, 2007; Shaw *et al.*, 2009), which increase the potential for apparent

competition among the species (van Veen *et al.*, 2006; Tylianakis, 2009).

Hence, the ongoing establishment of *A. levana* in southern Sweden makes this system a unique

opportunity to investigate the impact of such change on the distribution of native butterfly species

over a relatively short time period. If interspecific interactions are important determinants of

spatial distribution across the landscape, their signature should be detectable in the occupancy

pattern of the available environmental space. To model the realized niches of three sympatric

species and quantify the degree of niche overlap between them, we used the PCA-env method

applied within Broennimann's analysis framework (Broennimann *et al.*, 2012). We gathered high-

resolution data on land use and topography, and an index of nitrogen and phosphorus flows in the

soil, which reflects agricultural practices. We perform these analyses over two time-periods during

the establishment of *A. levana* and two geographically distinct regions (where *A. levana* has and

has not yet established), with the aim of assessing the potential effect of the establishment of *A.*

levana in the southern part of the country during the second period, while controlling for climatic

variability over time.

Specifically, as a recent colonizer of Sweden, *A. levana* is still in its initial phase of establishment

and, therefore, its distribution is expected to not be at equilibrium. Moreover, if changes in local

biotic interactions are important in this community, we expect the establishment of *A. levana* to

affect the distribution of the native species. We put forward three alternative mechanisms that, in

our study system, may explain the observed changes in distribution and density of the local

populations. We also discuss the opportunities and limitations of using occurrence records from

citizen-science databases and the reliability of the interpretation of the output results from the

method used.

MATERIALS AND METHODS

We studied the environmental space occupied by *A. urticae, A. io*, and *A. levana* in two regions,

corresponding to the county of Skåne in Southern Sweden and the Norrström drainage basin in

Central Sweden (including four counties: Södermanland, Stockholm, Uppsala, Västmanland, Fig.

1a), and two time-periods (first period 2001-2006 and second period 2009-2012). The regions are

separated by approximately 400 kilometers in a straight line. By aggregating the records over

specific time-periods, we aimed to have better estimates of species distributions, assuming these

patterns to remain relatively stable over short time-periods (i.e. to avoid yearly variation). Data

extraction was based on a regular grid covering both regions with a resolution of 1 km.

Species distribution is determined by the interplay of multiple variables operating at different

spatial scales (Pearson & Dawson, 2003). While at large scales, species distribution is limited by

physiological constraints mainly determined by climatic conditions, at small scales we expect their

distribution to be shaped by local variables such as land use and topography (Pearson & Dawson,

2003), as well as by direct and indirect biotic interactions (e.g. predation, competition, resources).

Considering the extent of our study areas, i.e. two regions of Sweden, we expected local-scale

variables, namely topography and land use, to be the most relevant factors for investigating niche

differences between species and potential niche shifts.

Occurrence Data

We extracted occurrence data for *A. urticae, A. io* and *A. levana* from the Swedish Species

Information Centre at SLU (www.artportalen.se, accessed on the 18/09/2015, average precision

reported for the data ± se of 168m ± 2m), a public database of species records in Sweden. The

Artportalen database gathers opportunistic occurrence data (presence only) collected at 90% by

amateurs with no specific required training in species identification (Gärdenfors *et al.*, 2014) and

who do not follow any specific protocol. For each species, we identify all grid cells for which the

species was recorded in each of the two time-periods and the two study regions.

For each period, we only considered species' presence across the selected grid cells, without

accounting for variation in the number of observations per cell as our main interest was to

determine species' occurrence patterns across the total available environment. In this way, we

reduced the potential bias associated with uneven sampling effort across sites (i.e. over-sampling

of more frequently visited sites). As a result of an overall growing interest for citizen science, both

the number of observations and the corresponding number of grid cells visited increased between

the first and the second period (Table S1 in Appendix S1). While the number of records is likely

to affect the extent of the environmental space sampled (surveyed), we are confident that the

sampling (reporting) effort for our study species was sufficiently high to limit this potential bias.

For the period 2009-2012, *A. urticae* and *A. io* are the first and fourth most reported butterfly

species in Artportalen, respectively, and they were in the top fifteen most reported species

according to the Swedish Butterfly Monitoring Scheme (records along transects, Petterson *et al.*,

2013). As a recently established species in Sweden, *A. levana* may still be in its initial phase of

establishment. Hence, its current distribution might not be at equilibrium yet or only reflect a

subset of its potential environmental niche. In our final dataset, occurrence of *A. urticae* was

recorded in 2935 grid cells, *A. io* in 2457, and *A. levana* in 599. Note that *A. levana* was only

recorded in Skåne.

Topographic Data

Aspect and slope were calculated based on a digital elevation map at an original 50m resolution

obtained from the Swedish University of Agricultural Science (https://maps.slu.se/get/). From the

original elevation map, we extracted the mean aspect and mean slope with the r.slope.aspect

function available in the GRASS GIS plugin for Quantum GIS 1.8 software (2012) and

recalculated both metrics for the 1km resolution grid, using PostgreSQL 9.4 and its spatial

extension PostGIS 2.1 (2014).

Land use Data

Land use data were collected from Naturvårdsverket at an original 25m resolution

(https://www.naturvardsverket.se/). The land use classification followed the Corine Land Cover.

We extracted the percentage of the different types of land use (forest, open land, arable land, non-

intensive agriculture, water body, and urban land) at 1km resolution grid cells. We also used

estimates of soil nutrient flow (in nitrogen and phosphorus) – which we used as a proxy for the

amount of nutrients available for plant root absorption – accessible from the SMED (Svenska Miljö

Emission Data, http://www.smed.se/) at an original resolution of the sub-catchment (municipality).

These estimates were modeled by means of simulation tools as well as measured data and

correspond to nitrogen and phosphorus loads to the water from diffuse sources across the whole

sub-catchment (Brandt *et al.*, 2009). Nutrient loads to the water from the sub-catchment result

from a combination of the run-off and leaching on the basis of assumptions on type-specific

concentrations for each type of land use. From these estimates, we calculated nutrient flow through

the 1km grid cells.

Realized Niche Shifts and Overlaps

We first extracted two axes that captured the maximum variation in land use and topography

available to the species, using the PCA-env ordination method described in Broennimann *et al*.

(2012). Second, we characterized, for each time-period and region, the realized niches of our study

species in the environmental space defined by the topographic and land use gradients extracted

above. Third, we quantified overlaps and differences for each species over time and between

species in each period in how they distributed themselves in this environment, using 100% of the

environmental space. All analyses were computed with the ecospat package (Broennimann *et al.*,

2015) in R 3.1.3 (R Core Team, 2015). To reduce the risk of generating false absences in un-

sampled areas due to the non-systematic sampling process that characterized our dataset, we

excluded from our analyses all grid cells where none of the focal species has been recorded, for

each time-period.

Following the framework proposed by Broennimann *et al.* (2012), we computed a weighted

Principal Components Analysis (PCA) on the environmental variables, after applying a kernel

density function to the number of sites of each specific environmental condition. The Gaussian

kernel density function is used to create a probability density function of each of the environmental

conditions available and of the occurrence of each species for each cell of the environmental space

(Broennimann *et al.*, 2012). We performed two distinct analyses in each region to prevent

environmental differences inherent to these regions (mostly in agricultural activity) to mask the

components of most interest for detecting niche partitioning occurring within region. A kernel

density function was also fitted on species occurrence records prior to projecting species

occupancy in the environmental space. Thereby, each cell of the environmental space is weighted

according to the availability of this specific environmental condition and species occurrence

records are weighed in a way that all species involved in the comparison are given similar total

weights.

We further tested for niche equivalency and niche similarity (*cf.* Warren *et al.*, 2008). For that, we

performed paired comparisons of the realized niches of each species over time and between species

within each period. The first test evaluates if the environmental conditions that define the niches

of two entities are identical. Specifically, niche equivalency is tested by comparing the overlap

between the two realized niches with the expected distribution of overlap obtained by randomly

reallocating the grid cells occupied by the two entities. The second test assesses the similarity in

the relative distribution over the environmental conditions defining the niches of two entities.

Niche similarity is tested by comparing the overlap between the two realized niches to the expected

distribution of overlap obtained by reallocating the density of occurrence of one entity across its

range of occupancy, while the occurrence of the other remains constant. In other words, this test

estimates the likelihood that niche centroids are significantly different from each other. For both

tests, expected distributions were based on 500 iterations of the randomization procedure.

RESULTS

Description of the available environment and realized niches

The environmental space sampled in each region was described by the first two axes of the

principal component analyses, capturing 43.0% and 41.6% of the environmental variation in

Norrström and Skåne respectively (Fig 1b & c, and Appendix S2). In both regions, the first PCA-

axis was strongly associated to a gradient defined by grid cells having a higher amount of arable

land at one end and more forest at the other. Note that in both cases the estimates of soil nitrogen

and phosphorus flow explained a large part of the variance captured by the first PCA-axis and were

associated with higher amount of arable land. In both regions, the variance along the second axis

reflected change in the mean aspect (varying from 0 degree North to 301 degrees at maximum).

Overall, the ecological niches occupied by *A. io* and *A. urticae* in each region between time-periods

were highly comparable (Figs 2 & 3).

Interspecific interactions

In spite of the overall similarities, we detected in each region and time-period significant differences among the species' realized niches. In most cases, the null hypothesis of niche equivalency and similarity was rejected (species interactions, $p<0.05$, Table 1), even though the three species displayed important overlap in their environmental space (Table 1, Figs 4).

In the two-dimensional environmental space in Skåne, *A. levana* overlapped with the two native species in both time-periods but the species' realized niches were neither equivalent nor similar ($p<0.05$, Table 1). Moreover, we observed a large decrease in the overlap of *A. levana* with the native species over time. In the first period it was 0.80 with *A. urticae* and *A. io*, while in the second period it was 0.68 with both native species. In the first period we did not observe a clear differentiation between *A. levana* and the resident species' niches (Fig. 4c & d top), but in the second period this partitioning appears clearly (Fig. 4c & d bottom). In the second period, *A. levana* preferentially occupied habitats with larger amount of forest per 1 km grid cell and the two native species were in comparison more present in mixed and agricultural habitats (Fig. 4c & d bottom).

The two native species (*A. urticae* and *A. io*) strongly overlapped in both periods and regions (overlap between 0.89 and 0.91, Table 1). This is not surprising considering that *A. io* and *A. urticae* are often observed in sympatry with ecological niches that are known to be difficult to differentiate. While in the first period, *A. urticae* and *A. io* displayed equivalent niches in Skåne (paired-species comparisons of niche equivalency, $p=0.327$, Table 1), the null hypothesis of niche equivalency was rejected in the first period in Norrström and in both regions in the second period ($p<0.05$, Table 1). Thus, although the realized niches of *A. urticae* and *A. io* significantly overlapped, they displayed non-random differences in their distributions. For all the paired-species comparisons

between *A. urticae* and *A. io*, we rejected the null hypothesis of niche similarity (tests of niche

similarity, *p*<0.05, Table 1); which suggests that these species occupied the available environment

differently in both time-periods and regions (shifts in their centroid). In Skåne in the first period,

A. urticae occupied proportionally more areas with larger amount of forest per 1 km grid cells than

A. io, whereas *A. io* was more abundant in mixed habitats (Fig. 4b top). In the second period, the

prevalence of *A. urticae* slightly shifted in comparison to *A. io* from forested areas toward habitats

with increasing amount of agricultural and urban lands (Fig. 4b bottom). In Norrström and for both

periods, *A. urticae* and *A. io* prevailed in environments with higher amount of forest (Fig. 2).

Niche shifts over time

The distribution of the three species shifted significantly over time in both regions, their

distribution in the environmental space being neither equivalent nor similar between the two time-

periods (tests, *p*<0.05, Table 1). The largest shift was observed for *A. levana* in Skåne, for which

the overlap in the realized niche between periods was of 0.75. For the two native species these

shifts were comparatively smaller, the overlap in the realized niche of each species between periods

varied between 0.82 and 0.87 (Fig. 5), but noticeable considering the short time scale of the study.

All shifts detected in the environmental space also corresponded to geographical shifts, as only

one layer of land use was available to cover both time-periods. For the recently established *A.*

levana, the observed shift in realized niche corresponded to a niche contraction as *A. levana* in the

first period seemed to have already colonized most of the available environmental conditions in

which the species occurred in the second period (Fig. 3c). In the second period, *A. levana*

preferentially occupied areas with larger amount of forest (Fig. 5c bottom). For the two native

species, the largest changes between the two periods were observed in Skåne (Fig. 5a & b).

Moreover, the extent of the environmental shift observed in Skåne was species-specific, being

larger for *A. urticae* (niche overlap=0.82, sd=0.052; Fig. 5a bottom) than for *A. io* (niche

overlap=0.83, sd=0.041; Fig. 5b bottom). The shift was also more clearly directed for *A. urticae*

than for *A. io* (Fig. 5a & b bottom). By shifting its distribution away from areas with larger amount

of forest, *A. urticae* reduced its overlap with *A. levana*, which over time has densified its

distribution in these environments across Skåne (niche overlap=0.68, sd=0.121; Fig. 4c bottom).

We observed the same pattern between *A. io* and *A. levana*, but to a lower extent (niche

overlap=0.68, sd=0.111; Fig. 4c bottom).

DISCUSSION

The distributions of our focal species were described and compared along the two first axes of the

environmental PCA. In both regions, the first axis was shaped along a gradient from arable land

to more forested land and the second axis correlated with topographic parameters (Fig.1a & b).

Based on our knowledge of the ecological niche of our focal species, we are confident in the

description of the species' environmental niche revealed by our models. The method developed by

Broennimann *et al.* (2012) gave results consistent with the overall characteristics of the realized

niches of the three butterfly species, which are known to occupy open habitats and woodlands,

wood hedges, and hedgerows (Asher *et al.*, 2001; Haahtela *et al.*, 2011).

Yet, the method also identified differences in realized niches among species in each region and

within species over time. We used the measure of overlap and the results from the niche

equivalency and similarity tests between realized niches to infer the role of interspecific

interactions in shaping species distributions. The large overlap in the realized niches of the two

native species in both regions and time periods suggests that no clear negative interaction is

involved in the partitioning of the niche between *A. urticae* and *A.io.* Nevertheless, the fact that

the realized niches of species were neither equivalent, nor similar, suggests some level of non-

random niche partitioning in the environmental space. Thus, while our results emphasized that the

two native species occupy the available environment differently, we could not identify a specific

environmental factor of niche differentiation. For example, we have observed in the field that *A.*

urticae tends to occupy more sun-facing slopes than *A. io*, but although we detected some effect

of aspect on species' distribution, the relatively coarse resolution of our data (1km grid cell)

prevents us to firmly support our field observations. Moreover, apart from the spatial component,

niche partitioning can also arise from, and in combination with, resource and/or temporal

partitioning (MacArthur & Levins, 1967). Regarding resource partitioning, both *A. urticae* and *A.*

io larvae feed on the leaves of stinging nettles, whose availability is not limited over the season.

Stinging nettle has even been shown to expand due to human land use and nitrogen pollution

(Taylor, 2009). Together with the lack of strong competition over resources, potential asynchrony

(temporal partitioning) in species phenology could also reduce the strength of a spatial signal in

the differentiation in species' realized niches.

The recently established *A. levana* overlaps with the realized niches of the native species and the

overlap decreased over time. In part, this decrease is likely a consequence of the described shifts

in realized niche of the three species over time. The largest shift was found for *A. levana* (in Skåne).

From the environmental conditions colonized by *A. levana* in the first period, we observed a

significant contraction in its realized niche in the second period toward habitats with a higher

amount of forest. This pattern suggests that the distribution of *A. levana* was not at equilibrium in

the first period and that it has progressively stabilized within habitats with a positive population

dynamic. We also found significant shifts in realized niches of the two native species over time, in

both regions. However, these shifts were most pronounced in Skåne where *A. urticae*, and to a

lower extent *A. io*, shifted their distributions away from forested areas towards habitats with higher

amount of agricultural and urban lands.

There are several processes that could explain these changes over time. The more pronounced shift

in the distribution of the native species in Skåne than in Norrström could partly reflect higher

regional climatic fluctuations over time. However, as the amplitudes of climatic variations over

time were comparable across the regions (Appendix S3), this factor does not appear to be a major

driver in our study system. The observed shifts in niche centroid over time could also be related to

the increase in the number of reported observations through time. Yet, this explanation also appears

rather unlikely, as the increases in the number of observation records were consistent across species

and regions, while all species did not show a consistent shift in their centroids. Thus, even if such

shifts may have been strengthened by the higher number of observation records over time, they

most likely represent genuine responses to other driving forces affecting the ecological niche of

native species.

Hence, in this community, changes in interspecific interactions appear to be a sensible explanation

for the described changes over time in species occupancy of the environmental space. One

contributing factor is probably the stabilization of the distribution of *A. levana,* as this resulted in

decreased overlap with the native species (Fig. 4c & d). This response may have been mediated

through the action of parasitoids (Dunn *et al.*, 2012). Indeed, studies have shown that the

community composition in herbivorous insects can largely be shaped by parasites (van Veen *et*

al., 2006; Tylianakis, 2009). All three species are heavily parasitized (personal observation) and

share many parasitoids (Hinz & Horstmann, 2007; Shaw *et al.*, 2009). Hence, the population

dynamics of these parasitoids might have been positively affected by the increase in potential hosts

(*A. levana*) between the two time-periods, as observed in tropical forest communities by Morris *et*

al. (2004). Moreover, the differences in phenology between *A. urticae, A. io*, and *A. levana* (Fig.

S1 in Appendix S1) may form a continuous breeding ground to stimulate parasitoid population

dynamics, and they may result in a prolonged temporal niche for parasitoids (Blitzer & Welter,

2011). In a scenario of parasitoid-driven apparent competition, the phenologically late species is

expected to be the more vulnerable as its life cycle will coincide with an increase in parasitoid

population size (Blitzer & Welter, 2011). In this study, the native species showed differences in

the magnitude of their shifts. The shift in the realized niche of *A. urticae* between periods was

stronger than for *A. io,* in terms of niche overlap. Niche partitioning was also stronger between *A.*

levana and *A. urticae* than with *A. io*. In the context of parasitoid buildup, this may seem

counterintuitive, as *A. urticae* is the earliest of the two native species. But because *A. urticae* is

bivoltine in Skåne, and produces a second brood later in the season (Fig. S1 in Appendix S1),

parasitoid populations may continue to build up over the season and result in the highest parasitoid

load for the second brood of *A. urticae*. Although we have no direct data to support the hypothesis

of parasitoid-driven niche differentiation, the circumstantial evidence is suggestive, and is further

reinforced by the observation that *A. levana* appears to share more parasitoids with *A. urticae* than

with *A. io,* based on the limited information available (Shaw *et al.*, 2009).

Differences in life history might be another possible explanation for the different shifts in

distribution among the three species. Based on experimental studies, Merckx *et al.* (2015)

suggested that because of its bivoltine life cycle, the plastic response of *A. urticae* to anthropogenic

environmental changes might be faster in comparison to the univoltine *A. io*. If this is true, *A.*

urticae would be expected to respond faster to the arrival of *A. levana* than *A. io*. This is consistent

with the described larger shift in distribution of *A. urticae* than *A. io* over time in Skåne. Likewise,

the bivoltine life cycle of *A. levana* may also participate to explain the large shift described in its

distribution over time.

It is also possible that differences in the way these species are able to adjust to the quality of their

host plants and the microclimate they offer may cause individuals of each species to choose

different nettle patches when coexisting in a region. The shift in niche centroid of *A. urticae*

reported here (toward agricultural land) is consistent with other studies conducted on *A. urticae*

and *A. io,* showing that *A. urticae* can to a larger extent be favored by the higher nutrient quality

of plants growing in agricultural land than *A. io* (Serruys & Van Dyck, 2014; Audusseau *et al.*,

2015; Merckx *et al.,* 2015). Changes in microclimate have been shown to differentially affect the

decline of butterfly species' following climate change (Wallisdevries & van Swaay, 2006) and to

modify butterflies' habitat associations (Davies *et al.*, 2006). We call for field experiments to

understand, at a finer spatial-scale, the differences among species in their optimal microclimatic

niche.

The method used was successful in handling the potential bias related to the non-standardized

sampling design associated to occurrence data extracted from the public database. Indeed, while

most observations were recorded in urban areas, the highest densities of occurrence of the two

native species based on the grid cell analysis were found in forest habitats (Figs 2 & 3). In addition,

the lower number of species records in the first period did not seem to have affected our capacity

to extract the main components of the realized niches as their shapes and centroids were consistent

through time. Hence, we consider that using occurrence records from citizen-science databases

offer great opportunities for investigating the realized niches of species and differences between

them, as long as the biological relevance of the modelling results is carefully considered. The

results of the method we used to characterize the realized niches of the focal butterfly species was

consistent with our knowledge of the species, and we recommend using occurrence data associated

with species and regions of high sampling effort in order to get more robust predictions. Such

regions often correspond to more densely populated areas as the reporting effort in citizen-based

monitoring programs is strongly related to the number of participants.

In conclusion, we found indications of niche partitioning among the three species and consider

that the observed shifts in species environmental niche were partly driven by the recent changes in

interspecific interactions. The establishment during range expansion of *A. levana* may have

modified biotic interactions, resulting in associated shifts in species distributions. Based on our

knowledge of the system, we suspect that the observed niche differentiation might be partly driven

by apparent competition mediated by shared parasitoids. Other factors such as interspecific

differences in life history could also have contributed to the observed pattern and lead to species-

specific change in habitat preference. The difference in the extent of the shift observed between *A.*

urticae and *A. io* during the establishment of *A. levana* can be related to their differences in

phenology, voltinism, and the higher number of shared parasitoids between *A. urticae* and *A.*

levana. Further investigations of the niche partitioning of these three nettle-feeding butterflies

where they all are native species would allow us to make predictions about the equilibrium this community may reach. More importantly, this framework is a promising tool for investigating the potential role of biotic interactions on species distributions, and for better predicting the outcome of their modifications on community composition and species-environment relationships.

ACKNOWLEDGMENTS

We thank the Strategic Research Program Ekoklim at Stockholm University for funding of this project. HA is acknowledging support from Helge Ax:son Johnsons stiftelse, RS from the FRB and EDF SA (FRB-CESAB project LOLA-BMS), and SN from the Swedish Research Council (2011-5636). We also wish to thank Vlad Dincã and Claudie Daniel for the photos presented in the Supporting Information, and Lars B. Pettersson for an insightful comment on the manuscript. We are also thankful for the comments from C. Stefanescu and two anonymous referees.

REFERENCES

Araújo, M.B. & Luoto, M. (2007) The importance of biotic interactions for modelling species distributions under climate change. *Global Ecology and Biogeography*, **16**, 743–753.

Asher, J., Warren, M., Fox, R., Harding, P., Jeffcoate, G. & Jeffcoate, S. (2001) *The millennium atlas of butterflies in Britain and Ireland*. Oxford University Press, Oxford.

Audusseau, H., Kolb, G. & Janz, N. (2015) Fertilization interacts with life history: variation in performance and stoichiometry in nettle-feeding butterfies. *Plos One*, **10**(5).

Berry, P., Dawson, T., Harrison, P. & Pearson, R. (2002) Modelling potential impacts of climate change on the bioclimatic envelope of species in Britain and Ireland. *Global Ecology and Biogeography*, **11**, 453–462.

Betzholtz, P.E., Pettersson, L.B., Ryrholm, N. & Franzén, M. (2013) With that diet, you will go
far: trait-based analysis reveals a link between rapid range expansion and a nitrogen-favoured
diet. *Proceedings of the Royal Society of London B: Biological Sciences*, **280**.

Blitzer, E. & Welter, S. (2011) Emergence asynchrony between herbivores leads to apparent
competition in the field. *Ecology*.

Brandt, M., Ejhed, H. & Rapp, L. (2009) Nutrient loads to the Swedish marine environment
in 2006. *Swedish Environmental Protection Agency Report*, **5815**, 93.

Broennimann, O., Fitzpatrick, M.C., Pearman, P.B., Petitpierre, B., Pellissier, L., Yoccoz, N.G.,
Thuiller, W., Fortin, M.J., Randin, C., Zimmermann, N.E., Graham, C.H. & Guisan, A. (2012)
Measuring ecological niche overlap from occurrence and spatial environmental data.
Global Ecology and Biogeography, **21**, 481–497.

Broennimann, O., Petitpierre, B., Randin, C., Engler, R., Cola, V.D., Breiner, F., D'Amen, M.,
Pellissier, L., Pottier, J., Pio, D., Mateo, R.G., Hordijk, W., Dubuis, A., Scherrer, D., Salamin,
N. & Guisan, A. (2015) ecospat: Spatial Ecology Miscellaneous Methods. R package version
1.1. URL http://CRAN.R-project.org/package=ecospat

Davis, A.J., Jenkinson, L.S., Lawton, J.H., Shorrocks, B. & Wood, S. (1998) Making mistakes
when predicting shifts in species range in response to global warming. *Nature*, **391**, 783–
786.

Davies, Z.G., Wilson, R.J., Coles, S. & Thomas, C.D. (2006) Changing habitat associations of a
thermally constrained species, the silver-spotted skipper butterfly, in response to climate
warming. *Journal of Animal Ecology*, **75**, 247-256.

Devictor, V., van Swaay, C., Brereton, T. *et al.* (2012) Differences in the climatic debts of birds
and butterflies at a continental scale. *Nature Climate Change*, **2**, 121–124.

Dunn, A.M., Torchin, M.E., Hatcher, M.J., Kotanen, P.M., Blumenthal, D.M., Byers, J.E., Coon,
C.A., Frankel, V.M., Holt, R.D., Hufbauer, R.A., Kanarek, A.R., Schierenbeck, K.A., Wolfe,
L.M. & Perkins, S.E. (2012) Indirect effects of parasites in invasions. *Functional Ecology*,
26, 1262–1274.

Eliasson, C. U., Ryrholm, N., Holmer, M., Jilg, K. & Gärdenfors, U. (2005) *Nationalnyckeln till*
Sveriges flora och fauna. Fjärilar. Dagfjärilar. Hesperiidae–Nymphalidae. Artdatabanken,
SLU, Uppsala, Sweden.

Gärdenfors, U., Jönsson, M., Obst, M., Wremp, A.M., Kindvall, O. & Nilsson, J. (2014) *Swedish*
LifeWatch – A Biodiversity Infrastructure Integrating and Reusing Data from Citizen
Science, Monitoring and Research.

Gilman, S.E., Urban, M.C., Tewksbury, J., Gilchrist, G.W. & Holt, R.D. (2010) A framework
for community interactions under climate change. *Trends in Ecology & Evolution*, **25**, 325–
331.

González-Megías, A., Menéndez, R., Roy, D., Brereton, T. & Thomas, C.D. (2008) Changes in
the composition of British butterfly assemblages over two decades. *Global Change Biology*,
14(7), 1464–1474.

Haahtela, T., Saarinen, K., Ojalainen, P., Aarnio, H. (2011). *Butterflies of Britain and Europe: A*
Photographic Guide. A. & C. Black Publishers Ltd.

Heikkinen, R.K., Luoto, M., Virkkala, R., Pearson, R.G. & Körber, J.H. (2007) Biotic
interactions improve prediction of boreal bird distributions at macro-scales. *Global*
Ecology and Biogeography, **16**, 754–763.

Hinz, R. & Horstmann, K. (2007) Uber Wirtsbeziehungen europaischer Ichneumon-Arten
(Insecta, Hymenoptera, Ichneumonidae, Ichneumoninae). *Spixiana*, **30**, 39.

IPCC (2014) *Climate Change 2013*. The physical science basis: Working group I contribution
to the fifth assessment report of the Intergovernmental Panel on Climate Change.
Cambridge University Press.

Leathwick, J.R. (1998) Are New Zealand's Nothofagus species in equilibrium with their
environment? *Journal of Vegetation Science*, **9**, 719–732.

MacArthur, R. & Levins, R. (1967) The limiting similarity, convergence, and divergence of
coexisting species. *American naturalist*, pp. 377–385.

Mason, T.H., Stephens, P.A., Apollonio, M. & Willis, S.G. (2014) Predicting potential

responses to future climate in an alpine ungulate: interspecific interactions exceed climate exceed. *Global change biology*, **20**, 3872–3882.

Merckx, T., Serruys, M. & Van Dyck, H. (2015) Anthropogenic host plant expansion leads a nettle-feeding butterfly out of the forest: consequences for larval survival and developmental plasticity in adult morphology. *Evolutionary Applications*.

Morris, R.J., Lewis, O.T. & Godfray, H.C.J. (2004) Experimental evidence for apparent competition in a tropical forest food web. *Nature*, **428**, 310–313.

Parmesan, C. (2006) Ecological and evolutionary responses to recent climate change. *The Annual Review of Ecology, Evolution, and Systematics*, **37**, 637–669.

Pearson, R.G. & Dawson, T.P. (2003) Predicting the impacts of climate change on the distribution of species: are bioclimate envelope models useful? *Global ecology and biogeography*, **12**, 361–371.

Pellissier, L., Pradervand, J.N., Pottier, J., Dubuis, A., Maiorano, L. & Guisan, A. (2012) Climate-based empirical models show biased predictions of butterfly communities along environmental gradients. *Ecography*, **35**, 684–692.

Pettersson, L., Mellbrand, K. & Ottvall, R. (2013) *Swedish Butterfly Monitoring Scheme, annual report for 2012*.

PostgreSQL (2014) *PostgreSQL, version 9.3.5*. PostgreSQL Global Development Group. URL http://www.enterprisedb.com/

Pöyry, J., Luoto, M., Heikkinen, R.K., Kuussaari, M. & Saarinen, K. (2009) Species traits explain recent range shifts of Finnish butterflies. *Global Change Biology*, **15**, 732–743.

Quantum GIS (2012) *Quantum GIS 1.8 Software*. QGIS Development Team. URL http://qgis.org/

R Core Team (2015) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. URL http://www.R-project.org/

Schweiger, O., Heikkinen, R.K., Harpke, A., Hickler, T., Klotz, S., Kudrna, O., Kühn, I., Pöyry, J. & Settele, J. (2012) Increasing range mismatching of interacting species under global

change is related to their ecological characteristics. *Global Ecology and Biogeography*, **21**, 88–99.

Serruys, M. & Van Dyck, H. (2014) Development, survival, and phenotypic plasticity in anthropogenic landscapes: trade-offs between offspring quantity and quality in the nettle-feeding peacock butterfly. *Oecologia*, **176**, 379–387.

Shaw, M.R., Stefanescu, C. & Van Nouhuys, S. (2009) *Parasitoids of European Butterflies*. Cambridge University Press, Cambridge.

Taylor, K. (2009) Biological Flora of the British Isles: *Urtica dioica* L. *Journal of Ecology*, **97**, 1436–1458.

Thuiller, W., Lavorel, S., Araújo, M.B., Sykes, M.T. & Prentice, I.C. (2005) Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences of the United States of America*, **102**, 8245–8250.

Tingley, R., Vallinoto, M., Sequeira, F. & Kearney, M.R. (2014) Realized niche shift during a global biological invasion. *Proceedings of the National Academy of Sciences*, **111**, 10233–10238.

Tylianakis, J.M. (2009) Warming up food webs. *Science*, **323**, 1300–1301.

Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D.A. (2008) Global change and species interactions in terrestrial ecosystems. *Ecology letters*, **11**, 1351–1363.

van Veen, F.F.J., Morris, R.J. & Godfray, H.C.J. (2006) Apparent competition, quantitative food webs, and the structure of phytophagous insect communities. *Annual Review of Entomology*, **51**, 187–208.

Wallisdevries, M.F. & van Swaay, C.A.M. (2006) Global warming and excess nitrogen may induce butterfly decline by microclimatic cooling. *Global Change Biology,* **12**, 1620-1626.

Warren, D.L., Glor, R.E. & Turelli, M. (2008) Environmental niche equivalency versus conservatism: quantitative approaches to niche evolution. *Evolution*, **62**, 2868–2883.

Wisz, M.S., Pottier, J., Kissling, W.D. *et al.* (2013) The role of biotic interactions in shaping distributions and realised assemblages of species: implications for species distribution

554   modelling. *Biological Reviews*, **88**, 15–30.

SUPPORTING INFORMATION

Appendix S1 Data description, phenology of the study species and consideration of biases.

Appendix S2 Table showing scores for each variable on the first three axes of the PCA-env and

the contribution of each axis expressed in percentage of the total variance (inertia).

Appendix S3 Average mean monthly temperature (Jan: 1, Feb: 2, etc.) in Skåne in comparison to

Norrström and in the first and second periods (mean±se).

BIOSKETCH Hélène Audusseau has been granted her PhD at Stockholm University. Her research

is concerned by the ecological and evolutionary responses of nettle-feeding butterflies to changes

in climate and land use.

Editor: Daniel Chapman.

AUTHORS' CONTRIBUTIONS

AH, ML, RS, NJ, SN participated in the design of the study. AH, ML, RS performed the analyses.

AH, ML, BK collected the data. AH wrote the first draft of the manuscript, which was substantially

improved through the contribution of RS, ML, NJ. All authors contributed to the final version of

the manuscript.

TABLE

Table 1 Table showing niche overlap (D) within species over time (time shift) and between species in each period (species interactions)

in Skåne and Norrström, respectively, with the associated *P*-values obtained from the tests of niche equivalency and similarity (no.

iterations=500).

Region		Period	Species	Test of niche equivalency		Test of niche similarity	
						Sp 1 -> Sp 2	Sp 2 -> Sp 1
			Sp 1 - Sp 2	Overlap D	*p*	*p*	*p*
Skåne	Time shift	P1 - P2	*A. urticae*	0.82	0.004	0.002	0.002
		P1 - P2	*A. io*	0.83	0.004	0.002	0.002
		P1 - P2	*A. levana*	0.75	0.004	0.002	0.002
	Species interactions	P1 - P1	*A. urticae - A. io*	0.89	0.327	0.002	0.002
		P1 - P1	*A. urticae - A. levana*	0.80	0.004	0.002	0.002
		P1 - P1	*A. io - A. levana*	0.80	0.004	0.002	0.002
		P2 - P2	*A. urticae - A. io*	0.89	0.032	0.002	0.002
		P2 - P2	*A. urticae - A. levana*	0.68	0.004	0.002	0.016
		P2 - P2	*A. io - A. levana*	0.68	0.004	0.002	0.004
Norrström	Time shift	P1 - P2	*A. urticae*	0.87	0.004	0.002	0.002
		P1 - P2	*A. io*	0.85	0.004	0.002	0.002
	Species interactions	P1 - P1	*A. urticae - A. io*	0.91	0.032	0.002	0.002
		P2 - P2	*A. urticae - A. io*	0.89	0.004	0.002	0.002

FIGURE LEGENDS

Figure 1 (a) Map of the two study-regions (in grey), corresponding to the county of Skåne in

Southern Sweden and the Norrström drainage basin in Central Sweden. (b), (c) Correlation circles

showing the contribution of the different variables to the two main axes of the PCA, which describe

the environmental space in Norrström and Skåne, respectively. S: Sweden, N: Norway, DK:

Denmark.

Figure 2 Density of species occurrence across the two-dimensional environmental space

describing the Norrström region. In columns are the densities of occurrence of (a) *Aglais urticae*

and (b) *Aglais io* for each period (in rows). The black gradient corresponds to the increase in the

density of occurrence of the species. The solid line corresponds to the limit of the environmental

space available. The dashed line corresponds to the 50% most frequently available environmental

conditions.

Figure 3 Density of species occurrence across the two-dimensional environmental space

describing the Skåne region. In columns are the densities of occurrence of (a) *Aglais urticae*, (b)

Aglais io and (c) *Araschnia levana* for each period (in rows). The black gradient corresponds to

the increase in the density of occurrence of the species. The solid line corresponds to the limit of

the environmental space available. The dashed line corresponds to the 50% most frequently

available environmental conditions.

Figure 4 Plots of the differences in density of occurrence, for each region and time-period, between

species. In columns are represented the differences in the density of occurrence in Norrström of

(a) *Aglais urticae* and *Aglais io* and in Skåne of (b) *A. urticae* and *A. io*, (c) *A. urticae* and *A. levana*

and (d) *A. io* and *A. levana*, for each period (in rows). Red cells indicate the prevalence of the first

species over the second species. Dark blue cells indicate the prevalence of the second species over

the first species. Grid cells with a color value of zero correspond to environmental conditions for

which the densities of occurrence of both species were similar. The boxplot and the standard

deviation (sd) associated with each graph show the variability of the differences of densities

observed across the available environment. The larger the box and the higher the sd are, the more

contrasting the species are in their distribution.

Figure 5 Plots of the differences in density of occurrence of each species over time, and for each

region. In columns are represented the differences in the density of occurrence of (a) *Aglais urtic*ae,

(b) *Aglais io* and (c) *A. levana* over time and for the regions Norrström and Skåne (in rows). Red

cells indicate the prevalence of the species in the second period. Dark blue cells indicate the

prevalence of the species in the first period. Grid cells with a color value of zero correspond to

environmental conditions for which the density of occurrence of the species did not vary over time.

The boxplot and the standard deviation (sd) associated with each graph show the variability of the

differences in densities observed across the available environment. The larger the box and the

higher the sd are, the more contrasting the species distributions are.

FIGURES

Figure 1

Figure 2

Figure 3

Figure 4

Species interactions

(a) *A. urticae - A. io*   (b) *A. urticae – A. io*   (c) *A. urticae - A. levana*   (d) *A. io- A. levana*

Norrström

Skåne

Period 1

Period 2

622   **Figure 5**

624