Widespread shortening of 3' untranslated regions and increased exon inclusion are evolutionarily conserved features of innate immune responses to infection

Athma A. Pai[a,1,2], Golshid Baharian[b,c,1], Ariane Pagé Sabourin[c], Jessica F. Brinkworth[b,c,d], Yohann Nédélec[b,c], Joseph W. Foley[e], Jean-Christophe Grenier[c], Katherine J. Siddle[f], Anne Dumaine[c], Vania Yotova[c], Zachary P. Johnson[g,h], Robert E. Lanford [i], Christopher B. Burge[a,j], Luis B. Barreiro[c,k,2]

[a]Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, USA

[b]Department of Biochemistry, Faculty of Medicine, University of Montreal, Montreal, QC, Canada

[c]Sainte-Justine Hospital Research Centre, Montreal, QC, Canada

[d]Department of Anthropology, University of Illinois, Urbana-Champaign, IL, USA

[e]Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA

[f]Department of Organismic and Evolutionary Biology, FAS Center for Systems Biology, Harvard University, Cambridge, MA, USA

[g]Yerkes National Primate Research Center, Emory University, Atlanta, GA 30322, USA

[h]Department of Human Genetics, Emory University School of Medicine, Atlanta, GA 30322, USA

[i]Texas Biomedical Research Institute, Department of Virology and Immunology, San Antonio, TX, USA.

[j]Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA

[k]Department of Pediatrics, Faculty of Medicine, University of Montreal, Montreal, QC, Canada

[1]These authors contributed equally to this work.

[2]To whom correspondence should be addressed. E-mail: athma@mit.edu and luis.barreiro@umontreal.ca

ABSTRACT

The contribution of pre-mRNA processing mechanisms to the regulation of immune

responses remains poorly studied despite emerging examples of their role as

regulators of immune defenses. Here, we used mRNA sequencing to quantify gene

expression and isoform abundances in primary macrophages from 60 individuals,

before and after infection with two live bacteria. In response to both bacteria we

identified thousands of genes that significantly change isoform usage in response to

infection, and found global shifts towards (*i*) the inclusion of cassette exons and (*ii*)

shorter 3' UTRs. Using complementary data collected in non-human primates, we

show that these features are evolutionarily conserved among primates. Finally, our

results suggest that the pervasive usage of shorter 3' UTRs is a mechanism for

particular genes to evade repression by immune-activated miRNAs. Collectively, our

results show that dynamic changes in RNA processing play a key role in the

regulation of innate immune responses.

INTRODUCTION

Innate immune responses depend on robust and coordinated gene expression programs involving the transcriptional regulation of thousands of genes (Huang et al. 2001; Smale 2010; Medzhitov and Horng 2009). These regulatory cascades start with the detection of microbial-associated products by pattern recognition receptors, which include Toll Like Receptors (TLRs), NOD-like receptors and specific C-type lectins (Medzhitov and Janeway 1998; Kawai and Akira 2010). These initial steps are followed by the activation of key transcription factors (e.g., NF-κB and interferon regulatory factors) that orchestrate the inflammatory and/or antiviral response signals involved in pathogen clearance and the subsequent development of appropriate adaptive immune responses (Medzhitov and Janeway 1998).

Although much attention has been devoted to characterizing transcriptional changes in response to infectious agents or other immune stimuli, we still know remarkably little about the contribution of post-transcriptional changes – specifically, changes in alternative pre-mRNA processing – to the regulation of the immune system (Lynch 2004; Martinez and Lynch 2013; Alasoo et al. 2015). Alternative splicing can impact cellular function by creating distinct mRNA transcripts from the same gene, which may encode unique proteins that have distinct or opposing functions. For instance, in human and mouse cells, several alternatively spliced forms of genes involved in the TLR pathway have been shown to function as negative regulators of TLR signaling in order to prevent uncontrolled inflammation (Rao et al. 2005; Wells et al. 2006; O'Connor et al. 2015). Additionally, several in-depth studies of individual genes suggest that alternative splicing plays an important role in increasing the diversity of transcripts encoding MHC molecules (Ishitani and Geraghty 1992) and in modulating intracellular signaling and intercellular interactions through the expression of various isoforms of key cytokines (Bihl et al. 2002; Nishimura et al. 2000), cytokine receptors (Rao et al. 2005; Goodwin et al. 1990; Koskinen et al. 2009), kinases (Jensen and Whitehead 2001) and adaptor proteins (Gray et al. 2010; Ohta et al. 2004; Janssens et al. 2002).

Yet, the extent to which changes in isoform usage are a hallmark of immune
responses to infection remains largely unexplored at a genome-wide level (Alasoo et
al. 2015; Rodrigues et al. 2013). Moreover, we still do not know which RNA
processing mechanisms contribute most to the regulation of immune responses, or
how these disparate mechanisms are coordinated.

To address these questions, we investigated genome-wide changes in transcriptome
patterns after independent infections with *Listeria monocytogenes* or *Salmonella*
typhimurium in primary human macrophages. Because of the distinct molecular
composition of these two pathogens and the way they interact with host cells, they
activate distinct innate immune pathways after infection (Haraga et al. 2008; Pamer
2004). Thus, our study design allowed us to evaluate the extent to which changes in
isoform usage are pathogen-specific or more generally observed in response to
bacterial infection. Furthermore, the large number of individuals in our study
allowed us to use natural variation in RNA processing to gain insight into fine-tuned
inter-individual regulation of immune responses. Our results provide a
comprehensive picture of the role of RNA processing in regulating early innate
immune responses to bacterial infection in human antigen-presenting cells.

RESULTS

Infection with either *Listeria* or *Salmonella* induces dramatic changes in mRNA
expression levels. Following 2 hours of infection with each bacteria, we collected
RNA-seq data from 60 matched non-infected and infected samples, with an average
of 30 million reads sequenced per sample (see Methods; Table S1). The first
principal component of the complete gene expression data set clearly separated
infected from non-infected samples, and both PC1 and PC2 clustered infected
samples by pathogen (i.e. *Listeria* or *Salmonella*; Figure 1A). Accordingly, we
observed a large number of differences in gene expression levels between infected
and non-infected cells, with 5,809 (39%) and 7,618 (51%) of genes showing
evidence for differential gene expression (DGE) after infection with *Listeria* and

Salmonella, respectively (using DESeq2, FDR ≤ 0.1% and |log$_2$(fold change)| > 0.5;

Figure S1A, Table S2). As expected, the sets of genes that responded to either

infection were strongly enriched (FDR ≤ 1.8×10^{-6}) for genes involved in immune-

related biological processes such as the regulation of cytokine production,

inflammatory responses, or T-cell activation (Table S3).

In order to uncover changes in isoform usage in response to infection using our

RNA-sequencing data, we applied two complementary approaches for analyzing

RNA processing. We first measured abundances of full isoforms to gain a holistic

perspective of the transcriptome landscape. Second, we used the relative

abundances of individual exons within a gene to gain a finer understanding of the

specific RNA processing mechanisms likely to be involved in the regulation of

immune responses.

Pervasive differential isoform usage in response to infection. We initially

sought to assess changes in isoform usage after infection using proportional

abundances of the different isoforms encoded by the same gene. To do so, we used

the transcripts per million (TPM) values reported by the RSEM software (Li et al.

2010) and calculated relative proportions by dividing isoform-level expression by

the overall expression level of the gene (i.e., TPM summed across all possible

isoforms). We then quantified differential isoform usage (DIU) between conditions

for genes with at least two annotated isoforms (*N = 11,353*) using a multivariate

generalization of the Welch's t-test that allowed us to test whether, as a group, the

proportional abundances of the different isoforms in each gene were significantly

different between infected and non-infected cells (Methods). We found that 1,456

(13%; after *Listeria* infection) and 2,862 (25%; after *Salmonella* infection) genes

showed evidence for significant DIU after infection (FDR ≤ 1%; Figure 1B for an

example of an DIU gene, Figure S1B, Table S4). Across all DIU genes, we observed a

mean of 10% (± 8% s.d.) and 13% (± 10% s.d.) change in relative isoform usage

upon infection with *Listeria* and *Salmonella*, respectively (defined as the maximum

change upon infection in relative isoform usage across transcripts of each gene,

Figure S1C). Following infection with *Listeria* and *Salmonella*, 6.2% and 10.2%,
respectively, of genes with a dominant isoform prior to infection (see methods) switch to
using an alternative dominant isoform (Figure S1E).

DIU genes were significantly enriched for genes involved in immune responses (FDR
≤ 0.01; Table S3), including several cytokines (e.g. *IL1B*, *IL7*, *IL20*), chemokines
(*CCL15*), regulators of inflammatory signals (e.g. *ADORA3*, *MAP3K14*) and genes
encoding co-stimulatory molecules required for T cell activation and survival (e.g.
CD28 in Figure 1B, *CD80*) (Figure S2). Notably, 86% of genes with DIU upon
infection with *Listeria* were also classified as having DIU after infection with
Salmonella, suggesting that changes in isoform usage are likely to be common across
a wide variety of immune triggers. To assess the robustness of our findings, we
tested for DIU using isoform-specific expression levels calculated with Kallisto, an
alignment-free quantification method (Bray et al. 2016) (see Supplementary
Methods). At an FDR of 5%, we observed a 74% (*Listeria*) and 78% (*Salmonella*)
agreement with DIU genes identified using RSEM, confirming that identification of
DIU genes is largely robust to the method used for isoform quantification.

Considering the large proportion of genes with significant DIU upon infection, we
sought to understand whether these changes arose from a shift in the usage of a
dominant isoform or increased isoform diversity within a given gene. To do so, we
calculated the Shannon diversity index, which measures the evenness of the isoform
usage distribution for each gene before and after infection (low values reflect usage
of one or few isoforms; high values reflect either the usage of a more diverse set of
isoforms or more equal representation of the same set of isoforms). $\Delta_{Shannon}$ thus
quantifies the change in isoform diversity after infection (Table S5). The majority of
genes (63% in *Listeria* and 68% in *Salmonella*) show an increase in isoform
diversity ($\Delta_{Shannon} > 0$) after infection (Figure S1D), indicating a shift away from
usage of the primary (pre-infection) isoform after infection and a general increase in
the diversity of transcript species following infection.

Previously, it had been reported that DGE and DIU generally act independently to shape the transcriptomes of different mammalian tissues (Barbosa-Morais et al. 2012). In contrast, we found that DIU genes were significantly enriched for genes with DGE (both up- and down-regulated genes, Fisher's exact test, $P \leq 7.5 \times 10^{-6}$ for both *Listeria* and *Salmonella*) (Figure S3A). Despite such enrichment, a substantial proportion of genes (47% (690 genes) in *Listeria* infected samples and 39% (1,105 genes) in *Salmonella* infected samples) with significant changes in isoform usage after infection do not exhibit DGE following infection (Figure 1C). Even after relaxing the FDR threshold for DGE by 100-fold, 685 (*Listeria*) and 1096 (*Salmonella*) DIU genes still exhibited no significant DGE. Thus, a considerable fraction of transcriptome changes in response to infection do occur solely at the level of RNA processing, and independently of changes in mean expression levels.

Directed shifts in 3' UTR length and alternative splicing following infection. Though useful in documenting the substantial transcriptome changes upon infection, aggregate isoform levels calculated by RSEM do not distinguish between different types of RNA processing changes. Thus, we next examined changes in five distinct categories of alternative RNA processing, allowing us to zoom in on particular molecular mechanisms that underlie differential isoform usage in response to infection. Specifically, we used human transcript annotations from Ensembl to quantify usage of (1) alternative first exons (AFEs), (2) alternative last exons (ALEs), (3) alternative polyadenylation sites, leading to tandem 3' untranslated regions (TandemUTRs), (4) retained introns (RIs), and (5) skipped exons (SEs). AFEs, ALEs, and TandemUTRs correspond to the alternative usage of terminal exons primarily affecting UTR composition, whereas RIs and SEs correspond to internal splicing events that usually affect the open reading frame.

For each gene or exon within each RNA processing category, we used the MISO software (Katz et al. 2010) to calculate a "percent spliced in" (PSI or Ψ) value (Table S6), defined as the proportion of transcripts from a gene that contain the "inclusion"

isoform (defined as the longer isoform for RIs, SEs, and TandemUTRs, or use of the exon most distal to the gene for AFEs and ALEs). ΔΨ values thus represent the difference between PSI values calculated for the infected versus non-infected samples. Overall, we observed many significant changes in RNA processing ($N \geq$ *1,098*) across all categories in both bacteria (significance was defined as Bayes Factor > 5 in at least 10% of individuals and |mean ΔΨ| > 0.05), compared to null expectations derived from measuring changes among pairs of non-infected samples (N=29) (Figure 2A, Table S7). Our criteria for determining significant changes also allows us to choose exons that are more likely to be consistently changing across many individuals; significantly changing exons have lower variance in ΔΨ values across individuals than exons that are not changing after infection (Figure S4). The greatest proportion of changes after infection occurred among retained intron and TandemUTR events. PSI values for 7.3% and 14% of retained introns and 7.6% and 16.7% of TandemUTRs significantly changed after infection with *Listeria* or *Salmonella*, respectively. When we considered the set of genes associated with at least one significant change in RNA processing, we observed an over-representation of Gene Ontology categories involved in immune cell processes and the cellular response to a stimulus (Figure 2B, Table S8).

Genome-wide shifts towards more prevalent usage of inclusion or exclusion isoforms have previously been observed as signatures of cellular stress responses or developmental processes (Sandberg et al. 2008; Mayr and Bartel 2009; Wong et al. 2013; Lackford et al. 2014a; Shalgi et al. 2014; Ji et al. 2009). To examine directional shifts in particular RNA processing mechanisms, we used mean ΔΨ values (defined as the mean ΔΨ across all individuals for each gene or exon) and observed a striking global shift towards the usage of upstream polyadenylation sites in tandem 3' UTR regions, indicating pervasive 3' UTR shortening following infection with either *Listeria* or *Salmonella* (Figure 2C; both $P < 2.2 \times 10^{-16}$ with Mann-Whitney U test; 94.4% and 98% of significant TandemUTRs changes in *Listeria* and *Salmonella*). Although less frequent, we also see a general trend towards usage of upstream polyadenylation sites in alternative last exons. Interestingly, we found a striking

correlation in the extent to which a given individual has a global shift towards 3'
UTR shortening and towards usage of an upstream ALE (Pearson R = 0.83, P = 4.97 ×
10^{-16} for Listeria and Pearson R = 0.78, P = 2.65 × 10^{-13} for Salmonella, Figure S6B).
These correlations suggest that the changes in ALEs and TandemUTRs are likely
being regulated by similar or coordinated mechanisms; rather they are likely
regulated through changes in the activities of cleavage and polyadenylation factors
(rather than splicing machinery), as suggested by previous reports (Takagaki and
Manley 1998; Di Giammartino et al. 2011; Taliaferro et al. 2016). Additionally, we
found a substantial increase in the inclusion of skipped exons after infection (both P
< $2.2×10^{-16}$ with Mann-Whitney U test; 79.8% and 77.3 % of significant SE changes
in *Listeria* and *Salmonella*, Figure S5). The consistency of these patterns suggests
that a dedicated post-transcriptional regulatory program might underlie genome-
wide shifts in RNA processing after bacterial infection.

Quantifying changes at the terminal ends of transcripts is challenging due to known
biases in standard RNA-seq protocols (Wang et al. 2009). Thus, we validated our
observation of 3' UTR shortening upon infection by sequencing RNA from 6
individuals after infection with *Listeria* and *Salmonella* with a 3'RNA-seq protocol,
which specifically captures the 3' most end of transcripts (see Methods). TPM
expression values calculated from 3' RNA-seq data highly correlate with TPMs from
RNA-seq data, establishing that this method is able to quantitatively measure of the
number of transcripts in a cell (R > 0.82 across all conditions, P < 2.2 × 10^{-16}, Figure
S6). Meta-gene plots around the polyadenylation sites of core and extended regions
of TandemUTRs show a notable increase after infection in sequencing coverage
around the upstream polyadenylation site relative to the downstream
polyadenylation site, specifically for genes with significantly shorter 3' UTRs upon
infection as assessed by the RNA-seq data (Figure 3A) Finally, to look at 3' UTR
shortening in specific genes, we calculate Ψ values for TandemUTRs using the
3'RNA-seq data (see Supplementary Methods) and find again that genes that are

significantly changing 3' UTR usage in the RNA-seq data exhibit a significant shift

towards negative $\Delta\Psi$ values in the 3'RNA-seq data ($P < 2.2 \times 10^{-16}$; Figure 3B).

RNA processing changes persist 24 hours after infection. To evaluate whether

the strong global shifts we observed were the result of a sustained immune

response to the bacteria rather than a short-lived cellular stress response, we

sequenced RNA from a subset of our individuals (*N=6*) after 24 hours (h) of

infection with *Listeria* or *Salmonella*. Samples sequenced after 24h were paired with

matched non-infected controls cultured for 24h to calculate $\Delta\Psi$ values for

alternative RNA processing categories using MISO (Katz et al. 2010). Across all

categories, we found many more significant changes in RNA processing and larger

$\Delta\Psi$ values after infection for 24h than after 2h (Figure S7). These results further

support a key role for RNA processing in controlling innate immune responses to

infection. For genes or exons that changed significantly after 2h of infection in this

set of 6 individuals (see Methods), the majority of changes (67%) remained

detectable at 24h, usually with similar magnitude and direction (Figure 4A). The one

notable exception is that introns that are preferentially retained at 2h in infected

cells tended to be spliced out at 24h. Our results thus indicate that the splicing of

alternative retained introns during the course of the immune response to infection

is temporally dynamic.

Shortening of 3' UTR regions and increased exon inclusion are evolutionarily

conserved responses to infection in primates. Since the innate immune response

is ancient, one might expect core aspects of this pathway to be evolutionarily

conserved among primates. To explore this question, we used high-depth RNA-seq

data from human, chimpanzee, and rhesus macaque primary whole blood cells (*N=6*

individuals from each species) before and after 4 hours of stimulation with

lipopolysaccharide (LPS), the major component of the outer membrane of Gram-

negative bacteria. Using an analysis pipeline similar to our analysis of infection after

24 hours of infection with *Listeria* or *Salmonella*, we calculated $\Delta\Psi$ values for

alternative RNA processing categories using MISO (Katz et al. 2010) (see

Supplementary Methods, Table S9). As with macrophages, we observed consistent

strong shifts towards increased exon inclusion and shorter 3' UTRs in human whole

blood cells stimulated with LPS. Furthermore, we observed similar trends toward

increased exon inclusion and significant 3' UTR shortening in macaques and

chimpanzees, supporting a conserved response of RNA processing pathways during

innate immune responses in primates (Figure 4B). In addition, these data shows

that the shortening of 3' UTRs and increased exon inclusion are not limited to

isolated macrophages but also observed as part of the immune response engaged by

interacting primary granulocytes and lymphocytes, more closely mimicking *in vivo*

cellular responses to infection.

Increased exon inclusion is associated with increased gene expression levels.

Since we observed strong connections between overall isoform usage and

differential gene expression, we explored the relationship between the occurrence

of particular RNA processing changes and the direction of gene expression changes

in response to infection. We found that genes with significant differences in skipped

exon usage pre- and post-infection (up to 80% of which have increased skipped

exon inclusion after infection) were more likely to be up-regulated after infection

(T-test, both $P \leq 1\times10^{-4}$ for *Listeria* and *Salmonella*; Figure 5A; Figure S8A). This

association remained when gene expression levels were calculated using reads

mapping to constitutive exons only (Figure S8A), eliminating concerns about

alternative spliced isoforms impacting gene expression estimates. Our observation

is consistent with a recent study that reported increased gene expression in tissues

with increased inclusion of evolutionarily novel SEs (Merkin et al. 2015). Genes with

other types of splicing changes showed no trend in expression changes.

Genes with significant skipped exon usage were enriched for Gene Ontology

categories such as "response to stimulus," and also for "RNA processing" and "RNA

stability" categories (Figure S9A). Included within these latter categories are several

members of two major splicing factor families, SR proteins and hnRNPs, which

showed increased inclusion of skipped exons and increased gene expression after

infection (Figure S9B). The SR protein and hnRNP families are known to function in

complex auto- and cross-regulatory networks (Huelga et al. 2012; Pandit et al.

2013), often antagonizing one another's effects on splicing (House and Lynch 2008;

Wang and Burge 2008; Busch and Hertel 2012). Consistent with the slightly higher

up-regulation of members of the hnRNP family, we observed a consistent

enrichment for intronic splicing silencers in upstream intronic regions around

significantly excluded skipped exons (Figure S9C). These elements are often bound

by hnRNPs to promote exclusion of a cassette exon (Wang and Burge 2008))

Variation across individuals provides insight into putative trans-regulatory

factors. A unique feature of our study design is our sampling of 60 individuals after

infection with both *Listeria* and *Salmonella*. By taking advantage of natural variation

in the regulation of RNA processing to infection, we aimed to gain further insight

into the connections between RNA processing and changes in gene expression levels

in response to infection. For each gene, we calculated the correlation between the

inter-individual variation in RNA processing changes ($\Delta\Psi$) and the fold changes in

gene expression levels upon infection. For most categories, we observed shifts in the

distribution of correlations between RNA processing and gene expression levels

relative to permuted controls (Figure S8B), with skipped exons and TandemUTRs

showing the most consistent patterns (Kolmogorov-Smirnov test, $P \leq 0.005$; Figure

5B). Increased skipped exon inclusion correlates with increased gene expression of

the gene across individuals in 69% of the genes with significant skipped exons. In

addition, preferential expression of shorter alternative 3' UTR isoforms tends to be

correlated with increased up-regulation of the associated genes in response to

infection. Our findings thus suggest that RNA processing changes may directly

impact gene expression levels, or at least are commonly regulated with changes in

transcription or mRNA stability.

Substantial inter-individual variation in the genome-wide average RNA processing

patterns also correlated strongly with average gene expression changes upon

infection, particularly for changes in 3' UTR usage. That is, individuals who exhibited more skipped exon inclusion or 3' UTR shortening also tended to exhibit larger global shifts towards up-regulation of gene expression levels upon infection (Figure S10B).

Next, we sought to exploit our relatively large sample size to identify candidate *trans*-factors impacting the amount of skipped exon and TandemUTR changes observed across individuals. Specifically, we examined the relationship between mean changes in exon inclusion and fold change in expression for all expressed genes, across the 60 individuals. When we focus on proteins with known RNA binding functions or domains – hypothesizing that these factors are more likely to regulate post-transcriptional mechanisms – we find many significant correlations (FDR ≤ 1%) for known regulators of the corresponding RNA processing category (Figure S11). For instance, genes whose changes in expression correlated with individual-specific mean 3' UTR shortening included several factors previously implicated in regulating polyadenylation site usage (such as cleavage and polyadenylation factors (Zheng and Tian 2014), hnRNP H proteins (Katz et al. 2010), and hnRNP F (Veraldi et al. 2001); Figure S11A). Additionally, many known regulators of alternative splicing – including a number of SR proteins and hnRNPs – have increased expression levels in individuals with larger extents of skipped exon inclusion (Figure S11B).

3' UTR shortening as a means to evade repression by immune-related miRNAs.
The largest global shift observed in alternative isoform abundance was for 3' UTR shortening, where as many as 98% of significantly changing TandemUTRs shifted towards usage of an upstream polyA site post-infection. Previously, global 3' UTR shortening has been associated with proliferating cells, particularly in the context of cellular transitions in development (Sandberg et al. 2008), cell differentiation (Lackford et al. 2014b), cancerous (Mayr and Bartel 2009) states, or global 3' UTR lengthening in embryonic development (Ji et al. 2009). However, macrophages do not proliferate, which we confirmed with a BrdU labeling assay in both resting and

infected conditions (Figure S12). This result indicates that the pervasive 3' UTR

shortening observed in response to infection is due to an active cellular process that

is independent of cell division and might be mechanistically distinct from that

leading to 3' UTR shortening in proliferating cells.

Previous studies in proliferating cells postulated that 3' UTR shortening can act as a

way to evade regulation by microRNAs (miRNAs), since crucial miRNA target sites

are most often found in 3' UTR regions (Sandberg et al. 2008). To evaluate this

hypothesis we performed small-RNA sequencing in 6 individuals after 2h and 24h of

infection with *Listeria* and *Salmonella*. When focusing specifically on miRNAs

expressed in macrophages (Table S10), we found that the extended UTR regions of

infection-sensitive 3' UTRs have a significantly higher density of miRNA target sites

compared to TandemUTRs that do not change in response to infection (Figure 6A).

This increased density of miRNA target sites was restricted to the extension region,

which is subject to shortening after infection: no difference in the density of target

sites was observed in the common "core" 3' UTR regions of the same genes (Figure

S13).

Next, we tested if the increased density of miRNA target sites was driven by the

enrichment of target sites for particular miRNAs expressed in macrophages. For

each miRNA that was expressed in either non-infected or infected macrophages, we

calculated an enrichment score assessing whether the target sites of that miRNA

were significantly enriched in the extended region of significantly shortened 3' UTRs

(with a background distribution matched for sequence composition; Figure S14; see

Supplemental Methods). We found 15 miRNAs with significantly enriched target

sites (FDR ≤ 10% and enrichment ≥ 1.5 fold) in either the *Listeria* or *Salmonella*

conditions, with 6 miRNAs over-represented in both bacterial conditions (Figure

6B). Interestingly, of the miRNAs with the highest enrichment after infection with

either *Listeria* or *Salmonella*, two (miR-146b and miR-125a) have previously been

shown to be important regulators of the innate immune response (Taganov et al.

2006; Perry et al. 2009; Boldin and Baltimore 2012; Kim et al. 2012; Banerjee et al.

2013). Accordingly, we found that 4 of the miRNAs with the highest enrichment
after infection (miR-125a, miR146b, miR-3661, and miR-151b) are all up-regulated
following infection, and strongly so after 24h of infection (Figure 6B). In the
majority of cases, greater 3' UTR shortening was associated with a stronger increase
in gene expression across individuals, suggesting that shifts towards shorter 3' UTRs
after infection allows these transcripts to escape from repression by specific
immune-induced miRNAs.

DISCUSSION

Taken together, our results provide strong evidence that RNA processing plays a key
role in the regulation of innate immune responses to infection. Despite known
differences in the regulatory pathways elicited in macrophages by *Listeria* and
Salmonella, our data revealed striking similarities in the overall patterns of RNA
processing induced in response to both pathogens. Given the strong overlaps, we
chose throughout this study to focus on consistent patterns across the two bacteria.

Though genes that have differential isoform usage in response to infection are more
likely to show changes in gene expression, we found that as many as 47% of genes
showing differential isoform usage have no evidence for differences in gene
expression upon infection. This observation suggests that a considerable proportion
of genes are regulated predominantly by post-transcriptional mechanisms, with
minimal changes in transcriptional regulation. Differential isoform usage is coupled
to a prominent increase in the diversity of isoforms following infection. Intriguingly,
genes showing the strongest increases in diversity upon infection were strongly
enriched among down-regulated genes (Figure S3B). This observation raises the
possibility that the increased isoform diversity could reflect a shift toward less
stable isoforms, e.g., those targeted by nonsense-mediated decay (NMD), thus
reducing transcript levels (Lewis et al. 2003; Green et al. 2003).

Upon infection of human macrophages with either *Listeria* or *Salmonella*, we
observed substantial shifts towards increased skipped exon inclusion and
shortening of 3' UTRs. These RNA processing patterns remained prominent when
interrogated in human whole blood after an alternative immune stimulus (LPS),
despite the fact the number of significant RNA processing changes in whole blood
was generally smaller than that observed in purified macrophages infected with live
bacteria. This reduced effect is likely explained by the fact that in whole blood, we
are interrogating a very heterogeneous cell population within which only a fraction
(∼3-6%) of the cell types are actually responding to LPS stimulation, such as
monocytes/macrophages (Zarember and Godowski 2002). Additionally,
transcriptional changes induced by live bacteria such as *Listeria* or *Salmonella*,
which concomitantly activates multiple innate immune pathways (Haraga et al.
2008; Pamer 2004), are grander than transcriptional changes that can induced by a
single purified ligand, such as LPS, which activates primarily the TLR4 pathway
(Poltorak et al. 1998). Despite this reduced statistical power, we still observed
conserved RNA processing signatures of immune stimulation through the primate
lineage, supporting the likely functional importance of these cellular responses and
the ubiquity of RNA processing changes after immune stimulation across evolution.
463

The most prominent hallmark of RNA processing changes after infection is a global
shortening of 3' UTRs. Interestingly, this effect resembles the global 3' UTR
shortening seen in proliferating activated T cells and transformed cells (Sandberg et
al. 2008) (Mayr and Bartel 2009) (Lackford et al. 2014b). Given that both T-cell and
macrophage activation result in the activation of shared signaling cascades (e.g., c-
Jun N-terminal kinase (JNK) – a key regulator of apoptosis – and nuclear factor κ-B
(NF-κB) activation), it is possible that, at least to some extent, 3' UTR shortening
observed in T-cells and macrophages stems from the activation of these shared
pathways. Previous studies hypothesized that 3' UTR shortening allows mRNAs to
escape regulation by dynamically changing miRNAs, suggesting that macrophages
may co-opt generally used mechanisms for the same purpose. (Sandberg et al. 2008;

Mayr and Bartel 2009). This shift towards shorter 3' UTRs could result from
increased miRNA-dependent degradation of transcripts with longer 3' UTRs post-
infection. However, genes that shift towards shorter 3' UTRs were more highly
expressed post-infection, making this explanation unlikely. Furthermore, the
increase in overall miRNA target site density is reminiscent of a previous
observation that genes with regulated 3' UTR usage across tissues have more target
sites for ubiquitously expressed miRNAs in their distal 3' UTRs (Lianoglou et al.
2013). However, in contrast to results across tissues, we observe that the target
sites in our distal 3' UTR regions are directly related to miRNAs whose expression
specifically changes after infection.

Specifically, our data support previous hypotheses and find evidence for that 3' UTR
shortening in macrophages is associated with the loss of target sites for a subset of
immune-regulated miRNAs, including miR-146b, miR-125a, and miR-151b (Figure
6B). These miRNAs target the extended 3' UTR regions of several critical activators
of the innate immune response – including the *IRF5* transcription factor (Lawrence
and Natoli 2011) and the MAP kinases *MAPKAP1* and *MAP2K4* (Arthur and Ley
2013) – all of which shift to usage of shorter 3' UTRs after infection. It is tempting to
speculate that without such 3' UTR shortening, the binding of one or more of
miRNAs to the 3' UTR region of these genes would compromise their up-regulation
and the subsequent activation of downstream immune defense pathways. Given that
many miRNAs are involved in either (or both) mRNA degradation and translational
efficiency, we cannot know for sure which of these processes is prevented by the 3'
UTR shortening of these genes.

The convergence towards similar isoform outcomes across many disparate genes
suggests the activation of factors acting in *trans* to drive global shifts towards
inclusion of cassette exons or usage of upstream proximal polyadenyalation sites.
Taking advantage of our relatively large sample size, we were able to identify
candidate *trans*-factors whose up-regulation upon infection might have widespread
impacts on RNA processing patterns, including some known regulators of

alternative splicing or 3'end processing (Figure S11). Sets of RNA binding factors

likely act in combination to influence final transcriptome states (Brooks et al. 2015).

For instance, a general up-regulation of splicing machinery following infection

(Munding et al. 2013), allowing the cell to recognize and use additional splice sites,

might contribute to increased exon inclusion and greater isoform diversity observed

following infection. Members of both the hnRNP and SR protein splicing factor

families show up-regulation of their overall gene expression levels (Figure S9B),

likely contributing to specific instances of exon inclusion as well as exclusion.

Correspondingly, changes in the regulation of many canonical cleavage and

polyadenylation factors were tied to individual-specific shifts in 3' UTR shortening.

The significant correlation between global shifts in ALE usage and TandemUTRs

across individuals further support the notion that cleavage and poladenylation

pathways are regulating the changing 3' UTR landscape upon infection (Figure

S10A). Intriguingly, fold changes in hnRNP H1 gene expression have high

correlations with both skipped exon inclusion and 3' UTR shortening, consistent

with its known splicing function and suggesting that this factor may play multiple

roles in shaping the transcriptome in response to bacterial infection.

An alternative mechanism that has been proposed for the global regulation of 3'

UTR shortening in T-cell activation and neuronal differentiation is a phenomenon

called *telescripting*. This is a phenomenon by which moderately lower U1 snRNA

levels relative to global increases in nascent mRNA production restrict U1's

secondary role in inhibiting premature cleavage and polyadenylation (Berg et al.

2012). While we did not directly measure mRNA levels, we observe a moderate but

significant increase in total RNA concentration after 2 hours of infection that is

consistent with the increase in nascent RNA observed by Berg et al. (Berg et al.

2012) (across 60 samples, paired t-test; P=0.006 for Listeria and P=4.57×10^{-6} for

Salmonella), combined with no changes in the levels of U1 snRNA (relative to 5S

rRNA, Supplemental Methods, Figure S15). Thus it is possible that a delayed

response of U1 snRNA could promote the usage of upstream polyadenylation sites

in both TandemUTRs and ALEs. Interestingly, Berg et al. also observe increased up-
regulation of individual internal exons upon moderate U1 knockdown in neurons
(Berg et al. 2012), which is consistent with our observation of increased skipped
exon inclusion.

Substantial inter-individual variation in the genome-wide average RNA processing
patterns also correlated strongly with average gene expression changes upon
infection, particularly for changes in 3' UTRs and skipped exons. That is, individuals
who exhibited more skipped exon inclusion or usage of upstream polyadenylation
sites also tended to exhibit larger global shifts towards up-regulation of gene
expression levels upon infection (Figure S10B). Though these patterns might result
from stochastic variation in levels of one or several *trans*-factors, they are also likely
to be influenced by genetic differences among individuals. It will be interesting to
ask whether global isoform differences might potentially be reflective of variation in
an individual's susceptibility to infection.

Materials and Methods

Complete details of the experimental and statistical procedures can be found in SI
Materials and Methods. Briefly, blood samples from 60 healthy donors were
obtained from Indiana Blood Center with informed consent and ethics approval
from the Research Ethics Board at the CHU Sainte-Justine (protocol #4022). All
individuals recruited in this study were healthy males between the ages of 18 and
55 y old. Blood mononuclear cells from each donor were isolated by Ficoll-Paque
centrifugation and blood monocytes were purified from peripheral blood
mononuclear cells (PBMCs) by positive selection with magnetic CD14 MicroBeads
(Miltenyi Biotec). In order to derive macrophages, monocytes were then cultured
for 7 days in RPMI-1640 (Fisher) supplemented with 10% heat-inactivated FBS
(FBS premium, US origin, Wisent), L-glutamine (Fisher) and M-CSF (20ng/mL; R&D
systems). After validating the differentiation/activation status of the monocyte-
derived macrophages we infected them at a multiplicity of infection (MOI) of 10:1

for *Salmonella typhimurium* and an MOI of 5:1 for *Listeria monocytogenes* for 2- and

24-hours. For primate comparisons, whole blood was drawn from 6 healthy adult

humans (CHU Sainte-Justine), 6 common chimpanzees, (Texas Biomedical Research

Institute), and 6 rhesus macaques (Yerkes Regional Primate Center). Human

samples were acquired with informed consent and ethics approval from the

Research Ethics Board (CHU Sainte-Justine, #3557). Non-human primate samples

were acquired in accordance with individual institutional IACUCC requirements. For

all species, 1ml of blood was drawn into a media-containing tube (Trueculture tube,

Myriad, US) spiked with ultrapure LPS (Invitrogen, USA) or endotoxin-free water

(control). Samples were stimulated with 1ug/ml LPS, at 37C for 4 hours before total

blood leukocytes were isolated and RNA collected. Genome-wide gene expression

profiles of untreated and infected/treated samples were obtained by RNA-

sequencing for both mRNA transcripts and small RNAs. After a series of quality

checks (SI Materials and Methods), mRNA transcript abundances were estimated

using RSEM (Li and Dewey 2011) and Kallisto (Bray et al. 2016) and microRNA

expression estimates were obtained as previously described(Siddle et al. 2014). To

detect genes with differential isoform usage between two groups of non-infected

and infected samples we used a multivariate generalization of the Welch's t-test.

Shannon entropy H_{sh} (also known as Shannon index) was applied to measure the

diversity of isoforms for each target gene before and after infection. Changes across

individual RNA processing events were quantified using the MISO software package

(v0.4.9) (Katz et al. 2010) using default settings and hg19 version 1 annotations.

Events were considered to be significantly altered post-infection if at least 10% (n

>= 6) of individuals had a BF >= 5 and the |mean $\Delta\Psi$| ≥ 0.05 (Table S6). To conduct a

targeting sequencing of 3' ends of transcripts, We used Smart-3SEQ (Foley et al,

manuscript in preparation), a modified version of the 3SEQ method (Beck et al.

2010) . We used a custom gene ontology script to test for enrichment of functional

annotations among genes that significantly changed isoform usage in response to

infection (SI Materials and Methods). All predicted miRNA target sites within

annotated TandemUTR regions were obtained using TargetScan (v6.2) (Friedman et

al. 2009).

Data Access

601 Data generated in this study have been submitted to the NCBI Gene Expression
602 Omnibus (GEO; http://www.ncbi.hlm.nih.gov/geo/) under the accession number
603 GSE73765, which comprises of the mRNA-seq data (GSE73502) and small RNA-seq
604 data (GSE73478).

Acknowledgements

We thank P Freese for a categorized list of RNA binding proteins, G McVicker for an
initial script to perform iterative gene ontology analyses and P Sudmant for help
parsing primate transcript annotations. We thank Y Katz, JM Taliaferro, P Sudmant, J
Tung, and members of the Barreiro and Burge labs for helpful discussions and
comments on the manuscript. This work was supported by grants from the elife
senio (232519), the Human Frontiers Science Program (CDA-00025/2012) and the
Canada Research Chairs Program (950-228993) (to L.B.B). We thank Calcul Québec
and Compute Canada for managing and providing access to the supercomputer
Briaree from the University of Montreal, used to do many computations reported in
the manuscript. Additional computational resources were performed on an MIT
computational cluster supported by the National Science Foundation (Grant No.
0821391). A.A.P. was supported by a Jane Coffin Childs postdoctoral fellowship. G. B.
was supported by a postdoctoral fellowship from the Fonds de la Recherche en
Santé du Québec.

References
626

Alasoo K, Estrada FM, Hale C, Gordon S, Powrie F. 2015. Transcriptional profiling of
macrophages derived from monocytes and iPS cells identifies a conserved
response to LPS and novel alternative transcription. *bioRxiv*.

Arthur JSC, Ley SC. 2013. Mitogen-activated protein kinases in innate immunity.
Nature Reviews Immunology **13**: 679–692.

Banerjee S, Cui H, Xie N, Tan Z, Yang S, Icyuz M, Thannickal VJ, Abraham E, Liu G.
2013. miR-125a-5p Regulates Differential Activation of Macrophages and
Inflammation. *Journal of Biological Chemistry* **288**: 35428–35436.

Barbosa-Morais NL, Irimia M, Pan Q, Xiong HY, Gueroussov S, Lee LJ, Slobodeniuc V,
Kutter C, Watt S, Çolak R, et al. 2012. The evolutionary landscape of alternative
splicing in vertebrate species. *Science* **338**: 1587–1593.

Beck AH, Weng Z, Witten DM, Zhu S, Foley JW, Lacroute P, Smith CL, Tibshirani R,
van de Rijn M, Sidow A, et al. 2010. 3'-End Sequencing for Expression
Quantification (3SEQ) from Archival Tumor Samples ed. I.O.L. Ng. *PLoS One* **5**:
e8768.

Berg MG, Singh LN, Younis I, Liu Q, Pinto AM, Kaida D, Zhang Z, Cho S, Sherrill-Mix S,
Wan L, et al. 2012. U1 snRNP Determines mRNA Length and Regulates Isoform
Expression. *Cell* **150**: 53–64.

Bihl MP, Heinimann K, Rudiger JJ, Eickelberg O, Perruchoud AP, Tamm M, Roth M.
2002. Identification of a novel IL-6 isoform binding to the endogenous IL-6
receptor. *American journal of respiratory cell and molecular biology* **27**: 48–56.

Boldin MP, Baltimore D. 2012. MicroRNAs, new effectors and regulators of NF-κB.
Immunological Reviews **246**: 205–220.

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq
quantification. *Nature Biotechnology*.

Brooks AN, Duff MO, May G, Yang L, Bolisetty M, Landolin J, Wan K, Sandler J,
Celniker SE, Graveley BR, et al. 2015. Regulation of alternative splicing in
Drosophila by 56 RNA binding proteins. *Genome Research* gr.192518.115.

Busch A, Hertel KJ. 2012. Evolution of SR protein and hnRNP splicing regulatory
factors. *WIREs RNA* **3**: 1–12.

Di Giammartino DC, Nishida K, Manley JL. 2011. Mechanisms and Consequences of
Alternative Polyadenylation. *Molecular Cell* **43**: 853–866.

Friedman RC, Farh KK-H, Burge CB, Bartel DP. 2009. Most mammalian mRNAs are

conserved targets of microRNAs. *Genome Research* **19**: 92–105.

Goodwin RG, Friend D, Ziegler SF, Jerzy R, Falk BA, Gimpel S, Cosman D, Dower SK,
March CJ, Namen AE, et al. 1990. Cloning of the human and murine interleukin-7
receptors: demonstration of a soluble form and homology to a new receptor
superfamily. *Cell* **60**: 941–951.

Gray P, Michelsen KS, Sirois CM, Lowe E, Shimada K, Crother TR, Chen S, Brikos C,
Bulut Y, Latz E, et al. 2010. Identification of a novel human MD-2 splice variant
that negatively regulates Lipopolysaccharide-induced TLR4 signaling. *The*
Journal of Immunology **184**: 6359–6366.

Green RE, Lewis BP, Hillman RT, Blanchette M, Lareau LF, Garnett AT, Rio DC,
Brenner SE. 2003. Widespread predicted nonsense-mediated mRNA decay of
alternatively-spliced transcripts of human normal and disease genes.
Bioinformatics **19**: i118–i121.

Haraga A, Ohlson MB, Miller SI. 2008. Salmonellae interplay with host cells. *Nature*
Reviews Microbiology **6**: 53–66.

House AE, Lynch KW. 2008. Regulation of Alternative Splicing: More than Just the
ABCs. *Journal of Biological Chemistry* **283**: 1217–1221.

Huang Q, Liu D, Majewski P, Schulte LC, Korn JM, Young RA, Lander ES, Hacohen N.
2001. The Plasticity of Dendritic Cell Responses to Pathogens and Their
Components. *Science* **294**: 870–875.

Huelga SC, Vu AQ, Arnold JD, Liang TY, Liu PP, Yan BY, Donohue JP, Shiue L, Hoon S,
Brenner S, et al. 2012. Integrative Genome-wide Analysis Reveals Cooperative
Regulation of Alternative Splicing by hnRNP Proteins. *Cell Reports* **1**: 167–178.

Ishitani A, Geraghty DE. 1992. Alternative splicing of HLA-G transcripts yields
proteins with primary structures resembling both class I and class II antigens.
Proceedings of the National Academy of Sciences **89**: 3947–3951.

Janssens S, Burns K, Tschopp J, Beyaert R. 2002. Regulation of interleukin-1-and
lipopolysaccharide-induced NF-κB activation by alternative splicing of MyD88.
Current Biology **12**: 467–471.

Jensen LE, Whitehead AS. 2001. IRAK1b, a novel alternative splice variant of
interleukin-1 receptor-associated kinase (IRAK), mediates interleukin-1
signaling and has prolonged stability. *Journal of Biological Chemistry* **276**:
29037–29044.

Ji Z, Lee JY, Pan Z, Jiang B, Bin Tian. 2009. Progressive lengthening of 3' untranslated
regions of mRNAs by alternative polyadenylation during mouse embryonic
development ed. J. Valcarcel. *PNAS* **106**: 7028–7033.

Katz Y, Wang ET, Airoldi EM, Burge CB. 2010. Analysis and design of RNA
sequencing experiments for identifying isoform regulation. *Nat Meth* **7**: 1009–
1015.
http://www.nature.com.proxy.uchicago.edu/nmeth/journal/v7/n12/pdf/nmet
h.1528.pdf.

Kawai T, Akira S. 2010. The role of pattern-recognition receptors in innate
immunity: update on Toll-like receptors. *Nat Immunol* **11**: 373–384.

Kim SW, Ramasamy K, Bouamar H. 2012. MicroRNAs miR-125a and miR-125b
constitutively activate the NF-κB pathway by targeting the tumor necrosis factor
alpha-induced protein 3 (TNFAIP3, A20).

Koskinen LL, Einarsdottir E, Dukes E, Heap GA, Dubois P, Korponay-Szabo IR,
Kaukinen K, Kurppa K, Ziberna F, Vatta S, et al. 2009. Association study of the
IL18RAP locus in three European populations with coeliac disease. *Hum Mol*
Genet **18**: 1148–1155.

Lackford B, Yao C, Charles GM, Weng L, Zheng X, Choi EA, Xie X, Wan J, Xing Y,
Freudenberg JM, et al. 2014a. Fip1 regulates mRNA alternative polyadenylation
to promote stem cell self-renewal. *The EMBO Journal* **33**: 878–889.

Lackford B, Yao C, Charles GM, Weng L, Zheng X, Choi EA, Xie X, Wan J, Xing Y,
Freudenberg JM, et al. 2014b. Fip1 regulates mRNA alternative polyadenylation
to promote stem cell self-renewal. *The EMBO Journal* **33**: 878–889.

Lawrence T, Natoli G. 2011. Transcriptional regulation of macrophage polarization:
enabling diversity with identity. *Nature Reviews Immunology* **11**: 750–761.

Lewis BP, Green RE, Brenner SE. 2003. Evidence for the widespread coupling of
alternative splicing and nonsense-mediated mRNA decay in humans. *PNAS* **100**:
189–192.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* **12**: 323.

Li B, Ruotti V, Stewart RM, Thomson JA, Dewey CN. 2010. RNA-Seq gene expression
estimation with read mapping uncertainty. *Bioinformatics* **26**: 493–500.

Lianoglou S, Garg V, Yang JL, Leslie CS, Mayr C. 2013. Ubiquitously transcribed genes
use alternative polyadenylation to achieve tissue-specific expression. *Genes &*
Development **27**: 2380–2396.

Lynch KW. 2004. Consequences of regulated pre-mRNA splicing in the immune
system. *Nature Reviews Immunology* **4**: 931–940.

Martinez NM, Lynch KW. 2013. Control of alternative splicing in immune responses:

many regulators, many predictions, much still to learn. *Immunological Reviews*
253: 216–236.

Mayr C, Bartel DP. 2009. Widespread Shortening of 3′UTRs by Alternative Cleavage
and Polyadenylation Activates Oncogenes in Cancer Cells. *Cell* **138**: 673–684.

Medzhitov R, Horng T. 2009. Transcriptional control of the inflammatory response.
Nature Reviews Immunology **9**: 692–703.

Medzhitov R, Janeway CA Jr. 1998. Innate immune recognition and control of
adaptive immune responses. *Seminars in Immunology* **10**: 351–353.

Merkin JJ, Chen P, Alexis MS, Hautaniemi SK. 2015. Origins and impacts of new
mammalian exons. *Cell Reports* **10**: 1992–2005.

Munding EM, Shiue L, Katzman S, Donohue JP, Ares M Jr. 2013. Competition between
Pre-mRNAs for the Splicing MachineryDrives Global Regulation of Splicing.
Molecular Cell **51**: 338–348.

Nishimura H, Yajima T, Naiki Y, Tsunobuchi H, Umemura M, Itano K, Matsuguchi T,
Suzuki M, Ohashi PS, Yoshikai Y. 2000. Differential roles of interleukin 15 mRNA
isoforms generated by alternative splicing in immune responses in vivo. *The*
Journal of experimental medicine **191**: 157–170.

Ohta S, Bahrun U, Tanaka M, Kimoto M. 2004. Identification of a novel isoform of
MD-2 that downregulates lipopolysaccharide signaling. *Biochem Biophys Res*
Commun **323**: 1103–1108.

O'Connor BP, Danhorn T, De Arras L, Flatley BR, Marcus RA, Farias-Hesson E, Leach
SM, Alper S. 2015. Regulation of Toll-like Receptor Signaling by the SF3a mRNA
Splicing Complex ed. X. Lin. *PLoS Genet* **11**: e1004932.

Pamer EG. 2004. Immune responses to Listeria monocytogenes. *Nature Reviews*
Immunology **4**: 812–823.

Pandit S, Zhou Y, Shiue L, Coutinho-Mansfield G, Li H, Qiu J, Huang J, Yeo GW, Ares M,
Fu X-D. 2013. Genome-wide analysis reveals SR protein cooperation and
competition in regulated splicing. *Molecular Cell* **50**: 223–235.

Perry MM, Williams AE, Tsitsiou E, Larner-Svensson HM, Lindsay MA. 2009.
Divergent intracellular pathways regulate interleukin-1β-induced miR-146a and
miR-146b expression and chemokine release in human alveolar epithelial cells.
FEBS Letters **583**: 3349–3355.

Poltorak A, He X, Smirnova I, Liu M-Y, Van Huffel C, Du X, Birdwell D, Alejos E, Silva
M, Galanos C, et al. 1998. Defective LPS Signaling in C3H/HeJ and C57BL/10ScCr
Mice: Mutations in Tlr4 Gene. *Science* **282**: 2085–2088.

Rao N, Nguyen S, Ngo K, Fung-Leung W-P. 2005. A novel splice variant of
interleukin-1 receptor (IL-1R)-associated kinase 1 plays a negative regulatory
role in Toll/IL-1R-induced inflammatory signaling. *Molecular and Cellular*
Biology **25**: 6521–6532.

Rodrigues R, Grosso AR, Moita L. 2013. Genome-wide analysis of alternative splicing
during dendritic cell response to a bacterial challenge. *PLoS One*.

Sandberg R, Neilson JR, Sarma A, Sharp PA, Burge CB. 2008. Proliferating Cells
Express mRNAs with Shortened 3' Untranslated Regions and Fewer MicroRNA
Target Sites. *Science* **320**: 1643–1647.

Shalgi R, Hurt JA, Lindquist S, Burge CB. 2014. Widespread inhibition of
posttranscriptional splicing shapes the cellular transcriptome following heat
shock. *Cell Reports* **7**: 1362–1370.

Siddle KJ, Deschamps M, Tailleux L. 2014. A genomic portrait of the genetic
architecture and regulatory impact of microRNA expression in response to
infection. *Genome Research*.

Smale ST. 2010. Selective Transcription in Response to an Inflammatory Stimulus.
Cell **140**: 833–844.

Taganov KD, Boldin MP, Chang KJ. 2006. NF-κB-dependent induction of microRNA
miR-146, an inhibitor targeted to signaling proteins of innate immune
responses.

Takagaki Y, Manley JL. 1998. Levels of Polyadenylation Factor CstF-64 Control IgM
Heavy Chain mRNA Accumulation and Other Events Associated with B Cell
Differentiation. *Molecular Cell* **2**: 761–771.

Taliaferro JM, Vidaki M, Oliveira R, Olson S, Zhan L, Saxena T, Wang ET, Graveley BR,
Gertler FB, Swanson MS, et al. 2016. Distal Alternative Last Exons Localize
mRNAs to Neural Projections. *Molecular Cell* **61**: 821–833.

Veraldi KL, Arhin GK, Martincic K, Chung-Ganster LH, Wilusz J, Milcarek C. 2001.
hnRNP F Influences Binding of a 64-Kilodalton Subunit of Cleavage Stimulation
Factor to mRNA Precursors in Mouse B Cells. *Molecular and Cellular Biology* **21**:
1228–1238.

Wang Z, Burge CB. 2008. Splicing regulation: From a parts list of regulatory
elements to an integrated splicing code. *RNA* **14**: 802–813.

Wang Z, Gerstein M, Snyder M. 2009. RNA-Seq: a revolutionary tool for
transcriptomics. *Nat Rev Genet* **10**: 57–63.

Wells CA, Chalk AM, Forrest A, Taylor D, Waddell N, Schroder K, Himes SR, Faulkner

G, Lo S, Kasukawa T, et al. 2006. Alternate transcription of the Toll-like receptor
signaling cascade. *Genome Biology* **7**: R10.

Wong JJL, Ritchie W, Ebner OA, Selbach M, Wong JWH, Huang Y, Gao D, Pinello N,
Gonzalez M, Baidya K, et al. 2013. Orchestrated Intron Retention Regulates
Normal Granulocyte Differentiation. *Cell* **154**: 583–595.

Zarember KA, Godowski PJ. 2002. Tissue Expression of Human Toll-Like Receptors
and Differential Regulation of Toll-Like Receptor mRNAs in Leukocytes in
Response to Microbes, Their Products, and Cytokines. *The Journal of*
Immunology **168**: 554–561.

Zheng D, Tian B. 2014. RNA-binding proteins in regulation of alternative cleavage
and polyadenylation. In *Systems Biology of RNA Binding Proteins*, pp. 97–127,
Springer.

815
816

Figure 1. Gene expression and isoform proportion differences in response to bacterial infection. (A) Principal component analysis of gene expression data from all samples (PC1 and PC2 on the *x-* and *y-axis*, respectively). (B) *ADORA3*, a gene with significant changes in isoform usage upon infection with *Listeria* and *Salmonella*. For each *ADORA3* isoform in response to infection, plotted are the average relative isoform usages across samples (*left panel*) and corresponding fold changes (*right panel*; $\log_2$ scale; with standard error bars). Isoforms are ordered by relative abundance in non-infected samples, and colored circles (*right panel*) correspond to colors in barplot (*left panel*). (C) Number of genes showing only DIU, only DGE, and both DIU and DGE upon infection with *Listeria* and *Salmonella*, (11,353 genes tested).

828
829

Figure 2. RNA processing changes in response to bacterial infection. (A)
Proportion of events for RNA processing category that are significantly changing
after infection with *Listeria* (*left*), *Salmonella* (*middle*), or variation between non-
infected samples as a control (*right*). Numbers indicate the number of significant
changes per category. (B) Significantly Gene Ontology categories for genes with any
significant RNA processing change (FDR ≤ 10%). (C) Distribution of ΔΨ values for
each RNA processing category. Negative values represent less inclusion, while
positive values represent more inclusion, as defined by the schematic exon
representations.

Figure 3. 3'RNA sequencing shows increased usage of upstream polyadenylation sites upon infection. (A) Meta-gene distributions of 3'RNA-seq read densities at the upstream polyA sites (core regions, *left*) and downstream polyA sites (extended regions, *right*) of Tandem 3' UTRs after infection with *Listeria* or *Salmonella* (*top* and *bottom*, respectively). Shown are the read distributions for non-infected samples across all Tandem 3' UTRs (*black*) and infected samples at Tandem 3' UTRs that significantly change after infection (*yellow*) or show no change after infection (*blue*), as called by the RNA-seq data. (B) Distribution of ΔΨ values calculated from 3'RNA-seq data for Tandem 3' UTRs. We observe significant shifts (*P* < 2.2 × 10⁻¹⁶ for both *Listeria* and *Salmonella*) towards negative ΔΨ values in Tandem UTRs that are identified as significantly changing in RNA-seq data (*yellow*) relative to Tandem UTRs without any change after infection (*blue*).

Figure 4. Directed shifts in RNA processing persist across time, stimulus conditions, and closely related species. (A) Correlations between ΔΨ values after 2 hours of infection (*x-axis*) and ΔΨ values after 24 hours of infection (*y-axis*), presented as density plots per RNA processing category in contrasting colors. Plotted are events that are significant after 2 hours of either *Listeria* or *Salmonella* infection. (B) Distributions of ΔΨ values for skipped exons (*purple*) and TandemUTRs (*blue*) following infection of whole blood cells with lipopolysaccharide, assessed in human (*top*), chimpanzee (*middle*) and rhesus macaque (*bottom*) individuals *(N=6* per species). We observe prominent global shifts in isoform distributions in human and macaque (SE$_{human}$ *P=0.003*, TandemUTR$_{human}$ *P=2.4×10^{-13}*; SE$_{macaque}$ *P=0.05*, TandemUTR$_{macaque}$ *P=8.8×10^{-6}*) with a more modest trend observed in chimpanzee, potentially due to poor transcript annotations in the chimpanzee genome (SE$_{chimpanzee}$ *P=0.26*, TandemUTR$_{chimpanzee}$ *P=0.002*). All p-values were calculated using a Student's t-test testing deviation from a mean of zero.

878
879

Figure 5. Relationship between RNA processing changes and gene expression
changes. (A) Distribution of fold changes in gene expression (*y-axis*, $\log_2$ scale) for
genes with significant skipped exon changes after infection (*purple*) and genes with
no change after infection (*gray*). (B) Distribution of Spearman correlations between
$\Delta\Psi$ and fold change in gene expression across 60 individuals per gene (*solid line*),
for genes with only one annotated alternative event and significantly changed SE
usage (*purple*; n_L = 46 genes and n_S = 97 genes) or tandem UTR usage (*blue*; n_L = 36
genes and n_S = 86 genes). Dotted lines show distribution of the correlation
coefficients after permuting $\Delta\Psi$ values.

Figure 6. Tandem 3' UTR shortening allows evasion of regulation by miRNAs.
(A) Distribution of frequency of miRNA target sites per nucleotide in the extended
regions of Tandem UTRs that either show no change after infection (*grey*) or
significantly change after infection (*blue*). (B) Significantly enriched miRNA target
sites (FDR ≤ 10%, |FC| > 1.5) in the extended regions of significantly changing
Tandem UTRs after infection with *Listeria*-only (*top*), with *Salmonella*-only (*middle*),
or both bacteria (*bottom*). For each bacteria, the barplots in the left panels show the
fold enrichment (*x-axis*, $\log_2$ scale) of target sites in the extended regions. White bars
represent non-significant enrichments. Panels on the right show the fold change in
miRNA expression (*x-axis*, $\log_2$ scale with standard error bars) after either 2 hours
of infection (*light colors*) or 24 hours of infection (*dark colors*).