1 **Title: Evolutionary assembly patterns of prokaryotic genomes**
2

3 **Authors:** Maximilian O. Press[1], Christine Queitsch[1], Elhanan Borenstein[1,2,3]*
4
Affiliations
[1]: Department of Genome Sciences, University of Washington, Seattle, WA, USA
[2]: Department of Computer Science and Engineering, University of Washington, Seattle, WA,
USA
[3]: External Faculty, Santa Fe Institute, Santa Fe, NM, USA
*Correspondence to: elbo@uw.edu

11
12 **Running title:** Evolutionary assembly of prokaryotic genomes
13

14 **Keywords**
15 Genome evolution, prokarya, prokaryote, constraint, epistasis, comparative method, evolutionary
16 predictability, RuBisCO, parallel evolution.

Abstract:

Evolutionary innovation must occur in the context of some genomic background, which limits

available evolutionary paths. For example, protein evolution by sequence substitution is

constrained by epistasis between residues. In prokaryotes, evolutionary innovation frequently

happens by macrogenomic events such as horizontal gene transfer (HGT). Previous work has

suggested that HGT can be influenced by ancestral genomic content, yet the extent of such gene-

level constraints has not yet been systematically characterized. Here, we evaluated the

evolutionary impact of such constraints in prokaryotes, using probabilistic ancestral

reconstructions from 634 extant prokaryotic genomes and a novel framework for detecting

evolutionary constraints on HGT events. We identified 8,228 directional dependencies between

genes, and demonstrated that many such dependencies reflect known functional relationships,

including, for example, evolutionary dependencies of the photosynthetic enzyme RuBisCO.

Modeling all dependencies as a network, we adapted an approach from graph theory to establish

chronological precedence in the acquisition of different genomic functions. Specifically, we

demonstrated that specific functions tend to be gained sequentially, suggesting that evolution in

prokaryotes is governed by functional assembly patterns. Finally, we showed that these

dependencies are universal rather than clade-specific and are often sufficient for predicting

whether or not a given ancestral genome will acquire specific genes. Combined, our results

indicate that evolutionary innovation via HGT is profoundly constrained by epistasis and

historical contingency, similar to the evolution of proteins and phenotypic characters, and

suggest that the emergence of specific metabolic and pathological phenotypes in prokaryotes can

be predictable from current genomes.

INTRODUCTION:

A fundamental question in evolutionary biology is how present circumstances affect future adaptation and phenotypic change (Gould and Lewontin 1979). Studies of specific proteins, for example, indicate that epistasis between sequence residues limits accessible evolutionary trajectories and thereby renders certain adaptive paths more likely than others (Weinreich et al. 2006; Gong et al. 2013; de Visser and Krug 2014; Harms and Thornton 2014). Similarly, both phenotypic characters (Ord and Summers 2015) and specific genetic adaptations (Christin et al. 2015; Conte et al. 2012) show strong evidence of parallel evolution rather than convergent evolution. That is, a given adaptation is more likely to repeat in closely related organisms than in distantly related ones. This inverse relationship between the repeatability of evolution and taxonomic distance implies a strong effect of lineage-specific contingency on evolution, also potentially mediated by epistasis (Orr 2005).

Such observations suggest that genetic adaptation is often highly constrained and that the present state of an evolving system can impact future evolution. Yet, the studies above are limited to small datasets and specific genetic pathways, and a more principled understanding of the rules by which future evolutionary trajectories are governed by the present state of the system is still lacking. For example, it is not known whether such adaptive constraints are a feature of genome-scale evolution or whether they are limited to finer scales. Moreover, the mechanisms that underlie observed constraints are often completely unknown. Addressing these questions is clearly valuable for obtaining a more complete theory of evolutionary biology, but more pressingly, is essential for tackling a variety of practical concerns including our ability to combat evolving infectious diseases or engineer complex biological systems.

Here, we address this challenge by analyzing horizontal gene transfer (HGT) in

prokaryotes. HGT is an ideal system to systematically study genome-wide evolutionary

constraints because it involves gene-level innovation, occurs at very high rates relative to

sequence substitution (Nowell et al. 2014; Puigbò et al. 2014a), and is a principal source of

evolutionary novelty in prokaryotes (Gogarten et al. 2002; Jain et al. 2003; Lerat et al. 2005;

Puigbò et al. 2014b). Clearly, many or most acquired genes are rapidly lost due to fitness costs

(van Passel et al. 2008; Baltrus 2013; Soucy et al. 2015), indicating that genes retained in the

long term are likely to provide a selective advantage. Moreover, not all genes are equally

transferrable (Jain et al. 1999; Sorek et al. 2007; Cohen et al. 2011), and not all species are

equally receptive to the same genes (Smillie et al. 2011; Soucy et al. 2015). However,

differences in HGT among species have been attributed not only to ecology (Smillie et al. 2011)

or to phylogenetic constraints (Nowell et al. 2014; Popa et al. 2011), but also to interactions with

the host genome (Jain et al. 1999; Cohen et al. 2011; Popa et al. 2011). Indeed, studies involving

single genes or single species support the influence of genome content on the acquisition and

retention of transferred genes (Pal et al. 2005; Iwasaki and Takagi 2009; Chen et al. 2011; Press

et al. 2013; Sorek et al. 2007; Johnson and Grossman 2014). For example, it has been

demonstrated that the presence of specific genes facilitates integration of others into genetic

networks (Chen et al. 2011), and that genes are more commonly gained in genomes already

containing metabolic genes in the same pathway (Pal et al. 2005; Iwasaki and Takagi 2009).

However, to date, a systematic, large-scale analysis of such dependencies has not been presented.

In this paper, we therefore characterize a comprehensive collection of genome-wide

HGT-based dependencies among prokaryotic genes, uncover potential rules of genome evolution

in prokaryotes, and demonstrate that the acquisition of genes is to some extent predictable based

on these rules. Overall, our study suggests that genetic innovation and adaptation are

substantially constrained through gene-level epistatic interactions such as those that we describe

influencing HGT.

RESULTS:

PGCE Inference

We first set out to detect pairs of genes for which the presence of one gene in the genome

promotes the gain of the other gene (though not necessarily *vice versa*) (Figure 1). Such "pairs of

genes with conjugated evolution" (PGCEs) represent putative epistatic interactions at the gene

level and may guide genome evolution. To this end, we obtained a collection of 634 prokaryotic

genomes, annotated by KEGG (Kanehisa et al. 2012), and linked through a curated phylogeny

(Dehal et al. 2010). For each of the 5801 genes that varied in presence across these genomes, we

reconstructed the probability of this gene's presence or absence on each branch of the

phylogenetic tree using a previously introduced method (Cohen and Pupko 2010), as well as the

probability that it was gained along these branches. We further confirmed that genes'

presence/absence was robust to the reconstruction method employed (99.5% agreement between

reconstruction methods used; Methods). From these reconstructions, we estimated the frequency

with which each gene was gained in the presence of each other gene, and followed previous

studies (Maddison 1990; Cohen et al. 2012) in using parametric bootstrapping (Figure S1) to

detect PGCEs – gene pairs for which one gene is gained significantly more often in the presence

of the other (Figure S2, SI Text). In total, we identified 8,415 PGCEs. We finally applied a

transitive reduction procedure to discard potentially spurious PGCEs, resulting in a final network

containing 8,228 PGCEs connecting a total of 2,260 genes (Figures S3, S4, SI Text). A detailed

description of the procedures used can be found in Methods, and the final list of PGCEs is

supplied as File S1.

PGCEs represent biologically relevant dependencies

Comparing this final set of PGCEs to known biological interactions, we confirmed that the

obtained PGCEs represent plausible biological dependencies. For example, genes sharing the

same KEGG Pathway annotations were more likely to form a PGCE (Figure 2A), as were genes

that are linked in an independently-derived network of bacterial metabolism (Levy and

Borenstein 2013) (Figure 2B). Moreover, PGCEs often linked genes in functionally related

pathways (Figure S5, SI Text). We similarly identified specific examples in which PGCEs

connected pairs of genes with well-described functional relationships. One such example is the

PGCE connecting *rbsL* and *rbsS* (sometimes written *rbcL*/*rbcS*), two genes that encode the large

and small subunits of the well-described photosynthetic enzyme ribulose-1-5-bisphosphate

carboxylase-oxygenase (RuBisCO), respectively. The rbsL subunit alone has carboxylation

activity in some bacteria, but the addition of rbsS increases enzymatic efficiency, consistent with

its PGCE dependency on *rbsL* (Figure 3A) (Andersson and Backlund 2008). Moreover, these

genes are known to undergo substantial horizontal transfer (Delwiche and Palmer 1996).

Multiple additional genes were found to promote *rbsS* gain (88 PGCEs in total, Table

S1), many of which, as expected, are associated with carbon metabolism. Other genes in this set,

however, unexpectedly implicated nitrogen acquisition, as well as other pathways (Table S2), in

promoting *rbsS* gain. For example, all components of the *urt* urea transport complex had a PGCE

link with *rbsS*, as shown by the reconstructed phylogenetic history of *urtA* and *rbsS* (Figure 3B).

This strict dependency could reflect nitrogen's role as a rate-limiting resource for primary

production in phytoplankton and other photosynthetic organisms (Eppley and Peterson 1979;

Sohm et al. 2011). In comparing the reconstructions from which *urtA-rbsS* and *rbsL-rbsS*

dependencies were inferred, we further observed that *rbsS* is gained only in lineages where both

dependencies were previously present. This indicates that while both *rbsL* and *urtA* may be

necessary for the acquisition of *rbsS*, neither *rbsL* nor *urtA* are independently sufficient for the

acquisition of *rbsS*. Other PGCEs may interact in similarly complex fashions in controlling the

acquisition of genes, and thus such relationships may be gene-specific and involve a variety of

biological mechanisms that may be difficult to generalize. For further analyses, we therefore

focused on analyzing large-scale patterns of PGCE connectivity and on exploring how the

dependencies between various genes structure the relationships between functional pathways.

PGCE network analyses reveal evolutionary assembly patterns

The *rbsS*-associated PGCEs described above show how PGCEs captured an assembly pattern

involving multiple pathways. Therefore, we next set out to infer global evolutionary assembly

patterns based on the complete set of PGCEs identified. Specifically, we used a network-based

topological sorting approach (SI Text) to rank all genes in the PGCE network. According to this

procedure, genes without dependencies occupy the first rank, genes in the second rank have

PGCE dependencies only on first rank genes, genes in the third rank have dependencies only on

first and second rank genes, and so on until all genes are associated with some rank. In other

words, the obtained ranking represents general patterns in the order by which genes are gained

throughout evolution, with the gain of higher-ranked genes succeeding the presence of the lower-

ranked genes on which they depend. Using this approach, we found that genes could be fully

classified into five ranks (Fig 4A). The first rank was by far the largest at 1,593 genes (most

genes do not have detectable dependencies), the second rank had 498 genes, and successive

ranks showed declining membership until the last (fifth) rank, with only 5 genes (Table S3).

To identify evolutionary assembly patterns from these ranks, we examined the set of genes in each rank and identified overrepresented functional categories (Table 1). These enriched functional categories indicate that certain functional groups of genes consistently occupy specific positions in these evolutionary assembly patterns, whether in controlling other genes' gain or in being controlled by other genes. For example, we found that the first rank was enriched for flagellar and piliar genes involved in motility, in addition to Type II secretion genes (many of which are homologous to or overlap with genes encoding piliar proteins) and certain two-component genes. The second rank was enriched for various metabolic processes, whereas later ranks were enriched for Type III and Type IV secretion systems and conjugation genes (Table 1). This finding suggests that habitat commitments are made early in evolution, mediated by motility genes that could underlie the choice and establishment of physical environments. This environmental choice is followed by a metabolic commitment to exploiting the new habitat. Last, genes for interaction with the biotic complement of these habitats are gained, and replaced frequently in response to evolving challenges. Considering two distinct but highly homologous pilus assembly pathways, one (fimbrial) was enriched in a low rank and one (conjugal) was enriched in a high rank, suggesting that the specific function of the gene rather than other sequence-level gene properties drove the ranking (Figure S6A). We additionally confirmed that the observed rank distribution for these functions is not explained by variation in the frequency of gene gain (Figure S6B). Furthermore, as expected, we observed that the gains of genes appearing late in the sort were overrepresented in later branches of the tree compared to the gains

of lower-ranked genes (Figures 4B, S7), suggesting that the chronology of gene acquisition

reflects the overall assembly patterns in gain order.

Evolution by HGT is predictable

The chronological ordering of ranks was relatively consistent across the tree (Figure 4B),

indicating that PGCE dependencies are universal across prokaryotes. Notably, this universality

also implies that gene acquisition is predictable from genome content. Put differently, if PGCEs

are universal, then PGCEs inferred in one clade of the tree are informative in making predictions

about gene acquisition in a different clade. Indeed, studies of epistasis-mediated protein

evolution indicate that the constriction of possible mutational paths should lead to predictability

in evolution, if epistasis is sufficiently strong (Weinreich et al. 2006). To explore this hypothesis

explicitly, we partitioned the tree into training and test sets (Figure 5A). As test sets, we selected

the Firmicutes phylum, and the Alphaproteobacteria/Betaproteobacteria subphyla. Choosing

whole clades as test sets (rather than randomly sampling species from throughout the tree)

guarantees that true predictions are based on universal PGCEs, rather than clade-specific PGCEs.

For each test set, we used a model phylogeny that excluded the test subtree as a training set, and

inferred PGCEs based on this pruned tree (Table S4, Figure S8A). We then used these inferred

PGCEs to score the likelihood of the gain of dependent genes on each branch in the test set,

based on the genome content of the branch's ancestor (Figure 5A, Table S4, SI Text). We used a

naïve and simplistic score: the proportion of genes upon which the gained gene depends that are

present in the reconstructed ancestor of each branch. In both test sets, we found that prediction

quality was surprisingly high (Figure 5B, Figure S8B), suggesting that PGCEs are taxonomically

universal and statistically robust in describing relationships between genes. This predictability is

consistent with the hypothesis that gene-gene dependencies constrain the evolution of genomes

by HGT. More broadly, this analysis and our finding that PGCEs predictably determines future evolutionary gains provide substantial evidence that the preponderance of parallel evolution over convergent evolution (Ord and Summers 2015; Conte et al. 2012) may be the result of specific, identifiable genetic dependencies that similarly impact the evolutionary trajectory taken by similar genomes.

DISCUSSION:

Combined, our findings provide substantial evidence to suggest that gene acquisitions in bacteria are governed by genome content through numerous gene-level dependencies. Our ability to detect these underlying dependencies is clearly imperfect, owing to various data and methodological limitations (SI Text, Figure S2). In reality the complete dependency network is therefore likely much denser than that described above and includes numerous dependencies and constraints that our approach may not be able to detect. Consequently, our estimates should be considered as a lower bound on the extent of gene-gene interactions and accordingly the predictability of HGT.

Notably, even considering such caveats, our observations dramatically expand our knowledge of the constraints on HGT. Previous studies of such constraints demonstrated that genes frequently acquired by HGT tend to occupy peripheral positions in biological networks (Jain et al. 1999; Cohen et al. 2011), are often associated with specific cellular functions, and are phylogenetically clustered. These observations suggested that properties of transferred genes are also important determinants of HGT regardless of recipient genome content (Jain et al. 1999; Cohen et al. 2011; Gophna and Ofran 2011) and that the acquisition of certain genes is clade-specific (Popa et al. 2011; Andam and Gogarten 2011). In contrast, our analysis demonstrates the importance of recipient genome content in strongly influencing the propensity of a new gene to

be acquired. In fact, to some extent, properties previously reported as determining the general

"acquirability" of genes across all species may reflect some average constraint across genomes.

By considering also variation in genomes acquiring genes, our analysis focused on specific

biological effects, whose strengths may vary from genome to genome.

Importantly, our model that gene acquisition is affected by recipient genome content is

consistent with the observed enrichment of HGT among close relatives, which presumably have

similar genome content (Gogarten et al. 2002; Andam and Gogarten 2011; Popa et al. 2011;

Popa and Dagan 2011). This taxonomic clustering of innovation by HGT is also in agreement

with previous studies that demonstrated that phenotypic (Gould and Lewontin 1979; Ord and

Summers 2015) and genetic (Conte et al. 2012; Christin et al. 2015) parallel evolution is more

common than convergent evolution, potentially due to the effects of contingency (Gould and

Lewontin 1979; Conte et al. 2012; Christin et al. 2015; Ord and Summers 2015). However, in

contrast to other studies, we present direct evidence that the mechanism by which contingency

controls evolution is epistasis. Furthermore the universality of the PGCEs shows that the

constraints underlying the effect of contingency operate outside the context of parallel evolution.

It should also be noted that while our analysis revealed several intriguing patterns, the

precise interpretation of some of these patterns remains unclear. For instance, the observed

correspondence of topological ranks of genes to chronology suggests that evolutionary age is a

potential contributor to such ranking, especially considering that our reconstructions likely lack

many genes that have not been retained in any extant genomes. However, the biological

plausibility and statistical robustness of PGCEs demonstrated above strongly argue that the

observed evolutionary patterns are the result of constraint-inducing dependencies. Future work

may therefore aim to quantify the trade-off between functional and chronological determinants in

apparent evolutionary constraints.

Finally, we demonstrate the predictability of genomic evolution by horizontal transfer

from current genomic content. As stated above, this finding also suggests that such dependencies

are fairly universal across the prokaryotic tree. It should be noted that our approach was designed

specifically to understand the PGCE network's significance and universality, rather than predict

gene acquisition. It is likely that an approach specifically engineered for gene acquisition

prediction would substantially outperform our approach. The estimates of predictability of

genomic evolution presented here are accordingly quite conservative.

The determinism and predictability of evolutionary patterns therefore appear to be an

outcome not only of intramolecular epistasis in proteins or phylogenetic constraints, but also of

genome-wide interactions between genes. This suggests that the evolution of medically,

economically, and ecologically important traits in prokaryotes depends on ancestral genome

content and is hence at least partly predictable, potentially informing research in the

epidemiology of infectious diseases, bioengineering, and biotechnology.

METHODS

All mathematical operations and statistical analyses were performed in R 2.15.3 (2012).

Probabilistic ancestral reconstructions were obtained using the *gainLoss* program (Cohen and

Pupko 2010). Phylogenetic simulations and plots were performed with the APE library (Paradis

et al. 2004). Network analyses and algorithms were implemented using either the *igraph* (Csardi

and Nepusz 2006) or *NetworkX* (Hagberg et al. 2013) libraries, and visualized using Cytoscape

v3.1.1 (Shannon et al. 2003).

Phylogenies

We used a pre-computed phylogenetic tree (Dehal et al. 2010) as a model of bacterial evolution. We mapped all extant organisms in this tree to organisms in the KEGG database by their NCBI genome identifiers, and pruned all tips that did not directly and uniquely map to KEGG. This yielded a phylogenetic tree connecting 634 prokaryotic species. For analyses involving subtrees of this phylogenetic tree, we used iTOL (Letunic and Bork 2011) to extract subtrees.

Inferring phylogenetic histories for genes

We used the *gainLoss* v1.266 software (Cohen and Pupko 2010), a set of presence/absence patterns of orthologous genes from KEGG (Kanehisa et al. 2012), and the phylogenetic tree described above to infer 1) the probabilities of presence and absence of genes at internal nodes of the tree, 2) gain and loss rates of each gene, and 3) tree branch lengths within a single model. We obtained a probabilistic ancestral reconstruction based on stochastic mapping for each of 5801 genes that were present in at least one species and absent in at least one species, and filtered out genes that were found to be gained less than twice throughout the tree, yielding 5031 genes which we further analyzed. We used the probabilities of presence and absence of each of these 5031 genes at each node and tip on the tree to compute the probability of each branch experiencing 1) gain (absent in ancestor and present in descendant) and 2) presence (present in both ancestor and descendant; Supporting Text). For a gene X on a branch with ancestor A and descendant B, we assume:

1. Pr(X *present* on branch) = Pr(X present in A ∩ X present in B) =

   Pr(X present in A) * Pr(X present in B)

2. Pr(X *gained* on branch) = Pr(X absent in A ∩ X present in B) =

Pr(X absent in A) * Pr(X present in B)

Note that these probability estimates are distinct from those obtained by using the *gainLoss*

continuous-time Markov chain on the same ancestral reconstruction, which consider also

hypothetical gains that are not retained and are thus not relevant to our analysis (Supporting

Text).

Robustness analysis of reconstruction method

We used a maximum-parsimony reconstruction as inferred by *gainLoss* to benchmark the

accuracy of the *gainLoss* reconstruction by stochastic mapping. In this analysis, only internal

node reconstructions were considered, as tip reconstructions (for which the states are known) are

not informative about algorithm performance. Since the maximum-parsimony reconstruction is

binary (presence/absence) and the stochastic mapping reconstruction is probabilistic, for

purposes of comparison we rounded the probabilities of the stochastic mapping reconstruction to

obtain a presence/absence reconstruction (*i.e.*, a probability >0.5 denotes presence and <=0.5

denotes absence). We computed the agreement between the two reconstructions as the

percentage of internal node reconstructions that agree on the state of the gene.

Quantifying PGCEs

We defined a "pair of genes with conjugated evolution" (PGCE) as a gene pair (i, j) for which

the presence of one gene i encourages the gain of the other, j Considering these genes as

phylogenetic characters, we therefore aim to detect pairs for which "gain" state transitions for

character j are enriched on branches where character i remains in the "present" state. This

problem is related to previous methods for detecting coevolution or correlation between

phylogenetic characters (Maddison 1990; Huelsenbeck et al. 2003; Cohen et al. 2012). Given N

branches and k genes, there are 2 N X k matrices, P and G, describing the probabilities,

respectively, of presence and gain of each gene along each branch. The test statistic for each

gene pair (i, j) is the probabilistic count of branches where the gain of gene j occurs, while

conditioning on the presence of gene i (cell C_{ij} in a k x k matrix C). To compute C across N

branches, we sum the conditional probabilities of the gain of gene j in the presence of gene i

across the tree, *i.e.* the products of the two N x k matrices, P (presence) and G (gain), for each

gene pair:

$$C_{ij} = \sum_{n=1}^{N} G_{ni} P_{nj}$$

Entries in C which are significantly larger than a null expectation of gains represent PGCEs

between the row and column genes of C.

Null distribution for PGCEs

For two independently evolving genes i and j, the counted gains of j in the presence of i, C_{ij}, will

be distributed under the null hypothesis (independent evolution) as some function of the

prevalence of i (the sum of P_i, the vector of probabilities of presence of i across branches of the

tree), the probabilistic count of gains experienced by j (the sum of G_j, the vector of probabilities

of gains of j across nodes of the tree), and the topology and branch lengths of the tree (τ):

$$C_{ij} \sim f(P_i, G_j, \tau)$$

As this distribution may be difficult to formalize for a specific dataset, we followed previous

studies (Cohen et al. 2012; Huelsenbeck et al. 2003; Maddison 1990) and approximated this null

distribution via parametric bootstrapping. Specifically, we simulated the evolution of 10^5 genes

along the tree using the APE library function *rTraitDisc()* (Paradis et al. 2004). For the gain and

loss rates used in these simulations, we used *gainLoss* gain and loss rates estimated for the 5801

empirical genes. We fit gamma distributions to these values by maximum likelihood using the

function *fitdistr()* from the MASS library (Venables and Ripley 2002). For both gains and losses,

we increased the shape parameter of the gamma distribution (by a factor of 3 for gains, 1.5 for

losses), to ensure that simulated genes showed sufficiently large numbers of gains. This was

necessary because parametric bootstrapping with the rates inferred by *gainLoss* resulted in left

skewed distributions of gene gains (compare Figures S1A, S1C, and S1E), which were likely to

confound null models, whereas for our null models to be applicable for this analysis, the

distribution of simulated gene gains should be roughly similar to the distribution of gains of

empirical genes (see Figure S1, Supplementary Text).

These simulated genes should evolve independently and thus represent a null model for

PGCEs. As above, we constructed matrices representing the probabilities of presence and gain of

these 10^5 genes across all of the branches of the phylogeny (P_{null} and G_{null}). We then multiplied

these matrices of simulated genes to compute a 10^5 x 10^5 matrix C_{null} of probabilistic counts. As

a null distribution for each pair of genes i and j with $C_{ij} > 1$ (those with $C_{ij} \leq 1$ are not

informative), we used the 1000 simulated genes with prevalence closest to gene i (rows of C_{null}),

and the 1000 simulated genes with a number of gains closest to gene j (columns of C_{null}). We

used the 10^6 simulated observations in the resulting submatrix of C_{null} as a null distribution for

C_{ij}. Notably, C_{ij} represents probabilistic counts, whereas C_{null} represents integer counts (the true

reconstruction is known). Consequently, we floored values in C_{ij}, such that all counts were

truncated at the decimal point. The comparison of C_{ij} to this null distribution yields an empirical

p-value; we rejected the null hypothesis of independence between genes i and j for the C_{ij}

observation at a 1% false discovery rate (Benjamini and Hochberg 1995) ($P < 7 \times 10^{-6}$).

Constructing a PGCE network.

For each entry in C_{ij} for which we observed a significant association, we recorded an edge from

gene i to gene j in a network of PGCEs. To focus purely on direct interactions, we subjected this

network to a transitive reduction (Hsu 1975). This reduction requires a directed acyclic graph

(DAG). To identify the largest possible DAG in our PGCE network, we identified and removed

the minimal set of edges inducing cycles (Supplementary Text). We performed a transitive

reduction of the resulting DAG using Hsu's algorithm (Hsu 1975) (Supplementary Text).

Mapping biological information to the network.

We used network rewiring (as implemented in the *rewire()* function of the *igraph* library (Csardi

and Nepusz 2006)) to generate null distributions of the PGCE network by randomly exchanging

edges between pairs of connected nodes, while excluding self-edges. In each permutation, we

performed 5N rewiring operations, where there are N edges in the network, to ensure sufficient

randomization. To estimate the relationship between the PGCE network and biological

information we calculated the number of edges shared between the PGCE network and a

metabolic network of all bacterial metabolism obtained from KEGG (Kanehisa et al. 2012; Levy

and Borenstein 2013), and the number of edges shared between members of the same functional

pathway as defined by KEGG, in both the original and randomized networks.

To determine whether genes with certain functional annotations were more likely to associate

with one another in the PGCE network, we examined the KEGG Pathway annotations of each

pair of genes in the network. We counted the number of edges leading from each pathway to

each other pathway, and obtained an empirical p-value for this count by comparing it to a null

distribution of the expected counts obtained by random rewiring as above.

Topological sorting of PGCE networks

To identify global patterns in our PGCE network, we performed topological sorting (Kahn 1962)

with grouping. Topological sorting finds an absolute ordering of nodes in a directed acyclic

graph (DAG), such that no node later in the ordering has an edge directed towards a node earlier

in the ordering. Grouping the sort allows nodes to have the same rank in the ordering if

precedence cannot be established between them, giving a unique solution. For a description of

the algorithm used, see Supplementary Text.

Prediction of HGT events on branches.

We used the PGCE network to predict the occurrence of specific HGT events (gene acquisitions)

on the tree in the following fashion. We used two test/training set partitions, with the clades of

Firmicutes and the Alpha/Betaproteobacteria as independent test sets, and the training sets as the

rest of the tree without these clades. To "train" PGCE networks, we performed ancestral

reconstruction of gene presence, PGCE inference, and network processing just as for the entire

tree. We only attempted to predict genes with at least one PGCE dependency ("predictable"

genes). We then considered each branch in the test set independently, attempting to predict

whether each predictable gene was gained on that branch based on the reconstructed genome at

the ancestor node. For each predictable gene-branch combination, our prediction score was the

proportion of the predictable gene's PGCE dependencies that are present in the ancestor. This is

the dot product of the gene presence/absence pattern of the ancestor node (A_i across i potentially

present genes) and a binary vector denoting which genes in the PGCE network the predictable

gene depends on (P_i across i genes in potential PGCEs), scaled by P_i:

$$score = \frac{\sum A_i P_i}{\sum P_i}$$

Note that this value ranges between 0 and 1 for each predicted gene. As true gains, we used our

reconstructed gene acquisition events for each branch in the test set. We arbitrarily called any

predictable gene-branch pair with a Pr(gain) > 0.5 as a gain, and any predictable gene-branch

pair with Pr(gain) <= 0.5 as no gain. We filtered out any gene-branch pair where the gene was

known to be present with Pr > 0.4, as in these cases the gene is probably already present. We

analyzed the accuracy of our prediction scores using receiver operating characteristic (ROC)

analysis and by comparing scores of the gain branches to those of the no-gain branches.

Data Access

Data are available at http://figshare.com/s/1f341994624c11e5b23706ec4bbcf141, along with

code for performing analyses.

Acknowledgements

We are obliged to members of the Borenstein and Queitsch laboratories, and to Evgeny

Sokurenko, Joe Felsenstein, and Willie Swanson for helpful discussions. We thank Ofir Cohen

for help with the *gainLoss* program. MOP was supported in part by National Human Genome

Research Institute Interdisciplinary Training in Genome Sciences Grant 2T32HG35-16. CQ is

supported by National Institute of Health New Innovator Award DP2OD008371. EB is

supported by National Institute of Health New Innovator Award DP2AT00780201. We thank

UW Genome Sciences Information Technology Services for high-performance computing

resources.

3
4

FIGURE LEGENDS

Figure 1. Workflow for deriving the PGCE network. (A): a model phylogeny and a set of

gene presence/absence patterns at the tips are used to generate an ancestral reconstruction, from

which gains are inferred. Filled circles represent the presence of a gene (distinguished by color),

empty circles represent absence of that gene. Inverted triangles represent points on the phylogeny

where the gene of the indicated color is inferred to be gained. (B): Based on inferred gain and

loss rates, many evolutionary scenarios are independently simulated and used as a null

expectation for evolutionary independence. Filled circles indicate presence of the simulated gene

and empty circles indicate absence, inverted triangles represent gains of the simulated gene on

the phylogeny. (C): A null distribution derived from simulated gene evolution is used to identify

dependencies between real genes. (D): These dependencies are modeled as a network. Filled

circles indicate genes (nodes), arrows indicate dependencies (edges).

Figure 2. PGCEs are enriched for biologically meaningful interactions. (A): The observed

number of PGCE edges connecting genes in the same pathway (dotted line), compared to the

expected distribution obtained from 1000 rewired networks with identical degree distributions.

(B): The observed number of PGCE edges that also appear in a bacteria-wide metabolic network,

compared to the expected distribution.

Figure 3. The phylogenetic history of *rbsL, urtA* and *rbsS*. The presence of each gene in each

branch in the phylogenetic tree is illustrated with a colored circle, with the circle's diameter

scaled to denote the probability of presence. (A): *rbsL* and *rbsS* evolutionary histories; (B): *urtA*

and *rbsS* evolutionary histories. The long branch leading to Archaea (bottom-most clade) was

reduced in size for graphical purposes.

Figure 4. Topological sorting of the PGCE dependency network reveals assembly patterns

that govern the evolutionary process. (A): Binned dependencies among the six ranks of genes

in the topological sort (left to right). Node size represents the number of genes in each rank

(using natural logarithm-scale). Edge width represents the number of PGCEs between genes in

different rank (natural logarithm-scale), all edges are directed to the right. (B) The gain of genes

from each rank in each branch of the phylogenetic tree is illustrated (circles). The different colors

represent different ranks. Circle sizes correspond to the proportion of gains on a branch

attributed to genes of that rank (e.g. a large red circle indicates that most gains on a branch

correspond to rank 1). The branch to Archaea (lower clade) has been reduced in size for

graphical purposes. See also Figure S7.

Figure 5. PGCE dependencies lead to taxonomically robust predictability of gene

acquisition. (A): Workflow for predicting gene acquisition between clades of the tree. A training

set is used to build a PGCE dependency model, which is then used to predict on which specific

branches genes are likely to be gained (green circles), based on dependencies inferred from the

training set (red and blue circles). (B): performance of PGCEs in predicting gene acquisitions in

two test sets (indicated clades of the prokaryotic tree). Areas under each curve: Firmicutes, 0.73;

Alpha/Beta-proteobacteria, 0.68. The diagonal dotted line represents the performance of a purely

random prediction. See also Figure S8.

23
24

1 FIGURES

Figure 1

Figure 2

Figure 3

A

Direction of edges

Ordered topological ranks

B

Topological rank
- 5
- 4
- 3
- 2
- 1

1
2

Figure 5

A

Test set

predict
gains

Training set

build
model

B

True positive rate

1.0
0.8
0.6
0.4
0.2
0.0

0.0 0.2 0.4 0.6 0.8 1.0
False positive rate

— test = Firmicutes
--- test = α/β-proteobacteria

1
2

1   **TABLES**

2   **Table 1.** Functional groups are enriched in different ranks of the topological sort.

Annotation label	P-value[1]	Enrichment Ratio[2]
Rank 1 Enrichments		
Cell motility	1.94E-07	1.40
Bacterial motility proteins	1.85E-11	1.41
Type II secretion system	2.61E-05	1.33
Two-component system	3.65E-04	1.25
Flagellar system	1.01E-09	1.43
Pilus system	2.11E-04	1.38
Metabolism[3]	3.37E-05	0.91
Xenobiotics biodegradation and metabolism[3]	1.07E-06	0.69
Carbohydrate metabolism[3]	0.00012	0.84
Type IV secretion system[3]	1.26E-09	0.20
Rank 2 Enrichments		
Metabolism	1.47E-04	1.23
Carbohydrate metabolism	3.08E-06	1.58
Rank 4 Enrichments		
Pathogenicity	1.88E-06	21.6
Conjugal transfer pilus assembly protein	1.08E-04	15.0
Type III protein secretion pathway protein	1.88E-06	21.6
ABC-2 type and other transporters	2.31E-04	12.5
Type IV secretion system	1.30E-03	8.04

3   1: from a hypergeometric test. All annotations displayed are significant at a 1% false discovery rate.

4   2: The ratio of the observed proportion of genes with this label in the indicated rank to the expected proportion

5   based on all genes in the network.

6   3: These annotations are depleted (i.e. enrichment ratio significantly less than one) in the first rank.

REFERENCES

Andam CP, Gogarten JP. 2011. Biased gene transfer in microbial evolution. *Nat Rev Microbiol* **9**: 543–55.

Andersson I, Backlund A. 2008. Structure and function of Rubisco. *Plant Physiol Biochem* **46**: 275–91.

Baltrus DA. 2013. Exploring the costs of horizontal gene transfer. *Trends Ecol Evol* **28**: 489–95.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc B* **57**: 289–300.

Chen HD, Jewett MW, Groisman E a. 2011. Ancestral genes can control the ability of horizontally acquired loci to confer new traits. *PLoS Genet* **7**: e1002184.

Christin P-A, Arakaki M, Osborne CP, Edwards EJ. 2015. Genetic enablers underlying the clustered evolutionary origins of C4 photosynthesis in angiosperms. *Mol Biol Evol* **32**: 846–58.

Cohen O, Ashkenazy H, Burstein D, Pupko T. 2012. Uncovering the co-evolutionary network among prokaryotic genes. *Bioinformatics* **28**: i389–i394.

Cohen O, Gophna U, Pupko T. 2011. The complexity hypothesis revisited: connectivity rather than function constitutes a barrier to horizontal gene transfer. *Mol Biol Evol* **28**: 1481–9.

Cohen O, Pupko T. 2010. Inference and characterization of horizontally transferred gene families using stochastic mapping. *Mol Biol Evol* **27**: 703–13.

Conte GL, Arnegard ME, Peichel CL, Schluter D. 2012. The probability of genetic parallelism and convergence in natural populations. *Proc Biol Sci* **279**: 5039–47.

Csardi G, Nepusz T. 2006. The igraph Software Package for Complex Network Research. *InterJournal* **Complex Sy**.

Dehal PS, Joachimiak MP, Price MN, Bates JT, Baumohl JK, Chivian D, Friedland GD, Huang KH, Keller K, Novichkov PS, et al. 2010. MicrobesOnline: an integrated portal for comparative and functional genomics. *Nucleic Acids Res* **38**: D396–400.

Delwiche CF, Palmer JD. 1996. Rampant horizontal transfer and duplication of rubisco genes in eubacteria and plastids. *Mol Biol Evol* **13**: 873–882.

Eppley RW, Peterson BJ. 1979. Particulate organic matter flux and planktonic new production in the deep ocean. *Nature* **282**: 677–680.

Gogarten JP, Doolittle WF, Lawrence JG. 2002. Prokaryotic evolution in light of gene transfer. *Mol Biol Evol* **19**: 2226–38.

Gong LI, Suchard MA, Bloom JD. 2013. Stability-mediated epistasis constrains the evolution of an influenza protein. *Elife* **2**: e00631.

Gophna U, Ofran Y. 2011. Lateral acquisition of genes is affected by the friendliness of their products. *Proc Natl Acad Sci U S A* **108**: 343–8.

Gould SJ, Lewontin RC. 1979. The Spandrels of San Marco and the Panglossian Paradigm: A Critique of the Adaptationist Programme. *Proc R Soc B Biol Sci* **205**: 581–598.

Hagberg A, Schult D, Swart P. 2013. NetworkX. High productivity software for complex networks. https://networkx.lanl.gov/.

Harms MJ, Thornton JW. 2014. Historical contingency and its biophysical basis in glucocorticoid receptor evolution. *Nature* **512**: 203–7.

Hsu HT. 1975. An Algorithm for Finding a Minimal Equivalent Graph of a Digraph. *J ACM* **22**: 11–16.

Huelsenbeck JP, Nielsen R, Bollback JP. 2003. Stochastic Mapping of Morphological Characters. *Syst Biol* **52**: 131–158.

Iwasaki W, Takagi T. 2009. Rapid pathway evolution facilitated by horizontal gene transfers across prokaryotic lineages. ed. I. Matic. *PLoS Genet* **5**: e1000402.

Jain R, Rivera MC, Lake J a. 1999. Horizontal gene transfer among genomes: the complexity hypothesis. *Proc Natl Acad Sci U S A* **96**: 3801–6.

Jain R, Rivera MC, Moore JE, Lake JA. 2003. Horizontal gene transfer accelerates genome innovation and evolution. *Mol Biol Evol* **20**: 1598–602.

Johnson CM, Grossman AD. 2014. Identification of host genes that affect acquisition of an integrative and conjugative element in Bacillus subtilis. *Mol Microbiol*.

Kahn AB. 1962. Topological Sorting of Large Networks. *Commun ACM* **5**: 558–562.

Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. 2012. KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic Acids Res* **40**: D109–14.

Lerat E, Daubin V, Ochman H, Moran NA. 2005. Evolutionary origins of genomic repertoires in bacteria. ed. D. Hillis. *PLoS Biol* **3**: e130.

Letunic I, Bork P. 2011. Interactive Tree Of Life v2: online annotation and display of phylogenetic trees made easy. *Nucleic Acids Res* **39**: W475–8.

Levy R, Borenstein E. 2013. Metabolic modeling of species interaction in the human microbiome elucidates community-level assembly rules. *Proc Natl Acad Sci U S A* **110**: 12804–9.

Maddison WP. 1990. A Method for Testing the Correlated Evolution of Two Binary Characters: Are Gains or Losses Concentrated on Certain Branches of a Phylogenetic Tree? *Evolution (N Y)* **44**: 539–557.

Nowell RW, Green S, Laue BE, Sharp PM. 2014. The extent of genome flux and its role in the differentiation of bacterial lineages. *Genome Biol Evol* **6**: 1514–29.

Ord TJ, Summers TC. 2015. Repeated evolution and the impact of evolutionary history on adaptation. *BMC Evol Biol* **15**: 137.

Orr HA. 2005. The Probability of Parallel Evolution. *Evolution (N Y)* **59**: 216–220.

Pal C, Papp B, Lercher MJ. 2005. Horizontal gene transfer depends on gene content of the host. *Bioinformatics* **21**: ii222–ii223.

Paradis E, Claude J, Strimmer K. 2004. APE: Analyses of Phylogenetics and Evolution in R language. *Bioinformatics* **20**: 289–290.

Van Passel MWJ, Marri PR, Ochman H. 2008. The emergence and fate of horizontally acquired genes in Escherichia coli. *PLoS Comput Biol* **4**: e1000059.

Popa O, Dagan T. 2011. Trends and barriers to lateral gene transfer in prokaryotes. *Curr Opin Microbiol* **14**: 615–23.

Popa O, Hazkani-Covo E, Landan G, Martin W, Dagan T. 2011. Directed networks reveal genomic barriers and DNA repair bypasses to lateral gene transfer among prokaryotes. *Genome Res* **21**: 599–609.

Press MO, Li H, Creanza N, Kramer G, Queitsch C, Sourjik V, Borenstein E. 2013. Genome-scale co-evolutionary inference identifies functions and clients of bacterial Hsp90. *PLoS Genet* **9**: e1003631.

Puigbò P, Lobkovsky AE, Kristensen DM, Wolf YI, Koonin E V. 2014a. Genomes in turmoil: Quantification of genome dynamics in prokaryote supergenomes. *BMC Biol* **12**: 66.

Puigbò P, Lobkovsky AE, Kristensen DM, Wolf YI, Koonin E V. 2014b. Genomes in turmoil: Quantification of genome dynamics in prokaryote supergenomes. *BMC Biol* **12**: 66.

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T. 2003. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* **13**: 2498–504.

Smillie CS, Smith MB, Friedman J, Cordero OX, David L a, Alm EJ. 2011. Ecology drives a global network of gene exchange connecting the human microbiome. *Nature* **480**: 241–4.

Sohm JA, Webb EA, Capone DG. 2011. Emerging patterns of marine nitrogen fixation. *Nat Rev Microbiol* **9**: 499–508.

Sorek R, Zhu Y, Creevey CJ, Francino MP, Bork P, Rubin EM. 2007. Genome-wide experimental determination of barriers to horizontal gene transfer. *Science* **318**: 1449–52.

Soucy SM, Huang J, Gogarten JP. 2015. Horizontal gene transfer: building the web of life. *Nat Rev Genet* **16**: 472–482.

Venables WN, Ripley BD. 2002. *Modern Applied Statistics with S.* Fourth Edi. Springer, Springer.

De Visser JAGM, Krug J. 2014. Empirical fitness landscapes and the predictability of evolution. *Nat Rev Genet* **15**: 480–490.

Weinreich DM, Delaney NF, Depristo MA, Hartl DL. 2006. Darwinian evolution can follow only very few mutational paths to fitter proteins. *Science* **312**: 111–4.

2012. R: A language and environment for statistical computing. R Development Core Team.