1   **Title:** Disruption of endosperm development is a major cause of hybrid seed inviability between
2   *Mimulus guttatus* and *M. nudatus*
3

4   Elen Oneal[1], John H. Willis[1], Robert G. Franks[2]
5
[1] Department of Biology, Duke University, 3319 French Family Science Center, 125 Science
Drive, Durham, NC, 27705, USA
[2] Department of Genetics, North Carolina State University, 2548 Thomas Hall, Raleigh, NC,
27695, USA
Email: eo22@duke.edu (EO), jwillis@duke.edu (JHW), rgfranks@ncsu.edu (RGF)
Corresponding author: Elen Oneal, email: eo22@duke.edu; phone (919) 684-3368

Summary
14
•   Divergence of developmental mechanisms within populations may lead to hybrid
developmental failure and inviability, and may be a factor driving speciation in flowering
plants.
•   We investigate the patterns of early and late embryo and endosperm development of seed
from *Mimulus guttatus* and the closely related, serpentine endemic *M. nudatus*, and compare
those patterns to that of hybrid seed derived from reciprocal, interspecific crosses. We
address whether disruption in hybrid seed development is the primary source of reproductive
isolation between these sympatric taxa.
•   *M. guttatus* and *M. nudatus* differ in the pattern and timing of endosperm and embryo
development. Some hybrid seed exhibit early disruption of endosperm development and are
completely inviable, while other hybrid seed exhibit comparatively normal early
development, but show impaired endosperm proliferation. These developmental patterns
underlie the phenotypes of mature seed, which are either very small and flat, (indicating little
to no endosperm), or shriveled, (indicating reduced endosperm volume). Both phenotypes
have low germination success. Hybrid seed inviability forms a potent reproductive barrier
between *M. guttatus* and *M. nudatus*.
•   We provide a partial developmental mechanism for the hybrid barrier between *M. guttatus*
and *M. nudatus*. In addition to illustrating the early stages of endosperm and embryo
development, we shed light on the extent of developmental variation between closely related
species within this important ecological model system.

35
36 **Keywords:** seed development, hybrid inviability, endosperm, postzygotic isolation, pollen-pistil
37 interactions, *Mimulus guttatus* (monkeyflower)
38
39
40

Introduction

The process of gradual evolution imposes a fundamental constraint on organismal development – each successful evolutionary shift, large or small, must allow for viable offspring (Smith *et al.,* 1985; Bonner, 1988; Beldade *et al.* 2002). This constraint is perhaps best visualized by the disruption of development frequently observed when two divergent populations hybridize, when both lineages themselves continue to produce viable offspring. In nature, these incompatibilities can keep species distinct by preventing gene flow. In the laboratory, we can make use of incompatibilities witnessed in hybrid offspring to investigate how development has evolved in isolation and how evolutionary constraint may shape developmental trajectories. Here we describe differences in seed development between a recently diverged species pair – *Mimulus guttatus* and *M. nudatus*. We further show that postzygotic failures of development are largely responsible for incompatibility in experimental crosses between this sympatric species pair. We propose that the *M. guttatus* sp. complex may serve as a new model to understand the evolution of development and developmental abnormalities in hybrid plants.

Development in multicellular organisms requires coordination across numerous cell lineages or types. The process of double fertilization in angiosperms is an extreme example as growth must be coordinated across two developing entities: the diploid embryo and the triploid, sexually-derived nutritive tissue called endosperm. Together these distinct entities comprise the angiosperm seed, a highly successful mode of reproduction employed by most vascular plants (Linkies *et al.,* 2010). While the developmental origins of embryo and endosperm have been known for over a century (Nawaschin, 1898; Guignard, 1899, Friedman, 2001), advances in genomic sequencing and gene expression analysis have only lately revealed the basic genetic details of embryogenesis and the development of endosperm (Girke *et al.,* 2000, Casson *et al.* 2005; Hsieh *et al.*, 2011). The developing endosperm and its interactions with the embryo is often responsible for hybrid seed failure (Brink & Cooper, 1947; Haig & Westoby 1991) emphasizing the critical and sensitive role it plays in promoting successful reproduction.

Despite the essential importance of endosperm, research on endosperm development has been largely restricted to the model system, *A. thaliana* and its close relatives (Scott et al., 1998; Josefsson et al. 2006; Burkart-Waco et al., 2013), even though several developmental and evolutionary peculiarities of the biology of *A. thaliana* may limit the applicability of research findings across the broader diversity of plants. For example, *A. thaliana* undergoes nuclear endosperm development, in which initial rounds of karyokinesis are not accompanied by cytokinesis. While this mode of development is shared by many groups of flowering plants, two other major modes of endosperm development, helobial and cellular, are also distributed widely among angiosperms (Bharathan, 2000). Indeed, *ab initio* cellular development, wherein karyokinesis is always followed by cytokinesis, is thought to be the ancestral state of endosperm development (Floyd & Friedman, 2000) and is characteristic of a few other model plant systems including *Solanum* (Lester & Kang, 1998) and *Mimulus* (Guilford & Fisk, 1951; Arekal, 1965). More broadly, the extent to which basic features of embryo and endosperm development may

83 vary among closely related taxa remains an open question and one which can only be addressed
84 by examining and comparing development between closely related species in other taxa.
85
Another factor limiting the applicability of research on endosperm in *A. thaliana* is that it is
predominantly self-fertilizing. Self-fertilization limits a major evolutionary pressure on seed
development by relaxing conflicts between maternal and paternal genomes over resource
provisioning to developing seeds. Relying on A. *thaliana* as a model of endosperm development
and failure potentially limits our ability to fully understand the evolutionary mechanisms shaping
plant development. It also limits our ability to investigate the role mating system may play in the
evolution of endosperm, a research topic that continues to garner increasing interest (Brandvain
& Haig, 2005; Friedman *et al.*, 2008; Köhler *et al.,* 2012; Haig, 2013). Moreover, while seed
inviability may be a potent hybrid barrier and potential driver of plant speciation (Tiffin *et al.*,
2001), nearly all research on hybrid seed lethality in *A. thaliana* is focused on lethality resulting
from interploidy crosses (Scott *et al.,* 1998; Köhler *et al.,* 2003), potentially limiting its
application to divergence among diploid taxa.
98
To extend our understanding of seed development and seed failure beyond *A. thaliana* we turn to
the *Mimulus guttatus* species complex. The genus *Mimulus* (Phrymaeceae) has emerged as a
model system in which to investigate the genetic basis of ecological adaptation and the role of
mating system evolution in promoting species divergence (Lowry & Willis, 2010; Martin &
Willis, 2010; Wright *et al.,* 2013). Knowledge of the pattern of seed development in this
increasingly important genus is limited to two papers published over 50 years ago on the species
Mimulus ringens (Arekal, 1965) and the cultivar *M. tigrinus* (Guilford & Fisk, 1951), likely a
hybrid between *M. luteus* and the Chilean species *M. cupreus* (Cooley & Willis, 2009). Seed
inviability is a common outcome of crosses between members of the *M. guttatus* species
complex, a highly diverse group of populations, ecotypes and species distributed across western
North America (Vickery 1966, 1978). *M. guttatus* is the most geographically widespread and
genetically diverse member of the complex (Wu *et al.*, 2007; Oneal et al. 2014), and exhibits
varying interfertility with other members of the complex (Vickery, 1978; Wu *et al.,* 2007),
however, hybridization between the closely related members of this complex is frequently
accompanied by varying levels of hybrid seed failure (Vickery, 1978).
114
While the edaphic endemic, *M. nudatus* is likely recently derived from a *M. guttatus*-like
ancestor (Oneal *et al.*, 2014), this species pair exhibits the highest level of sequence divergence
of any within the *M. guttatus* sp. complex (~3% genomic sequence divergence; L. Flagel,
personal communication). Populations of serpentine-adapted *M. guttatus* overlap with those of *M.*
nudatus at multiple serpentine soil sites in the California Coastal Ranges. Despite their close
physical proximity and recent divergence, *M. guttatus* and *M. nudatus* rarely form hybrids. The
absence of naturally occurring hybrids is all the more striking given that *M. guttatus* and *M.*
nudatus also overlap substantially in flowering time and share multiple pollinators (Gardner &
Macnair, 2000; J. Selby, unpublished data). Gardner and Macnair (2000) found that controlled
field and greenhouse crosses recovered very few normal seed but instead produced seed that
were shriveled and comparatively flattened, and that failed to germinate (Gardner, 2000).
Together, these findings raise the possibility that the recent divergence between *M. guttatus* and

M. nudatus has been accompanied by a shift in the pattern of embryo and endosperm
development, that in turn contributes to their reproductive isolation.
129
Here we investigate early embryo and endosperm development within *M. guttatus*, a species
which has emerged as an important model system in ecological and evolutionary genetics, and
contrast it with development of *M. nudatus*, a serpentine soil endemic. Furthermore, we
investigate whether seed inviability is the primary reproductive barrier between *M. guttatus* and
M. nudatus (Gardner & Macnair, 2000). We first address whether interspecific pollen can
successfully germinate and penetrate the ovary when either species serves as pollen donor. We
then compare the development of hybrid seed with that of the normal pattern of development in
both species, and attempt to determine at what point in development hybrid seed failure arises.
Finally, we connect the development of hybrid seed to the phenotypes of mature seed collected
from hybrid fruits and confirm that hybrid seed are largely inviable.
140
We find that *M. guttatus* and *M. nudatus* exhibit divergent trajectories of early embryo and
endosperm development, and suggest that early disruption of endosperm development and a later
failure of endosperm proliferation are the major causes of hybrid seed failure and comprise major
isolating mechanism between these species. Our work is the first to examine the pattern of seed
development in *M. guttatus* and the first to examine the extent to which early seed development
varies between *M. guttatus* and a closely related species, *M. nudatus*. Finally, since seed lethality
is a common outcome of hybridization between multiple members of the *M. guttatus* sp. complex
(Vickery, 1978), our results suggest that *M. guttatus*, which is already emerging as a model
system in ecological genetics, could also provide valuable insight into the genetic basis of
fundamental developmental processes and their importance for speciation in this group.
151
152
153   **Materials and Methods**
154
155   *Growth and pollination of Mimulus spp.*
156
M. guttatus is the most widespread member of the *M. guttatus* sp. complex and is adapted to a
variety of soil conditions (Lowry *et al.* 2009, Wright *et al.* 2014), including serpentine soil.
Serpentine-adapted *M. guttatus* and *M. nudatus* individuals were collected in 2008 from
sympatric populations located at two serpentine soil sites in the Donald and Sylvia McLaughlin
Natural Reserve in Lake County, California, and brought back to the Duke Research
Greenhouses where they were self-fertilized for at least 2 generations to produce inbred lines.
We use one inbred line per population in this study (see Table 1 for a list of accessions), and two
populations each of serpentine-adapted *M. guttatus* and *M. nudatus*. The lines CSS4 (*M.
guttatus*), CSH10 (*M. nudatus*) and DHRo22 (*M. nudatus*) were inbred for 2 generations. Most
data generated using the *M. guttatus* accession DHR14 was acquired from a line that was inbred
for 3 generations; however, we were unable to complete the study with these individuals, as they
died prematurely in the greenhouse; we completed the study with a 6-generation inbred line of
DHR14. All plants used in this study were grown from seeds that were first cold-stratified for 10
days at 4ºC, then placed in a greenhouse with 30% relative humidity and a light/temperature

regime of 18-hour days at 21 °C and 6-h nights at 16 °C. Following germination, individuals
were placed in 2.5-inch square pots where they were maintained for the duration of the study.
173
M. guttatus and *M. nudatus* are both self-compatible and self-fertilize regularly in the field,
although *M. nudatus* is primarily outcrossing (Ritland & Ritland, 1989). Both species
produce hermaphroditic, chasmogamous flowers with four anthers, and invest similarly in male
(e.g., stamens) vs. female (e.g., pistil) structures, however, *M. nudatus* flowers are smaller than
those of *M. guttatus* and produce proportionately fewer ovules and pollen grains (~20% as many
ovules and pollen grains) (Ritland & Ritland, 1989). To account for this imbalance in pollen
production, we used 4 new flowers (i.e., 16 anthers) whenever *M. nudatus* served as pollen donor
to *M. guttatus*. All crosses and self-pollinations were performed in the morning and using the
same protocol. Pollen recipients were emasculated 1-3 days prior. The day of pollination, mature
pollen was obtained by tapping the stamens of a fresh flower onto a glass slide, and then
collected with a pair of sterile forceps and placed directly on the open, receptive stigma of the
pollen recipient.
186
187   *Pollen tube growth assay*
188
Pollinated styles and ovaries were fixed in Farmer's solution (3:1 95% EtOH:acetic acid) for at
least 12 hours, then softened in 8N NaOH for 24 hours before being left to stain overnight in a
decolorized aniline blue solution (0.1% in 0.1 M K_3PO_4) (Kearns & Inouye, 1993) that
differentially stains pollen tubes callose plugs. Stained styles and ovaries were mounted on a
slide and examined with a Zeiss Axio Observer equipped with a fluorescent lamp. We performed
10 reciprocal crosses for each sympatric population pair (CSS4 x CSH10; CSH10 x CSS4;
DHR14 x DHRo22; DHRo22 x DHR14) and then collected the styles and ovaries after 24 hours,
a period sufficient to allow pollen from self-pollinations to penetrate the ovary in each
population (Oneal, personal observation). For each pollination event we noted whether the pollen
had successfully germinated and whether pollen tubes were observed within the ovary.
199
200   **Seed set and viability**
201
We performed 8 reciprocal crosses for each sympatric population pair of *M. guttatus*/*M. nudatus*
and 8 self-pollinations of each accession, collected the mature fruits and counted the resulting
seeds under a dissection scope. Throughout, we use the term "self-fertilization" to describe
fertilizations performed with pollen from the same accession (i.e., inbred line), but not
necessarily the same individual plant. Normal *M. guttatus* and *M. nudatus* seeds are round, fully
filled and unbroken with a light brown, reticulate coat (Searcy & Macnair, 1990). We counted
the number of seed found in self-fertilized and hybrid fruits and categorized them by outward
morphology. We also took pictures of mature seed using a Zeiss Lumar.V12 stereoscope
outfitted with a AxioCam MRM firewire monocrome camera and measured the length of up to
25 seed morphs for each self-fertilized accession and reciprocal cross. We sowed round,
shriveled, and flat self-fertilized and hybrid seeds (see below) to compare germination rates. All
seeds were cold-stratified, placed in the Duke Greenhouses as above, and examined over the
course of 14-days for signs of germination.

215
Seed development
Fruits resulting from interspecific crosses consistently contained seeds that fell into one of three different morphological categories (see below). We used microscopy to connect early embryo and endosperm development with the seed morphologies found in mature fruits and to compare the growth, and embryo and endosperm development, of self-fertilized seeds to those of reciprocal, sympatric hybrid seed. Self-fertilized and hybrid fruits were collected from 1-5 days after pollination (DAP), and then at 9 DAP. Emasculated, but unpollinated ovaries were collected at 1-2 DAP.

We first examined whole-mounted fruits to get an initial sense of the pattern of embryo and endosperm development in self-fertilized and hybrid seeds from 1 to 5 DAP. Plant material was fixed in a solution of 9:1 EtOH:acetic acid for at least 2 hours and up to 48 hours, then washed twice in 90% EtOH for a minimum of 30 minutes per wash. Tissue was subsequently cleared in Hoyer's solution (70% chloral hydrate, 4% glycerol and 5% gum arabic) for at least 12 hours. A final dissection of the fruit in Hoyer's solution allowed unfertilized ovules and immature seed to be separated from the ovary or fruit and then mounted on a glass slide. We collected 3 replicate fruits per DAP for each hybrid cross or self-fertilization, as well as 3 unpollinated ovaries from each accession. Mounted specimens were observed with a Zeiss Axioskop2 or Zeiss Axio Imager using differential interference contrast (DIC) microscopy. We took pictures of up to 10 ovules per unfertilized ovary, 10 immature seed per self-fertilized fruit, and up to 10 of each seed morph (see below) per hybrid fruit, and used these images to measure the size of seed morphs.

We used laser confocal microscopy (LCM) to better visualize the pattern of early seed failure observed in whole mounted fruits (see below). Tissue was stained with propidium iodide with according to Running (2007). Plant material was fixed under a vacuum with a solution containing 3.7% [v/v] formaldehyde, 5% [v/v] propionic acid, 70% [v/v] ethanol, and then subjected to a graded ethanol series to remove residual chorophyll. Tissue was then subjected to a decreasing ethanol series, stained with propidium iodide dissolved in 0.1M L-arginine (pH 12.4) for 2-4 days (stain time depended upon the size of the plant material), rinsed for 2-4 days in a 0.1M L-arginine buffer (pH 8.0), subjected to another graded ethanol series and then a final graded xylene series. Unfertilized ovules and immature seeds were dissected out and mounted in Cytoseal XYL. Images were acquired with a Zeiss 710 inverted scanning confocal microscope equipped with an argon laser. Some images (e.g., seed at 4-5 DAP) required the collection of extended z-stacks, which were assembled into composite 3-D images using the Zeiss Zen software.

We examined cross sections of self-fertilized and hybrid seeds collected at 9 DAP. Fruits were fixed under a vacuum with a solution containing 3.7% [v/v] formaldehyde, 50% [v/v] EtOH, and 5% [v/v] glacial acetic acid, subjected to a graded ethanol series, then stained overnight in a 0.1% Eosin solution. Stained tissue was subjected to a graded xylene series and then infused with and mounted in paraplast parafin. Fruits were sliced with a microtome into 0.8 micron sections, stained with toluidine blue, and sealed with Cytoseal XYL for imaging. Slides were examined

258 and photographed with a Zeiss Axio Imager outfitted with a QImaging MicroPublisher 5.0 MP
259 color camera.
260
To determine whether hybrid seeds that appeared to fail early in the course of development (see
below) represented fertilized seeds (as opposed to unfertilized, aborted ovules), we used three
lines of evidence. First, we used a vanillin stain to test for seed coat development in immature
hybrid seed. In *A. thaliana*, vanillin in acidic solution (i.e., a 1% [w/v] vanillin solution in 6 N
HCL) turns red or brown upon binding to proanthocyanidins in the seed coat; a positive stain is
indicative of seed coat development and suggests that fertilization has occurred (Deshpande *et al.,*
1986; Roszak & Köhler, 2011). We tested for seed coat development in 5 DAP reciprocal hybrid
seed, and unpollinated ovaries (negative control) collected 5 days after emasculation. Second,
we measured the length, from micropylar to chalazal end, of hybrid seed and compared their
growth trajectory to that of self-fertilized seed. Third, using our LCM images, we compared the
width of the central cell of unfertilized ovules from a subset of our accessions (DHR14 and
DHRo22) to the width of the putative primary endosperm cell of DHR14 x DHRo22 reciprocal
hybrid seed that exhibited signs of arrest at 2 DAP. An increase in size of the putative primary
endosperm cell over the central cell is suggestive of successful fertilization (Williams, 2009).
Throughout, measurements of size were taken using ImageJ software (Rasband, 1997). All
crosses are given with the female parent listed first (i.e., female x male). Unless accessions
differed significantly (noted in the text), data are pooled across accessions for both *M. guttatus*
and *M. nudatus.*
279
280 **Results**
281
282 ***Pollen germination and tube growth***
283
The inability of pollen from one species to successfully germinate, tunnel down the style, and
penetrate the ovary of another species is a common prezygotic barrier to hybridization in
flowering plants (Galen & Newport, 1988; Boavida *et al.,* 2001; Campbell *et al.,* 2003; Ramsey
et al., 2003). In the *M. guttatus* sp. complex, pollen failure contributes to transmission distortion
in crosses between *M. guttatus* and the closely related *M. nasutus* (Fishman *et al.,* 2008).
Gardner and Macnair (2000) reported that mature hybrid fruits contained few viable seeds but
were filled with "dust"; however, they did not specify whether these particles were aborted seeds
or unfertilized ovules. To clarify this, we investigated whether *M. guttatus* pollen could
germinate and successfully penetrate the ovary of *M. nudatus* and vice versa, a precondition for
fertilization. We found that interspecific pollen successfully germinated in all crosses and that
some pollen grains were nearly always successful in tunneling down to the ovary within 24-
hours (Table 2; Fig. 1). Identity of the female parent did not affect ability of interspecific pollen
to penetrate the ovary (Wilcox rank-sum test, p > 0.1, data pooled across accessions).
Furthermore, fruits resulting from *M. guttatus* x *M. nudatus* crosses typically swell and increase
in size in a manner similar to fruits resulting from self-fertilization in either species (Fig. 2).
299
300 **Seed set and germination success**
301

The majority of seeds from mature self-fertilized fruits of *M. guttatus* and *M. nudatus* are round
and unbroken with a light brown, reticulate coat (Searcy & Macnair, 1990) (Fig. 2, Fig. 3;
termed "round" in this work). Most self-fertilized fruits (30 of 32) also contained a minority of
seeds which were shriveled and irregularly shaped (Fig 3. termed "shriveled"), and were
significantly smaller than the usual, round seed for both species (seed length: *M. guttatus* $F_{1,98}$ =
58.647, $p < 0.001$; *M. nudatus* $F_{1,95}$ = 108.14, $p < 0.001$). For both species, the size difference
between round and shriveled seeds varied with accession (two-way ANOVA: *M. guttatus* $F_{1,98}$ =
4.24, $p = 0.002$; *M. nudatus* $F_{1,95}$ = 10.0, $p = 0.042$). In addition, several self-fertilized fruits (1 *M.*
nudatus and 5 *M. guttatus*) contained a few seeds (16 total) that were considerably smaller than
round, wild type seed and of a brown, flat appearance (Fig 3. termed "flat"). The proportion of
round seed was lower in *M. guttatus* than *M. nudatus* (*M. guttatus:* round mean = 69.3% ± 19.2
SD; *M. nudatus:* round mean = 80.7% ± 21.8 SD) (Fig. 3). Total seed set per fruit of self-
fertilized accessions was 77.3 (± 40.9 SD) for *M. guttatus* and 84.0 (± 19.7 SD) for *M. nudatus*.
(Fig. 4).
316
Total seed set did not differ between self-fertilized fruits and fruits resulting from interspecific
crosses (two-way ANOVA, $p > 0.1$ for *M. guttatus* or *M. nudatus* female) (Fig 4; Supplementary
Fig. 1). Of 16 interspecific *M. guttatus* x *M. nudatus* crosses, only one produced one round seed.
Instead, most hybrid seeds (mean = 76.8% ± 24.8 SD) were dark brown and shriveled (termed
"shriveled"), resembling the shriveled seed present at lower frequency in self-fertilized fruits
(Fig. 2e, Fig. 3, Supplementary Fig. 2). The remaining hybrid seeds (mean = 23.2% ± 24.8 SD)
were very small, dark brown and flattened in appearance (Fig. 2e, Fig. 3, Supplementary Fig. 2)
(termed "flat"). These latter seeds were clearly distinguishable from unfertilized ovules, which
are smaller and light pink in color (due to a lack of seed coat) (Searcy & Macnair, 1990). We
found one round seed with endosperm that had exploded through the seed coat in one of 16
mature fruits where *M. nudatus* was the female (CSH10 x CSS4). Otherwise, *M. nudatus* x *M.*
guttatus crosses produced hybrid seed that were either shriveled or small and flat (mean
shriveled = 40.0% ± 19.2 SD; mean flat = 59.6% ± 19.1 SD) (Fig. 3).
330
331   *Germination Success*
332
Averaging across accessions, 92% (± 5.6 SD; *N*=50) of seed from self-fertilized *M. guttatus*
accessions germinated, while 62.5% (± 26.5 SD; $N = 32$) of self-fertilized *M. nudatus* seed
germinated (see Table 3 for germination success by accession and cross); the difference between
the species was not significant (Fisher exact test, $p = 0.484$). Shriveled seeds from self-fertilized
fruits germinated at a lower rate for both species (*M. guttatus* = 13.6%; *M. nudatus* = 3.5%, $p <$
0.001 for both comparisons, Fisher's exact test). Hybrid seed germinated only when *M. guttatus*
was the female, and then at significantly lower rates (6.01% overall; Fisher's exact test, $p <$
0.0001). None of the flat seeds germinated, including flat seeds from self-fertilized fruits.
341
342   **Seed development**
343
344   *M. guttatus* seed development
345

The *Mimulus* mature female gametophyte is the Polygonum type, which posesses two haploid
synergid cells, a haploid egg cell and two antipodal cells, one of which is binucleate (as
described for *M. ringens* (Arekal, 1965)) (Fig. 5a. Within 24-hours after pollination, many *M.*
guttatus seeds can be seen undergoing the first transverse division of the primary endosperm cell.
At 2 DAP, endosperm development consists of 2 to 8 evenly spaced endosperm nuclei (Fig. 5b),
and the establishment of the chalazal and micropylar domains. The micropylar domain is
anchored by two cells whose nuclei accumulate multiple nucleoli—signs of endoreduplication, a
phenomenon commonly observed in plant tissue (Galbraith et al., 1991). The chalazal
haustorium, also containing two very large nuclei, differentiates from the central endosperm,
occupying the chalazal domain. At times, the cells micropylar haustorium can be seen to
penetrate beyond the base of the micropylar domain towards the chalazal domain. By 3 DAP,
cellularized endosperm continues to proliferate, the embryo is at the 2/4-cell stage, and the
chalazal haustorium has already begun to degrade (Fig. 5c). Between 4-5 DAP, the embryo
progresses rapidly from the 8-celled stage with suspensor to the late globular stage (Fig. 5d,e).
The seed contains regularly dispersed endosperm (Fig. 5e), which becomes densely packed by 9
DAP (Fig. 6e); at this point the micropylar haustorium has largely degenerated, but remains
visible as a darkly stained element in the micropylar domain (Fig 6e).
363
M. nudatus seed development
The female gametophyte of *M. nudatus* is significantly smaller than that of *M. guttatus* (one-
tailed t-test, *p* < 0.001). Development of *M. nudatus* seeds parallels that of *M. guttatus* but with
some important differences. Most notably, endosperm and embryo development proceed more
slowly in *M. nudatus* than in *M. guttatus* (Fig. 5f-i). In addition, *M. nudatus* endosperm is less
compact and regularly spaced. Division of the primary endosperm nucleus is initiated by 2 DAP
and completed by 3DAP (Fig. 5f,g). The chalazal and micropylar haustoria emerge by 3 DAP;
both are more prominent and persist longer in *M. nudatus* than *M. guttatus* (Fig. 5g,h). At 5 DAP,
the *M. nudatus* embryo is a 16-cell embryo (Fig. 5i). At 9 DAP, embryo development ranges
from heart stage to torpedo stage (Fig. 6h).
374
375 *Hybrid seed development*
376
Earlier work suggested that the primary barrier to hybridization between *M. guttatus* and *M.*
nudatus was the formation of nonviable hybrid seed (Gardner & Macnair, 2000). To test this, we
compared the development of seeds from reciprocal, interspecific crosses to that of seeds
resulting from self-fertilizations for each accession of *M. guttatus* and *M. nudatus*, respectively.
We found that reciprocal *M. guttatus* x *M. nudatus* crosses produced broadly similar
developmental trajectories for hybrid seed: an early stage of arrested development (Fig. 7), a
pattern of delayed embryo development visible by 5 DAP (Fig. 6a-d), and retarded endosperm
proliferation evident at 9 DAP (Fig. 6e-i).
385
Seeds that fail early are distinguishable as early as 2 DAP. At this stage, *M. guttatus* and *M.*
nudatus self-fertilized seed have undergone at least one and often a few divisions of the primary
endosperm cell. In hybrid seed at 2 DAP, regardless of which species serves as maternal parent,
the putative primary endosperm cell widens and becomes significantly larger than the central cell

of the female gametophyte of the maternal parent (t-test, $p < 0.001$ for both crossing directions).
These seeds are also significantly longer than unfertilized ovules (t-test, $p < 0.0001$ for both
crossing directions) but do not increase substantially in size over time (Fig. 8a,b).
393
Confocal microscopy of these hybrid seed indicates that at 2 DAP, transverse division of the
primary endosperm cell has failed to occur; cells walls are not evident, and the cell is filled with
multiple vacuoles (Fig. 7b,d). Fruits at 4-5 DAP may show some signs that the primary
endosperm cell of these early arrested seed may have undergone one or a few divisions (Fig.
7c,e), including the presence of a few cell walls (Fig. 7e). Also at this stage, one or a few
endosperm nuclei appear to contain multiple nucleoli, potentially due to endoreduplication (Fig.
7c,e) . Intriguingly, for both cross directions the primary endosperm cell of 2 DAP hybrid seeds
appears to filled with nucleic acids (either DNA or RNA) bound to the propidium iodide stain.
This fluorescence often, but not always, diminishes by 4-5 DAP (Fig 7c,e). As in self-fertilized
seed, micropylar haustoria are evident by 2 DAP and at later stages exhibit multiple nucleoli. By
5 DAP, a few arrested seeds may even show evidence of embryo growth (Fig 7c), albeit at a very
early stage. Exposure to vanillin stain of hybrid fruit at 5 DAP reveals two distinct sizes of dark
seed. We suggest that the smaller seeds, which are darker than ovules from unpollinated ovaries
(Fig. 9), represent the early arrested hybrid seeds (i.e., flat seeds), while the larger dark seeds
represent hybrid seed that develop embryos and proliferating endosperm, but have a later,
shriveled appearance.
410
411 *Later hybrid seed development*
412
In contrast to these early arresting seeds, many hybrid seed undergo development that at the
earliest stages (2-4 DAP) closely resembles that of the maternal parent (as described above),
including in the initial pattern of endosperm division and cellularization and the timing of the
emergence and eventual degeneration of chalazal and micropylar haustoria. Notably, these seeds
are typically slightly smaller than seed from the self-fertilized maternal parent (Fig. 8a,b; see
Supplementary Fig. 3 for growth by accession). For both crossing directions, however, by 5 DAP
hybrid embryo development is slightly delayed: *M. guttatus* x *M. nudatus* embryos range from 8-
to 16-cell embryos, as compared to *M. guttatus* embryos, which are at the globular stage (Fig.
7a,b). Similarly, *M. nudatus* x *M. guttatus* hybrid embryos are at the 8-cell stage while the *M.
nudatus* embryo is more typically at the 16-cell to early globular stage (Fig. 7c,d). This delay
persists at later stages and moreover, is accompanied by defects in endosperm proliferation at 9
DAP (Fig. 7e-i). Compared to *M. guttatus* and *M. nudatus* self-fertilized seed, at 9 DAP, hybrid
seed exhibit endosperm that is patchily distributed and less dense. Connecting these patterns of
early and late endosperm development in hybrid seed with the phenotypes of hybrid seeds found
in mature fruits leads us to conclude that the early arrested seed most likely become the small,
flat seeds recovered in mature fruits, while the hybrid seed that continue to develop, but in a
delayed fashion, likely mature to become the shriveled seed of mature fruits (Fig. 2e,f).
430
431
Discussion

Comparing the pattern of embryo development of *M. guttatus* and *M. nudatus* with that of *M.*
ringens (Arekal, 1965), and to a lesser extent, the cultivar *M. tigrinus* (Guilford & Fisk, 1951)
enables us to shed light on the variation in seed development within the Phrymaceae. Like *M.*
ringens and *M. tigrinus*, both *M. guttatus* and *M. nudatus* exhibit the Polygonum-type of female
gametophyte and *ab initio* cellular endosperm. They also share the development of micropylar
and chalazal haustoria, organs that appear to funnel nutrients from the maternal plant to the
developing seed (Raghavan, 1997; Nguyen *et al.,* 2000, Płachno *et al.*, 2013) (Fig. 5).
Intriguingly, Arekal (1965) The chalazal haustorium appears to be more prominent and persist
longer in *M. nudatus* than in *M. guttatus* or *M. ringens* (Arekal, 1965). Another notable
developmental difference between *M. guttatus* and *M. nudatus* is the pattern of endosperm
development: endosperm cellularization appears to produce cells that are more regularly
dispersed in *M. guttatus* than in *M. nudatus* at 5 DAP (Fig. 6a,c). Finally, the pace of embryo
development also differs, with the *M. guttatus* embryo at the globular stage by 5 DAP, while that
of *M. nudatus* is still at the 16-cell stage (Fig. 6a,c).

Dysfunctional endosperm development is associated with reduced hybrid seed viability in many
groups of flowering plants, including *Solanum* (Johnston and Hanneman, 1982; Lester & Kang,
1998), *Amaranthus* (Pal et al., 1972), *Lilium* (Dowrick & Brandham, 1970), *Oryza* (Fu *et al.,*
2009), and *Arabidopsis* (Scott *et al.,* 1998), and now, *Mimulus*, where disrupted endosperm
development manifests itself as one of two phenotypes. In the first phenotype, division of the
primary endosperm cell almost never occurs. The arrested seed enlarges slightly and seed coat
development occurs, indicating that fertilization has occurred, but initial growth quickly plateaus
and by 5 DAP, these arrested seeds are less than 1/3 the size of developing hybrid seeds. These
early arrested seed likely eventually become the small flat seeds found in mature hybrid fruits,
and never successfully germinate.

The second dysfunctional endosperm phenotype is represented by hybrid seed that appear to
develop relatively normally from 1 to 5 DAP, but exhibit impaired endosperm proliferation by 9
DAP, and are ultimately deficient in total endosperm volume as demonstrated by their shriveled,
phenotype at maturity. Embryo development is also impaired, with a slight delay evident at 5
DAP that continues to accumulate by 9 DAP (Fig. 6). The development of these seeds suggests
that even when endosperm cellularization initially proceeds, transfer of resources from the
maternal plant to this nutritive tissue may yet be limited. In addition to its primary role of
providing nutrition to the embryo, endosperm tissue actively regulates and modulates embryo
growth (Lester & Kang, 1998, Costa *et al.,* 2004, Hehenberger *et al.,* 2012). Disruption in
endosperm development may be accompanied by arrested or reduced embryo development
(Lester & Kang, 1998, Scott *et al.*, 1998). Future experiments, such as embryo rescue (e.g.,
Rebernig *et al.* 2015), would be needed to tease apart the relative contributions of endosperm vs.
embryo inviability due to the strong hybrid incompatibility *M. guttatus* and *M. nudatus*.
Intriguingly, both small, flat seeds and shriveled seeds have been previously described in crosses
involving copper-adapted *M. guttatus* (Searcy & Macnair, 1990), suggesting a common
developmental mechanism underlying failed seed development in the *M. guttatus* species
complex.
474

475   *Speciation*
476
By visualizing the progress of pollen tubes and examining development of seeds in hybrid fruits,
we conclude that interspecific pollen is functionally capable of fertilization regardless of which
species serves as maternal parent, and also that under controlled conditions, fertilization
produces large numbers of hybrid seed in both directions. Nevertheless, *M. guttatus* and *M.*
nudatus are strongly reproductively isolated. Field experiments, as well as microsatellite and
genomic sequencing data suggest that introgression between these species is rare (Gardner &
Macnair, 2000; Oneal *et al.,* 2014; L. Flagel, unpublished data). Few if any seeds of round
appearance are produced from crosses between *M. guttatus* and *M. nudatus*. Early dysfunctional
endosperm development results in small, flat hybrid seed that never germinate, and the fraction
of these apparently inviable seed can be substantial, ranging from 23% of hybrid seed when *M.*
guttatus is female, to 60% when *M. nudatus* serves as female (Fig. 3). Germination success of
shriveled hybrid seed is very low (6.0% averaged across *M. guttatus* accessions; 0% for *M.*
nudatus accessions).
490
We conclude, like Gardner and Macnair (2000), that postzygotic seed inviability forms the
primary barrier between *M. guttatus* and *M. nudatus*, but differ with their conclusions in some
respects. Most notably, we cannot rule out that subtle pollen-pistil interactions may yet serve as a
partial prezygotic isolating mechanism, since we did not explicitly test whether interspecific
pollen suffers a competitive disadvantage in fertilization success. We note that hybrid crosses
where *M. nudatus* served as female had substantially lower seed set than self-fertilized *M.*
nudatus, which is suggestive of a relative deficiency of *M. guttatus* pollen when fertilizing *M.*
nudatus. Second, speculating that bee pollinators would find landing on the *M. guttatus* stigma
more difficult, Gardner and Macnair (2000) concluded that any gene flow was likely asymmetric
and would flow more in the direction of *M. guttatus* to *M. nudatus* than the reverse. We found,
however, that none of the hybrid seed in which *M. nudatus* was the female parent germinated.
Gardner and Macnair (2000) also found that only rounded hybrid seed germinated, and at a very
low rate (< 1%), while shriveled seeds did not germinate at all. We found instead that none of the
few round, hybrid seed germinated, but up to 6.74% of shriveled hybrid seed germinated. We
attribute this difference to the fact that we recovered very few round hybrid seed to assay for
germination (*N*=3) and to differences in our categorization of hybrid seed: we found that mature
hybrid seed lie on a continuum of endosperm fullness, and that distinguishing between round and
shriveled hybrid seed was somewhat subjective. Since the shriveled appearance of these hybrid is
indicative of incomplete endosperm development (Lester & Kang, 1998), in the future, seed
weight may be a better measure of the completeness of endosperm development in hybrid seed.
511
Studies of aberrant seed development in interploidy crosses between *A. thaliana* accessions
suggest that dosage imbalances in the expression of imprinted paternally and maternally
expressed alleles and/or their regulatory targets causes dysfunctional embryo and endosperm
development and ultimately, aborted seeds (Birchler, 1993, Köhler *et al.,* 2003, Reyes and
Grossniklaus 2003, Josefsson *et al.* 2006, Erilova *et al.* 2009, Kradolfer *et al.* 2013). In theory,
such dosage imbalances could underlie failed diploid crosses as well, for example via changes in
imprinting status, sequence divergence or gene duplication of involved loci in one or both

species, all of which might alter the critical balance of dosage-sensitive genes necessary for
normal seed development (Johnston and Hanneman, 1982; Birchler & Veitia, 2010; Köhler *et al.,*
2010, Köhler *et al.,* 2012, Birchler 2014). While the vast majority of reciprocal *M. guttatus* x *M.*
nudatus seeds appear deficient in endosperm development, it is intriguing that one *M. nudatus* x
M. guttatus cross produced two rounded seeds with exploded endosperm, a phenotype associated
with excessive expression of paternally imprinted alleles in failed *A. thaliana* interploidy crosses
(Scott *et al.*, 1998). This raises the possibility that imbalances in the dosages of genes involved in
mediating maternal investment in endosperm of developing seeds may contribute to postzygotic
isolation between *M. guttatus* and *M. nudatus*. Mapping the genes associated with interspecific
endosperm failure will enable us to test this possibility.
529
While hybrid seed lethality has long been recognized as a common postzygotic isolating
mechanism among members of the ecologically and genetically diverse *M. guttatus* sp. complex
(Vickery, 1978; Gardner and Macnair, 2000), our work is the first to provide a partial
developmental mechanism—early arrested endosperm development and later failures of
endosperm proliferation—for that outcome, and to provide insight into the early stages of
endosperm and embryo development for members of the complex. We find that despite the fact
that *M. nudatus* is likely recently derived from a *M. guttatus*-like ancestor (Oneal *et al.*, 2014),
these species exhibit different patterns of embryo and endosperm development. The temporal
coordination of development across cell types with different developmental roles is increasingly
recognized as a critical aspect of achieving normal development (Del Toro-De León *et al.,* 2014;
Gillmor *et al.,* 2014). We suggest that divergence in the timing of development between *M.*
guttatus and *M. nudatus* may partly underlie the near complete hybrid barrier between them. Our
work provides a framework for further investigation of the role of this fundamental
developmental feature in the divergence between these closely related species. The extensive
genomic tools already developed for the *M. guttatus* sp. complex, including an annotated genome
sequence for *M. guttatus*, extensive Illumina re-sequence data from *M. nudatus*, and the
continued development of transgenic experimental methods (Yuan *et al.,* 2014) will only
enhance future work on this important aspect of plant evolution and speciation.
548

549   **Acknowledgements**
550
The authors would like to thank Phillip Benfey for allowing us to use a Zeiss Axio Observer to
obtain pictures of pollen-tube fluorescence, and Louise Lieberman for her assistance in obtaining
images of pollen tube growth. We thank Yasheng Gao of the Duke Microscopy Facility for
assistance with DIC imaging of unfertilized ovules and developing seed, as well as Eva Johannes
at NC State University, and Benjamin Carlson at Duke with assistance with confocal microscopy.
Jessica Selby initially collected the populations used in this study. We thank Yaniv Brandvain,
Clément Lafon-Placette, and Claudia Köhler and three anonymous referees for their insightful
comments an earlier manuscript. Funding was provided by the National Science Foundation
grants EF-0328636 and EF-0723814 to J.W., as well an NIH NRSA fellowship
(1F32GM097929-01) to E.O.

References

Arekal GD. 1965. Embryology of *Mimulus ringens*. *Botanical Gazette* **1965**: 58-66.

Beldade P, Koops K, Brakefield PM. 2002. Developmental constraints versus flexibility in morphological evolution. *Nature* **416**: 844-847.

Bharathan G. 1999. Endosperm development in angiosperms: a phylogenetic analysis. *Biodiversity, taxonomy, and ecology*, 167-180.

Birchler JA. 1993. Dosage analysis of maize endosperm development. Annual Review of Genetics **27**: 181-204.

Birchler JA. 2014. Interploidy hybridization barrier of endosperm as a dosage interaction. Frontiers in Plant Science **5**: 1-4.

Birchler JA & Veitia RA. 2010. The gene balance hypothesis: implications for gene regulation, quantitative traits and evolution. New Phytologist **186**: 54-62.

Boavida LC, Silva JP, Feijó JA. 2001. Sexual reproduction in the cork oak (*Quercus suber* L). II. Crossing intra-and interspecific barriers. *Sexual Plant Reproduction* **14**: 143-152.

Bomblies K, Lempe J, Epple P, Warthmann N, Lanz C, Dangl JL, Weigel D. 2007. Autoimmune response as a mechanism for a Dobzhansky-Muller-type incompatibility syndrome in plants. *PLoS biology* **5**: e236.

Bonner JT. 1988. *The evolution of complexity*. Princeton Univ. Press, Princeton, NJ.

Burkart-Waco D, Ngo K, Dilkes B, Josefsson C, Comai L. 2013. Early disruption of maternal–zygotic interaction and activation of defense-like responses in *Arabidopsis* interspecific crosses. *The Plant Cell Online*, *25*(6), 2037-2055.

Brandvain Y, Haig D. 2005. Divergent mating systems and parental conflict as a barrier to hybridization in flowering plants. *The American Naturalist* **166**: 330-338.

Brink RA, & Cooper DC. 1947. The endosperm in seed development. *The Botanical Review* **13**: 479-541.

Campbell DR, Alarcon R, Wu CA. 2003. Reproductive isolation and hybrid pollen disadvantage in *Ipomopsis*. *Journal of evolutionary biology* **16**: 536-540.

Casson S, Spencer M, Walker K, & Lindsey K. 2005. Laser capture microdissection for the analysis of gene expression during embryogenesis of Arabidopsis. *The Plant Journal* **42**: 111-123.

Cooley AM, Willis JH. 2009. Genetic divergence causes parallel evolution of flower color in
Chilean Mimulus. *New Phytologist* **183**: 729-739.

Costa CB, Costa JA, de Queiroz LP, Borba EL. (2013). Self-compatible sympatric *Chamaecrista*
(Leguminosae-Caesalpinioideae) species present different interspecific isolation mechanisms
depending on their phylogenetic proximity. *Plant Systematics and Evolution*, *299*(4), 699-711.

Costa LM, Gutierrez-Marcos JF, Dickinson HG. 2004. More than a yolk: the short life and
complex times of the plant endosperm. *Trends in plant science* **9**: 507-514.

Coyne JA, Orr HA. 2004. *Speciation*. Sunderland, MA: Sinauer Associates.

Del Toro-De León G, García-Aguilar M, Gillmor CS. 2014. Non-equivalent contributions of
maternal and paternal genomes to early plant embryogenesis. *Nature* **514**, 624-627.

Deshpande SS, Cheryan M, Salunkhe DK, & Luh BS. 1986. Tannin analysis of food products.
Critical Reviews in Food Science & Nutrition **24:**, 401-449.

Dowrick GJ, Brandram SN. 1970. Abnormalities of endosperm development in *Lilium* hybrids.
Euphytica **19**: 433-442.

Erilova A, Brownfield L, Exner V, Rosa M, Twell D, Scheid OM, Hennig L, Köhler C. 2009.
Imprinting of the Polycomb Group gene *MEDEA* srves as a ploidy sensor in *Arabidopsis*. *PLoS*
Genetics **5**: e1000663

Fishman L, Aagaard J, Tuthill JC. 2008. Toward the evolutionary genomics of gametophytic
divergence: patterns of transmission ratio distortion in monkeyflower (*Mimulus*) hybrids reveal a
complex genetic basis for conspecific pollen precedence. *Evolution* **62**: 2958-2970.

Floyd SK, Friedman WE. 2000. Evolution of endosperm developmental patterns among basal
flowering plants. *International Journal of Plant Sciences* **161**: S57-S81.

Friedman WE. 2001. Comparative embryology of basal angiosperms. *Current Opinion in Plant*
Biology **4**: 14-20

Friedman, WE, Madrid, EN, Williams, JH. 2008. Origin of the fittest and survival of the fittest:
relating female gametophyte development to endosperm genetics. *International Journal of Plant*
Sciences **169**: 79-92.

Fu XL, Lu YG, Liu XD, Li JQ. 2009. Crossability barriers in the interspecific hybridization
between *Oryza sativa* and *O. meyeriana*. *Journal of integrative plant biology* **51**: 21-28.

Galbraith DW, Harkins KR, Knapp S. 1991. Systemic endopolyploidy in *Arabidopsis thaliana*.
Plant Physiology **96**: 985-989.
650
651 Galen C, Newport MEA. 1988. Pollination quality, seed set, and flower traits in *Polemonium*
652 *viscosum*: complementary effects of variation in flower scent and size. *American Journal of*
653 *Botany*, 900-905.
654
655 Gardner M, Macnair M. 2000. Factors affecting the co-existence of the serpentine endemic
656 Mimulus nudatus Curran and its presumed progenitor, *Mimulus guttatus* Fischer ex DC.
657 *Biological Journal of the Linnean Society* **69**: 443-459.
658
659 Gillmor CS, Silva-Ortega CO, Willmann MR, Buendía-Monreal M, Poethig RS. 2014. The
660 Arabidopsis Mediator CDK8 module genes CCT (MED12) and GCT (MED13) are global
661 regulators of developmental phase transitions. *Development* **141**: 4580-4589.
662
663 Girke T, Todd J, Ruuska S, White J, Benning C, & Ohlrogge J. 2000. Microarray analysis of
664 developing Arabidopsis seeds. *Plant Physiology* **124**: 1570-1581.
665
666 Grossniklaus U, Spillane C, Page DR, Köhler C. 2001. Genomic imprinting and seed
667 development: endosperm formation with and without sex. *Current opinion in plant biology*, **4**:
668 21-27.
669
670 Guignard L. 1899. Sur les anthérozoides et la double copulation sexuelle chez les végétaux
671 angiospermes. *Comptes Rendus de l'Aca- demie des Sciences de Paris* **128**: 864-871.
672
673 Haig D. 2013. Kin conflict in seed development: an interdependent but fractious collective.
674 *Annual review of cell and developmental biology* **29**: 189-211.
675
676 Haig D, Westoby M. 1991. Genomic imprinting in endosperm: its effect on seed development in
677 crosses between species, and between different ploidies of the same species, and its implications
678 for the evolution of apomixis. *Philosophical Transactions of the Royal Society, B* **333**: 1-13.
679
680 Hehenberger E, Kradolfer D, Köhler C. 2012. Endosperm cellularization defines an important
681 developmental transition for embryo development. *Development* **139**: 2031-2039.
682
683 Hoballah ME, Gübitz T, Stuurman J, Broger L, Barone M, Mandel T, Dell'Olivo A, Arnold M,
684 Kuhlemeier C. 2007. Single gene–mediated shift in pollinator attraction in *Petunia*. *The Plant*
685 *Cell Online* **19**: 779-790.
686
687 Hsieh TF, Shin J, Uzawa R, Silva P, Cohen S, Bauer MJ, Hashimoto M, Kirkbride RC, Harada
688 JJ, Zilberman D, Fischer RL. 2011. Regulation of imprinted gene expression in Arabidopsis
689 endosperm. *Proceedings of the National Academy of Sciences* **108**: 1755-1762.
690

Johnston SA, Hanneman RE. 1982. Manipulations of endosperm balance number overcome
crossing barriers between diploid *Solanum* species. Science **217**: 446-448.

Josefsson C, Dilkes B, Comai L. 2006. Parent-dependent loss of gene silencing during
interspecies hybridization. *Current Biology* **16**: 1322-1328.

Kearns CA, Inouye DW. 1993. *Techniques for Pollination Biologists*. University Press of
Colorado, CO. pp. 77–151

Köhler, C, L. Hennig, C. Spillane, P. Stephane, W. Gruissem, and U. Grossniklaus. 2003. The
Polycomb-group protein *MEDEA* regulates seed development by controlling expression of the
MADS-box gene *PHERES1*. *Genes and Development* **17**: 1540-1553

Kradolfer D, Hennig L, Köhler C. 2013. Increased maternal genome dosage bypasses the
requirement of the *FIS* polycomb repressive complex 2 in *Arabidopsis* seed development. *PLoS*
genetics **9**: e1003163.

Köhler C, Mittelsten Scheid O, Erilova A. 2010. The impact of the triploid block on the origin
and evolution of polyploid plants. *Trends in Genetics* **26**: 142-148.

Köhler C, Wolff P, Spillane C. 2012. Epigenetic mechanisms underlying genomic imprinting in
plants. *Annual review of plant biology* **63**: 331-352.

Maynard Smith J, Burian R, Kauffman S, Alberch P, Campbell J, Goodwin B, Lande R, Raup D,
Wolpert L. 1985. Developmental constraints and evolution: a perspective from the Mountain
Lake conference on development and evolution. *Quarterly Review of Biology* **60:** 265-287.

Lester RN, Kang JH. 1998. Embryo and endosperm function and failure in *Solanum* species and
hybrids. *Annals of Botany* **82**: 445-453.

Linkies A, Graeber K, Knight C, Leubner-Metzger G. 2010. The evolution of seeds. *New*
Phytologist **186**: 817-831.

Lowry DB, Willis JH. 2010. A widespread chromosomal inversion polymorphism contributes to
a major life-history transition, local adaptation, and reproductive isolation. *PLoS Biology* **8**:
e1000500.

Martin NH, Willis JH. 2010. Geographical variation in postzygotic isolation and its genetic basis
within and between two *Mimulus* species. *Philosophical Transactions of the Royal Society B*
365:3469-2478.

Nawaschin SG. 1898. Resultate einer Revision der Befruchtungs- vorgänge bei Lilium martagon
und Fritillaria tenella. *Bulletin de l'Academie des Sciences de Saint Petersbourg* **9**: 377-382.

Nguyen H, Brown RC, Lemmon BE. 2000. The specialized chalazal endosperm in *Arabidopsis*
thaliana and *Lepidium virginicum* (Brassicaceae). *Protoplasma* **212**: 99-110.
737
738   Oneal E, Lowry DB, Wright KM, Zhu Z, Willis JH. 2014. Divergent population structure and
739   climate associations of a chromosomal inversion polymorphism across the *Mimulus guttatus*
740   species complex. *Molecular ecology* **23**: 2844-2860.
741
742   Pal M, Pandey RM, Khoshoo TN. 1982. Evolution and improvement of cultivated amaranths.
743   *Journal of Heredity* **73**: 353-356.
744
745   Płachno BJ, Świątek P, Sas-Nowosielska H, Kozieradzka-Kiszkurno M. 2013. Organisation of
746   the endosperm and endosperm–placenta syncytia in bladderworts (Utricularia, Lentibulariaceae)
747   with emphasis on the microtubule arrangement. *Protoplasma* **250**: 863-873.
748
749   Ramsey J, Bradshaw HD, Schemske DW. 2003. Components of reproductive isolation between
750   the monkeyflowers Mimulus lewisii and M. cardinalis (Phrymaceae). *Evolution* **57**: 1520-1534.
751
752   Rasband WS. ImageJ. U. S. National Institutes of Health, Bethesda, Maryland, USA,
753   http://imagej.nih.gov/ij/, 1997-2014.
754
755   Reyes JC, Grossniklaus U. 2003. Diverse functions of *Polycomb* group proteins during plant
756   development. *Seminars in Cell and Developmental Biology* **14**: 77-84
757
758   Rieseberg LH, Willis JH. 2007. Plant speciation. *Science* **317**: 910-914.
759
760   Ritland C, Ritland K. 1989. Variation of sex allocation among eight taxa of the *Mimulus guttatus*
761   species complex (Scrophulariaceae). *American Journal of Botany* **1989**: 1731-1739.
762
763   Roszak, P, & Köhler, C. 2011. Polycomb group proteins are required to couple seed coat
764   initiation to fertilization. *PNAS* **108**: 20826-20831.
765
766   Running, M. 2007. Nuclear Staining of Plants for Confocal Microscopy. *Cold Spring Harbor*
767   *Protocols*, **2007**: pdb-prot4685.
768
769   Schemske DW, Bradshaw HD. 1999. Pollinator preference and the evolution of floral traits in
770   monkeyflowers (*Mimulus*). *PNAS* **96**: 11910-11915.
771
772   Scott RJ, Spielman M, Bailey J, Dickinson HG. 1998. Parent-of-origin effects on seed
773   development in *Arabidopsis thaliana*. *Development* **125**: 3329-3341.
774
775   Searcy KB, Macnair MR. 1990. Differential seed production in Mimulus guttatus in response to
776   increasing concentrations of copper in the pistil by pollen from copper tolerant and sensitive
777   sources. *Evolution* **1990**: 1424-1435.
778

779   Soltis PS, Soltis DE. 2004. The origin and diversification of angiosperms. *American Journal of*
780   *Botany* **91**: 1614-1626.
781
782   Tiffin P, Olson S, Moyle LC. 2001. Asymmetrical crossing barriers in angiosperms. *Proceedings*
783   *of the Royal Society of London, B* **268**: 861-867.
784
785   Vickery RK. 1966. Speciation and isolation in section Simiolus of the genus *Mimulus*. *Taxon* **15:**
786   55-63.
787
788   Vickery RK. 1978. Case studies in the evolution of species complexes in *Mimulus*. In: Hecht
789   MK, Steere WC, Wallace B., *eds. Evolutionary Biology*, Volume 11, pp 405-507.
790
791   Williams, JH. 2009. *Amborella trichopoda* (Amborellaceae) and the evolutionary developmental
792   origins of the angiosperm progamic phase. *American Journal of Botany*, **96:** 144-165
793
794   Wright KM, Lloyd D, Lowry DB, Macnair MR, Willis JH. 2013. Indirect evolution of hybrid
795   lethality due to linkage with selected locus in *Mimulus guttatus*. *PLoS biology* **11:** e1001497.
796
797   Wu CA, Lowry DB, Cooley AM, Wright KM, Lee YW, Willis JH. 2007. *Mimulus* is an
798   emerging model system for the integration of ecological and genomic studies. *Heredity* **100**: 220-
799   230.
800
801   Yuan YW, Sagawa JM, Frost L, Vela JP, Bradshaw HD. 2014. Transcriptional control of floral
802   anthocyanin pigmentation in monkeyflowers (*Mimulus*). *New Phytologist* **204**: 1013-1027.
803
804

804   **Figure 1**. Images of pollen tubes penetrating the ovaries within 24-hours of pollination for
805   interspecific crosses of *M. guttatus* and *M. nudatus*. Pollen tubes are stained with aniline blue
806   and visualized with a Zeiss Axio Observer equipped with a fluorescent lamp and DAPI filter. (a)
807   An unpollinated *M. guttatus* ovary. (b) Pollen tubes from *M. nudatus* pollen growing down a *M.*
808   *guttatus* style. (c) Pollen tubes from *M. guttatus* pollen growing down a *M. nudatus* style.
809
Figure 2. Developing fruits resulting from *M. guttatus* and *M. nudatus* self-fertilizations and
reciprocal crosses of *M. guttatus* x *M. nudatus*, as well as images of mature seed recovered from
self-fertilized and hybrid fruits. All crosses are female x male. (a) *M. guttatus* self-fertilized fruit.
(b) *M. nudatus* self-fertilized fruit. (c) *M. guttatus* x *M. nudatus* fruit. (d) *M. nudatus* x *M.*
guttatus. (e) round *M. guttatus* seed; shriveled and flat *M. guttatus* x *M. nudatus* seed. (f) **d**:
round *M. nudatus* seed; shriveled and flat *M. nudatus* x *M. guttatus* seed. All seed types appear
to have developed a seed coat.
817
818   **Figure 3.** Percentages of each mature seed phenotype resulting from self-fertilized *M. guttatus*
819   and *M. nudatus* self-fertilizations and reciprocal *M. guttatus* x *M. nudatus* crosses, pooled across
820   accessions. All crosses are female x male.
821
822   **Figure 4.** Seed set for self-fertilized and reciprocal crosses of *M. guttatus* and *M. nudatus*,
823   pooled across accessions. For seed set by accession, see Supplementary Figure 1. All crosses are
824   female x male.
825
Figure 5. Development of normal *M. guttatus* and *M. nudatus* seeds. **an**: antipodal nuclei; **ccn**:
central cell nucleus; **c**: chalazal end; **ch**: chalazal haustorium; **ec**: egg cell; **en**: endosperm; **em**:
embryo; **m**: micropylar end; **mh**: micropylar haustorium; **mn**: micropylar nucleus; **pen**: primary
endosperm nucleus; **sn**: synergid nucleus; (a) *M. guttatus* female gametophyte. (b) *M. guttatus*, 2
days after pollination (DAP); endosperm (**en**) has undergone at least three divisions. (c) *M.*
guttatus, 3 DAP with faintly visible chalazal haustorium (**ch**), an embryo (**em**) and micropylar
nuclei **(mn)** with multiple nucleoli. (d) *M. guttatus*, 4 DAP with a 16-cell embryo (**em**) with
suspensor, and a micropylar nucleus (**mn**) with multiple nucleoli. (e) *M. guttatus*, 5 DAP.
Endosperm (**en**) is densely packed and surrounds the globular embryo (**em**). (f) *M. nudatus*, 2
DAP with one nucleus (**en**) in the chalazal domain and one nucleus in the micropylar domain
(**mn**). (g) *M. nudatus*, 3 DAP: a chalazal haustorium (**ch**) is plainly visible, a micropylar
haustorium (**mh**) with nuclei has been established, and transverse division of endosperm nuclei
(en) is occurring. (h) *M. nudatus*, 4 DAP, displaying a prominent chalazal haustorium (**ch**) with
two heavily nucleolated nuclei, proliferating endosperm (**en**), an 8-cell embryo (**em**) and a
micropylar haustorium (**mh**) (the cells of the mh are not visible in this microscopic plane). (i) *M.*
nudatus, 5 DAP. The chalazal haustorium (**ch**) has largely degraded, and a 16-cell embryo (**em**)
is visible.
843
Figure 6. Compared to self-fertilized *M. guttatus* and *M. nudatus* seed at 5 DAP (a-d) and 9
DAP (e-i), reciprocal *M. guttatus* x *M. nudatus* hybrid seed exhibit delayed embryo growth and
impaired endosperm development. **c**: chalazal end; **en**: endosperm; **em**: embryo: **m**: micropylar
end; **mn**: micropylar nuclei; **sc**: seed coat. (a) *M. guttatus* self-fertilized seed at 5 DAP.

Endosperm (**en**) has nuclei with regularly dispersed cell walls and an embryo (**em**) is at the
globular stage. (b) *M. guttatus* x *M. nudatus* seed at 5 DAP. Endosperm (**en**) resembles that of
self-fertilized *M. guttatus*, but embryo (**em**) is delayed at 8-cell stage. (c) *M. nudatus* self-
fertilized seed at 5 DAP. Endosperm (**en**) has nuclei with regularly dispersed cell walls and
embryo (**em**) is at the 16-cell stage; micropylar nuclei (**mn**) are also visible. (d) *M. nudatus* x *M.*
guttatus seed at 5 DAP. Endosperm (**en**) is poorly developed and embryo (**em**) is at the 8-cell
stage. (e) *M. guttatus* seed at 9 DAP. These seeds have a well-developed seed coat (**sc**), densely
packed endosperm (**en**), and a torpedo stage embryo (**em**). (f and g) *M. guttatus* x *M. nudatus*
seed at 9 DAP, with loosely packed and irregularly deposited endosperm (**en**) and a heart stage
embryo (**em**). (h) *M. nudatus* seed at 9 DAP with densely packed endosperm (**en**) and a heart
stage endosperm (**em**). (i) *M. nudatus* x *M. guttatus* seed with irregularly developed endosperm
(**en**) and late globular stage embryo (**em**).

Figure 7. Images of the *M. guttatus* female gametophyte (a) and early arrested hybrid seed from
reciprocal, sympatric crosses between *M. guttatus* and *M. nudatus* (b-f). Early arrested hybrid
seed show clear signs of fertilization, including a widened primary endosperm cell (**pec**)
compared to the width of the gametophytic central cell (a, b, d), signs of mitosis within the
primary endosperm nucleus (**pen**) (i.e., visible nucleoli), the development of micropylar nuclei
(**mn**) and, occasionally, a visible embryo (**em**). **c**: chalazal end; **ccn**: central cell nucleus; **cw**: cell
wall; **em**: embryo; **en**: endosperm; **enn**: endosperm nucleus; **m**: micropylar end; **mh**: micropylar
haustorium; **mn**: micropylar nucleus; **pec**: primary endosperm cell; **pen**: primary endosperm
nucleus; All crosses are female x male. (a) *M. guttatus* female gametophyte. (b) *M. guttatus* x *M.*
nudatus, 2 days after pollination (DAP) with an undivided primary endosperm nucleus (**pen**)**;** a
micropylar nucleus is visible. (c) arrested *M. guttatus* x *M. nudatus* seed, 5 DAP; endosperm
nucleus (**enn**) has multiple nucleoi evident, but there is little to no endosperm present; a
micropylar nucleus (**mn**) with multiple nucleoli and an embryo (**em**) are presented. (d) *M.*
nudatus x *M. guttatus* seed, 2 DAP, with an undivided primary endosperm nucleus (**pen**). (e)
arrested *M. nudatus* x *M. guttatus* seed, 5 DAP. One endosperm nucleus (**enn**) with at least one
nucleolus is visible, as is a possible endosperm cell wall (**cw**). (f) *M. nudatus* x *M. guttatus*
arrested seed, 9 DAP. A primary endosperm cell (**pec**) and micropylar haustorium (**mh**) are
visible.
878
879

Figure 8. Growth trajectories of self-fertilized and hybrid seed, pooled across accessions. For
seed size increase by accession, see Supplementary Figure 2. All crosses are female x male. (a)
M. guttatus self-fertilized seed and *M. guttatus* x *M. nudatus* hybrid seed. (b) *M. nudatus* self-
fertilized seed and *M. nudatus* x *M. guttatus* seed.

Figure 9. Vanillin stain test for developed seed coat. (a) Ovule from *M. nudatus* unpollinated
ovaries collected 5 days after emasculation. (b) *M. guttatus* x *M. nudatus* seed, 5 DAP (C) *M.*
nudatus x *M. guttatus* seed, 5DAP. Two hybrid seed types are visible: small, arrested seed. Both
appear to have seed coat.
889

890   **Supplementary Figure 1**. (a) Seed set for self-fertilized *M. guttatus* accessions and sympatric *M.*
891   *guttatus* x *M. nudatus* crosses. (b) Seed set for self-fertilized *M. nudatus* accessions and
892   sympatric *M. nudatus* x *M. guttatus* crosses. All crosses are female x male.
893
894   **Supplementary Figure 2. (A)** Percentages of each mature seed phenotype resulting from self-
895   fertilized *M. guttatus* and sympatric *M. guttatus* x *M. nudatus* crosses. (B) Percentages of each
896   mature seed phenotype resulting from self-fertilized *M. nudatus* and sympatric *M. nudatus* x *M.*
897   *guttatus* crosses. All crosses are female x male.
898
899   **Supplementary Figure 3.** Growth trajectories of self-fertilized seed and hybrid seed from
900   sympatric, pairwise crosses between *M. guttatus* x *M. nudatus*, broken down by accession. All
901   crosses are female x male. (a) CSS4 self-fertilized seed and CSS4 x CSH10 hybrid seed. (b)
902   DHR14 self-fertilized seed and DHR14 x DHRo22 hybrid seed. (c) CSH10 self-fertilized seed
903   and CSH10 x CSS4 hybrid seed. (d) DHRo22 self-fertilized seed and DHRo22 x DHR14 seed.
904
905

905
906
907
908 **Table 1** Sampling localities and accession names for sympatric serpentine-adapted *M. guttatus*
909 and *M. nudatus* populations.
910

Accessions	Latitude (N)	Longitude (W)
M. guttatus/*M. nudatus*		
CSS4/CSH10	38.861	-122.415
DHR14/DHRo22	38.859	-122.411

917
918

919 **Table 2** Pollen germination and ovary penetration data for sympatric interspecific crosses
920 between *M. guttatus* and *M. nudatus*. All crosses are female x male.
921

Cross	N	Pollen germinated	Ovary Penetration
M. guttatus x *M. nudatus*			
CSS4 x CSH1	10	100%	90%
DHR14 x DHRo22	10	100%	80%
M. nudatus x *M. guttatus*			
CSH10 x CSS4	10	100%	90%
DHRo22 x DHR14	10	100%	80%

932
933
934

Table 3 Germination success of self-fertilized and hybrid seed. All crosses are female x male.

	Seed type	% Germinated	N
Self-fertilized			
M. guttatus CSS4	round	96.0	25
M. guttatus CSS4	shriveled	11.3	106
M. guttatus CSS4	small, flat	0.0	14
M. guttatus DHR14	round	88.0	25
M. guttatus DHR14	shriveled	15.6	115
M. nudatus CSH10	round	81.2	16
M. nudatus CSH10	shriveled	4.40	91
M. nudatus CSH10	flat	0.0	2
M. nudatus DHRo22	round	43.8	16
M. nudatus DHRo22	shriveled	0.0	30
M. guttatus x *M. nudatus*			
CSS4 x CSH10	round	0.0	1
CSS4 x CSH10	large, flattened	10.7	75
CSS4 x CSH10	small, flattened	0.0	50
DHR14 x DHRo22	large, flattened	2.78	108
DHR14 x DHRo22	small, flattened	0.0	17
M. nudatus x *M. guttatus*			
CSH10 x CSS4	round, exploded	0.0	2
CSH10 x CSS4	large, flattened	0.0	104
CSH10 x CSS4	small, flattened	0.0	50
DHRo22 x DHR14	large, flattened	0.0	98
DHRo22 x DHR14	small, flattened	0.0	138

967
968
969   Figure 1
970
971

972
973
974   Figure 2
975
976

977
978   Figure 3
979
980

981
982
983   Figure 4
984
985

986
987
988   Figure 5
989
990

990
991

992
993
994 Figure 6
995

998 Figure 7

1000
1001
1002 Figure 8a
1003

1003

1004
1005
1006   Figure 8b
1007
1008

1008

1009
1010   Figure 9
1011
1012
1013