Variation in linked selection and recombination drive genomic divergence during allopatric speciation of European and American aspens

Jing Wang[1], Nathaniel R. Street[2], Douglas G. Scofield[1,3,4], Pär K. Ingvarsson[1]

[1] Department of Ecology and Environmental Science, Umeå University, SE-90187, Umeå, Sweden

[2] Umeå Plant Science Centre, Department of Plant Physiology, Umeå University, SE-90187, Umeå, Sweden

[3] Department of Ecology and Genetics: Evolutionary Biology, Uppsala University, Uppsala, Sweden

[4] Uppsala Multidisciplinary Center for Advanced Computational Science, Uppsala University, Uppsala, Sweden

Corresponding author:

Dr Pär K. Ingvarsson, Department of Ecology and Environmental Science, Umeå University, Umeå, SE 90187, Sweden. Phone: +46907867414; Fax: +46-(0)-90-786-6705; E-mail: par.ingvarsson@emg.umu.se

Keywords: *Populus tremula*, *P. tremuloides*, Whole-genome re-sequencing, demographic histories, heterogeneous genomic differentiation, linked selection, recombination

Abstract

Despite the global economic and ecological importance of forest trees, the genomic basis of differential adaptation and speciation in tree species is still poorly understood. *Populus tremula* and *P. tremuloides* are two of the most widespread tree species in the Northern Hemisphere. Using whole-genome re-sequencing data of 24 *P. tremula* and 22 *P. tremuloides* individuals, we find that the two species diverged ~2.2-3.1 million years ago, coinciding with the severing of the Bering land bridge and the onset of dramatic climatic oscillations during the Pleistocene. Both species have experienced substantial population expansions following long-term declines after species divergence. We detect widespread and heterogeneous genomic differentiation between species, and in accordance with the expectation of allopatric speciation, coalescent simulations suggest that neutral evolutionary processes can account for most of the observed patterns of genomic differentiation. However, there is an excess of regions exhibiting extreme differentiation relative to those expected under demographic simulations, which is indicative of the action of natural selection. Overall genetic differentiation is negatively associated with recombination rate in both species, providing strong support for a role of linked selection in generating the heterogeneous genomic landscape of differentiation between species. Finally, we identify a number of candidate regions and genes that may have been subject to positive and/or balancing selection during the speciation process.

Introduction

Understanding how genomes diverge during the process of speciation has received a great deal of attention in the evolutionary genetics literature in recent years (Nosil et al. 2009; Strasburg et al. 2012; Seehausen et al. 2014). Under strict neutrality, differentiation is expected to accumulate as a result of the stochastic fixation of polymorphisms by genetic drift (Coyne and Orr 2004). Demographic processes, such as population bottlenecks or expansions, can accelerate or decelerate the rate of differentiation through changes in the effective population sizes of nascent daughter species (Avise 2000). Random genetic drift and demographic processes are both expected to affect the entire genome (Luikart et al. 2003). Natural selection, however, only influence loci involved in ecological specialization and/or reproductive isolation, resulting in patterns of polymorphisms and divergence that deviate from neutral predictions (Luikart et al. 2003; Via 2009). The functional architectures of genomes, e.g. mutation and recombination rates, are also important factors in determining genomic landscape of differentiation (Noor and Bennett 2009; Nachman and Payseur 2012; Renaut et al. 2013). For example, suppressed recombination could increase genetic differentiation either by limiting inter-species gene flow to prevent the break-up of co-adapted alleles, or through the diversity-reducing effects of linked selection (Noor and Bennett 2009). However, disentangling the relative importance of these evolutionary forces when interpreting patterns of genomic divergence remains a challenge in speciation genetics.

With the advance of next generation sequencing (NGS) technologies, a growing number of studies have found highly heterogeneous patterns of genomic differentiation between recently diverged species (Turner et al. 2005; Ellegren et al. 2012; Renaut et al. 2013; Carneiro et al. 2014; Feulner et al. 2015). A common explanation for these patterns is that levels of gene flow between species differ across the genome. Increased genetic divergence is usually observed in a small number of regions containing loci involved in reproductive isolation ('speciation islands'), where as the remainder of the genome is still permeable to ongoing gene flow and therefore shows lower levels of differentiation (Nosil et al. 2009; Sousa and Hey 2013). However, some recent studies have argued that highly differentiated regions represent 'incidental islands' that are not tied to the speciation processes per se. Rather they are

seen simply as a result of the diversity-reducing effects of linked selection that
accelerate lineage sorting of ancestral variation and increase interspecific
differentiation, especially in regions of reduced recombination (Turner and Hahn
2010; Cruickshank and Hahn 2014). In addition, long-term balancing selection is
supposed to maintain stable trans-species polymorphisms and leaves signatures of
unusually low genetic differentiation between species (Charlesworth 2006). Under
these scenarios, natural selection alone is sufficient to generate patterns of
heterogeneous genomic differentiation even under complete allopatry (Noor and
Bennett 2009; Turner and Hahn 2010). Finally, strictly neutral forces, such as
stochastic genetic drift and complex demographic processes, can also create
heterogeneous genomic divergence and generate patterns of divergence and
polymorphism that mimic the effects of selection (Nosil et al. 2009; Campagna et al.
2015). In general, the three hypotheses listed above are not mutually exclusive and
exhaustive examination of these hypotheses requires detailed information on the
speciation process, such as the timing of speciation, the geographic and demographic
context in which it occurred (Nosil and Feder 2012).

Although largely understudied compared to other model species, forest trees
represent a promising system to understand the genomic basis of species divergence
and adaptive evolution; as a group they have developed diverse strategies to adapt and
thrive across a wide range of climates and environments (Neale and Kremer 2011).
Populus tremula (European aspen) and *P. tremuloides* (American aspen) are two of
the most ecologically important and geographically widespread tree species of the
Northern Hemisphere (Figure 1a). Both are keystone species, display rapid growth,
with high tolerance to environmental stresses and long-distance pollen and seed
dispersal via wind (Eckenwalder 1996; Müller et al. 2012). In addition, they both
harbor among the highest level of intraspecific genetic diversity reported in plant
species so far (Wang et al. forthcoming). Based on their morphological similarity and
close phylogenetic relationships, they are considered to be sister species, or less
commonly, conspecific subspecies (Eckenwalder 1996; Wang et al. 2013). They can
readily cross and artificial hybrids usually show high heterosis (Hamzeh and
Dayanandan 2004; Tullus et al. 2012). A recent study based on a handful of nuclear
and chloroplast loci suggests that the first opening of the Bering land bridge may have
driven the allopatric speciation of the two species (Du et al. 2015).

Due to their continent-wide distributions, extraordinary levels of genetic and

phenotypic diversity, along with the availability of a high-quality reference genome in

the congener, *P. trichocarpa* (Tuskan et al. 2006), *P. tremula* and *P. tremuloides*

represent a promising system for evaluating how various evolutionary processes have

shaped the patterns of genomic divergence during speciation. In this study, we use

whole-genome re-sequencing data from both species to estimate and infer their

divergence time and historical demographic processes of the two species. Explicit

characterizations of the demographic history not only allow us to estimate historical

population size fluctuations in both species, but also increase the accuracy of

identifying regions or genes that have been under natural selection. By incorporating

the inferred demographic scenarios into the null model, we investigate the extent to

which demographic and selective events have contributed to the overall patterns of

genomic differentiation between the two species. We also identify a number of outlier

regions and genes that likely have evolved in response to positive and/or balancing

selection during the speciation process.

Results

We generated whole-genome resequencing data for 24 *P. tremula* and 22 *P.*

tremuloides. The high extent of conserved synteny between the genomes of aspen and

P. trichocarpa (Pakull et al. 2009; Robinson et al. 2014) allowed us to map all reads

to the *P. trichocarpa* reference genome (v3.0) (Tuskan et al. 2006) after adapter

removal and quality trimming (see Materials and Methods). More than 88% of

sequenced reads were aligned and the mean coverage of uniquely mapped reads per

site was 25.1 and 22.5 in samples of *P. tremula* and *P. tremuloides*, respectively

(Table S1). Two complementary bioinformatics approaches were used in this study

(Figure S1): (1) For those population genetic statistics that relied on inferred site-

frequency-spectrum (SFS), estimation was performed directly from genotype

likelihoods without calling genotypes (Nielsen et al. 2011) as implemented in

ANGSD (Korneliussen et al. 2014). (2) For those estimations that required accurate

genotype calls, single nucleotide polymorphisms (SNPs) and genotypes were called

with HaplotypeCaller in GATK (Danecek et al. 2011). In total, we identified

5,894,205 and 6,281,924 SNPs passing filtering criteria (see Materials and Methods)

across the 24 *P. tremula* samples and 22 *P. tremuloides* samples, respectively.

Population structure

We used NGSadmix (Skotte et al. 2013) to infer individual ancestry based on

genotype likelihoods, which takes the uncertainty of genotype calling into account. It

clearly sub-divided all sampled individuals into two species-specific groups when the

number of clusters (*K*) was 2 (Figure 1b). When *K* = 3, there was evidence for further

population sub-structuring in *P. tremuloides*, where individuals from populations of

Alberta and Wisconsin clustered into two subgroups. With *K* = 4, most individuals of

P. tremula were inferred to be a mixture of two genetic components, showing slight

clinal variation with latitude. No further structure was found when *K* = 5 (Figure 1b).

A principal component analysis (PCA) further supported these results (Figure 1c).

Only the first two components were significant based on the Tracy-Widom test (Table

S2), which explained 21.4% and 2.1% of total genetic variance, respectively (Figure

1c). Among the total number of polymorphisms in the two species, fixed differences

between *P. tremula* and *P. tremuloides* accounted for 1.1%, whereas 16.7% of

polymorphisms were shared between species, with the remaining polymorphic sites

being private in either of the two species (Figure 1d).

To further examine the extent of population subdivision in *P. tremuloides*, we

measured F_{ST} and d_{xy} between the two subpopulations (Alberta and Wisconsin) along

individual chromosomes (Table S3). We found low levels of genetic differentiation

(average F_{ST}: 0.0443±0.0325) between the two subpopulations (Table S3). Total

sequence differentiation in the inter-population comparison (mean d_{xy} =

0.0165±0.0083) was similar to mean sequence differences in intra-population

comparisons ($\pi_{Alberta}$: 0.0161±0.0081; $\pi_{Wisconsin}$: 0.0157±0.0080, Table S3), indicating

that individuals of the two populations were genetically not more different from each

other than individuals within each population. Based on the summaries of site

frequency spectrum (Tajima's *D* and Fay & Wu's *H*), both populations exhibited

strong skews toward low-frequency variants (negative *D*) and intermediate skews

toward high frequency-derived variants (negative *H*) (Table S3), suggesting that they

likely experienced similar species-wide demographic events.

Demographic histories

We used *fastsimcoal2* (Excoffier et al. 2013), a coalescent simulation-based method,

to infer the past demographic histories of *P. tremula* and *P. tremuloides* from the joint

site frequency spectrum. Eighteen divergence models were evaluated (Figure S2;

Table S4), and all models began with the split of the ancestral population into two

derived populations and differed in terms of (i) whether post-divergence gene flow

was present or not, (ii) levels and patterns of gene flow between the two species, and

(iii) how population size changes occurred, either at the time of species divergence or

afterwards (Figure S2). The best-fitting model was a simple isolation-with-migration

model where, after the two species diverged, *P. tremuloides* experienced exponential

growth and whereas a stepwise population size change occurred in *P. tremula* (Figure

2a). The exact parameter estimates of divergence time, gene flow, population sizes

and their associated 95% confidence intervals (CIs) are given in Table 1. The

estimated divergence time between *P. tremula* and *P. tremuloides* (T_{DIV}) was ~2.3

million years ago (Mya) (bootstrap range [BR]: 2.2-3.1 Mya). The contemporary

effective population sizes (N_e) of *P. tremula* ($N_{P.tremula}$) and *P. tremuloides*

($N_{P.tremuloides}$) were 102,814 (BR: 93,688-105,671) and 309,500 (BR: 247,321-310,105)

respectively, with both being larger than the effective population size of their

common ancestor (N_{ANC} = 56,235 [48,012-69,492])). Gene flow ($2N_em$, where N_e is

the effective population size and m is the migration rate) from *P. tremuloides* to *P.*

tremula was higher (0.202 migrants per generation [0.156-0.375]) than in the opposite

direction (0.053 [0.052-0.117]), most likely reflecting the higher N_e in *P. tremuloides*

than in *P. tremula* (Slatkin 1985). Overall, the migration rates in both directions were

fairly low given the large N_e of both species (Morjan and Rieseberg 2004), which is

not unexpected given the large geographical distance and disjunct distributions

between the two species.

We employed the multiple sequential Markovian coalescent (MSMC) method

to investigate changes of N_e over time based on inferring the time to the first

coalescence between pairs of haplotypes (Schiffels and Durbin 2014). Higher

resolution of recent population size changes is expected when more haplotypes are

used (Schiffels and Durbin 2014). We therefore applied MSMC to phased whole-

genome sequences from one (two haplotypes), two (four haplotypes) and four (eight

haplotypes) individuals in each species, respectively. We did not include more

haplotypes because of the high computational cost of larger samples. The MSMC-

based estimates of N_e for both *P. tremula* (60,796) and *P. tremuloides* (49,701) at the

beginning of species divergence (around 2.3 Mya) were very similar to the

fastsimcoal2-based estimates of N_e for their ancestral population (Figure 2). The two

species experienced similar magnitudes of population decline following their initial

divergence (Figure 2b). Population expansion in *P. tremuloides* occurred around

50,000-70,000 years ago and continued up to the present (Figure 2b), whereas *P.*

tremula experienced a population expansion following a substantially longer periods

of bottleneck (Figure 2b).

To assess the possible confounding effects of population subdivision and

biased sampling scheme on demographic inferences in both species, we first applied

MSMC analysis using *P. tremuloides* individuals originating from populations in

Alberta and Wisconsin separately and compared them to the result obtained from the

pooled samples (Figure S3). Although the Wisconsin population was found to have

undergone a decline in population size during the last 2000-3000 years ago (Figure

S3), both local populations of *P. tremuloides* show evidence for a longer period of

species-wide expansion when compared with *P. tremula*, which was in accordance

with the results observed from the pooled samples. Second, demographic inferences

of both species were also supported by the summary statistics based on the nucleotide

diversity (θ_π) and site frequency spectrum (Tajima's *D* and Fay & Wu's *H*) (Figure

S4). The θ_π in the two subpopulations of *P. tremuloides* were all marginally higher

than in *P. tremula*, suggesting that the large effective population size found in *P.*

tremuloides is not influenced by the presence of intraspecific population subdivision

(Figure S4a). In addition, the signal of more negative values of Tajima's *D* in both

local and pooled samples of *P. tremuloides* (Figure S4b) suggest that it may have

gone through a more pronounced and/or longer period of population expansion

compared to *P. tremula*. The lower values of the genome-wide Fay &Wu's *H* in *P.*

tremula (Figure S4c), on the other hand, might reflect the relatively longer period of

low population size during the bottleneck. Taken together, these results suggest that

population subdivision of *P. tremuloides* and the unbalanced sampling schemes

between the two species have negligible effects on our demographic inferences.

Genome-wide patterns of differentiation and identification of outlier regions

against the best-fitting demographic model

To investigate patterns of interspecific genetic differentiation across the genome, we
calculated the standard measure of genetic divergence, F_{ST}, between *P. tremula* and *P.*
tremuloides over non-overlapping 10 Kbp windows (Figure 3). Levels of genetic
differentiation varied greatly throughout the genome, with the majority of windows
showing high genetic differentiation (mean $F_{ST} = 0.386$) between species (Figure 3).

In order to test the extent to which historical demographic events can explain
the observed patterns of genetic divergence between the two species, we used
coalescent simulations performed in *msms* (Ewing and Hermisson 2010) to compare
the observed patterns of differentiation to that expected under three demographic
models (Figure S5). The demographic scenario in model 1 was same as the best-
fitting model inferred by *fastsimcoal2* (Figure S5a; Table S5). In another two models,
we incorporated the population subdivision of *P. tremuloides* into the best-fitting
demographic model. In model 2 (Figure S5b; Table S5), we assume there was no gene
flow between the two subpopulations of *P. tremuloides* and explored different values
of their divergence time until the simulated F_{ST} values between the two
subpopulations matched those observed (Figure S6). The same procedure was applied
to model 3 (Figure S5c; Table S5), except that we there assume the per-generation
gene flow between the two subpopulations of *P. tremuloides* ($4N_e m$) was equal to 10
and increased their divergence time in tandem with gene flow. To assess the fit of
these models, we compared two summary statistics, θ_π and Tajima's D, between the
simulated and observed data for both species. As can be seen from Figure S7 and
Figure S8, there was generally a good agreement between observed and simulated
data sets for all three models. In addition, the above three models showed consistent
distributions of simulated F_{ST} values between *P. tremula* and *P. tremuloides*,
indicating that the presence of population subdivision in *P. tremuloides* has little
effect on the overall patterns of genomic divergence that we observe between the two
species (Figure S9).

Comparing the observed distribution of inter-specific F_{ST} with that obtained
from simulations based on the best-fitting demographic model, we found that the
observed distribution was flatter and contained greater proportions of regions falling
in the extremes of distribution (Figure 4a). We used the distribution of inter-specific
F_{ST} based on 500,000 coalescent simulations to identify outlier windows that were
likely targets of natural selection by performing a false discovery rate (FDR) of 1% as

our cut-off. With this approach, 674 and 262 windows (FDR<0.01) were, respectively,

identified as showing exceptionally high and low F_{ST} between the two species (Figure

4a). After examining the genomic distribution, physical sizes and overlaps of these

outlier windows, we found that both highly and lowly differentiated regions were

randomly distributed across the genome (Figure 3) and that the sizes of these regions

appeared to be rather small, with the majority occurring on a physical scale smaller

than 10 Kbp (Figure S10).

Signatures of selection in outlier regions

As F_{ST} is a relative measure of differentiation and is thus sensitive to any processes

that alter intra-species genetic variation (Charlesworth 1998; Cruickshank and Hahn

2014), we quantified and compared inter-specific genetic differentiation between two

unions of outlier windows and the rest of the genome using three additional

approaches: (i) pairwise nucleotide divergence between species (d_{xy}), which is a

measure that is independent of within-species diversity (Nei 1987); (ii) relative node

depth (RND) (Feder et al. 2005), which takes into account possible variation in the

mutation rate among genomic regions by dividing d_{xy} of the two aspen species with

d_{xy} between aspens and a third more distantly related species (*P. trichocarpa*); and

(iii) the proportion of inter-specific shared polymorphisms. Compared to the genomic

background average, both d_{xy} and RND revealed much greater divergence between

the two species in regions of high differentiation (Figure 4b; Table S6) and, in

accordance with these patterns, the proportion of inter-specific shared polymorphisms

was significantly lower in these regions (Figure 4b; Table S6). In addition, these

regions are characterized by multiple signatures of positive selection within one or

both species, including significantly reduced levels of polymorphism (θ_π), skewed

allele frequency spectrum towards rare alleles (more negative Tajima's D), increased

high-frequency derived alleles (more negative Fay & Wu's H), and stronger signals of

linkage disequilibrium (LD) ($P<0.001$, Mann-Whitney U test) (Figure 4c; Table S6).

Relative to genome-wide averages, these regions also contained significantly higher

proportions of fixed differences that were caused by derived alleles fixed in either

species (Figure 4c; Table S6).

In contrast to patterns found in regions of high differentiation, regions of low

differentiation showed significantly higher levels of polymorphism, excesses of

intermediate-frequency alleles (higher Tajima's D and Fay & Wu's H values), higher

proportions of inter-specific shared polymorphisms and negligible proportions of

fixed differences compared to the genomic background (Figure 4b,c; Table S6). It is

therefore likely that some of these regions have been targets of long-term balancing

selection in both species (Charlesworth 2006). Consistent with this prediction, we

found slightly lower or comparable levels of LD in these regions (Figure 4c; Table

S6), which is likely due to the long-term effects of recombination on old balanced

polymorphisms (Leffler et al. 2013). The higher d_{xy} and RND values we observe in

these regions may, however, be a consequence of the higher levels of ancestral

polymorphisms that were maintained predating the split of the two species (Figure 4b;

Table S6) (Cruickshank and Hahn 2014).

Impact of recombination rate on patterns of genetic differentiation

We examined relationships between the scaled recombination rates ($\rho=4N_e c$) and

levels of inter-species divergence over non-overlapping 10 Kbp windows across the

genome (Figure S11). We found a significant negative correlation between relative

divergence, measured as F_{ST} that depends on genetic diversity within species, and

recombination rates in both *P. tremula* (Spearman's ρ=-0.121, *P*-value<0.001) and *P.*

tremuloides (Spearman's ρ=-0.157, *P*-value<0.001) (Figure S11a). In contrast to F_{ST},

we observed significantly positive correlations between absolute divergence d_{xy} and

recombination rates in both species (*P. tremula*: Spearman's ρ=0.199, *P*-value<0.001;

P. tremuloides: Spearman's ρ=0.140, *P*-value<0.001) (Figure S11b).

Because $\rho=4N_e c$, where c is the per-generation recombination rate and N_e is

the effective population size, a reduction of N_e in regions linked to selection will

lower local estimates of ρ even if local c is identical to other regions in the genome. In

order to account for such effects and to obtain a measure of recombination that is

independent of local N_e, we compared ρ/θ_π between regions with extreme genetic

differentiation and the remainder of the genome. Relative to the genomic background,

our results showed significantly suppressed recombination in outlier regions

displaying either exceptionally high or low inter-specific differentiation (Figure 4c).

Genes under selection

The availability of the annotated *P. trichocarpa* genome enabled functional analyses
of candidate target genes within regions that were likely under selection. In total, 722
and 391 genes were located in outlier windows displaying exceptionally high and low
differentiation (Table S7 and S8), respectively. Compared to the genome overall, we
did not find significantly higher gene density in these outlier windows ($P>0.05$,
Mann-Whitney U test; Figure S12). We used the Gene Ontology (GO) assignments
of those candidate genes to assess whether specific GO terms were significantly over-
represented. After accounting for multiple comparisons, we did not detect over-
representation of any functional category among the candidate genes within regions of
high differentiation. However, we identified 60 significantly overrepresented GO
terms for genes located within regions showing significantly low genetic
differentiation and that were likely candidates for being under the influence of
balancing selection. Most of these GO categories were associated with immune and
defense responses, signal transduction or apoptosis (Table S9). Nevertheless, some
caution should be applied when interpreting these results since we observed a skewed
pattern of low coverage breadth in outlier windows displaying significantly low
differentiation compared to either the genomic background or to those highly
differentiated windows (Figure S13). Such unequal coverage breadth likely results
from the inherent technical hurdle of short-read sequencing technologies and likely
represents difficulties of mapping short reads to a reference genome in highly
polymorphic regions (Brandt et al. 2015). After stringent quality filtering, more reads
were discarded in these regions, which further decreased the amount of usable
information. Therefore, future studies, incorporating a combination of careful
experimental design and long-read sequencing technologies, are needed to evaluate
the accuracy of the evolutionary significance of balancing selection found in the
candidate genes shown here.
390

391   **Discussion**

393   We use a population genomic approach to resolve the evolutionary histories of two
394   widespread and closely related forest tree species, and to highlight how genome-wide
395   patterns of differentiation have been influenced by a variety of evolutionary
396   processes. Our simulation-based analyses indicate that *P. tremula* and *P. tremuloides*

diverged around 2.2-3.1 Mya during the Late Pliocene and/or Early Pleistocene. This timing corresponds closely with the first opening of the Bering Strait, which occurred 3.1-5.5 Mya and broke up the overland intercontinental migration route of terrestrial floras between Eurasia and North America (Marincovich and Gladenkov 1999; Gladenkov et al. 2002). This may have been less of an immediate barrier to wind-dispersed *Populus* than some other tree species, but the severing of the Bering land bridge associated with the onset of dramatic climatic oscillations throughout the Pleistocene were likely the principal drivers for initial divergence between *P. tremula* and *P. tremuloides* (Comes and Kadereit 1998; Milne and Abbott 2002; Du et al. 2015). Given the modern-day geographic isolation, disjunct distributions and extremely low rates of gene flow, our results support an allopatric model of speciation for these two aspen species (Morjan and Rieseberg 2004). The coalescent-based, intra-specific demographic analyses using MSMC demonstrate that both species have experienced substantial population expansions following long-term declines after species divergence. The population expansion of *P. tremuloides* has occurred over the last 50,000-70,000 years, following the retreat of the penultimate glaciation and has continued up to the present (Kaufman and Manley 2004). *P. tremula*, in contrast, experienced a more extended population contraction and, consistent with many other forest trees in Europe, the initiation of the population expansion in *P. tremula* coincided with the end of the Last Glacial Maximum (Hewitt 2000; Hewitt 2004).

Consistent with the expectation for allopatric speciation, where the absence of gene flow allowed for the accumulation of inter-specific differentiation due to stochastic genetic drift (Coyne and Orr 2004), we detect widespread genomic differentiation between the two species. Although neutral processes are likely responsible for the majority of the genetic differentiation we observe between the two species on a genome-wide scale (Coyne and Orr 2004; Strasburg et al. 2012), a number of regions exhibit more extreme genetic differentiation compared to expectations based on demographic simulations and thses regions also show multiple evidences of the action of natural selection (Nielsen et al. 2009). If natural selection has truly been one of the dominant evolutionary forces shaping genetic differentiation between the two species, regions of low recombination would be expected to show increased F_{ST} values, but not increased d_{xy} values (Noor and Bennett 2009; Cruickshank and Hahn 2014). This occurs because natural selection (through either selective sweeps and/or background selection) removes neutral variation over longer

distances in regions of low recombination (Begun and Aquadro 1992). As a
consequence, relative measures of divergence (e.g. F_{ST}) that rely on within-species
diversity are expected to be higher in regions with restricted recombination (Noor and
Bennett 2009; Nachman and Payseur 2012). In contrast, increased absolute
divergence (e.g. d_{xy}) is only expected if reduced gene flow occurred in regions of low
recombination (Nachman and Payseur 2012). In accordance with this view, we
observe a significant negative relationship between population-scaled recombination
rates (ρ) and F_{ST}, but not with d_{xy}, in both species (Noor and Bennett 2009; Keinan et
al. 2010). Contrary to expectations of heterogeneous gene flow, our findings highlight
a significant effect of linked selection in generating the heterogeneous differentiation
landscape we observe between the two *Populus* species (Turner and Hahn 2010;
Cruickshank and Hahn 2014; Burri et al. 2015).

Rather than being physically clustered into just a few large, discrete genomic
'islands' as expected when species diverge in the presence of gene flow (Turner et al.
2005), differentiation islands in our study system are distributed throughout the
genome, being narrowly defined and mostly located in regions with substantially
suppressed recombination. Combined with the multiple signatures of positive
selection we observe in these regions, these islands of divergence likely represent
candidate regions harboring loci closely tied to species-specific adaptation rather than
loci that are "resistant" to gene flow (Turner and Hahn 2010). In addition, we find no
functional over-representation for candidate genes located in these regions, suggesting
that a wide range of genes and functional categories have likely been involved in the
adaptive evolution of these two species (Wolf et al. 2010).

In addition to the highly differentiated regions that show signs of species-
specific positive selection, we also identify a number of lowly differentiated regions
that are candidates for being affected by balancing selection in both species
(Charlesworth 2006). Apart from low inter-specific divergence and high intra-specific
diversity, these regions also contain an excess of sites at intermediate frequencies, a
greater proportion of shared polymorphisms between the two species and lack of fixed
inter-specific differences. Genes within these regions are enriched for immune and
defense response, signal transduction and apoptosis, highlighting the influence of co-
evolutionary arms races between hosts and natural enemies on the persistence of
functional genetic diversity in immunity and defense-related genes (Tiffin and

Moeller 2006; Salvaudon et al. 2008). That said, caution is required when interpreting

the functional properties of the regions identified here, and future studies of these

candidate genes are clearly needed to better assess the adaptive genetic potential of

these widespread forest tree species to current and future climate change.

A number of factors may have influenced our demographic inferences and the

detection of natural selection in the two species. First, the presence of within-species

population subdivision could have magnified the inference of demographic expansion

in *P. tremuloides*, because pooling samples from populations of Alberta and

Wisconsin skews the SFS toward low-frequency polymorphism (more negative

Tajima's D) (Figure S4) and results in larger estimates of effective population sizes

than estimates obtained from local samples (Figure S3). However, all our analyses

suggest that this confounding effect is very weak (see Results) and the divergence

between the two subpopulations of *P. tremuloides* likely is too recent to have any

major effects on the demographic reconstruction and tests of selection in these species

(Chikhi et al. 2010). Another caveat concerns the sampling scheme used in the two

species. Local samples in *P. tremula* may not adequately reflect species-wide

demography compared to the pooled samples in *P. tremuloides*. However, the extent

to which this might influence the estimates of inter-specific F_{ST} deserves further

study. More generally, sampling should likely be more extensive in both species to

capture a greater proportion of the species-wide diversity, although local sampling is

expected to only have small effects in species with high gene flow like *Populus*

(Wakeley 2000). Finally, inter-specific hybridization in either species could

potentially bias our results. However, there are no other species of *Populus* occurring

in regions where the *P. tremula* individuals were sampled. For *P. tremuloides*,

naturally occurring hybridization is only known to occur with *P. grandidentata* in

central and eastern North America where the two species co-occur (Pregitzer and

Barnes 1980). Therefore, any possible hybridization in our study would be limited to

samples from the Wisconsin population of *P. tremuloides* but, as noted above, we did

not detect any major differences in patterns of genetic variation between the two

subpopulations, suggesting little or no effect of hybridization.

Conclusion

Here we provide insights into the speciation and recent evolutionary histories of two
closely related forest tree species, *P. tremula* and *P. tremuloides*. Our study supports
an allopatric model of speciation for the two species, which are estimated to have
diverged around 2.2-3.1 Mya as a result of the first opening of Bering Strait.
Coalescent simulations suggest that stochastic genetic drift and historical
demographic processes can largely explain the genome-wide patterns of
differentiation between species. However, there is an excess of regions displaying
extreme inter-specific genetic differentiation in the observed data compared with
demographic simulations, which is likely indicative of the action of natural selection.
In addition, we find that heterogeneous genomic divergence is strongly driven by
linked selection and variation in recombination rate in the two species. Instead of
being clustered into a few large genomic "islands" as is expected under a model of
speciation with gene flow, regions of pronounced differentiation are characterized by
multiple signatures of positive selection in one or both species, and are distributed
throughout the genome at many small, independent locations. Regions displaying
exceptionally low differentiation are likely candidate targets of long-term balancing
selection, which are strongly enriched for genes involved in immune and defense
response, signal transduction and apoptosis, suggesting a possible link to long-term
co-evolutionary arms races with pest and pathogens. Our study highlights that future
work should integrate more information on the natural histories of speciation, such as
divergence time, geographical context, magnitudes of gene flow, demographic
histories and sources of adaptation, when interpreting the meaning of observed
patterns of genomic divergence between closely related species.
520

521 **Materials and Methods**
522

Population samples, sequencing, quality control and mapping
A total of 24 individuals of *P. tremula* and 22 individuals of *P. tremuloides* were
collected and sequenced (Figure 1a and Table S1). For each individual, genomic DNA
was extracted from leaf samples and paired-end sequencing libraries with an insert
size of 600bp were constructed according to the Illumina library preparation protocol.
Sequencing was carried out on the Illumina HiSeq 2000 platform at the Science for
Life Laboratory in Stockholm, Sweden. All samples were sequenced to a target

coverage of 25$_X$. The sequencing data has been deposited in the Short Read Archive

(SRA) at NCBI under accession numbers SRP065057 and SRP065065 for samples of

P. tremula and *P. tremuloides*, respectively.

For raw sequencing reads (Wang et al. 2015), we used Trimmomatic (Lohse et

al. 2012) to remove adapter sequences and cut off bases from either the start or the

end of reads when the base quality was lower than 20. Reads were completely

discarded if there were fewer than 36 bases remaining after trimming. We then

mapped all reads to the *P. trichocarpa* reference genome (v3.0) (Tuskan et al. 2006)

with default parameters implemented in bwa-0.7.10 using the BWA-MEM algorithm

(Li unpublished data, http://arxiv.org/abs/1303.3997, last accessed May 26, 2013).

Local realignment was performed to correct for the mis-alignment of bases in regions

around insertions and/or deletions (indels) using RealignerTargetCreator and

IndelRealigner in GATK v3.2.2 (DePristo et al. 2011). In order to account for the

occurrence of PCR duplicates introduced during library construction, we used

MarkDuplicates in Picard (http://picard.sourceforge.net) to remove reads with

identical external coordinates and insert lengths. Only the read with the highest

summed base quality was kept for downstream analyses.

Filtering sites

Prior to variant and genotype calling, we employed several filtering steps to exclude

potential errors caused by paralogous or repetitive DNA sequences. First, after

investigating the empirical distribution, we removed sites showing extremely low

(<100 reads across all samples per species) or high (>1200 reads across all samples

per species) read coverage. Second, as a mapping quality score of zero is assigned for

reads that could be equally mapped to multiple genomic locations, we removed sites

containing more than 20 such reads among all samples in each species. Third, we

removed sites that overlapped with known repeat elements as identified by

RepeatMasker (Tarailo-Graovac and Chen 2009). After all filtering steps, there were

42.8% of sites across the genome left for downstream analyses. Among them, 54.9%

were found within gene boundaries, and the remainder (45.1%) was located in

intergenic regions.

SNP and genotype calling

We employed two complementary approaches for SNP and genotype calling (Figure S1): (i) Direct estimation without calling genotypes was implemented in the software ANGSD v0.602 (Korneliussen et al. 2014). Only reads with a minimal mapping quality of 30 and bases with a minimal quality score of 20 were considered. For all filtered sites in both species, we defined the alleles that were the same as those found in the *P. trichocarpa* reference genome as the ancestral allelic state. We used the -doSaf implementation to calculate the site allele frequency likelihood based on the SAMTools genotype likelihood model at all sites (Li et al. 2009), and then used the –realSFS implementation to obtain a maximum likelihood estimate of the unfolded SFS using the Expectation Maximization (EM) algorithm (Kim et al. 2011). Several population genetic statistics were then calculated based on the global SFS (Figure S1). (ii) Multi-sample SNP and genotype calling was implemented in GATK v3.2.2 with HaplotypeCaller and GenotypeGVCFs (Figure S1) (DePristo et al. 2011). A number of filtering steps were performed to reduce false positives from SNP and genotype calling: (1) Remove SNPs that were located in regions not passing all previous filtering criteria; (2) Removed SNPs with more than 2 alleles in both species; (3) Removed SNPs at or within 5bp from any indels; (4) Assigned genotypes as missing if their quality scores (GQ) were lower than 10, and then removed SNPs with more than two missing genotypes in each species; (5) Removed SNPs showing significant deviation from Hardy-Weinberg Equilibrium (*P*<0.001) in each species.

Population structure

Population genetic structure was inferred using the program NGSadmix (Skotte et al. 2013), with only sites containing lower than 10% of missing data being used. We used the SAMTools model (Li et al. 2009) in ANGSD to estimate genotype likelihoods and then generated a beagle file for the subset of the genome that was determined as being variable using a likelihood ratio test (*P*-value $<10^{-6}$) (Kim et al. 2011). We predefined the number of genetic clusters *K* from 2 to 5, and the maximum iteration of the EM algorithm was set to 10,000.

As another method to visualize the genetic relationships among individuals, we performed principal component analysis (PCA) using ngsTools accounting for sequencing errors and uncertainty in genotype calls (Fumagalli et al. 2014). The

expected covariance matrix across pairs of individuals in both species was computed

based on the genotype posterior probabilities across all filtered sites. Eigenvectors and

eigenvalues from the covariance matrix were generated with the R function eigen, and

significance levels were determined using the Tracy-Widom test as implemented in

EIGENSOFT version 4.2 (Patterson et al. 2006).

Demographic history

We inferred demographic histories associated with speciation for *P. tremula* and *P.*

tremuloides using a coalescent simulation-based method implemented in *fastsimcoal*

2.5.1 (Excoffier et al. 2013). Two-dimensional joint site frequency spectrum (2D-SFS)

was constructed from posterior probabilities of sample allele frequencies by ngsTools

(Fumagalli et al. 2014). 100,000 coalescent simulations were used for the estimation

of the expected 2D-SFS and log-likelihood for a set of demographic parameters in

each model. Global maximum likelihood estimates for each model were obtained

from 50 independent runs, with 10-40 conditional maximization algorithm cycles.

Model comparison was based on the maximum value of likelihood over the 50

independent runs using the Akaike information criterion (AIC) and Akaike's weight

of evidence (Excoffier et al. 2013). The model with the maximum Akaike's weight

value was chosen as the optimal one. We assumed a mutation rate of 2.5×10^{-9} per site

per year and a generation time of 15 years in *Populus* (Koch et al. 2000) when

converting estimates to units of years and individuals. Parameter confidence intervals

of the best model were obtained by 100 parametric bootstraps, with 50 independent

runs in each bootstrap.

We then employed multiple sequential Markovian coalescent (MSMC)

method to estimate variation of scaled population sizes (N_e) over historical time in

both species (Schiffels and Durbin 2014), which is an extension of a pairwise

sequential Markovian coalescent (PSMC) method (Li and Durbin 2011). Prior to the

analysis, all segregating sites within each species were phased and imputed using

fastPHASE v1.4.0 (Scheet and Stephens 2006). A generation time of 15 years and a

rate of 2.5×10^{-9} mutations per nucleotide per year (Koch et al. 2000) were used to

covert the scaled times and population sizes into real times and sizes.

Genome-wide patterns of differentiation

We have previously shown that linkage disequilibrium (LD) decays within 10
kilobases (Kbp) in both *P. tremula* and *P. tremuloides* (Wang et al. forthcoming), and
we thus divided the genome into 39,406 non-overlapping windows of 10 Kbp in size
to investigate patterns of genomic differentiation between species. For a window to be
included in the downstream analyses, we required there to be at least 1000 bases left
after all above filtering steps. Levels of genetic differentiation between species at each
site were estimated using method-of-moments F_{ST} estimators implemented in ngsFST
from the ngsTools package (Fumagalli et al. 2014), which calculates indices of the
expected genetic variance between and within species from posterior probabilities of
sample allele frequencies, without relying on SNP or genotype calling (Fumagalli et
al. 2013). We then averaged F_{ST} values across all sites within each 10 Kbp non-
overlapping window.
640
Coalescent simulations for detecting outlier windows
In order to examine thresholds for detection of outlier windows that may have been
targets of natural selection, we conducted coalescent simulations to compare observed
patterns of genetic differentiation (F_{ST}) to those expected under different demographic
models (see Results). All simulations were performed using the program *msms* v3.2rc
(Ewing and Hermisson 2010) based on demographic parameters derived from the
best-fitting model inferred by *fastsimcoal2.5.1* (Excoffier et al. 2013). Population-
scaled recombination rates (ρ) were assumed to be between 1 Kbp^{-1} and 5 Kbp^{-1} given
the large variation we found in both species (Wang et al. forthcoming). We simulated
genotypes corresponding to a 10 Kbp region with the same sample size as the real
data for 100,000 replications, from where we simulate genotype likelihoods using the
program msToGlf in ANGSD (Korneliussen et al. 2014) by assuming a mean
sequencing depth of 20_X and an error rate of 0.5%. We estimated two summary
statistics, nucleotide diversity (θ_π) and Tajima's D, from sample allele frequency
likelihoods in ANGSD for all simulation replicates to test whether the simulated data
matches the observed data. To assess whether any of the observed windows display
F_{ST} values deviating significantly from neutral expectations, we determined the
conditional probability (*P*-value) of observing more extreme inter-specific F_{ST} values
among simulated data sets than among the observed data. Our significance was based
on running 500,000 coalescent simulations of the most acceptable demographic null

model (see Results). We then corrected for multiple testing by using the False

Discovery Rate (FDR) adjustment, and only windows with FDR lower than 1% were

considered as candidate regions affected by selection (Storey 2002).

Molecular signatures of selection in outlier regions

To assess the occurrence of selection in outlier windows displaying either

exceptionally high or low differentiation, we compared these two unions of outlier

windows to the remaining portion of the genome by a variety of additional population

genetic statistics in both species. First, θ_π, Tajima's D (Tajima 1989) and Fay & Wu's

H (Fay and Wu 2000) were calculated from sample allele frequency likelihoods in

ANGSD over non-overlapping 10 Kbp windows. Second, levels of LD and

population-scaled recombination rates (ρ)were estimated and compared. To evaluate

levels of LD within each 10 Kbp window, the correlation coefficients (r^2) between

SNPs with pairwise distances larger than 1 Kbp were calculated using VCFtools

v0.1.12b (Danecek et al. 2011). Population-scaled recombination rates (ρ) were

estimated using the Interval program of LDhat 2.2 (McVean et al. 2004) with

1,000,000 MCMC iterations sampling every 2,000 iterations and a block penalty

parameter of five. The first 100,000 iterations of the MCMC iterations were discarded

as burn-in. Resulting estimates of r^2 and ρ were averaged over each 10 Kbp window.

In both species, windows were discarded in the estimation of r^2 and ρ if there were

less than 3 Kbp and/or 10 SNPs left from previous filtering steps. Finally, we used the

program ngsStat (Fumagalli et al. 2014) to calculate several additional measures of

genetic differentiation: (1) with *P. trichocarpa* as an outgroup, the proportion of fixed

differences that is caused by derived alleles fixed in either *P. tremula* or *P.*

tremuloides among all segregating sites; (2) the proportion of inter-species shared

polymorphisms among all segregating sites; (3) d_{xy}, which was calculated from

sample allele frequency posterior probabilities at each site and was then averaged over

each 10 Kbp window; and (4) the relative node depth (RND), which was calculated

by dividing the d_{xy} of the two aspen species with d_{xy} between aspen (represented by

24 samples of *P. tremula* in this study) and *P. trichocarpa* (24 samples; see Wang et

al. forthcoming). Significance of the differences between outlier windows and the

genome-wide averages for all above mentioned population genetic statistics were

examined using one-sided Wilcoxon ranked-sum tests.

Gene ontology (GO) enrichment

To determine whether any functional classes of genes were overrepresented among
regions that were candidates for being under selection, we performed functional
enrichment analysis of GO using Fisher's exact test by agriGO's Term Enrichment
tool (http://bioinfo.cau.edu.cn/agriGO/index.php) (Du et al. 2010). GO groups with
fewer than two outlier genes were excluded from this analysis. *P*-values of Fisher's
exact test were further corrected for multiple testing with Benjamini-Hochberg false
discovery rate (Benjamini and Hochberg 1995). GO terms with a corrected *P*-value
<0.05 were considered to be significantly enriched.

Acknowledgements

We are grateful to Rick Lindroth for providing access to the samples of *P.*
tremuloides used in this study. We thank Carin Olofsson for extracting DNA for all
samples used in this study. We thank both the editor and two anonymous referees for
their useful comments on the manuscript. The research has been funded through
grants from Vetenskapsrådet and a Young Researcher Award from Umeå University
to PKI. JW was supported by a scholarship from the Chinese Scholarship Council.
The authors also would like to acknowledge support from Science for Life
Laboratory, the National Genomics Infrastructure (NGI), and Uppmax for providing
assistance in massive parallel sequencing and computational infrastructure.

References

718
Avise JC. 2000. Phylogeography: the history and formation of species. Cambridge:
Harvard University Press.
Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and
powerful approach to multiple testing. *J R Stat Soc Series B Methodol*. 57:289-
300.
Brandt DY, Aguiar VR, Bitarello BD, Nunes K, Goudet J, Meyer D. 2015. Mapping
bias overestimates reference allele frequencies at the HLA genes in the 1000
genomes project phase I data. *G3*. 5:931-941.
Burri R, Nater A, Kawakami T, Mugal CF, Olason PI, Smeds L, Suh A, Dutoit L,
Bureš S, Garamszegi LZ, et al. 2015. Linked selection and recombination rate
variation drive the evolution of the genomic landscape of differentiation across
the speciation continuum of *Ficedula* flycatchers. *Genome Res*. 25:1656-1665.

Campagna L, Gronau I, Silveira LF, Siepel A, Lovette IJ. 2015. Distinguishing noise
from signal in patterns of genomic divergence in a highly polymorphic avian
radiation. *Mol Ecol.* 24:4238-4251.
Carneiro M, Albert F, Afonso S, Pereira R, Burbano H, Campos R, Melo-Ferreira J,
Blanco-Aguiar J, Villafuerte R, Nachman M, et al. 2014. The genomic
architecture of population divergence between subspecies of the European
Rabbit. *PLoS Genet.* 10:e1003519.
Charlesworth B. 1998. Measures of divergence between populations and the effect of
forces that reduce variability. *Mol Biol Evol.* 15:538-543.
Charlesworth D. 2006. Balancing selection and its effects on sequences in nearby
genome regions. *PLoS Genet.* 2:e64.
Chikhi L, Sousa VC, Luisi P, Goossens B, Beaumont MA. 2010. The confounding
effects of population structure, genetic diversity and the sampling scheme on
the detection and quantification of population size changes. *Genetics* 186:983-
995.
Comes HP, Kadereit JW. 1998. The effect of Quaternary climatic changes on plant
distribution and evolution. *Trends Plant Sci.* 3:432-438.
Coyne JA, Orr HA. 2004. Speciation. Sunderland: Sinauer Associates.
Cruickshank TE, Hahn MW. 2014. Reanalysis suggests that genomic islands of
speciation are due to reduced diversity, not reduced gene flow. *Mol Ecol.*
23:3133-3157.
Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE,
Lunter G, Marth GT, Sherry ST, et al. 2011. The variant call format and
VCFtools. *Bioinformatics* 27:2156-2158.
DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis
AA, Del Angel G, Rivas MA, Hanna M, et al. 2011. A framework for
variation discovery and genotyping using next-generation DNA sequencing
data. *Nat Genet.* 43:491-498.
Du S, Wang Z, Ingvarsson PK, Wang D, Wang J, Wu Z, Tembrock LR, Zhang J.
2015. Multilocus analysis of nucleotide variation and speciation in three
closely related *Populus* (Salicaceae) Species. *Mol Ecol.* 24:4994-5005.
Du Z, Zhou X, Ling Y, Zhang Z, Su Z. 2010. agriGO: a GO analysis toolkit for the
agricultural community. *Nucleic Acids Res.* 38:W64-W70.
Eckenwalder JE. 1996. Systematics and evolution of *Populus*. In Stettler RF,
Bradshaw HD, Heilman PE, Hinckley TM, editors. Biology of *Populus* and its
Implications for Management and Conservation. Ottawa: NRC Research
Press.p.7-32.
Ellegren H, Smeds L, Burri R, Olason PI, Backström N, Kawakami T, Künstner A,
Mäkinen H, Nadachowska-Brzyska K, Qvarnström A, et al. 2012. The
genomic landscape of species divergence in *Ficedula* flycatchers. *Nature*
491:756-760.
Ewing G, Hermisson J. 2010. MSMS: a coalescent simulation program including
recombination, demographic structure and selection at a single locus.
Bioinformatics 26:2064-2065.
Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa V, Foll M. 2013. Robust
demographic inference from genomic and SNP data. *PLoS Genet.* 9:e1003905.
Fay JC, Wu C-I. 2000. Hitchhiking under positive Darwinian selection. *Genetics*
155:1405-1413.

Feder JL, Xie X, Rull J, Velez S, Forbes A, Leung B, Dambroski H, Filchak KE, Aluja M. 2005. Mayr, Dobzhansky, and Bush and the complexities of sympatric speciation in *Rhagoletis*. *Proc Natl Acad Sci USA*. 102:6573-6580.

Feulner P, Chain F, Panchal M, Huang Y, Eizaguirre C, Kalbe M, Lenz T, Samonte I, Stoll M, Bornberg-Bauer E, et al. 2015. Genomics of divergence along a continuum of parapatric population differentiation. *PLoS Genet.* 11:e1004966.

Fumagalli M, Vieira FG, Korneliussen TS, Linderoth T, Huerta-Sánchez E, Albrechtsen A, Nielsen R. 2013. Quantifying population genetic differentiation from next-generation sequencing data. *Genetics* 195:979-992.

Fumagalli M, Vieira FG, Linderoth T, Nielsen R. 2014. ngsTools: methods for population genetics analyses from next-generation sequencing data. *Bioinformatics* 30:1486-1487.

Gladenkov AY, Oleinik AE, Marincovich L, Barinov KB. 2002. A refined age for the earliest opening of Bering Strait. *Palaeogeogr Palaeoclimatol Palaeoecol.* 183:321-328.

Hamzeh M, Dayanandan S. 2004. Phylogeny of *Populus* (Salicaceae) based on nucleotide sequences of chloroplast trnT-trnF region and nuclear rDNA. *Am J Bot.* 91:1398-1408.

Hewitt G. 2004. Genetic consequences of climatic oscillations in the Quaternary. *Philos Trans R Soc Lond B Biol Sci.* 359:183-195.

Hewitt G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* 405:907-913.

Kaufman D, Manley W. 2004. Quaternary glaciations—Extent and chronology Part II: North America. In Ehlers J, Gibbard PL. editors. Developments in Quaternary Science.p.1-440.

Keinan A, Reich D, Begun DJ. 2010. Human population differentiation is strongly correlated with local recombination rate. *PLoS Genet.* 6:e1000886.

Kim SY, Lohmueller KE, Albrechtsen A, Li Y, Korneliussen T, Tian G, Grarup N, Jiang T, Andersen G, Witte D, et al. 2011. Estimation of allele frequency and association mapping using next-generation sequencing data. *BMC bioinformatics* 12:231.

Koch MA, Haubold B, Mitchell-Olds T. 2000. Comparative evolutionary analysis of chalcone synthase and alcohol dehydrogenase loci in *Arabidopsis*, *Arabis*, and related genera (Brassicaceae). *Mol Biol Evol.* 17:1483-1498.

Korneliussen TS, Albrechtsen A, Nielsen R. 2014. ANGSD: analysis of next generation sequencing data. *BMC bioinformatics* 15:356.

Leffler EM, Gao Z, Pfeifer S, Ségurel L, Auton A, Venn O, Bowden R, Bontrop R, Wall JD, Sella G, et al. 2013. Multiple instances of ancient balancing selection shared between humans and chimpanzees. *Science* 339:1578-1582.

Li H, Durbin R. 2011. Inference of human population history from individual whole-genome sequences. *Nature* 475:493-496.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* 25:2078-2079.

Lohse M, Bolger A, Nagel A, Fernie AR, Lunn JE, Stitt M, Usadel B. 2012. RobiNA: a user-friendly, integrated software solution for RNA-Seq-based transcriptomics. *Nucleic Acids Res.* 40: W622-W627.

Luikart G, England PR, Tallmon D, Jordan S, Taberlet P. 2003. The power and promise of population genomics: from genotyping to genome typing. *Nat Rev Genet.* 4:981-994.

Marincovich L, Gladenkov AY. 1999. Evidence for an early opening of the Bering Strait. *Nature* 397:149-151.

McVean GA, Myers SR, Hunt S, Deloukas P, Bentley DR, Donnelly P. 2004. The fine-scale structure of recombination rate variation in the human genome. *Science* 304:581-584.

Milne RI, Abbott RJ. 2002. The origin and evolution of Tertiary relict floras. *Adv Bot Res.* 38:281-314.

Morjan CL, Rieseberg LH. 2004. How species evolve collectively: implications of gene flow and selection for the spread of advantageous alleles. *Mol Ecol.* 13:1341-1356.

Müller A, Leuschner C, Horna V, Zhang C. 2012. Photosynthetic characteristics and growth performance of closely related aspen taxa: on the systematic relatedness of the Eurasian *Populus tremula* and the North American *P. tremuloides*. *Flora* 207:87-95.

Nachman MW, Payseur BA. 2012. Recombination rate variation and speciation: theoretical predictions and empirical results from rabbits and mice. *Philos Trans R Soc Lond B Biol Sci* 367:409-421.

Neale DB, Kremer A. 2011. Forest tree genomics: growing resources and applications. *Nat Rev Genet.* 12:111-122.

Nei M. 1987. Molecular evolutionary genetics. New York: Columbia University Press.

Nielsen R, Hubisz MJ, Hellmann I, Torgerson D, Andrés AM, Albrechtsen A, Gutenkunst R, Adams MD, Cargill M, Boyko A, et al. 2009. Darwinian and demographic forces affecting human protein coding genes. *Genome Res.* 19:838-849.

Nielsen R, Korneliussen T, Albrechtsen A, Li Y, Wang J. 2011. SNP calling, genotype calling, and sample allele frequency estimation from New-Generation Sequencing data. *PloS One* 7:e37558.

Noor MA, Bennett SM. 2009. Islands of speciation or mirages in the desert? Examining the role of restricted recombination in maintaining species. *Heredity* 103:439-444.

Nosil P, Feder JL. 2012. Genomic divergence during speciation: causes and consequences. *Philos Trans R Soc Lond B Biol Sci* 367:332-342.

Nosil P, Funk DJ, Ortiz☐Barrientos D. 2009. Divergent selection and heterogeneous genomic divergence. *Mol Ecol.* 18:375-402.

Pakull B, Groppe K, Meyer M, Markussen T, Fladung M. 2009. Genetic linkage mapping in aspen (*Populus tremula* L. and *Populus tremuloides* Michx.). *Tree Genet Genomes* 5:505-515.

Patterson N, Price A, Reich D. 2006. Population structure and eigenanalysis. *PLoS Genet.* 2:e190.

Pregitzer KS, Barnes BV. 1980. Flowering phenology of *Populus tremuloides* and *P. grandidentata* and the potential for hybridization. *Can J Forest Res.* 10:218-223.

Renaut S, Grassa C, Yeaman S, Moyers B, Lai Z, Kane N, Bowers J, Burke J, Rieseberg L. 2013. Genomic islands of divergence are not affected by geography of speciation in sunflowers. *Nature Commun.* 4:1827.

Robinson KM, Delhomme N, Mähler N, Schiffthaler B, Önskog J, Albrectsen BR, Ingvarsson PK, Hvidsten TR, Jansson S, Street NR. 2014. *Populus tremula* (European aspen) shows no evidence of sexual dimorphism. *BMC Plant Biol.* 14:276.

Salvaudon L, Giraud T, Shykoff JA. 2008. Genetic diversity in natural populations: a
fundamental component of plant–microbe interactions. *Curr Opin Plant Biol.*
11:135-143.
Scheet P, Stephens M. 2006. A fast and flexible statistical model for large-scale
population genotype data: applications to inferring missing genotypes and
haplotypic phase. *Am J Hum Genet.* 78:629-644.
Schiffels S, Durbin R. 2014. Inferring human population size and separation history
from multiple genome sequences. *Nat Genet.* 46:919-925.
Seehausen O, Butlin RK, Keller I, Wagner CE, Boughman JW, Hohenlohe PA,
Peichel CL, Saetre G-P, Bank C, Brännström Å, *et al.* 2014. Genomics and the
origin of species. *Nat Rev Genet.* 15:176-192.
Skotte L, Korneliussen TS, Albrechtsen A. 2013. Estimating individual admixture
proportions from next generation sequencing data. *Genetics* 195:693-702.
Slatkin M. 1985. Gene flow in natural populations. *Annu Rev Ecol Syst.* 16:393-430.
Sousa V, Hey J. 2013. Understanding the origin of species with genome-scale data:
modelling gene flow. *Nat Rev Genet.* 14:404-414.
Storey JD. 2002. A direct approach to false discovery rates. *J R Stat Soc Series B Stat*
Methodol. 64:479-498.
Strasburg JL, Sherman NA, Wright KM, Moyle LC, Willis JH, Rieseberg LH. 2012.
What can patterns of differentiation across plant genomes tell us about
adaptation and speciation? *Philos Trans R Soc Lond B Biol Sci.* 367:364-373.
Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by
DNA polymorphism. *Genetics* 123:585-595.
Tarailo-Graovac M, Chen N. 2009. Using RepeatMasker to Identify Repetitive
Elements in Genomic Sequences. *Curr Protoc Bioinformatics* 4:1-4.10.
Tiffin P, Moeller DA. 2006. Molecular evolution of plant immune system genes.
Trends Genet. 22:662-670.
Tullus A, Rytter L, Tullus T, Weih M, Tullus H. 2012. Short-rotation forestry with
hybrid aspen (*Populus tremula* L.× *P. tremuloides* Michx.) in Northern
Europe. *Scand J Forest Res.* 27:10-29.
Turner T, Hahn M, Nuzhdin S. 2005. Genomic islands of speciation in *Anopheles*
gambiae. PLoS Biol. 3:e285.
Turner TL, Hahn MW. 2010. Genomic islands of speciation or genomic islands and
speciation? *Mol Ecol.* 19:848-850.
Tuskan G, Difazio S, Jansson S, Bohlmann J, Grigoriev I, Hellsten U, Putnam N,
Ralph S, Rombauts S, Salamov A, et al. 2006. The genome of black
cottonwood, *Populus trichocarpa* (Torr. & Gray). *Science* 313:1596-1604.
Via S. 2009. Natural selection in action during speciation. *Proc Natl Acad Sci USA.*
106:9939-9946.
Wakeley J. 2000. The effects of subdivision on the genetic divergence of populations
and species. *Evolution* 54:1092-1101.
Wang J, Scofield D, Street N, Ingvarsson P. 2015. Variant calling using NGS data in
European aspen (*Populus tremula*). In: Sablok G, Kumar S, Ueno S, Kuo J,
Varotto C, editors. Advances in the Understanding of Biological Sciences
Using Next Generation Sequencing (NGS) Approaches. Springer. p.43-61.
Wang J, Street NR, Scofield DG, Ingvarsson PK. forthcoming. Natural selection and
recombination rate variation shape nucleotide polymorphism across the
genomes of three related *Populus* species. *Genetics* doi: genetics.115.183152.
Wang Z, Du S, Dayanandan S, Wang D, Zeng Y, Zhang J. 2013. Phylogeny
reconstruction and hybrid analysis of *Populus* (salicaceae) based on nucleotide

sequences of multiple single-copy nuclear genes and plastid fragments. *PloS*
One 9:e103645.
Wolf JB, Lindell J, Backström N. 2010. Speciation genetics: current status and
evolving approaches. *Philos Trans R Soc Lond B Biol Sci.* 365:1717-1733.

Figures and Tables

Figure 1. **Geographic distribution and genetic structure of 24 *Populus tremula***
and 22 *P. tremuloides*. (a) Map showing the current geographic distribution of *P.*
tremula (red) and *P. tremuloides* (blue). Yellow circles and triangles indicate the
locations where the 24 individuals of *P. tremula* and 22 individuals of *P. tremuloides*
were sampled. (b) Genetic structure of the two species inferred using NGSadmix. The
y-axis quantifies subgroup membership, and the x-axis shows the sample ID for each
individual. (c) Principal component analysis (PCA) plot based on genetic covariance
among all individuals of *P. tremula* (red circle) and *P. tremuloides* (green square and
blue triangle). The first two principle components (PCs) are shown, with PC1
explaining 21.04% (P=2.51×10^{-19}, Tacey-Widom test) of the overall genetic variation
and separating the two species and PC2 explaining 2.09% (P=9.65×10^{-4}, Tracy-
Widom test) of the overall variation and separating samples from Wisconsin (blue
triangle) and Alberta (green square) in *P. tremuloides*. (d) Pie chart summarizing the
proportion of fixed, shared and exclusive polymorphisms of the two species.

Figure 2. **Demographic history of *Populus tremula* and *P. tremuloides*.** (a)
Simplified graphical summary of the best-fitting demographic model inferred by
fastsimcoal2. (b) Multiple sequential Markovian coalescent (MSMC) estimates of the
effective population size (N$_e$) changes for *P. tremula* (red line) and *P. tremuloides*
(blue line) based on the inference from two (dashed), four (dotted) and eight (solid)
phased haplotypes in each species. Time scale on the x-axis is calculated assuming a
neutral mutation rate per generation (μ) = 3.75×10^{-8} and generation time (g) = 15
years. The grey bar indicates the speciation time inferred by *fastsimcoal2*.

Figure 3. Genome-wide divergence.   Chromosomal distribution of genetic
differentiation (F_{ST}) between *Populus tremula* and *P. tremuloides*. The small, light
blue dots indicate F_{ST} values estimated over 10 Kbp non-overlapping windows. Grey
lines indicate F_{ST} values estimated over 100 Kbp non-overlapping windows.
Locations for windows displaying extreme differentiation relative to demographic
simulations are highlighted with colored bars above the plot. Among them, candidate
windows displaying significantly high differentiation (orange bars) are located on the

topside; candidate windows displaying significantly low differentiation (green bars)

are located at the bottom.

Figure 4. Identification of outlier windows that are candidates for being affected by

natural selection. (a) Distribution of genetic differentiation (F_{ST}) between *P. tremula*

and *P. tremuloides* from the observed (blue bar) and simulated data sets (black line).

The dashed lines indicate the thresholds for determining significantly (False

Discovery Rate <1%) high (orange bars) and low (green bars) inter-specific

differentiation based on coalescent simulations. (b) Comparisons of d_{xy}, relative node

depth (RND) and the proportion of inter-specific shared polymorphisms among

regions displaying significantly high (orange boxes) and low (green boxes)

differentiation versus the genomic background (blue boxes). (c) Comparisons of

multiple population genetic statistics, nucleotide diversity (θ_π), Tajima's D, Fay

&Wu's H, linkage disequilibrium (r^2), recombination rate (ρ/θ_π), the proportion of

fixed differences caused by derived alleles fixed in either *P. tremula* or *P. tremuloides*,

among regions displaying significantly high (orange boxes) and low differentiation

(green boxes) versus the genomic background (blue boxes). Asterisks designate

significant differences between the outlier and the rest of genomic regions by Mann-

Whitney U test (* *P*-value < 0.05; ** *P*-value < 1e-4; ****P*-value <2.2e-16).

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024 **Table 1**. Inferred demographic parameters of the best-fitting demographic model

1025 shown in Figure 2a.

1026

Parameters	Point estimation	95% CI[a]	
		Lower bound	Upper bound
N_{ANC}	56235	48012	69492
$N_{P.tremula}$	102814	93688	105671
$N_{P.tremuloides}$	309500	247321	310105
$2Nm_{P.tremuloides->P.tremula}$	0.202	0.156	0.375
$2Nm_{P.tremula->P.tremuloides}$	0.053	0.052	0.117
T_{DIV}	2332410	2186760	3113520

1027

Parameters are defined in Figure 2a. N indicates the effective population size of *P.*

tremula, *P. tremuloides* or their ancestral population, m indicates the migration rates

between species on either direction, T_{DIV} indicates the estimated divergence time

between the two species obtained from *fastsimcoal2*.

[a]Parametric bootstrap estimates obtained by parameter estimation from 100 data sets

simulated according to the overall maximum composite likelihood estimates shown in

point estimation columns. Estimations were obtained from 100,000 simulations per

likelihood.

1036

1037

(a)

ANC $N_{ANC} = 56235$

0.05

0.20

2.33MYA

P. tremula

P. tremuloides

$N_{P.tremula} = 102814$ $N_{P.tremuloides} = 309500$

(b)

Effective population size

P.tremula   *P.tremuloides*
- - - 2 haplotypes   - - - 2 haplotypes
······ 4 haplotypes   ······ 4 haplotypes
——— 8 haplotypes   ——— 8 haplotypes

Time (years ago)

F_{ST}