# Adaptation to heavy-metal contaminated environments proceeds via selection on pre-existing genetic variation.

**Kevin M. Wright 1\*, Uffe Hellsten 2, Chenling Xu 3, Annie L. Jeong 4,5, Avinash Sreedasyam 6, Jarrod A. Chapman 2, Jeremy Schmutz 6, Graham Coop 3, Daniel S. Rokhsar 2,7, John H. Willis 4**

1.    Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA.
2.    Department of Energy Joint Genome Institute, Walnut Creek, CA 94598, USA.
3.    Department of Evolution and Ecology, University of California, Davis, CA, 95616, USA.
4.    Department of Biology, Duke University, Durham, NC 27708, USA.
5.    University Program in Genetics and Genomics, Duke University Medical Center, Durham, North Carolina, USA
6.    Hudson Alpha Institute for Biotechnology, Huntsville, Alabama 35806, USA.
7.    Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA

\* Corresponding Author: wright@fas.harvard.edu

**Across a species range, islands of stressful habitats impose similar selection pressures on isolated populations. It is as yet unclear, when populations respond to these selective pressures, the extent to which this results in convergent genetic evolution and whether convergence is due to independent mutations or shared ancestral variation. We address these questions investigating a classic example of adaptation by natural selection - the colonization of plant species to heavy metal contaminated soils. We use field-based reciprocal transplant experiments to demonstrate that mine alleles at a major copper tolerance QTL, *Tol1*, are strongly selected in the mine environment, but are neutral, or nearly so, in the off-mine environment. To identify scaffolds in genetic linkage with this locus, we assemble the genome of a mine adapted *M. guttatus* genotype and sequence near isogenic lines (NILs) homozygous for tolerant or non-tolerant alleles at *Tol1*. We identify genes with differential expression between NILs and differences in allele frequency between independent pairs of mine and off-mine populations to identify *Tol1* candidate genes. We identify a single gene, a multicopper oxidase, with large differences in expression between NILs and allele frequency between populations. Furthermore, we find patterns of genetic variation at *Tol1*, and four additional candidate adaptation loci, are consistent with selection acting upon beneficial haplotypes that predates the existence of the copper mine habitat. We estimate the age of selected *Tol1* haplotype to be at least 1700 years old and was at a frequency of 0.4-0.6% in the ancestral population when mining was initiated 150 years ago. These results suggest that adaptation to the mine habitat routinely occurs via selection on ancestral variation, rather than independent *de-novo* mutations or migration between populations.**

The most spectacular demonstrations of evolution often involve independent populations repeatedly evolving similar traits in response to similar environmental pressures (Endler 1977; Joron and Mallet 1998; Schluter, 2000). Convergent evolution of similar phenotypes suggests convergence in genotype, as may occur when selection operates on the same ancestral allele in independent populations or species (Colosimo *et al.* 2005; The Heliconius Genome Consortium, 2012). Additionally, there are multiple instances of the similar phenotypes evolving via independent genetic mutations at the same locus (Tishkoff *et al.* 2007, Karasov *et al.* 2010). When evolution occurs via selection on pre-existing genetic variation, adaptation to novel environments will proceed quite quickly. Conversely, when evolution proceeds via independent genetic mutations, adaptation will be limited by the waiting time for beneficial mutations to occur and go to high frequency. In order to understand the circumstances under which either of these evolutionary scenarios is likely to occur, and make predictions about how species will respond to rapid environmental changes such as global climate change, we investigate the genomic basis of a classic example of convergent phenotypic evolution.

Investigations into the repeated colonization of stressful soil environments by plants species have a long history in the development of evolutionary biology (Kruckeberg 1951; Turner 1969; Antonovics *et al.* 1971; Macnair 1987). With the initiation of the first, and still on-going, field experiments at Park Grass, UK,

scientists demonstrated that soil nutrient levels and pH impose strong selection on the local plant communities causing some plant species to rapidly evolve tolerance, whereas other species, despite ample migration opportunities, never colonize these stressful habitats (Brenchley 1958). This result has been replicated in investigations of plants colonizing heavy metal contaminated soils (Walley *et al.* 1974, Bradshaw 1984, Symeonidis *et al.* 1985; Al-Hiyaly *et al.* 1990, 1993). Interestingly, the only species to colonize contaminated soils also have tolerant individuals in populations unexposed to soil contamination. Whereas species that did not evolved tolerance, lack tolerant individuals in unexposed populations (*summarized in* Bradshaw, 1991). Given these results, Bradshaw concluded that tolerant species possessed the necessary genetic variation in ancestral populations, and species which remain non-tolerant lack the necessary genetic variation to adapt to novel environmental pressures (1991). However these results do not eliminate the possibility that tolerance was absent in the ancestral population and spread, by migration, to uncontaminated sites following adaptation. To distinguish between these two hypotheses, we need additional evidence on the source of adaptive genetic variation in natural populations.

Here, we investigate the recent and repeated adaptation of the annual wildflower, *Mimulus guttatus*, to copper contaminated sites near Copperopolis, CA (Macnair *et al.* 1993). Contaminated sites have 10-100 fold increase in the concentration of copper in the soil compared to uncontaminated sites (Allan and Sheppard 1971; Macnair et al. 1993; S. Table 1). Copper mining in this region of the Sierra Nevada foothills started in 1861 (Aubury 1908). Mine populations are located in close geographic proximity, within 40km of each other, and *M. guttatus* is very common in uncontaminated soils adjacent to mine sites (Allan and Sheppard 1973; Macnair *et al.* 1993). Copper tolerance is nearly fixed at four mine populations (99.77%, N =2796); at intermediate frequency (56.0%; N=1121) at eleven sites, mostly located within one mile of contaminated soil, and at low frequency (2.2%; N=1056;) in 11 populations located 1-10 miles from contaminated soil (Macnair *et al.* 1993). This suggests that the copper tolerance phenotype, measured as root growth in a hydroponic solution, is under strong selection in contaminated habitats. It remains to be determined whether each mine was colonized independently and whether the low frequency of tolerance at uncontaminated sites represents ancestral variation or back migration from the mines. Previously, we identified a single large effect QTL for copper tolerance, *Tol1*, in line from the Copperopolis mine population (Wright *et al.* 2013). It is unknown whether this same locus contributes to copper tolerance in other mine adapted populations. We use field based transplant experiments, population genomic sequencing, and coalescence modeling to address three key questions: Does *Tol1* affect fitness in mine habitat, and if so, is it the primary determinate of survival in this habitat? Did convergent phenotypic evolution in independent mine populations occur via convergent genetic evolution? Did mine alleles at the candidate adaptation loci evolve from new beneficial mutations or via selection on pre-existing standing variation?

**Tol1 affects fitness in copper mine habitat**

We directly measured whether the mine and off-mine genotypes are locally adapted
to their respective habitats using a reciprocal transplant experiment across multiple
years (S. Figure 1). We found that plants from the mine population have greater
fitness than off-mine genotypes in mine habitat, supporting the hypothesis of strong
selection in this environment (Figure 1; S. Table 2-9). In the off-mine habitat, mine
genotypes were also slightly favored compared to off-mine genotypes (Figure 1; S.
Table 2-9). This suggests that high fitness in the mine environment does not cause a
detectable trade-off in fitness in the off-mine environment. In order to measure
whether the copper tolerance locus affects fitness, we conducted a reciprocal
transplant experiment with introgression lines possessing alternate alleles at *Tol1*.
We constructed introgression lines by reciprocally backcrossing lines with mine and
off-mine alleles at the tolerance locus (*Tol1*$_{CC}$ or *Tol1*$_{MM}$) to mine (CCC$_{BC}$) and off-
mine (MED$_{BC}$) genotypes (S. Figure 2). In the off-mine habitat, we found no
significant effect of the *Tol1* locus in our CCC$_{BC}$ introgression lines, (95% CI: 0.14, -
0.36, $p$ = 0.68, S. Table 6 ) which suggests that tolerant alleles at *Tol1* are neutral (or
only weakly selected) in off-mine populations. Conversely, we found that mine
alleles at *Tol1* greatly increases the probability of survival to flowering in the mine
habitat (Figure 1; S. Table 2). The strength of selection (s) on *Tol1*, defined as the
relative fitness of each allele, was significantly stronger in the off-mine genomic
background (MED$_{BC}$ s=0.69) compared to the mine genomic background (CCC$_{BC}$
s=0.30). Additionally, the difference in fitness between parental lines, (s=0.90; S.
Table 2) is greater than the fitness effect of *Tol1*. These two results support the
hypothesis that additional loci contribute to fitness in the mine habitat. Our finding
that the *Tol1* mine allele does not impose a measureable fitness cost in the off-mine
habitat is consistent with either back-migration or ancestral variation as possible
scenarios explaining the low frequency of copper tolerant plants in this
environment. To distinguish between these two scenarios we need to measure
segregating variants at *Tol1* in mine and off-mine populations.
141
**Targeted assembly of *Tol1* region**
To accurately measure genetic variation at *Tol1*, we need to first improve the
assembly of this region. *Tol1* is located in a poorly assembled pericentromeric
region of linkage group 9 in the M *guttatus* reference assembly, notably
characterized by large amounts repetitive DNA and low recombination rates
(Wright *et. al.* 2013). We sought to map all genomic scaffolds linked to *Tol1* by
resequencing plants derived from our original Near Isogenic Line (NIL) mapping
population (Wright *et al.* 2013). To accomplish this, we sequenced two pools of
plants each composed of 100 plants homozygous for either tolerant (NIL_*Tol1*$_{TT}$) or
nontolerant (NIL_*Tol1*$_{NTNT}$) alleles at the genetic marker, *Sc84_37kb*, located 0.32 cM
from *Tol1*  (Wright *et al.* 2013). We reason that that scaffolds in this region should,
except in the case of a rare recombination event, be fixed for different haplotypes in
the NIL_*Tol1*$_{TT}$ and NIL_*Tol1*$_{NTNT}$ pools. We estimate the recombination distance
between genomic scaffolds and the *Sc84_37kb* marker as the frequency of SNPs in
the two pools to used to define the two pools (S. Text). We identified eight scaffolds,
comprising 4.9 Mb, within a somewhat enlarged search space of 0.80 cM (S. Figure
3; S. Table 10). We also identified small portions of other scaffolds that are in tight

linkage to *Tol1* (data not shown), suggesting problems with the v1.1 reference
assembly and/or translocations between the reference and mapping lines. To
address these assembly issues we sequenced a Copperopolis inbred line, CCC52, to
220X coverage and constructed a *de novo* genome assembly (S. Text, S. Table 11).
We repeated our mapping effort using this *de novo* assembly and identified 53
CCC52 scaffolds, totaling 3.1Mb in length, within 0.80 cM of the *Sc84_37kb* marker
(S. Text, S. Table 12). While most of the scaffolds are homologous to reference
assembly *Tol1* linked scaffolds, eight of them are unique to the CCC52 assembly.
Using this improved assembly, we sought to identify genes that exhibit significant
differences in expression between NIL_*Tol1*$_{TT}$ and NIL_*Tol1*$_{NTNT}$ lines.
169
**Differential expression of *Tol1* link genes**
We measured gene expression in the NIL_*Tol1*$_{TT}$ and NIL_*Tol1*$_{NTNT}$ lines  grown in a
full-strength Hoaglands solution and Hoaglands enriched with 1.25 ppm CuSO4. We
extracted mRNA from root tissue, sequenced each mRNA library to 140X coverage,
and aligned all reads to a *de novo* transcriptome assembly (S. Text). We find that the
majority of differentially expressed genes are due to allelic variation between the
NIL_*Tol1*$_{TT}$ and NIL_*Tol1*$_{NTNT}$ lines, the copper treatment had negligible effect on
gene expression. We identified 277 genes that are differentially expressed (DE)
(Benjamini & Hockberg adjusted p-value < 0.1 and log$_2$ fold change > 1.0) between
lines in either treatment, 37 of these genes are located on the 53 *Tol1* linked
scaffolds (S. Figure 4, S. Table 13).  Some *Tol1* link DE genes - including a heavy
metal ion transporter, a potassium ion transporter, and a multicopper oxidase –
function to maintain intracellular ion homeostasis and may underlie the copper
tolerance phenotype. Next, we sought to determine whether *Tol1* linked DE genes
exhibit large differences in allele frequency in mine/off-mine population
comparison.
186
**Identification of candidate adaptation loci at *Tol1***
We sought to identify loci at *Tol1* that may underlie adaptation to the copper mine
environment by measuring the genetic variation within and between two mine and
two offmine populations. In a classic hard selective sweep, selection drives a single
new beneficial mutation to near fixation, causing linked sites to hitchhike to high
frequency, leading to a local reduction in genetic variation and large differences in
allele frequency between the ancestral and derived populations (Maynard Smith &
Haigh 1974, Barton 2000, Berg and Coop 2015). Alternatively, selection on pre-
existing adaptive variants, which reside on multiple haplotypes, is predicted to
produce a much smaller change in allele frequency at surrounding sites because no
single haplotype goes to fixation (Innan and Kim 2004, 2008; Prezeworski *et al.*
2005). This event is sometimes termed a soft selective sweep (Messer and Petrov
2013). By comparing independent populations that have both adapted to similar
environments, we can distinguish between selection on newly derived beneficial
mutations or shared ancestral variation.
202
We sequenced pooled DNA from 20-31 natural isolates from two mine and two off-
mine populations to 34-72X genome-wide coverage (S. Text, S. Table 1, S. Figure 5).

Genome wide nucleotide diversity is similar across all populations, ($\pi$ = 0.0254 –
0.0268 ; S. Table 14). Genome-wide estimates of genetic differentiation between
populations, measured with $F_{st}$ and the $G$ statistic (accounts for difference in
coverage among sites and populations, S. Text) are higher between mine/off-mine
compared to off-mine/off-mine population comparisons ($F_{st}$ M/OM= 0.07-0.14; $F_{st}$
OM/OM=0.02; S. Table 15). Elevated genome-wide estimates of differentiation
between mine/off-mine populations is caused by a few, large effect genomic regions
(ie. S. Figure 6) whereas the majority of the genome shows little differentiation.
These results suggest that the mine populations did not undergo a strong bottleneck
and/or gene flow readily occurs between populations in contrasting habitats.
215
We identified genomic regions with elevated difference in allele frequency at *Tol1*,
as well as at other sites across the genome. Differentiation at *Tol1* is not elevated
across all scaffolds in tight linkage to this locus (S. Figure 7, 8), as would be expected
from a single hard selective sweep (S. Text), instead we find multiple loci with
nearly fixed differences in allele frequency between mine/off-mine populations
surrounded by large regions of shared genetic variation. We focus on loci that
exceed the 0.5% genome-wide threshold of allelic differentiation in both mine/off-
mine population comparisons (allelic differentiation is measured using the $G$
statistic in 500bp bins, S. Text). We identify five distinct candidate adaptation loci,
harboring seven genes, within the 53 *Tol1* link CCC52 scaffolds (S. Table 16). The
signal of differentiation at these loci is relatively narrow (<20kb) with no significant
differentiation at physically adjacent regions. To identify which, if any, of these
genes maybe a *Tol1* candidate, we compared these genes to the 37 differential
expression genes from the RNA-seq experiment. We find, one gene, a multicopper
oxidase (MCO) is significantly differentiated in both experiments. MCO is located on
a small scaffold (*s47895 - 1.7kb*) that is absent from the reference genome.
Furthermore, we find that mine populations exhibit a 6X fold enrichment of aligned
reads at MCO (S. Table 17), suggesting a recent tandem duplication of this gene is
responsible for the 12X increase in expression observed in copper tolerant lines (S.
Table 13). This gene represents an interesting *Tol1* candidate, but additional
functional studies are required to determine whether it is responsible for the copper
tolerance phenotype. Building upon the results from our transplant experiment, that
*Tol1* accounts for some but not all of the variation in fitness in the mine
environment (Figure 1), we next sought to identify additional loci with a population
genetic signature of selection.
241
**Genome-wide survey of candidate adaptation loci**
We identified multiple genomic regions with elevated levels of differentiation
between mine and offmine populations. As with *Tol1*, we focused on candidate
adaptation loci that exceed the 0.5% genome-wide $G$ statistic threshold in mine/off-
mine population comparisons. Some loci are unique to one mine/off-mine
population comparison (S. Table 18, 19), however, loci with the largest difference in
allele frequency, 0.5% and 0.1% outliers, are much more likely to be shared
between the mine/off-mine population comparisons than would be expected from
random chance (S. Table 20; S. Figure 9). We rank the loci by their degree

differentiation (the quantity of physically adjacent 0.5% outlier bins) in both
mine/off-mine population comparisons and find that *Tol1*, although it has a large
effect on fitness, ranks as the fifth most differentiated region of the genome (Table
1). The identify of the top candidate adaptation loci in our screen are largely robust
to changes in the genome-wide $G$ statistic threshold (1%, 0.5%, 0.1%; S. Table 21),
clustering of adjacent outlier bins (5kb, 10kb, 100kb; S. Table 22), the minimum
number of SNPs per bin and measures of population differentiation (5 or 10 SNPs; $G$
or $F_{st}$; S. Table 23). Nearly all candidate adaptation loci are restricted to single genes
which encode proteins that function to maintain intracellular ion homeostasis
(copper ion transporter, iron-zinc ion transporter, transmembrane ferric reductase,
and potassium ion transporter;Table 1) and, in some cases, they have been
implicated in heavy-metal tolerance (ie. Hanikenne *et al.* 2008 ; Grennan *et al.*
2009). The young age of mine populations and narrow interval of divergence
(<30kb) suggests that beneficial variants pre-date the establishment of mine
populations.
266
**Selection on pre-existing genetic variation**
The narrow interval of genetic differentiation at each candidate adaptation locus is
inconsistent with expectations from a hard selective sweep. Given the intense
selection on *Tol1* (s=0.69, Figure 1), a hard selective sweep would cause the mine
allele to reach near fixation in ~20 generations and reduce genetic diversity by
>50% for ~3Mb from the selected site (S. Text). In contrast, we observe that for *Tol1*
outlier loci, heterozygosity recovers to 50% of background within a maximum of
20kb (S. Table 22). This quick recovery, despite strong selection, strongly suggests
that the selected allele at *Tol1* was segregating in the ancestral population prior to
the establishment of the mine populations. If this is true, we predict that each mine
population would share the same haplotype at the candidate loci. In order to assess
this hypothesis for the *Tol1* region, and all other candidate adaptation loci, we
measured genetic differentiation between mine populations.
280
We find that three loci in tight linkage to *Tol1* and four additional unlinked
candidate adaptation loci exhibit regions of very low genetic differentiation between
mine populations (Table 2; i.e. Figure 2, S. Figure 9-15). This is inconsistent with
independent sweeps of new mutations, but rather with selection on shared standing
variation. Mine populations share the same *core haplotype* at these loci (extending
from 1-18kb; Table 2) and outside this core, genetic differentiation between mine
populations rises dramatically to levels comparable to differentiation between mine
and off-mine populations (Figure 2). This peak-valley-peak pattern is additional
evidence that the *core haplotype* was present in the ancestral population and
recombined onto multiple genomic backgrounds prior to the establishment of the
copper mines (Roesti *et al.* 2014). How old must the *core haplotypes* be to have an
opportunity to recombine onto different genomic backgrounds prior to the
initiation of mining in the region?
294
**Age of core haplotypes**

We estimate the age of the *core haplotypes* using the rate at which genetic diversity
around the mine allele recovers to background levels (S. Text). We find that all *core*
*haplotypes* predate the establishment of the copper mines (Table 2). Not
surprisingly, the locus with the strongest signature of selection, MET, has the
youngest *core haplotype* (639 generations), but even this haplotype existed in the
ancestral Copperopolis population long before the establishment of the mine (150
years; Table 2). In contrast to the four unlinked *core haplotype* regions, we find
three tightly linked *core haplotypes* at *Tol1*. This suggests these haplotypes were in
linkage disequilibrium (LD) in the ancestral population, a hypothesis consistent
with the result that each scaffold maps to <0.05 cM of the same position (S. Table
12), and LD was maintained as they rose to high frequency in each mine population.
We estimate the age of the *core haplotypes* at the three *Tol1* linked genes to predate
the establishment of the copper mines by hundreds-thousands of years (Table 2).
The CDK and PUM haplotypes have been present in the ancestral Copperopolis
population for 1.7-2.7K generations, whereas the MCO haplotype is much older
(12.5K generations). Using the age of these haplotypes and the strength of selection
on *Tol1*, we calculate that the CDK and PUM haplotypes were at 0.4-0.6% and the
MCO at 4.5% frequency when the copper mines were first established (S. Table 24).
These values are inflated for MCO because it is on a small scaffold with few
informative SNPs (S. Figure 15), thus we consider the CDK and PUM estimates to
better reflect the age and frequency of the selected allele at *Tol1*.
317
**Conclusion**
In this study, we investigate a classic example of rapid adaptation, colonization of
heavy metal contaminated environments. We investigate the genetic control of a
major copper tolerance QTL, *Tol1*. This locus has a significant effect on fitness in the
mine environment (s = 0.69). We improve the assembly of the pericentromeric
region harboring *Tol1*. Next we use whole genome RNA expression measurements
and population genomic sequencing to identify candidate genes at this locus. We
identify a single gene, a multicopper oxidase, with large differences in expression
between tolerant and nontolerant lines and allele frequency between mine and off-
mine populations. Interestingly, three tightly linked genes at *Tol1*, including MCO,
possess *core haplotypes* that are strongly differentiated from the off-mine
populations and highly similar between mine populations. This pattern suggests
these *core haplotypes* predates the establishment of the mine populations. We
develop a novel method to estimate the age of these haplotypes and find that they
predate the establishment of the copper mines by 1000s of generations.
Furthermore, we estimate the frequency of this haplotype in the ancestral
population at the onset of mining (< 1%) is consistent with estimates of the
frequency of tolerant plants in current populations inhabiting uncontaminated sites
(Macnair *et al.* 1993). Looking throughout the genome we find that the top four
candidate adaptation loci exhibit a *core haplotype* pattern similar to what we
observe at *Tol1*. The function of the majority of these genes, to maintain intracellular
ion homeostasis, suggests they contributed to adaptation in these copper
contaminated environments. This work combining field-based estimates of
selection, population genomics, and theoretical analyses provides thorough support

for the classic hypothesis that rapid adaptation in natural populations primarily
proceeds via selection on standing genetic variation (Bradshaw 1991).
344
345

345    **Figures**
346
**Figure 1 (A)** Mean probability of survive to flowering for all genotypes in mine plots
in 2012 reciprocal transplant experiment (error bars = standard deviation). The
probability a plant survives to flower, and the p-values testing whether there is a
significant effect of genotype, is calculated from the predicted values using a logistic
regression. The different genotypes are: parental mine, parental off-mine, F1 hybrid,
and backcross lines (BC). Backcross lines are homozygous for Copperopolis mine
allele at *Tol1* (red bars) or MED off-mine allele at *Tol1*, blue bars. **(B)** Mine plot,
replicate 1. **(C)** Survival to flowering for all genotypes in offmine plots in 2012
reciprocal transplant experiment. Due to low germination, we have no data for the
BC MED with off-mine allele at *Tol1*. **(D)** Off-mine plot, replicate 1.
357
**Figure 2** Genetic differentiation at top ranked candidate adaptation locus on
*Scaffold8*. (**A**) Genetic differentiation measured with G statistic for individual SNPs .
Arrows depict annotated genes from *M. guttatus* genome V1.1. – Black arrows are
genes within *core haplotype* **(B)** ratio: $\pi_{Mine}/ \pi_{Off-mine}$ measured in 500bp windows.
Orange dots/lines: Mine1 vs. Off-mine1. Blue dots/lines: Mine2 vs. Off-mine2. Green
dots in **(A)**: Mine1 vs. Mine2. Green line in **(B)** ratio of pairwise $\pi$: $\pi_{Mine1+Mine2}/ \pi_{Mine-}$
$_{Off-mine}$. Grey dashed line indicates genomic intervals with no SNP data.
365
**Supplemental Figure 1** Local adaptation of parental genotypes in the mine habitat.
A. Experimental plot 1 at the copper mine, soil Cu is 142ppm. B. Experimental off-
mine plot 1, soil Cu is 12.9ppm. C. Copperopolis parental line flowering on May12,
2012. D. Probability of survival to flowering of parental genotypes in two years of
reciprocal transplant experiments. Mine genotypes orange=2007; red=2012. Off-
mine genotypes: purple=2007; blue=2012. E. Map of experimental plots at Napoleon
Mine, 15 km from Copperopolis, CA. Orange shaded region is approximate outline of
mine. Orange diamonds are two mine plots, and blue triangles are two off-mine
plots.
375
**Supplemental Figure 2** Generation of outbred BC lines of known genotype at *Tol1*
locus.  Backcrossing to Copperopolis parental lines, CCC, proceeded for four
generations and backcross to Off-mine parental line, MED, proceeded for one
generation. Afterwards heterozygous *Tol1* plants were identified with *Tol1* linked
markers and selfed to produce plants homozygous for alternate alleles at *Tol1*.
Independent backcross lines were then intercrossed to produce outbred plants of
known genotype at *Tol1* locus.
383
**Supplemental Figure 3** Assembly of *Tol1* region.
Plot of the recombination distance between a focal gene and the *Sc84_37kb* marker
(indicated with blue arrow). Recombination distance estimated using all
informative SNPs within focal gene (S. Text). *Tol1* linkage threshold (recombination
distance of 0.008, or 0.8cM) is denoted with blue line. Black arrows at top indicate
approximate location of v1.1 scaffolds.
390

**Supplemental Figure 4** Differentially expressed transcripts in tolerant and non-
tolerant lines.
Plot of log2 fold changes over mean expression. Blue horizontal lines represent fold
change in expression of 1.0. Red dots indicate transcripts that have adjusted p-value
less than 0.1. Triangles indicate transcripts that exceed 10 or -10 log2 fold change in
expression. **A.** Difference in expression between NIL_$Tol1_{TT}$ and NIL_$Tol1_{NTNT}$ lines
in Hoaglands + Cu treatment. We identified 93 and 107 transcripts down- and up-
regulated respectively at an FDR of 0.1. **B.** Hoaglands treatment. We identified 119
and 105 transcripts to be down- and up-regulated respectively at an FDR of 0.1.
400
401
402 **Supplemental Figure 5** Map of Copperopolis, Calaverous Cty, CA, USA.
403 Red circles are mine populations and blue circles denote off-mine populations.
404 Population key: CCC – Copperopolis mine; MCN - McNulty mine; OM1 –Off-Mine1/
405 O'Byrnes Ferry Rd.; and OM2 Off-mine2/ Stage Coach Rd.
406
407 **Supplemental Figure 6** Scaffold wide plots of allelic divergence between
408 Copperopolis and Off-mine 1.
409 Plot of rank $G$ statistic between Copperopolis and Off-mine 1 for scaffold 8. Each dot
410 is 500bp bin - minimum 10 SNPs per bin- sliding window step size of 100bp. Black
411 line is 0.5% genome wide threshold used to identify outlier bins.
412
413
**Supplemental Figure 7** Genetic Differentiation at *Tol1* Linked Reference Genome
Scaffolds
Order of scaffolds (reference genome v1.1) is same for S. Figure 3. Scaffold wide
plots of allelic divergence, genome wide rank $G$ statistic, between (A) Copperopolis /
Off-mine 1, orange dots and (B) McNulty Mine / Off-mine 2, blue dots. Each dot is
500bp bin - minimum 10 SNPs per bin- sliding window step size of 100bp. Black line
is 0.5% genome wide threshold used to identify outlier bins.
421
422 **Supplemental Figure 8** Genetic Differentiation at *Tol1* Linked CCC52 Genome
423 Scaffolds.
424 Population comparisons same as presented in S. Figure 7. Scaffolds are in numerical
425 order from 1-53 according to name ID in S Table 12.
426
**Supplemental Figure 9** Plot of allelic divergence between Copperopolis-Off-mine1
and McNulty-Off-mine2.
Candidate adaptation loci exceed 0.1% (dark blue), 0.5% (blue), 1.0% (light blue)
rank G statistic threshold in both mine/off-mine comparisons. Allelic divergence
measured as average G score for SNPs within 500bp window, each dot is a single
500bp bin with data for both population comparisons.
433
**Supplemental Figure 10** Allelic divergence for candidate region: rank 2, scaffolds
148 and 198. These two scaffolds are adjacent in M. guttatus genome V. 2.0. (**A**)
Genetic differentiation measured with G statistic for individual SNPs . Arrows depict

annotated genes from *M. guttatus* genome V1.1. – Black arrows are genes within
*core haplotype* **(B)** ratio: $\pi_{Mine}$/ $\pi_{Off\text{-}mine}$ measured in 500bp windows. Orange
dots/lines: Mine1 vs. Off-mine1. Blue dots/lines: Mine2 vs. Off-mine2. Green dots in
**(A)**: Mine1 vs. Mine2. Green line in **(B)** ratio of pairwise $\pi$: $\pi_{Mine1+Mine2}$/ $\pi_{Mine\text{-}Off\text{-}mine}$ .
Grey dashed line indicates genomic intervals with no SNP data.
442
443  **Supplemental Figure 11** Allelic divergence for candidate region: rank 3, scaffold 4.
444  Figure details are provided in legend of S Figure 9.
445
446  **Supplemental Figure 12** Allelic divergence for candidate region: rank 4, scaffold 1.
447  Figure details are provided in legend of S Figure 9.
448
449  **Supplemental Figure 13** Allelic divergence for *Tol1* link candidate locus CDK,
450  scaffold 7467, CCC52 genome. Figure details are provided in legend of S Figure 9.
451
452  **Supplemental Figure 14** Allelic divergence for *Tol1* link candidate locus PUM,
453  scaffold 10201, CCC52 genome. Figure details are provided in legend of S Figure 9.
454
455  **Supplemental Figure 15** Allelic divergence for *Tol1* link candidate locus MCO,
456  scaffold 47895, CCC52 genome. Figure details are provided in legend of S Figure 9.

461
462
463 **Tables**
464
465 **Table 1. Top ranked candidate adaptation loci**

| Rank | Scaffold | Length (kb) | N - Genes | N - Bins Exceed 0.5% | N - SNPs Fixed Diff. | Locus ID | Annotation |
|---|---|---|---|---|---|---|---|
| 1 | 8 | 32.9 | 10 | 97 | 70\65 | MET | Cytochrome P450; Methytransferase; Protease Inhibitor |
| 2 | 148/198 | 24.1 | 2 | 39 | 0\32 | FER | Transmembrane Ferric Reductase |
| 3 | 4 | 7.7 | 2 | 25 | 0\0 | ZNC | Zinc/Iron Ion Transporter |
| 4 | 1 | 8.1 | 1 | 19 | 0\5 | COP | Copper Ion Transporter |
| 5A | 84 | 1.8 | 1 | 11 | 0\0 | MND | **Tol1 Link** - Mandelate Racemase |
| 5B | 84 | 5.8 | 1 | 8 | 3\0 | CDK | **Tol1 Link** - Cyclin Dependent Kinase |
| 6 | 44 | 2.5 | 1 | 9 | 0\0 | SHD | Protein Binding, SH3 domain |
| 7 | 80 | 6.2 | 1 | 6 | 0\0 | DUF | Unknown Function |
| 8 | 115 | 0.9 | 1 | 5 | 0\0 | KIT | Potassium Ion Transporter |
| 9 | 47 | 2.1 | 0 | 5 | 0\0 | - | Intergenic |
| 10 | 51 | 0.8 | 1 | 4 | 0\0 | HSP | Heat Shock Protein, HSP90. |

466
467
468
Candidate adaptation loci are composed on 500bp bins that exceed 0.5% rank G
statistic threshold in both mine/off-mine population comparisons. Outlier bins are
grouped together using the 10kb clustering algorithm. Loci are ranked by the
number of outlier bins at a locus. The physical length of a candidate locus is defined
by the region of overlap between both mine-off-mine population comparisons. The
FER locus spans two adjacent scaffolds: 148 and 198. The number of SNPs with fixed
differences is presented for CCC vs. Off-mine1 \ MCN vs. Off-mine2 comparisons.
Gene annotation from *M. guttatus* genome v1.1, phytozome.net. Annotation of loci
that span multiple genes is restricted to genes with strongest signal of
differentiation.
479
480
481
482

**Table 2 –Age of core haplotypes at candidate adaptation loci.**

| Locus ID | Core Haplotype | | | Coalescence Time | | | | Annotation |
|---|---|---|---|---|---|---|---|---|
| | Scaffold | Length (kb) | Num. Genes | CCC | MCN | CCC/ OM1 | MCN/ OM2 | |
| MET | 8 | 19.6 | 4 | 639 | 1191 | 1901 | 5699 | Cytochrome P450; Methytransferase; Protease Inhibitor; Unknown Func. |
| FER | 148/198 | 7.06 | 1 | 1533 | 2022 | 2285 | 1725 | Transmembrane Ferric Reductase |
| ZNC | 4 | 2.4 | 1 | 17191 | 960 | 1622 | 1537 | Zinc/Iron Ion Transporter |
| COP | 1 | 1.27 | 1 | 5062 | 3119 | 3956 | 3931 | Copper Ion Transporter |
| *Tol1* **Linked Genes** | | | | | | | | |
| CDK | 7467* | 1.26 | 1 | 1768 | 7840 | 19121 | 15778 | Cyclin Dependent Kinase |
| PUM | 10201* | 8.51 | 1 | 2789 | 10162 | 19264 | 23900 | Pumilio-family, RNA binding |
| MCO | 47895* | 1.06 | 1 | 12543 | 90664 | 340560 | 56920 | Multicopper oxidase |

The length and age of the core haplotypes are estimated by modeling the rate at
which $\pi_{M1-M2}$ (pairwise $\pi$ between mine populations ) increases with distance from
the selected site using the program, MSSEL, see supplemental text for additional
details. The length of the core haplotype is estimated using equation 1 in S. Text.
Coalesence time in presented in generations for within population (CCC; MCN) and
between population (CCC/OM1; MCN/OM2) analyses. *Scaffold ID corresponds to
CCC52 de novo assembly.

**References**

Al-Hiyaly, S. A. K., McNeilly, T., & Bradshaw, A. D. (1990). The Effect of Zinc Contamination from Electricity Pylons. Contrasting Patterns of Evolution in Five Grass Species. New Phytologist, 114(2), 183–190.

Al-Hiyaly, S. A. K., McNeilly, T., Bradshaw, A. D., & Mortimer, A. M. (1993). The effect of zinc contamination from electricity pylons. Genetic constraints on selection for zinc tolerance. Heredity, 70, 22-32.

Allen, W. R., & Sheppard, P. M. (1971). Copper Tolerance in Some Californian Populations of the Monkey Flower, *Mimulus guttatus*. Proc. R. Soc. B., 177(1047), 177–196.

Antonovics, J., Bradshaw, A. D., & Turner, R. G. (1971). Heavy metal tolerance in plants. Adv. Ecol. Res., 7:1–85.

Aubury, L. E. (1905). The copper resources of California (No. 23). WW Shannon, Superintendent State Print.

Barton, N. H. (2000). Genetic hitchhiking. Phil. Tran. R. Soc. B., 355(1403), 1553-1562.

Berg, J. J., & Coop, G. (2015). A Coalescent Model for a Sweep of a Unique Standing Variant. Genetics. 201.

Bradshaw, A. D. (1984). The importance of evolutionary ideas in ecology and vice versa. Evolutionary ecology, 1-25.

Bradshaw, A. D. (1991). The Croonian Lecture, 1991. Genostasis and the limits to evolution. Phil. Tran. R. Soc. B., 333(1267), 289–305.

Brenchley, W. E. (1958). The Park Grass Plots at Rothamsted 1856-1949. Rep. Rothamsted Exp. Stn.

Colosimo, P. F., Hosemann, K. E., Balabhadra, S., Villarreal, G., Dickson, M., Grimwood, J., et al. (2005). Widespread parallel evolution in sticklebacks by repeated fixation of Ectodysplasin alleles. Science, 307(5717), 1928–1933.

Endler, J. A. (1977). Geographic Variation, Speciation and Clines (p. 268). Princeton University Press.

Grennan, A. K. (2009). Identification of genes involved in metal transport in plants. Plant Physiology, 149(4), 1623–4.

539     Hanikenne, M., Talke, I. N., Haydon, M. J., Lanz, C., Nolte, A., Motte, P., et. al. (2008).
540     Evolution of metal hyperaccumulation required cis-regulatory changes and
541     triplication of HMA4. Nature, 453(7193), 391–395.
542
543     Heliconius Genome Consortium. (2012). Butterfly genome reveals promiscuous
544     exchange of mimicry adaptations among species. Nature, 487(7405), 94-98.
545
546     Innan, H., & Kim, Y. (2004). Pattern of polymorphism after strong artificial selection
547     in a domestication event. Proc. Nat. Acad. Sci.101(29), 10667-10672.
548
549     Innan, H., & Kim, Y. (2008). Detecting local adaptation using the joint sampling of
550     polymorphism data in the parental and derived populations. Genetics, 179(3), 1713-
551     1720.
552
553     Joron, M., & Mallet, J. L. (1998). Diversity in mimicry: paradox or paradigm? Trends
554     Eco. & Evol, 13(11), 461-466.
555
556     Karasov, T., Messer, P. W., & Petrov, D. A. (2010). Evidence that Adaptation in
557     *Drosophila* Is Not Limited by Mutation at Single Sites. PLoS Genetics, 6(6),
558     e1000924.
559
560     Kruckeberg, Arthur R. (1951) Intraspecific variability in the response of certain
561     native plant species to serpentine soil. Amer. Jour. Botany (1951): 408-419
562
563     Macnair, M. R. (1987). Heavy metal tolerance in plants: A model evolutionary
564     system. Trends Eco. & Evol, 2(12), 354–359.
565
566     Macnair, M. R., Smith, S. E., & Cumbes, Q. J. (1993). Heritability and distribution of
567     variation in degree of copper tolerance in Mimulus guttatus at Copperopolis,
568     California. Heredity, 71(5), 445–455.
569
570     Maynard-Smith, J., & Haigh, J. (1974). The hitch-hiking effect of a favourable gene.
571     Genetics Research, 23(01), 23–35.
572
573     Messer, P. W., & Petrov, D. A. (2013). Population genomics of rapid adaptation by
574     soft selective sweeps. Trends Eco. & Evol, 28(11), 659–69.
575
576     Prezeworski, M., Coop, G., & Wall, J. D. (2005). The signature of positive selection on
577     standing genetic variation. Evolution, 59(11), 2312-2323.
578
579     Roesti, M., Gavrilets, S., Hendry, A. P., Salzburger, W., & Berner, D. (2014). The
580     genomic signature of parallel adaptation from shared genetic variation. Molecular
581     Ecology, 23(16), 3944-3956.
582
583     Schluter, D. (2000). The Ecology of Adaptive Radiation. Oxford, England: Oxford
584     University Press.

586  Symeonidis, L., McNeilly, T., & Bradshaw, A. D. (1985). Differential tolerance of three
587  cultivars of Agrostis capillaries l. to cadmium, copper, lead, nickel and zinc. New
588  Phytologist, 101(2), 309–315.
589

590  Tishkoff, S. A., Reed, F. A., Ranciaro, A., Voight, B. F., Babbitt, C. C., Silverman, J. S., ...
591  Deloukas, P. (2007). Convergent adaptation of human lactase persistence in Africa
592  and Europe. Nature Genetics, 39(1), 31–40.
593

594  Turner, R. G. (1969). Heavy metal tolerance in plants. Vol. 9, 399-410. Blackwell
595  Oxford.
596

597  Walley, K. A., Khan, M. S. I., & Bradshaw, A. D. (1974). The potential for evolution of
598  heavy metal tolerance in plants. I. Heredity, 32, 309-319.
599

600  Wright, K. M., Lloyd, D., Lowry, D. B., Macnair, M. R., & Willis, J. H. (2013). Indirect
601  evolution of hybrid lethality due to linkage with selected locus in *Mimulus guttatus*.
602  PLoS Biol, 11(2), e1001497.

1A.

1B.

1C.

1D.

Off-Mine$_P$  F1  Mine$_P$  Mine$_{BC4}$  Off-Mine$_{BC1}$

# S Figure 1

S Figure 2

S Figure 3

# S Figure 4

S Figure 5

# S Figure 6

# S. Figure 7

# S. Figure 8

S Figure 9

**A.** S Figure 10

**Scaffold 148**

**Scaffold 198**

S Figure 11

S Figure 12

S Figure 13

S Figure 14

S Figure 15

Scaffold 47895 (kb)