Crop diversity enhances disease suppressive potential in soils

Ariane L. Peralta,[a*]YanmeiSun,[a*]Marshall D. McDaniel,[b]and Jay T. Lennon[c]

Department of Biology, East Carolina University, Greenville, NC, USA[a]; Department of

Agronomy, Iowa State University, USA[b]; Department of Biology, Indiana University,

Bloomington, IN, USA[c]

*A.L.P and Y.S. contributed equally to this work.

E-mail: peraltaa@ecu.edu; sunya14@ecu.edu; marsh@iastate.edu, lennonj@indiana.edu

Running Head: Crop diversity enhances disease suppression

Corresponding author: Ariane L. Peralta, e-mail: peraltaa@ecu.edu

ABSTRACT

Biodiversity is thought to regulate a wide range of agroecosystem processes including plant

production and disease suppression. Farmers have used crop rotations, a form of biodiversity, for

thousands of years and this may be due, in part, to early observations of "disease prevention" in

the form of increased yield.   However, the evidence for a mechanistic link between crop

rotations and disease suppression has not yet been elucidated. Disease suppressive soils are

characterized by the biocontrol properties provided by resident soil microorganisms. Biocontrol

properties include antibiosis via production of antifungal or antibacterial compounds known to

suppress the growth of soil-borne pathogens. In this study, we investigated the impact of long-

term crop diversity (via rotation) on microbial communities and disease suppressive functional

potential in soils. We hypothesized that plant and microbial biodiversity provide disease

suppressive functions in soils. To address these hypotheses, we collected soil samples from a 12-

year crop rotation experiment at the Kellogg Biological Station Long-Term Ecological Research

(KBS LTER) site. We sampled seven treatments along a crop diversity gradient (monoculture to

five crop species) and a spring fallow (naturally regenerating plants) treatment to examine the

influence of crop diversity on total bacterial community composition (16S rRNA gene

sequencing) and a subset of microorganisms capable of producing antifungal compounds (2,4-

diacetylphloroglucinol: *phlD* gene fingerprint analysis; pyrrolnitrin: *prnD* gene quantitative

PCR). Our study revealed that crop diversity significantly influenced bacterial community

composition, and crop rotations decreased bacterial diversity by 4% on average compared to

monocultures. Crop rotations did, however, increase disease suppressive functional group *prnD*

gene abundance in the more diverse rotation (corn-soybean-wheat + cover crops) by about 9%

compared to monocultures. Variation in plant inputs to soil organic matter pools may be a

possible mechanism driving shifts in microbial community patterns and disease suppressive

functional potential.

Keywords

Crop rotation; disease suppression; microbial diversity; structure-function relationships

Abbreviations

2,4-diacetylphloroglucinol (DAPG); plant growth promoting rhizobacteria (PGPR); plant

pathogen suppression (PPS); pyrrolnitrin (PRN)

1. Introduction

Agricultural intensification has led to declines in biodiversity and in associated ecosystem functions (Tilman et al., 2002). However, crop diversification within agricultural landscapes is a strategy used to alleviate this loss in biodiversity. Increases in crop biodiversity on the farm can take many forms, such as crop rotations, cover crops, inter-cropping, and cover crop mixtures. From a management or conservation perspective, crop rotations are not the traditional form of increasing biodiversity. Instead of managing species in space, crop rotations increase diversity through time. This is because at any given time the species richness on a farm using crop rotations is often one (same as monocultures), but there is a diverse suite of biochemical inputs from crops planted at different times to soil microorganisms. There is mounting evidence that this form of 'temporal biodiversity' may provide some of the same beneficial ecosystem functions as traditional spatial biodiversity, such as carbon sequestration, pest control, and nutrient cycling (Ball et al., 2005; McDaniel et al., 2014b; Tiemann et al., 2015; Venter et al., 2016).

Different mechanisms confer plant pathogen suppression (PPS) in the soil. First, crops and the soil microbial community are linked by the plant inputs of carbon and nutrients to the soil. Studies have shown the quantity and quality, and likely even the diversity of crop inputs (residues and rhizodeposits) to soil can alter the microbial community and functioning (Zak et al., 2003; Hättenschwiler et al., 2005; Dijkstra et al., 2010; van der Putten et al., 2016). Second, each crop will affect the physical characteristics of the soil environment whether it is with differences in water use (Tilman et al., 2002), shading (Liebman and Dyck, 1993), aggregation (Tiemann et al., 2015), root morphology affecting porosity (Smucker, 1993), or all of the above. Both of these chemical and physical mechanisms of crop influence on soil microbial

communities might alter the functions important to crops, such as nutrient mineralization from

soil organic matter (Barness et al., 1991), N_2 fixation (Reed et al., 2010), plant growth promotion

(Lugtenberg and Kamilova, 2009), and plant pathogen suppression (Bever et al., 1997; Haas and

Defago, 2005; Perez et al., 2008). The stimulation of these functions is what likely controls

plant-soil feedbacks in agroecosystems (Bever et al., 1997; Kulmatiski et al., 2008; van der

Putten et al., 2016).

Greater plant pathogen suppression has been associated with soil microbial communities

known to exhibit antimicrobial properties, which ultimately provide plants with protection from

soil-borne plant pathogens. Greater soil microbial diversity can provide more opportunity for

PPS potential microorganisms to be maintained although may not be required. Maintenance of

these PPS microorganisms in the community are thought to occur due to competition for iron,

antibiosis, lytic enzymes, and induction of system resistance with host plant (Doornbos et al.,

2012). Specifically, antibiosis has been linked to disease suppressive capacity, whereby the

abundance of antagonistic bacteria has been associated reductions in fungal pathogens through

competitive inhibition (Weller et al., 2002; Haas and Defago, 2005). Bacterial production of

secondary metabolites 2,4-diacetylphloroglucinol (DAPG) and pyrrolnitrin (PRN) are two potent

toxins known to suppress fungal pathogens in agricultural soils (Garbeva et al., 2004a; Garbeva

et al., 2004b; Haas and Defago, 2005). In a previous study, increased plant diversity was

associated with enhanced soil disease suppressiveness measured by increased DAPG and PRN

producers (Latz et al., 2012). In other studies, streptomycetes (a well-known group of bacteria

possessing antibiotic inhibitory effects) were found to contribute to disease suppression in

agricultural soils (Wiggins and Kinkel, 2005; Perez et al., 2008). However, the diversity,

composition, and disease suppressive activity among streptomycetes communities has also been

found to be unrelated to plant diversity treatments (Bakker et al., 2010). Thus, the relationship

between biodiversity and disease suppression in agricultural soils remains unclear. By focusing

on the disease suppressive capacity of soil, we can evaluate how agricultural land-use strategies

and subsequent changes in the soil environment and resident microorganisms impact plant

growth (Bakker et al., 2010; Kulmatiski and Beard, 2011).

Given the unknown effect of crop diversity, via rotations on microbial communities and

plant pathogen suppression, we used a long-term (12 y) crop rotation study at the Kellogg

Biological Station LTER to test the effect of crop diversity on soil bacterial biodiversity and PPS

potential. Specifically, our research addresses the following questions: (1) what is the

relationship between crop diversity and soil microbial community composition and disease

suppressive functional potential? and (2) what is the relationship between changes in soil

physicochemical properties, soil microbial community composition, and disease suppressive

functional potential in response to a crop diversity gradient? We tested the hypothesis that plant

and microbial biodiversity provide disease suppressive functions in soils. We predicted that soils

from high crop diversity (i.e., long crop rotations) would have greater soil bacterial diversity and

have greater PPS capacity compared to soil microbial communities developed under low crop

diversity (i.e., monoculture and short crop rotations like corn-soybean).

2. Methods

2.1. Site description & experimental design

We   collected   soils   from   the   Biodiversity   Gradient   Experiment

(http://lter.kbs.msu.edu/research/long-term-experiments/biodiversity-gradient/) at W.K. Kellogg

Biological Station Long-Term Ecological Research (KBS LTER) site in southwest, Michigan,

USA. Mean annual temperature is 9.7 °C and mean annual precipitation is 890 mm. The soils are

Kalamazoo (fine-loamy) and Oshtemo (coarse-loamy) mixed, mesic, Typic Hapluadalfs formed

under glacial outwash (Crum and Collins, 1995). The crop rotation treatments at the Biodiversity

Gradient Experiment included: monoculture corn (*Zea maize*, mC), corn with 1 red clover

(*Trifolium pretense* L.) cover crop (C_{1cov}), corn-soy (*Glycine max*, CS), corn-soy-wheat (*Triticum*

aestivum, CSW), CSW with red clover (CSW_{1cov}), CSW with red clover and cereal rye (*Secale*

cereal L., CSW_{2cov}), and a spring fallow treatment that was just plowed every spring but contains

7-10 naturally-occurring plant species in the region (Table 1). This spring fallow treatment is

considered the benchmark for plant diversity in the region, and under same tillage. Plantings of

cover crop were dependent on the main crop in rotation (Smith and Gross, 2006). The

experiment was in a randomized complete block design, which included four blocks or replicates

of each treatment. All plots received the same tillage at 15 cm depth, and no fertilizer or

pesticides were applied to these plots.

2.2. Soil sampling

We sampled soil from six crop diversity treatments, but to eliminate any immediate crop

effect all the treatments were sampled in the corn phase and a spring fallow treatment (Table 1)

on November 1, 2012. In each plot, we collected five soil cores (5 cm diameter, 10 cm depth)

and then homogenized the cores in the field. A subsample from each composite sample was

sieved through 4 mm in the field, flash frozen in the field in liquid nitrogen, and stored at -80 °C

prior to molecular-based microbial analyses. On the remaining sample, soil chemical properties

(total carbon, total nitrogen, ammonium, nitrate, pH, texture) were analyzed as originally

reported elsewhere (McDaniel et al., 2014a; McDaniel et al., *In press*). Labile C was measured as

permanganate oxidizable C (POXC). Overall biological activity and amount of potentially

mineralizable carbon (PMC) and nitrogen (PMN) were analyzed using the aerobic incubation

method (McDaniel et al., *In press*).

2.3. Bacterial community sequencing

We extracted DNA using the MoBio Power Soil DNA Isolation Kit (MO BIO

Laboratories, Inc., Carlsbad, CA). DNA concentration was adjusted to a standard concentration

of 20 ng μl^{-1} and used as template. To characterize bacterial taxonomic diversity, we used

barcoded primers (515f/806r primer set) developed by the Earth Microbiome Project to target the

V4-V5 region of the bacterial 16S subunit of the ribosomal RNA gene (16S rRNA) (Caporaso et

al., 2012). For each sample, PCR product combined from three 50 μl reactions, concentration

quantified, and PCR product from each soil sample was combined in equimolar concentrations

for paired-end 250×250 sequencing using the Illumina MiSeq platform according to details in

Muscarella et al. (2014). Briefly, we assembled the paired-end 16S rRNA sequence reads using

the Needleman algorithm (Needleman and Wunsch, 1970). All sequences were subjected to

systematic checks to reduce sequencing and PCR errors. High quality sequences (i.e., >200 bp in

length, quality score of >25, exact match to barcode and primer, and contained no ambiguous

characters) were retained. In addition, we identified and removed chimeric sequence using the

UCHIME algorithm (Edgar et al., 2011). We aligned our sequence data set with the bacterial

SILVA-based bacterial reference database (Yilmaz et al., 2013). During data analysis,

operational taxonomic units (OTUs) were binned at 97% sequence identity and phylogenetic

classifications of bacterial sequences performed. Sequences were processed using the software

package *mothur* v.1.35.1 (Schloss et al., 2009; Kozich et al., 2013).

2.4. Composition and abundance of disease suppression genes

We classified disease suppressive taxa as the subset of soil microorganisms possessing

genes that are required for the production of antifungal compounds 2,4-diacetylphloroglucinol

(DAPG) and pyrrolnitrin (PRN) (Garbeva et al., 2004b; Haas and Defago, 2005). We targeted

phlD and *prnD*, which are known to code for a subset of DAPG producers and PRN producers,

respectively in environmental samples (according to methods in Latz et al. 2012). We assessed

the composition of disease suppressive microorganisms by targeting the *phlD* gene using

terminal restriction fragment length polymorphism (T-RFLP) (von Felten et al., 2011). For *phlD*

gene amplification, the forward primer B2BF (5′-ACCCACCGCAGCATCGTTTATGAGC-3′)

and reverse primer FAM-BPR4 (5′-CCGCCGGTATGGAAGATGAAAAAGTC-3′) yielded a

629 bp product. In each 25 μL PCR reaction, we combined 5% dimethylsulfoxide, 0.8 mg ml^{-1}

bovine serum albumin, 1× GoTaq Colorless Master Mix (Promega, Madison, WI), 0.2 μM of

each primer and 5 μL of template DNA. Reactions were cycled with an initial denaturation at 94

°C for 2 min, followed by 35 cycles of 94 °C for 1 min, 60 °C for 1 min, and 72 °C for 2 min,

with a final extension carried out at 72 °C for 10 min (von Felten et al., 2011). The amplified

PCR products were purified with QIAquick PCR purification kit (Qiagen, Valencia, CA). After

purification, amplicons generated from each sample were digested in multiple restriction

enzymes overnight in 12 μL reaction mixtures containing 4 μL of PCR product, 1×enzyme buffer

(von Felten et al., 2011). After digestion, the enzymes were inactivated for 5 min at 80 °C, and

the digested products were purified according to the purification kit protocol (Qiagen, Valencia,

CA). For T-RFLP analysis, we combined 1.5 μL of the digested product with 9 μL of HiDi

formamide (Applied Biosystems, Foster City, CA) and 0.4 μL of internal size standard ABI

GeneScan LIZ 600 (Applied Biosystems, Foster City, CA). The samples were incubated for 3

min at 96 °C and then stored on ice prior to fragment analysis. We determined the length and

relative abundance of terminal restriction fragments (T-RFs) using an ABI 3130×1 Genetic

Analyzer (Applied Biosystems, Foster City, CA). Electrophoresis conditions were 60 °C and 15

kV with a run time of 41 min using POP-7 polymer. The resulting data was analyzed using

GeneMapper Software 4.0 (Applied Biosystems, Foster City, CA). The peak detection limit was

set to 50 fluorescence intensity units.

We assessed the relative abundance of disease suppressive functional genes by targeting

prnD using quantitative PCR (qPCR) (Garbeva et al., 2004b). The partial *prnD* gene abundance

was quantified using a SYBR green assay with primers prnD-F (5′-

TGCACTTCGCGTTCGAGAC-3′) and prnD-R (5′-GTTGCGCGTCGTAGAAGTTCT-3′)

(Garbeva et al., 2004b). The 25 μL PCR reaction contained 1× GoTaq Colorless Master Mix

(Promega, Madison, WI), 0.4 μM of each primer, and 5 μL of template DNA. Cycling conditions

were as following: initial cycle 95 °C for 10 min, and 30 cycles of 95 °C for 15 s and 60 °C for 1

min. For the qPCR standard curve, *prnD* gene was amplified from soil genomic DNA. PCR

fragments were cloned to pGEM-T Easy Vector System according to the manufacturer's manual

(Promega, Madison, WI). Plasmids were extracted using the QIAprep Spin Miniprep kit (Qiagen,

Valencia, CA), and cloned fragments were verified by PCR and agarose gel electrophoresis.

Dilutions of plasmid DNA containing *prnD* gene were used to generate standard curves in

quantities ranging from 5.0×10^2 to 5.0×10^7 copies. We quantified the *prnD* gene in 25 μL

reaction volumes containing about 20 ng DNA template, 1×TaqMan Environmental Master Mix

2.0 (Applied Biosystems, Valencia, CA), 1× SYBR green I, and 0.4 μM of each primer.

Fragments were amplified with an initial denaturation step at 95 °C for 10 min, followed by 40

cycles of 95°C for 15 s, 60 °C for 1 min. For each sample, PCR reactions were run in triplicate.

We obtained standard curves based on serial dilutions of mixed PCR product amplified from soil

samples. Reactions were analyzed on a BIO-RAD CFX-96Real-Time System (Bio-Rad,

Hercules, California, USA).

2.5. Statistical analyses

We tested for differences in total bacterial diversity (based on Shannon Diversity index

H′) and *prnD* gene abundance in response to crop diversity treatment using analysis of variance

(ANOVA). We checked that data met assumptions of analyses, and we treated crop diversity

treatment as a fixed factor and block as a random effect. We used Tukey's Honestly Significant

Difference (HSD) tests to identify between-group differences in bacterial diversity and *prnD*

gene abundance.

To visualize patterns of microbial community composition, we used Principal

Coordinates Analysis (PCoA) of the microbial community composition based on the Bray-Curtis

dissimilarity coefficient for each possible pair of samples using the R statistical package (R Core

Development Team 2015). To test for differences in total bacterial communities and a subset of

previously identified biocontrol bacterial taxa (i.e., *Psuedomonas* spp. and *Streptomyces* spp.)

among crop diversity treatments, we used non-parametric permutational multivariate analysis of

variance (PERMANOVA) implemented with the *adonis* function in the R Statistics Package R

version 3.0.2 (R Development Core Team 2015). PERMANOVA was also used to assess the

contribution of soil factors to the variation in bacterial community composition. The R^2 value

reported refers to the treatment sums of squares divided by the total sums of squares for each soil

factor in the model. Because the *adonis* function carries out sequential tests (similar to Type I

sums of squares) (Oksanen et al., 2010), the effect of the last soil factor or soil biological activity

factor of the model was included in the final PERMANOVA model summary (Peralta et al.,

2012). We also performed a similarity percentage analysis (SIMPER) using the *simper* function

(R Statistics Package R version 3.0.2) (Clarke, 1993; Warton et al., 2012) to identify the bacterial

OTUs responsible for community differences between monoculture corn and other crop diversity

treatments and is based on the contribution of individual taxa to the average Bray-Curtis

dissimilarity. We also performed multiple linear regression (gene abundance ~ crop number +

total soil carbon + soil moisture + soil ammonium + soil nitrate) to test the influence of soil

factors and crop diversity number on abundance of disease suppression/biocontrol gene *prnD*

using the *lm* function in the R Statistics Package R version 3.0.2 (R Core Development Team

2015).

3. Results

3.1. Bacterial community composition along a crop diversity gradient

A total of 12,539,359 sequence reads were generated, and we analyzed 47,261 OTUs for

bacterial community "patterns". A summary of soil attributes is presented in Table S1 and

elsewhere (McDaniel and Grandy, 2016). The crop diversity treatment significantly influenced

bacterial community composition (R^2 = 0.37, p < 0.001; Table S2a, Fig. 1). Bacterial

communities from the fallow plots and the most diverse crop rotations (CSW, CSW_{1cov},

CSW_{2cov}) were more similar to each other than the lower crop diversity treatments (C_{1cov}, CS)

(Fig. 1). The monoculture corn (mC) treatment was more distinct in bacterial community

composition than all other crop diversity treatments (Fig. 1). Bacterial diversity, as measured

using Shannon Diversity Index (H′), was surprisingly greater under lower crop diversity systems

than higher crop diversity systems, but highest in fallow treatments the most diverse non-

cropping system (crop rotation: $F_{6,20}$ = 10.16, p <0.0001; block: $F_{1,20}$ = 0.20, p = 0.6600; Fig. 2).

Among, the corn cropping systems, mC had the highest Shannon Diversity Index and decreased

by up to as much as 4 % in the most diverse rotation of corn-soybean-wheat with two cover

crops (CSW_{2cov}).

Bacterial community composition was best explained by soil texture (R^2 = 0.066, p<0.05,

Table 3a). However, bacterial community composition was marginally affected by soil moisture

(R^2 = 0.048, p < 0.10, Table 2). Labile C had an effect on bacterial community composition (R^2

= 0.074, p < 0.05), but potentially mineralizable C did not. Potentially mineralizable nitrogen

(PMN), however, which is produced in the same aerobic incubation as PMC (a biologically-

available N pool), explained significant variation in bacterial community composition (R^2 =

0.063, p<0.05, Table 3).

The bacterial taxa primarily responsible for treatment differences between mC and the

other crop diversity treatments are *Sphingomonadales* spp. and *Acidobacteria* subgroup Gp6

(Table S3). When we compared a subset of taxa representing broad biocontrol bacterial

community (composed of *Streptomyces* spp. and *Pseudomonas* spp.), there was no significant

pattern in community composition across the crop diversity treatment (PERMANOVA; crop

rotation: $R^2 = 0.321$, $p = 0.132$; Table S4).

3.2. Disease suppression functional potential in soils along a crop diversity

Crop diversity affected the composition and abundance of disease-suppression soil

microorganisms. We observed a significant shift in the composition of disease-suppression

microorganisms (represented by *phlD* gene T-RFLP) along the crop diversity gradient

(PERMANOVA; crop rotation: $R^2 = 0.52$, $p = 0.037$; Fig. 3, Table S2b). The *phlD* community

composition in the fallow treatment was different from other cropping systems (Fig. 3). The

relative abundance of dominant T-RF 280 bp fragment group, previously identified as an

important disease suppressive bacterial population (von Felten et al., 2011), accounted for about

70% of the disease suppressive community under fallow conditions. In addition, the disease

suppressive functional group T-RF 582 bp was a dominant group, representing about 31-97%

relative abundance across all crop diversity treatments. In addition, *prnD* gene abundances in

cropping systems were higher than under fallow conditions (crop rotation: $F_{6,20} = 7.51$, $p =$

0.0003; Fig. 4). In cropping systems, the *prnD* gene in CSW$_{2cov}$ treatment was the most

abundant, and the gene abundance was significantly higher than in CSW and fallow treatments

(Fig. 4). Our diversity benchmark, the fallow treatment (i.e., lowest crop diversity), showed the

lowest *prnD* gene abundances (Fig. 4). Based on multiple linear regression analysis, plant and

soil factors significantly influenced *prnD* abundance (Adjusted $R^2=0.40$, $F=4.571$, $p=0.005$).

Crop species number (p=0.003), soil carbon (p=0.002), and soil moisture (p=0.0005)

significantly influenced *prnD* gene abundance (Table 4).

4. Discussion

We found that crop rotation history changed bacterial diversity and disease suppression

potential in agricultural soils in the current study. Contrary to our prediction, bacterial diversity

decreased with increasing cropping diversity (Fig. 2). In contrast, disease suppressive potential

of the soil microbial community increased with crop diversity, with the lowest suppressive

potential in the no crop fallow treatments (Fig. 4). A possible explanation for this pattern in

belowground biodiversity is the contribution of cover crop species to the rotation and the

interaction with weedy plant species associated with these plots and. We observed that without

crop plants (as reflected in the no crop fallow treatment), disease suppressive potential was

significantly diminished compared to crop treatments, possibly due to reduced selection for soil

microorganisms with disease suppression traits. The composition of the soil microbial

community may be more important than diversity to soil suppressive function. Thus, crop

rotation has the potential to impact diseases suppressive function, providing evidence for

sustainable biocontrol of soil-borne pathogens.

4.1. Crop diversity decreases belowground (bacterial) diversity

Crop rotation history decreased bacterial diversity and increased disease suppression

potential in this 12-year crop diversity study. Contrary to our prediction, however, crop diversity

decreased soil bacterial diversity (Fig. 2). A recent meta-analysis showed that the crop rotation

effect increased soil bacterial diversity (i.e., Shannon's diversity index H′) most notably in the

first 5 years of treatment, but crop rotations occurring in greater than 5 years were more variable

in diversity (Venter et al., 2016). Although, a few of studies included in the meta-analysis were

based on high throughput sequencing approaches (i.e., pyrosequencing) also found decreases in

bacterial diversity with increasing crop diversity (Alvey et al., 2001; Yin et al., 2010; Reardon et

al., 2014; Venter et al., 2016).

The pattern of reduced bacterial diversity (based on 16S rRNA gene sequencing) was

lower in soils with higher cropping diversity did not align with our initial predictions. There are a

number of potential reasons for this pattern. One explanation for this pattern in belowground

biodiversity is the presence of weedy plant species. Increasing crop diversity within a rotation is

used as a strategy for reducing reliance on synthetic herbicides and been correlated with reduced

weed diversity. Early observations at this long-term crop diversity study revealed decreasing

weed abundance with increasing crop diversity during the 2002-2004 seasons (Smith and Gross,

2007). Specifically, the monoculture treatments (including mC) had an average of 13 weed

species per m^{-2}, but the most diverse cropping systems (CSW$_{2cov}$) had only 5 or 6 in 2003. Thus,

while our crop diversity is lower in monocultures, they actually may have greater total plant

diversity compared to more diverse rotations when weeds are included. Last, instead of diversity,

the Shannon Index might also be looked at as an indicator of a shift in microbial carbon usage.

More specifically, a study by McDaniel and Grandy (McDaniel and Grandy, 2016), using the

very same soils we used in this study, found that catabolic evenness (a diversity measure of the

catabolism of a suite of 31 carbon compounds) also decreased with increasing crop diversity.

This indicates that this trend is not just structural, but also functional, and may indicate

specialization.

4.2. Crop diversity enhances plant pathogen suppression

The diversity of plant pathogen suppressive (PPS) microbial community increased with

crop diversity treatment (Fig. 4). In addition, we found that the increased crop diversity, via

rotation, increased the abundance and composition of a specific plant pathogen suppression gene.

Together, these results suggest that cropping diversity may increase the disease suppressive

functional potential of agricultural soils. These findings are consistent with previous studies

suggesting that plant diversity can enhance protection against soil-borne pathogens by fostering

antagonistic soil bacterial communities (Latz et al., 2012; van der Putten et al., 2016). One

potential explanation for this effect is changes in plant root exudation, which may lead to

enrichment of plant growth promoting rhizobacteria (PGPRs) (Badri et al., 2009; Chaparro et al.,

2012). In previous studies, interactions among the total microbial community and soil-borne

pathogens in the plant rhizosphere have influenced both plant growth and productivity (Bakker et

al., 2010; Penton et al., 2014).

The addition of cover crops to rotations, in particular, strongly increased disease

suppressive potential. This along with evidence from previous studies shows that crop rotations

may prevent many forms of crop disease caused by *Fusarium* spp., *Phytophthora*, and

Rhizoctonia spp. (Raaijmakers et al., 2009; van der Putten et al., 2016). Soil microbial diversity

has been implicated as important for soil disease suppression; sterilized soils lose suppressive

capacity, and adding soil microorganisms to sterilized soil facilitates disease suppression

functional capacity (Garbeva et al., 2006; Brussaard et al., 2007; Postma et al., 2008). Biocontrol

bacteria can also provide disease suppression against plant pathogens by way of the following

mechanisms: competition for iron, antibiosis, lytic enzymes, induction of system resistance of

host plants (Doornbos et al., 2012). Plants can also recruit specific biocontrol microorganisms in

some cases. A previous study suggests that beneficial pseudomonads are recruited depending on

the most dominant soil-borne pathogen infecting crop species. Specifically, *Pseudomonas* spp.

capable of DAPG production are more effective at controlling *G. faminis var. tritici,* while

Pseudomonas spp. capable of producing phenazines were enhanced in fields where phenazine-

sensitive *Rhizoctonia* plant pathogen were most dominant (Berendsen et al., 2012; Mavrodi et

al., 2012). In the present study, we analyzed a subset of previously reported biocontrol bacterial

taxa (e.g., *Pseudomonas* spp. and *Streptomyces* spp.) across the crop diversity gradient; however,

we did not detect distinct changes in putative biocontrol community composition (Table S4).

The gene abundance results suggest that incorporation of cover crop in rotations increases

prnD gene abundance, a gene associated disease suppressive microbial taxa capable of producing

antifungal compound pyrrolnitrin (PRN) (Garbeva et al., 2004b; Haas and Defago, 2005).

Specifically, cover crop species may have important effects on the *prnD* gene abundance and

disease suppressive functional potential in soils. The *prnD* gene abundance in cropping systems

is higher than in fallow treatment. In addition, there are no reports linking the functional group

T-RF 280 to a specific genotype of laboratory strains (von Felten et al., 2011). In our study, this

bacterial group is capable of producing 2,4-diacetylphloroglucinol (DAPG) and affects DAPG

community composition (Fig. 3). The T-RF 582 bp was also reported to be a major DAPG-

producing functional *Pseudomonas kilonensis* in a previous study (von Felten et al., 2011). The

abundance of DAPG and PRN producers increasing with plant diversity has been previously

observed (Latz et al., 2012). Compared to agricultural soils, the PRN producers were more

frequently detected in grassland or grassland-derived plots (Garbeva et al., 2004a; Garbeva et al.,

2004b). The *prnD* gene abundance increased in the presence of grasses, but the legume species

tended to decrease the DAPG and PRN producer abundance (Latz et al., 2012). Without crop

plants (as reflected in the fallow treatment), we observed that disease suppressive potential

significantly declined. Disease suppression traits such as antifungal production may not be

needed and are not maintained in the community. When agricultural management is absent, there

is reduced selection for soil microorganisms with disease suppression traits. Together, our

findings combined with previous studies suggest that the land-use regime, plant diversity, and

plant species are involved in structuring disease suppressive microbial communities.

4.3 Crop-to-bacterial feedback mechanisms and links to the "rotation effect"

Disease suppression may have a major role in what is colloquially referred to as "the

rotation effect." The increases in yields seen by farmers over the millennia (Karlen et al., 1994)

may be due in large part to disease suppression. Crop rotations may also provide other important

benefits like enhanced nutrient provisioning to plants, improvement of soil physical properties,

increases in soil C, and increases in soil microbial and faunal activity that also could be

responsible for the increased yields responsible for the rotation effect (Ball et al., 2005; van der

Putten et al., 2016). Our study provided evidence that crop rotations alter soil bacterial

community composition, but the mechanisms through which this occurs can include chemical,

physical, and biological changes to the soil environment. Crops can influence soil properties in a

variety of ways, including chemically and physically. Chemically, plants provide carbon to the

soil environment through root exudation of recently assimilated photosynthate, composed of

soluble, low molecular weight organic compounds (Neumann and Romheld, 2007). As a

consequence, the increased C flow from root exudates can stimulate soil microbial activity.

Changes in plant inputs through variation in either root exudation rates or chemical composition

are likely a major factor to how crops and crop rotations, can alter belowground microbial

communities. Our study focused on soil bacterial community composition. It has been identified

that crop rotation also influences soil fungal and faunal communities, which are also important

members of the soil food web (McLaughlin and Mineau, 1995). Changes in root exudates have

been observed to shift microbial community composition and stimulate a diverse microbial

community (Hooper et al., 2000; Stephan et al., 2000; Paterson et al., 2009; Dijkstra et al., 2010).

Physically, crop diversity (especially rotations) can enhance soil properties like improving plant

water availability by lowering bulk density, increasing soil pore space, and increasing soil

aggregate formation (Tilman et al., 2002; McDaniel et al., 2014b; Tiemann et al., 2015), which

could have indirect influence over the soil bacterial community as well. Biologically, soil

microorganisms can provide disease suppression against plant pathogens through competition for

nutrients, antibiosis, and induction of system resistance of host plants (Doornbos et al., 2012).

Cover crops are the most salient feature of these crop rotations affecting the soil bacterial

community in general. This is not surprising, since cover crops have been shown to influence

several soil properties, which likely have indirect effects on the soil bacterial community

composition. Soil properties like total C, total N, pH, and bulk density and porosity have all been

shown to increase with cover crops (Bullock, 1992; Liebman and Dyck, 1993; Tilman et al.,

2002; McDaniel et al., 2014a; McDaniel et al., 2014b; Tiemann et al., 2015). In our study, higher

crop diversity but lower soil bacterial diversity supported higher disease-suppression functional

potential in soils. We posit that rotation has a filtering effect on soil microorganisms, whereby

crop diversity selects for antagonistic microorganisms with disease suppressive potential. Our

study further supports the hypothesis that plant diversity can support biocontrol functional

potential by enhancing antagonistic properties of resident soil microorganisms against soil-borne

pathogens. The soil microbial community composition may be more important than soil

microbial diversity to soil ecological function. Increasing evidence supports that crop species and

soil type have been shown to shape the soil microbial composition and function (McDaniel et al.,

2014b; Tiemann et al., 2015; Venter et al., 2016); thus, land management can have a large effect

on soil microbial processes and microbial community composition (Jangid et al., 2008; Lauber et

al., 2008; Orr et al., 2015).

Acknowledgements

This work was supported by the U.S. Department of Agriculture National Institute of Food and

Agriculture Postdoctoral Fellowship (2012-67012-19845 to A.L.P.) and the National Science

Foundation (DEB 1442246 to J.T.L.). The funders had no role in study design, data collection

and interpretation, preparation of the manuscript, or decision to submit the work for publication.

Support was also provided by the NSF Long-term Ecological Research Program (DEB 1027253)

at the Kellogg Biological Station and by Michigan State University AgBioResearch. We would

like to thank the Kellogg Biological Station LTER for logistical support and use of sampling

sites. We also thank M. Muscarella, J. Ford, S. Krahnke, and M. Brewer for microbial analyses

support and B. O'Neill, A.S. Grandy and T.M. Schmidt Labs for field and soil analyses support.

All sequence data and metadata have been submitted to NCBI and are available at XXXX.

References

Alvey, S., Bagayoko, M., Neumann, G., Buerkert, A., 2001. Cereal/legume rotations affect
chemical properties and biological activities in two West African soils. Plant Soil 231,
45-54.
Badri, D.V., Quintana, N., El Kassis, E.G., Kim, H.K., Choi, Y.H., Sugiyama, A., Verpoorte, R.,
Martinoia, E., Manter, D.K., Vivanco, J.M., 2009. An ABC transporter mutation alters
root exudation of phytochemicals that provoke an overhaul of natural soil microbiota.
Plant Physiol. 151, 2006-2017.
Bakker, M.G., Glover, J.D., Mai, J.G., Kinkel, L.L., 2010. Plant community effects on the
diversity and pathogen suppressive activity of soil streptomycetes. Appl. Soil Ecol. 46,
35-42.
Ball, B., Bingham, I., Rees, R., Watson, C., Litterick, A., 2005. The role of crop rotations in
determining soil structure and crop growth conditions. Can. J. Soil Sci. 85, 557-577.
Barness, E., Chen, Y., Hadar, Y., Marschner, H., Romheld, V., 1991. Siderophores of
Pseudomonas putida as an iron source for dicot and monocot plants. Plant Soil 130, 231-
241.
Berendsen, R.L., Pieterse, C.M., Bakker, P.A., 2012. The rhizosphere microbiome and plant
health. Trends Plant Sci. 17, 478-486.
Bever, J.D., Westover, K.M., Antonovics, J., 1997. Incorporating the soil community into plant
population dynamics: the utility of the feedback approach. J. Ecol. 85, 561-573.
Brussaard, L., De Ruiter, P.C., Brown, G.G., 2007. Soil biodiversity for agricultural
sustainability. Agric. Ecosyst. Environ. 121, 233-244.
Bullock, D.G., 1992. Crop rotation. Crit. Rev. Plant Sci. 11, 309-326.
Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens,
S.M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J.A., Smith, G., Knight, R.,
2012. Ultra-high-throughput microbial community analysis on the Illumina HiSeq and
MiSeq platforms. ISME J. 6, 1621-1624.
Chaparro, J.M., Sheflin, A.M., Manter, D.K., Vivanco, J.M., 2012. Manipulating the soil
microbiome to increase soil health and plant fertility. Biol. Fertility Soils 48, 489-499.
Clarke, K.R., 1993. Non‐parametric multivariate analyses of changes in community structure.
Aust. J. Ecol. 18, 117-143.
Crum, J.R., Collins, H.P., 1995. KBS Soils [Online], W. K. Kellogg Biological Station Long-
Term Ecological Research Project, Michigan State University, Hickory Corners, MI.
Dijkstra, F.A., Morgan, J.A., Blumenthal, D., Follett, R.F., 2010. Water limitation and plant
inter-specific competition reduce rhizosphere-induced C decomposition and plant N
uptake. Soil Biol. Biochem. 42, 1073-1082.
Doornbos, R.F., van Loon, L.C., Bakker, P.A., 2012. Impact of root exudates and plant defense
signaling on bacterial communities in the rhizosphere. A review. Agron. Sustainable Dev.
32, 227-243.
Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., Knight, R., 2011. UCHIME improves
sensitivity and speed of chimera detection. Bioinformatics 27, 2194-2200.
Garbeva, P., Postma, J., Van Veen, J., Van Elsas, J., 2006. Effect of above‐ground plant species
on soil microbial community structure and its impact on suppression of Rhizoctonia
solani AG3. Environ. Microbiol. 8, 233-246.

Garbeva, P., van Veen, J.A., van Elsas, J.D., 2004a. Microbial diversity in soil: Selection of
microbial populations by plant and soil type and implications for disease suppressiveness.
Annu. Rev. Phytopathol. 42, 243-270.
Garbeva, P., Voesenek, K., van Elsas, J.D., 2004b. Quantitative detection and diversity of the
pyrrolnitrin biosynthetic locus in soil under different treatments. Soil Biol. Biochem. 36,
1453-1463.
Haas, D., Defago, G., 2005. Biological control of soil-borne pathogens by fluorescent
pseudomonads. Nature Rev. Microbiol. 3, 307-319.
Hättenschwiler, S., Tiunov, A.V., Scheu, S., 2005. Biodiversity and litter decomposition in
terrestrial ecosystems. Annu. Rev. Ecol. Evol. Syst., 191-218.
Hooper, D.U., Bignell, D.E., Brown, V.K., Brussaard, L., Dangerfield, J.M., Wall, D.H., Wardle,
D.A., Coleman, D.C., Giller, K.E., Lavelle, P., Van der Putten, W.H., De Ruiter, P.C.,
Rusek, J., Silver, W.L., Tiedje, J.M., Wolters, V., 2000. Interactions between
aboveground and belowground biodiversity in terrestrial ecosystems: Patterns,
mechanisms, and feedbacks. Bioscience 50, 1049-1061.
Jangid, K., Williams, M.A., Franzluebbers, A.J., Sanderlin, J.S., Reeves, J.H., Jenkins, M.B.,
Endale, D.M., Coleman, D.C., Whitman, W.B., 2008. Relative impacts of land-use,
management intensity and fertilization upon soil microbial community structure in
agricultural systems. Soil Biol. Biochem. 40, 2843-2853.
Karlen, D., Varvel, G., Bullock, D.G., Cruse, R., 1994. Crop rotations for the 21st century.
Advances in Agronomy 53.
Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., Schloss, P.D., 2013. Development of
a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence
data on the MiSeq Illumina sequencing platform. Appl. Environ. Microbiol. 79, 5112-
5120.
Kulmatiski, A., Beard, K.H., 2011. Long-term plant growth legacies overwhelm short-term plant
growth effects on soil microbial community structure. Soil Biol. Biochem. 43, 823-830.
Kulmatiski, A., Beard, K.H., Stevens, J.R., Cobbold, S.M., 2008. Plant–soil feedbacks: a meta-
analytical review. Ecol. Lett. 11, 980–992.
Latz, E., Eisenhauer, N., Rall, B.C., Roscher, C., Scheu, S., Jousset, A., 2012. Plant diversity
improves protection against soil-borne pathogens by fostering antagonistic bacterial
communities. J. Ecol. 100, 597-604.
Lauber, C.L., Strickland, M.S., Bradford, M.A., Fierer, N., 2008. The influence of soil properties
on the structure of bacterial and fungal communities across land-use types. Soil Biol.
Biochem. 40, 2407-2415.
Liebman, M., Dyck, E., 1993. Crop rotation and intercropping strategies for weed management.
Ecol. Appl. 3, 92-122.
Lugtenberg, B., Kamilova, F., 2009. Plant-growth-promoting rhizobacteria. Annu. Rev.
Microbiol. 63, 541–545.
Mavrodi, O.V., Mavrodi, D.V., Parejko, J.A., Thomashow, L.S., Weller, D.M., 2012. Irrigation
differentially impacts populations of indigenous antibiotic-producing Pseudomonas spp.
in the rhizosphere of wheat. Appl. Environ. Microbiol. 78, 3214-3220.
McDaniel, M.D., Grandy, A.S., 2016. Soil microbial biomass and function are altered by 12
years of crop rotation. SOIL Discuss. 2016, 1-39.

McDaniel, M.D., Grandy, A.S., Tiemann, L.K., Weintraub, M.N., 2014a. Crop rotation
572   complexity regulates the decomposition of high and low quality residues. Soil Biol.
573   Biochem. 78, 243-254.
574   McDaniel, M.D., Grandy, A.S., Tiemann, L.K., Weintraub, M.N., *In press*. Eleven years of crop
575   diversification alters the decomposition dynamics of litter mixtures incubated with soils.
576   Ecosphere.
577   McDaniel, M.D., Tiemann, L.K., Grandy, A.S., 2014b. Does agricultural crop diversity enhance
578   soil microbial biomass and organic matter dynamics? a meta-analysis. Ecol. Appl.
579   McLaughlin, A., Mineau, P., 1995. The impact of agricultural practices on biodiversity. Agric.
580   Ecosyst. Environ. 55, 201-212.
581   Muscarella, M., Bird, K., Larsen, M., Placella, S., Lennon, J., 2014. Phosphorus resource
582   heterogeneity in microbial food webs. Aquat. Microb. Ecol. 73, 259-272.
583   Needleman, S.B., Wunsch, C.D., 1970. A general method applicable to the search for similarities
584   in the amino acid sequence of two proteins. J. Mol. Biol. 48, 443-453.
585   Neumann, G., Romheld, V., 2007. The Release of Root Exudates as Affects by the Plant
586   Physiological Status, In: Pinton, R., Varanini, Z., Nannipieri, P. (Eds.), The Rhizosphere:
587   Biochemistry and Organic Substances at the Soil-plant Interface, 2nd ed. CRC Press, pp.
588   23-72.
589   Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., O'Hara, R.B., Simpson, G.L., Solymos, P.,
590   Stevens, M.H.H., Wagner, H., 2010. Community Ecology Package 'vegan'.
591   Orr, C., Stewart, C., Leifert, C., Cooper, J., Cummings, S., 2015. Effect of crop management and
592   sample year on abundance of soil bacterial communities in organic and conventional
593   cropping systems. J. Appl. Microbiol.
594   Paterson, E., Midwood, A.J., Millard, P., 2009. Through the eye of the needle: a review of
595   isotope approaches to quantify microbial processes mediating soil carbon balance. New
596   Phytol. 184, 19-33.
597   Penton, C.R., Gupta, V., Tiedje, J.M., Neate, S.M., Ophel-Keller, K., Gillings, M., Harvey, P.,
598   Pham, A., Roget, D.K., 2014. Fungal community structure in disease suppressive soils
599   assessed by 28S LSU gene sequencing. PloS One 9, e93893.
600   Peralta, A.L., Matthews, J.W., Flanagan, D.N., Kent, A.D., 2012. Environmental factors at
601   dissimilar spatial scales influence plant and microbial Communities in restored wetlands.
602   Wetlands 32, 1125-1134.
603   Perez, C., Dill-Macky, R., Kinkel, L.L., 2008. Management of soil microbial communities to
604   enhance populations of Fusarium graminearum-antagonists in soil. Plant Soil 302, 53-69.
605   Postma, J., Schilder, M.T., Bloem, J., van Leeumen-Haagsma, W.K., 2008. Soil suppressiveness
606   and functional diversity of the soilmicroflora in organic farming systems. Soil Biol.
607   Biochem. 40, 2394–2406.
608   Raaijmakers, J.M., Paulitz, T.C., Steinberg, C., Alabouvette, C., Moënne-Loccoz, Y., 2009. The
609   rhizosphere: a playground and battlefield for soilborne pathogens and beneficial
610   microorganisms. Plant Soil 321, 341-361.
611   Reardon, C., Gollany, H., Wuest, S., 2014. Diazotroph community structure and abundance in
612   wheat–fallow and wheat–pea crop rotations. Soil Biol. Biochem. 69, 406-412.
613   Reed, S.C., Townsend, A.R., Cleveland, C.C., Nemergut, D.R., 2010. Microbial community
614   shifts influence patterns in tropical forest nitrogen fixation. Oecologia 164, 521-531.
615   Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski,
616   R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B., Thallinger, G.G.,

617   Van Horn, D.J., Weber, C.F., 2009. Introducing mothur: Open-Source, Platform-
618   Independent, Community-Supported Software for Describing and Comparing Microbial
619   Communities. Appl. Environ. Microbiol. 75, 7537-7541.
620   Smith, R.G., Gross, K.L., 2006. Weed community and corn yield variability in diverse
621   management systems. Weed Sci. 54, 106-113.
622   Smith, R.G., Gross, K.L., 2007. Assembly of weed communities along a crop diversity gradient.
623   J. Appl. Ecol. 44, 1046-1056.
624   Smucker, A., 1993. Soil environmental modifications of root dynamics and measurement. Annu.
625   Rev. Phytopathol. 31, 191-218.
626   Stephan, A., Meyer, A.H., Schmid, B., 2000. Plant diversity affects culturable soil bacteria in
627   experimental grassland communities. J. Ecol. 88, 988-998.
628   Tiemann, L.K., Grandy, A.S., Atkinson, E.E., Marin-Spiotta, E., McDaniel, M.D., 2015. Crop
629   rotational diversity enhances belowground communities and functions in an
630   agroecosystem. Ecol. Lett. 18, 761-771.
631   Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R., Polasky, S., 2002. Agricultural
632   sustainability and intensive production practices. Nature 418, 671-677.
633   van der Putten, W.H., Bradford, M.A., Pernilla Brinkman, E., Voorde, T.F., Veen, G., 2016.
634   Where, when and how plant–soil feedback matters in a changing world. Funct. Ecol.
635   Venter, Z.S., Jacobs, K., Hawkins, H.-J., 2016. The impact of crop rotation on soil microbial
636   diversity: A meta-analysis. Pedobiologia.
637   von Felten, A., Meyer, J.B., Defago, G., Maurhofer, M., 2011. Novel T-RFLP method to
638   investigate six main groups of 2,4-diacetylphloroglucinol-producing pseudomonads in
639   environmental samples. J. Microbiol. Methods 84, 379-387.
640   Warton, D.I., Wright, S.T., Wang, Y., 2012. Distance‐based multivariate analyses confound
641   location and dispersion effects. Methods Ecol. Evol. 3, 89-101.
642   Weller, D.M., Raaijmakers, J.M., Gardener, B.B.M., Thomashow, L.S., 2002. Microbial
643   populations responsible for specific soil suppressiveness to plant pathogens. Annu. Rev.
644   Phytopathol. 40, 309-+.
645   Wiggins, E., Kinkel, L.L., 2005. Green manures and crop sequences influence alfalfa root rot and
646   pathogen inhibitory activity among soil-borne streptomycetes. Plant Soil 268, 271-283.
647   Yilmaz, P., Parfrey, L.W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J.,
648   Ludwig, W., Glöckner, F.O., 2013. The SILVA and "All-species living tree project
649   (LTP)" taxonomic frameworks. Nucleic Acids Res. 42, D643-D648.
650   Yin, C., Jones, K.L., Peterson, D.E., Garrett, K.A., Hulbert, S.H., Paulitz, T.C., 2010. Members
651   of soil bacterial communities sensitive to tillage and crop rotation. Soil Biol. Biochem.
652   42, 2111-2118.
653   Zak, D.R., Holmes, W.E., White, D.C., Peacock, A.D., Tilman, D., 2003. Plant diversity, soil
654   microbial communities, and ecosystem function: are there any links? Ecology 84, 2042-
655   2050.

FIGURES

Figure 1

Figure 1.Ordination from Principal Coordinates Analysis depicting soil bacterial communities

along a cropping diversity gradient. Symbols are colored according to cropping diversity

treatment   (mC=monoculture   corn; C_{1cov} =corn/1   cover   crop;   CS=corn/soy;

CSW=corn/soy/wheat; CSW_{1cov} =corn/soy/wheat/1 cover crop; CSW_{2cov}=corn/soy/wheat/2 cover

crops; fallow=spring fallow, tilled annually).

674 Figure 2

676 Figure 2. Total bacterial diversity (mean ± SEM based on Shannon Diversity Index H′) in

677 response to long-term crop diversity treatment. Different letters above points reflect significant

678 differences in gene abundance along crop diversity gradient at *p*< 0.05 (Tukey's HSD *post-hoc*

679 analysis).

688   Figure 3

690   Figure 3. Ordination from Principal Coordinates Analysis of disease suppressive community

691   based on T-RFLP of *phlD* gene (DAPG producers) T-RF relative abundance along crop diversity

692   gradient.

703   Figure 4

Crop Divesity

705   Figure 4. Abundance of *prn*D gene (PRN producers) in response to crop diversity treatment

706   analyzed using quantitative PCR and expressed as log copy number of *prn*D gene. Different

707   letters above points reflect significant differences in Different letters above boxplots considered

708   significantly different in gene abundance at $p < 0.05$ (Tukey's HSD *post-hoc* analysis).

 TABLES

719 Table 1. Cropping diversity treatments at the Kellogg Biological Station Long-term Ecological

720 Research (KBS LTER) Biodiversity Gradient Experiment Plots. Plant treatments were

721 established in 2000. Treatments were composed of monoculture, two-crop rotation, three-crop

722 rotation +/- cover crops, and fallow plots (early successional) and soil collected during the corn

723 phase of the rotation. Treatment abbreviations are in parentheses.

Crop diversity treatment description	Number of crop species
(1) Continuous monoculture (mC)	1
(2) Continuous monoculture, one cover crop (C_{1cov})	2
(3) Two-crop rotation (CS)	2
(4) Three-crop rotation (CSW)	3
(5) Three-crop rotation, one cover crop (CSW_{1cov})	4
(6) Three-crop rotation, two cover crops (CSW_{2cov})	5
(7) Spring Fallow/early successional field (fallow)	10

Table 2. Summary of the contribution of (A) soil factors (original data from McDaniel et al.

2014) and (B) soil biological activity (original data from McDaniel et al. XXX) on bacterial

community variation at the KBS Biodiversity Gradient Experimental Plots based on

permutational MANOVA (PERMANOVA). Soil factor effects were considered to significantly

contribute to community variation at $P < 0.05$.

(a) Soil Factors

Effect	df	SS	MS	F	R^2	p-value
Sand	1	0.088	0.088	2.243	0.066	0.014
Silt	1	0.088	0.088	2.239	0.066	0.020
Clay	1	0.087	0.087	2.207	0.065	0.024
pH	1	0.057	0.057	1.444	0.043	0.143
Nitrate	1	0.023	0.023	0.593	0.018	0.893
Ammonium	1	0.019	0.019	0.496	0.015	0.966
Nitrogen	1	0.043	0.043	1.086	0.032	0.326
Carbon	1	0.036	0.036	0.921	0.027	0.491
Moisture	1	0.064	0.064	1.622	0.048	0.078
Residuals	18	0.707	0.039		0.534	
Total	27	1.325			1	

744   (b) Soil Biological Activity

Effect	df	SS	MS	F	R^2	p-value
PMN	1	0.083	0.083	1.821	0.063	0.049
PMC	1	0.062	0.062	1.358	0.047	0.146
POXC	1	0.097	0.097	2.125	0.074	0.028
Residuals	24	1.100	0.046		0.830	
Total	27	1.325			1	

760   Table 3. Summary of multiple linear regression to test the influence of disease suppressive

761   functional potential (*prnD* gene abundance) on soil factors and crop diversity.

Factor	Estimate	Std error	t-value	*p*-value
Intercept	7.444	0.420	17.728	< 0.001
Crop_number	-0.085	0.025	-3.355	0.003
Carbon	0.180	0.050	3.618	0.002
Moisture	-11.564	2.817	-4.105	< 0.001
Ammonium	-0.701	0.948	-0.739	0.468
Nitrate	0.093	0.136	0.684	0.501

1   Table S1 Soil properties measured along the KBS cropping diversity gradient averaged over four replicate blocks. We acknowledge

2   M.D. McDaniel and A.S. Grandy for these results (*McDaniel et al. 2014).

Treatment	Total C (g C kg^{-1} soil)	Total N (g N kg^{-1} soil)	NH$_4^+$ (mg N kg^{-1} soil)	NO$_3^-$ (mg N kg^{-1} soil)	pH	Clay (%)	Silt (%)	Sand (%)
Fallow	8.74 ± 2.41	0.77 ± 0.16	0.07 ± 0.04	1.50 ± 0.31	6.70 ± 0.28	25 ± 9	39 ± 19	36 ± 11
CSW-2cov	8.98 ± 1.81	0.91 ± 0.12	0.09 ± 0.04	1.24 ± 0.50	6.25 ± 0.11	21 ± 10	51 ± 22	29 ± 12
CSW-1cov	9.63 ± 1.29	0.91 ± 0.04	0.06 ± 0.02	1.09 ± 0.51	6.37 ± 0.25	24 ± 10	46 ± 17	31 ± 7
CSW	7.43 ± 1.58	0.70 ± 0.10	0.07 ± 0.03	1.26 ± 0.62	6.54 ± 0.21	21 ± 10	45 ± 21	34 ± 12
CS	7.70 ± 2.11	0.73 ± 0.26	0.10 ± 0.09	1.16 ± 0.74	6.66 ± 0.12	23 ± 11	41 ± 24	36 ± 14
C-1cov	9.09 ± 1.86	0.93 ± 0.16	0.06 ± 0.02	1.50 ± 0.65	6.46 ± 0.25	27 ± 8	36 ± 19	38 ± 12
C	8.09 ± 1.24	0.71 ± 0.15	0.16 ± 0.16	1.31 ± 0.36	6.58 ± 0.38	25 ± 9	36 ± 19	38 ± 11

5   *McDaniel MD, Grandy AS, Tiemann LK, Weintraub MN. 2014. Crop rotation complexity regulates the decomposition of high and

6   low quality residues. Soil Biol Biochem 78:243-254.

Table S2 Effects of crop rotation on total bacterial community composition based on 16S rRNA amplicon sequencing (a) and disease suppressive community composition based on *phlD* gene T-RFLP (b) at the KBS Biodiversity Gradient Experimental Plots based on permutational MANOVA (PERMANOVA) results. Rotation effect was considered to significantly contribute to community variation at $P < 0.05$.

(a) 16S rRNA gene

	df	SS	MS	F	R^2	p-value
Rotation	6	0.493	0.082	2.08	0.372	<0.001
Residuals	21	0.832	0.040		0.628	
Total	27	1.32			1	

(b) *phlD* gene T-RFLP

	df	SS	MS	F	R^2	p-value
Rotation	8	3.024	0.378	1.748	0.518	0.037
Residuals	13	2.811	0.216		0.482	
Total	21	5.835			1.000	

Table S3 Summary of similarity percentages (SIMPER) of the top 10 bacterial taxa responsible for differences in bacterial community composition between monoculture corn and each of the crop diversity/fallow treatments.

CS-C

OTU ID	average	sd	ratio	ava	avb	cusum	taxonomy
Otu000001	0.0031	0.0019	1.6782	0.0291	0.0267	0.0117	Bacteria;Proteobacteria;Alphaproteobacteria;Sphingomonadales;Sphingomonadaceae;unclassified
Otu000002	0.0028	0.0016	1.7098	0.0361	0.0305	0.0221	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000004	0.0024	0.0010	2.4510	0.0190	0.0183	0.0312	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000003	0.0023	0.0019	1.2195	0.0192	0.0174	0.0398	Bacteria;Acidobacteria;Acidobacteria_Gp4;Acidobacteria_Gp4_order_incertae_sedis;Acidobacteria_Gp4_family_incertae_sedis;Gp4
Otu000005	0.0016	0.0011	1.4928	0.0126	0.0126	0.0457	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000028	0.0012	0.0004	3.4057	0.0051	0.0051	0.0504	Bacteria;unclassified;unclassified;unclassified;unclassified;unclassified
Otu000064	0.0012	0.0013	0.9627	0.0017	0.0030	0.0550	Bacteria;Firmicutes;Bacilli;Bacillales;unclassified;unclassified
Otu000017	0.0011	0.0010	1.1353	0.0052	0.0061	0.0593	Bacteria;Verrucomicrobia;Spartobacteria;Spartobacteria_order_incertae_sedis;Spartobacteria_family_incertae_sedis;Spartobacteria_genera_incertae_sedis
Otu000007	0.0011	0.0009	1.2244	0.0063	0.0081	0.0635	Bacteria;Proteobacteria;Alphaproteobacteria;Rhizobiales;Bradyrhizobiaceae;unclassified
Otu000006	0.0011	0.0008	1.4008	0.0067	0.0076	0.0675	Bacteria;Verrucomicrobia;Spartobacteria;Spartobacteria_order_incertae_sedis;Spartobacteria_family_incertae_sedis;Spartobacteria_genera_incertae_sedis

C-C1cov

OTU ID	average	sd	ratio	ava	avb	cusum	taxonomy
Otu000001	0.0051	0.0033	1.5290	0.0267	0.0358	0.0178	Bacteria;Proteobacteria;Alphaproteobacteria;Sphingomonadales;Sphingomonadaceae;unclassified
Otu000002	0.0027	0.0015	1.7418	0.0305	0.0292	0.0271	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000004	0.0025	0.0013	1.8750	0.0183	0.0178	0.0358	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000008	0.0020	0.0013	1.4620	0.0032	0.0071	0.0427	Bacteria;Actinobacteria;Actinobacteria;Actinomycetales;Micrococcaceae;unclassified
Otu000005	0.0016	0.0010	1.5955	0.0126	0.0104	0.0484	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000009	0.0015	0.0011	1.3491	0.0054	0.0080	0.0535	Bacteria;Proteobacteria;Betaproteobacteria;Burkholderiales;Oxalobacteraceae;unclassified
Otu000003	0.0014	0.0010	1.3998	0.0174	0.0169	0.0585	Bacteria;Acidobacteria;Acidobacteria_Gp4;Acidobacteria_Gp4_order_incertae_sedis;Acidobacteria_Gp4_family_incertae_sedis;Gp4
Otu000011	0.0014	0.0008	1.7199	0.0061	0.0089	0.0634	Bacteria;Proteobacteria;Betaproteobacteria;Burkholderiales;Comamonadaceae(97);unclassified(93)
Otu000028	0.0013	0.0009	1.4952	0.0051	0.0036	0.0681	Bacteria;unclassified;unclassified;unclassified;unclassified;unclassified;
Otu000049	0.0013	0.0012	1.0088	0.0027	0.0047	0.0725	Bacteria;Acidobacteria;Acidobacteria_Gp7;Acidobacteria_Gp7_order_incertae_sedis;Acidobacteria_Gp7_family_incertae_sedis;Gp7

CSW -C

OTU ID	average	sd	ratio	ava	avb	cusum	taxonomy
Otu000001	0.0037	0.0025	1.4701	0.0256	0.0267	0.0127	Bacteria;Proteobacteria;Alphaproteobacteria;Sphingomonadales;Sphingomonadaceae;unclassified
Otu000003	0.0034	0.0026	1.3177	0.0238	0.0174	0.0242	Bacteria;Acidobacteria;Acidobacteria_Gp4;Acidobacteria_Gp4_order_incertae_sedis;Acidobacteria_Gp4_family_incertae_sedis;Gp4
Otu000004	0.0031	0.0024	1.3097	0.0194	0.0183	0.0349	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000005	0.0025	0.0016	1.5315	0.0119	0.0126	0.0433	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000002	0.0024	0.0018	1.3352	0.0302	0.0305	0.0513	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000006	0.0019	0.0016	1.2433	0.0111	0.0076	0.0579	Bacteria;Verrucomicrobia;Spartobacteria;Spartobacteria_order_incertae_sedis;Spartobacteria_family_incertae_sedis;Spartobacteria_genera_incertae_sedis
Otu000064	0.0018	0.0020	0.9285	0.0037	0.0030	0.0642	Bacteria;Firmicutes;Bacilli;Bacillales;unclassified;unclassified
Otu000025	0.0013	0.0015	0.8854	0.0057	0.0040	0.0687	Bacteria;Bacteroidetes;Sphingobacteria;Sphingobacteriales;Chitinophagaceae;Flavisolibacter
Otu000016	0.0013	0.0008	1.6594	0.0069	0.0069	0.0732	Bacteria;Acidobacteria;Acidobacteria_Gp4;Acidobacteria_Gp4_order_incertae_sedis;Acidobacteria_Gp4_family_incertae_sedis;Gp4
Otu000028	0.0013	0.0008	1.6126	0.0042	0.0051	0.0777	Bacteria;unclassified;unclassified;unclassified;unclassified;unclassified

CSW1cov-
C

OTU ID	average	sd	ratio	ava	avb	cusum	taxonomy
Otu000001	0.0080	0.0029	2.7324	0.0427	0.0267	0.0233	Bacteria;Proteobacteria;Alphaproteobacteria;Sphingomonadales;Sphingomonadaceae;unclassified
Otu000002	0.0053	0.0017	3.2004	0.0198	0.0305	0.0388	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000008	0.0037	0.0026	1.4427	0.0107	0.0032	0.0497	Bacteria;Actinobacteria;Actinobacteria;Actinomycetales;Micrococcaceae;unclassified
Otu000004	0.0029	0.0020	1.4488	0.0143	0.0183	0.0582	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000025	0.0021	0.0018	1.1963	0.0074	0.0040	0.0644	Bacteria;Bacteroidetes;Sphingobacteria;Sphingobacteriales;Chitinophagaceae;Flavisolibacter
Otu000006	0.0018	0.0014	1.2519	0.0108	0.0076	0.0697	Bacteria;Verrucomicrobia;Spartobacteria;Spartobacteria_order_incertae_sedis;Spartobacteria_family_incertae_sedis;Spartobacteria_genera_incertae_sedis
Otu000003	0.0018	0.0012	1.4489	0.0151	0.0174	0.0749	Bacteria;Acidobacteria;Acidobacteria_Gp4;Acidobacteria_Gp4_order_incertae_sedis;Acidobacteria_Gp4_family_incertae_sedis;Gp4
Otu000005	0.0016	0.0009	1.7331	0.0115	0.0126	0.0796	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000064	0.0014	0.0009	1.5663	0.0038	0.0030	0.0838	Bacteria;Firmicutes;Bacilli;Bacillales;unclassified;unclassified
Otu000028	0.0014	0.0010	1.3996	0.0032	0.0051	0.0879	Bacteria;unclassified;unclassified;unclassified;unclassified;unclassified

CSW2cov-C

OTU ID	average	sd	ratio	ava	avb	cumsum	Taxonomy
Otu000001	0.0135	0.0041	3.3287	0.0538	0.0267	0.0354	Bacteria;Proteobacteria;Alphaproteobacteria;Sphingomonadales;Sphingomonadaceae;unclassified
Otu000002	0.0059	0.0022	2.7222	0.0188	0.0305	0.0507	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000004	0.0036	0.0025	1.4686	0.0118	0.0183	0.0602	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000008	0.0036	0.0020	1.7843	0.0104	0.0032	0.0697	Bacteria;Actinobacteria;Actinobacteria;Actinomycetales;Micrococcaceae;unclassified
Otu000003	0.0035	0.0022	1.6243	0.0168	0.0174	0.0789	Bacteria;Acidobacteria;Acidobacteria_Gp4;Acidobacteria_Gp4_order_incertae_sedis;Acidobacteria_Gp4_family_incertae_sedis;Gp4
Otu000006	0.0033	0.0032	1.0351	0.0130	0.0076	0.0876	Bacteria;Verrucomicrobia;Spartobacteria;Spartobacteria_order_incertae_sedis;Spartobacteria_family_incertae_sedis;Spartobacteria_genera_incertae_sedis
Otu000035	0.0026	0.0024	1.0971	0.0072	0.0031	0.0944	Bacteria;Acidobacteria;Acidobacteria_Gp1;Acidobacteria_Gp1_order_incertae_sedis;Acidobacteria_Gp1_family_incertae_sedis;Gp1
Otu000021	0.0022	0.0015	1.4605	0.0086	0.0042	0.1001	Bacteria;Proteobacteria;Alphaproteobacteria;Rhizobiales;unclassified;unclassified
Otu000009	0.0020	0.0010	2.0627	0.0094	0.0054	0.1054	Bacteria;Proteobacteria;Betaproteobacteria;Burkholderiales;Oxalobacteraceae;unclassified
Otu000027	0.0020	0.0014	1.4213	0.0073	0.0039	0.1106	Bacteria;Proteobacteria;Alphaproteobacteria;unclassified;unclassified;unclassified

fallow-C

OTU ID	average	sd	ratio	ava	avb	cusum	taxonomy
Otu000001	0.0029	0.0018	1.6218	0.0237	0.0267	0.0101	Bacteria;Proteobacteria;Alphaproteobacteria;Sphingomonadales;Sphingomonadaceae;unclassified
Otu000004	0.0024	0.0015	1.5848	0.0197	0.0183	0.0185	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000002	0.0022	0.0009	2.3624	0.0261	0.0305	0.0262	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000005	0.0015	0.0009	1.5931	0.0125	0.0126	0.0314	Bacteria;Acidobacteria;Acidobacteria_Gp6;Acidobacteria_Gp6_order_incertae_sedis;Acidobacteria_Gp6_family_incertae_sedis;Gp6
Otu000009	0.0015	0.0007	2.1148	0.0083	0.0054	0.0365	Bacteria;Proteobacteria;Betaproteobacteria;Burkholderiales;Oxalobacteraceae;unclassified
Otu000028	0.0013	0.0009	1.4890	0.0034	0.0051	0.0411	Bacteria;unclassified;unclassified;unclassified;unclassified;unclassified
Otu000003	0.0013	0.0008	1.5674	0.0171	0.0174	0.0456	Bacteria;Acidobacteria;Acidobacteria_Gp4;Acidobacteria_Gp4_order_incertae_sedis;Acidobacteria_Gp4_family_incertae_sedis;Gp4
Otu000016	0.0013	0.0007	1.7643	0.0046	0.0069	0.0500	Bacteria;Acidobacteria;Acidobacteria_Gp4;Acidobacteria_Gp4_order_incertae_sedis;Acidobacteria_Gp4_family_incertae_sedis;Gp4
Otu000020	0.0013	0.0009	1.3536	0.0069	0.0044	0.0544	Bacteria;Proteobacteria;Alphaproteobacteria;Sphingomonadales;Sphingomonadaceae;unclassified
Otu000006	0.0011	0.0008	1.4200	0.0076	0.0076	0.0584	Bacteria;Verrucomicrobia;Spartobacteria;Spartobacteria_order_incertae_sedis;Spartobacteria_family_incertae_sedis;Spartobacteria_genera_incertae_sedis

Table S4. Effects of crop rotation on biocontrol bacterial community (composed of

Streptomyces spp. and *Pseudomonas spp.*) at the KBS Biodiversity Gradient Experimental

Plots based on permutational MANOVA (PERMANOVA) results. Rotation effect was

considered to significantly contribute to community variation at $P < 0.05$.

	df	SS	MS	F	R^2	p-value
Rotation	6	0.768	0.128	1.654	0.321	0.132
Residuals	21	1.626	0.077		0.679	
Total	27	2.394			1	