Title: Fitness costs of herbicide resistance across natural populations of the common

morning glory, *Ipomoea purpurea*

Authors: Megan L. Van Etten[1], Adam Kuester[1], Shu-Mei Chang[2], Regina S Baucom[1]

Author affiliation:

[1] Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor,

MI 48103

[2] Plant Biology Department, University of Georgia, Athens, GA 30602

[3] Corresponding author: Department of Ecology and Evolutionary Biology, University of

Michigan, Ann Arbor, MI 48103; 734-763-8297; mvanette@umich.edu

Running head: Fitness cost of herbicide resistance

Keywords: cost of herbicide resistance, fitness cost, glyphosate, *Ipomoea purpurea*,

germination, early life history, seed quality, trade-offs

Data is archived on Dryad doi:xxx

Abstract

Although fitness costs associated with plant defensive traits are widely expected, they are

not universally detected, calling into question their generality. Here we examine the

potential for life history trade-offs associated with herbicide resistance by examining seed

germination, root growth, and above-ground growth across 43 naturally occurring

populations of *Ipomoea purpurea* that vary in their resistance to RoundUp®, the most

commonly used herbicide worldwide. We find evidence for life history trade-offs

associated with all three traits; highly resistant populations had lower germination rates,

shorter roots and smaller above-ground size. A visual exploration of the data indicated

that the type of trade-off may differ among populations. Our results demonstrate that

costs of adaptation may be present at stages other than simply the production of progeny

in this agricultural weed. Additionally, the cumulative effect of costs at multiple life

cycle stages can result in severe consequences to fitness when adapting to novel

environments.

Introduction

Plant defense is generally hypothesized to involve a cost. This expectation stems

from the surprising observation of genetic variation underlying plant defense traits in

many natural systems, whether the elicitor of damage is an herbivore, a pathogen, or an

herbicide (Simms and Rausher 1987, 1989; Stahl et al. 1999; Baucom and Mauricio

2004; Bakker et al. 2006; Menchari et al. 2006; Délye et al. 2010; Kuester et al. 2015). If

there were no costs associated with defense, traits conferring either resistance or

tolerance to damage should increase to fixation rendering all individuals in the population

highly defended (Rausher and Simms 1989). Despite our expectations of a trade-off

between fitness and defense, however, reviews of the literature consistently show that

costs are not ubiquitous regardless of the elicitor of selection or the study organism at

hand (Bergelson and Purrington 1996; Coustau and Chevillon 2000).

Three main ideas have been proposed to explain the absence of such costs. First,

there are a diverse number of potential mechanisms responsible for adaptation to a

damaging agent, only some of which may incur a cost (Powles and Yu 2010; Vogwill et

al. 2012). A single gene nucleotide substitution that leads to herbicide resistance, for

example, may not alter the efficiency of translated proteins and therefore not incur a cost

(e.g. Yu et al. 2007; Yu et al. 2010). On the other hand, a mechanism that provides

resistance to a range of different herbicides through changes in growth may be more

likely to impose fitness costs. Second, costs may not be detected if the genetic

background is not properly controlled (Bergelson and Purrington 1996; Vila-Aiub et al.

2009b; Vila-Aiub et al. 2011). Control of the genetic background, either by performing

crosses (Baucom and Mauricio 2004; Menchari et al. 2008; Giacomini et al. 2014) or

ensuring replication across multiple genetic backgrounds (Cousens et al. 1997; Strauss et

al. 2002) increases the likelihood that a cost will be detected (Bergelson and Purrington

1996). Third, researchers often examine only a portion of the life cycle (i.e., seed

production or fecundity) and may do so in artificial and/or non-competitive conditions

(Vila-Aiub et al. 2009b; Vila-Aiub et al. 2011). Studies that examine a range of traits are

more likely to identify potential growth and/or fitness differences associated with plant

defense compared to those that focus solely on measures of fecundity (Vila-Aiub et al.

2009b).

The phenomenon of herbicide resistance in plant weeds provides a particularly

useful system to investigate the nature and types of costs associated with plant defense,

since we know when selection by the herbicide began, the strength of selection, and often

the frequency of herbicide use. However, as in other systems examining the evolution of

plant defense, fitness costs of herbicide resistance are often not detected (Bergelson and

Purrington 1996; Gemmill and Read 1998; Vila-Aiub et al. 2009b). Despite

recommendations to control/increase the number of genetic backgrounds (Bergelson and

Purrington 1996), and to examine multiple life history stages when determining if

resistance incurs a cost (Primack and Kang 1989; Vila-Aiub et al. 2009b), only 25% of

herbicide resistance studies control for background effects; further, only 7-10% of cost

studies examine multiple stages of the life cycle (Vila-Aiub et al. 2009b). Fewer still

examine the potential for fitness costs using a large number of naturally occurring

populations sampled from a species' range, an approach suggested almost 20 years ago

(Cousens et al. 1997; Strauss et al. 2002). Just as the mechanism of resistance can vary

among species, populations of the same weed have been shown to harbor different

mechanisms of resistance to the same herbicide (Christopher et al. 1991; Christopher et

al. 1992; Christopher et al. 1994; Preston and Powles 1998; Yu et al. 2008; Délye et al.

2010), thus increasing the likelihood that costs may likewise vary among populations. It

is also possible, though rarely tested, that fitness costs have been ameliorated in some

herbicide resistant populations relative to other populations due to the evolution of

modifier loci (i.e. compensatory evolution, Darmency et al. 2015). The above hypotheses

for the lack of costs are all interrelated: because resistance could be due to a variety of

mechanisms (Délye et al. 2013a), costs may be apparent at only certain life history

stages, expressed in particular environments (Vila-Aiub et al. 2009b), or apparent in

some populations but not others. Thus, there remain crucial gaps in our understanding of

where in the life history of a plant tradeoffs between fitness-enhancing traits and

resistance might be apparent, and further, how ubiquitous such trade-offs may be across a

species' range (Vila-Aiub et al. 2011; Neve et al. 2014).

92   The common morning glory, *Ipomoea purpurea*, a noxious weed of US

agriculture (Webster and MacDonald 2001), provides an excellent system to examine the

strength and type of potential costs that may be present in natural populations. This

species exhibits variability in resistance to glyphosate (Baucom and Mauricio 2008;

Kuester et al. 2015), which is the main ingredient in the herbicide RoundUp®.

RoundUp® is currently the most widely used herbicide in agriculture (Fernandez-

Cornejo et al. 2014), and of the approximately 30 resistant weeds that have been

examined (Heap 2015), only a third are reported to express fitness costs (Ismail et al.

2002; Pedersen et al. 2007; Brabham et al. 2011; Giacomini et al. 2014; Shrestha et al.

2014; Vila-Aiub et al. 2014; Glettner and Stoltenberg 2015; Goh et al. 2015). *I. purpurea*

has long been considered to exhibit low-level resistance to glyphosate (Culpepper

(2006)), and previously we have shown that this low-level resistance (estimated as

proportion leaf damage) has an additive genetic basis and is under positive selection in

the presence of the herbicide (Baucom and Mauricio 2008). Further, a recent replicated

dose-response experiment of 43 populations sampled from the southeastern and Midwest

US showed that some populations of *I. purpurea* exhibit ~100% survival after application

of the field dose of RoundUp® (*i.e.,* resistance), whereas other populations exhibit high

susceptibility (Kuester et al. 2015). Although we find variability in resistance across

natural populations, it is unclear if this defense trait involves a cost. We investigated this

question within one population using artificial selection for increased/decreased

resistance and discovered that the seed production of individuals from the increased

resistance lines was not significantly lower than that of susceptible lines in the absence of

the herbicide, suggesting that there may not be a fecundity cost associated with resistance

in this species. However, there was some indication that progeny quality may be lower in

resistant individuals – resistant lines exhibited a trend for reduced seed viability

compared to susceptible lines (Debban et al. 2015). This finding suggests that trade-offs

between fitness enhancing traits (e.g., germination and resistance) may be present within

this species, which could manifest as a cost by reducing the overall fitness of resistant

compared to susceptible lineages in the absence of herbicide.

Here we determine if there are trade-offs associated with resistance by examining

germination, early root growth and above-ground growth across 43 populations of *I.*

purpurea. We specifically ask the following: (1) are there potential trade-offs associated

with resistance across this species' range in the US, manifest in the form of (i) lower

germination and/or (ii) smaller size at early life history stages (i.e., early germinant,

young plant)?, and (2) do resistant populations exhibit the same type of potential trade-

off, which may indicate the nature and expression of fitness costs may vary across

populations?

Materials and Methods

Seed collection and control of maternal/environmental effects

Multiple fruits were collected from up to 79 individuals separated by at least 2 m from 43

populations located across the Midwest and Southeastern US (Table S1; Fig S1). These

seeds (hereafter field-collected seeds) were used in several experiments to determine

resistance, germination and early growth characteristics. To homogenize the effects of

maternal environment on seed quality, we chose a subset of the populations (N=18), grew

them in a common greenhouse for one generation and collected the autonomously self-

pollinated seeds from a similar growing and mating system environment (hereafter once-

selfed seeds).

Estimate of herbicide resistance

To determine glyphosate resistance across populations, a dose-response experiment was

conducted by planting a single field-collected seed from 10 randomly chosen maternal

lines from each population in six glyphosate treatments (including a non-herbicide

control treatment) in each of two greenhouse rooms. Full details of the dose-response

experiment are presented in Kuester *et al.* (2015) - for simplicity, we present resistance as

the percent survival per population at 1.70 kg a.i./ha of glyphosate, a rate which is

slightly higher than the suggested field rate of 1.54 kg a.i./ha. Individual seeds were

scarified, planted, allowed to grow for three weeks, and then treated with the herbicide

(PowerMax Roundup; Monsanto, St. Louis, Missouri) using a hand-held CO_2 pressurized

sprayer (Spraying Systems Co., Wheaton, IL). Survival was scored three weeks after

treatment application, and the population estimate of resistance was determined as the

proportion of individuals that survived glyphosate.

Germination

We performed three germination experiments to determine if resistance influenced seed

traits. First, we examined germination using field-collected seeds in a petri-dish assay in

the laboratory; second, we examined germination of the field-collected seeds in the soil in

the greenhouse; and third we performed a petri-dish assay in the lab using seeds

generated via selfing in the greenhouse (once-selfed seeds) to examine the potential for

maternal field environmental effects. For the first experiment using field-collected seeds,

we measured seed weight and germination characteristics using field-collected seeds

from each population (N=43). Up to five (ave 4.6) seeds from 8-79 maternal lines per

population (ave 38, total 1621, see Table S1 for exact sample sizes per population) were

randomly chosen for the germination test. From this pool of seeds we randomly chose a

subset of families per population (8-49 maternal lines per population; Table S1) for

which the selected seeds were weighed (as a group) to determine the average seed

weight. All of the selected seeds were placed in a small petri dish (one dish per family),

submerged in filtered water and allowed to germinate in the lab under ambient light and

temperature. Water was added as necessary every three days to prevent drying out. Petri

dishes were completely randomized across lab benches. Germination was scored

periodically until no further germination was recorded. Final pre-scarification

germination was scored after 16 days, with successful germination considered the

emergence of a normal radicle. At this time, seeds that had not imbibed water (by visual

determination) were scarified and germination was again scored after 1 week. We

recorded the final number of seeds exhibiting normal germination, the number of seeds

needing scarification, the number of scarified seeds that germinated, and the number that

had abnormal germination. For the second germination assay, we examined germination

data from one replicate (housed in a single greenhouse room) of the dose-response

experiment mentioned earlier in which seeds were scarified and planted in conetainers

with 1 seed per pot (10 maternal families per population). Germination was scored after

three weeks and used to calculate the percentage of seeds that germinated.

For the third and final germination assay, we used seeds from maternal lines that

were selfed once in the greenhouse. Two sets of five seeds for up to 8 maternal lines

(randomly selected) for each of 18 populations were placed in petri dishes with water.

Germination was scored after 11 days. If seeds had not imbibed water they were scarified

and scored again in one week. We calculated the percentage of seeds with normal

germination prior to scarification, the percentage of seeds that germinated after

scarification, and the percentage that had abnormal germination.

Early root and above-ground growth

To examine early root growth, we again used the once-selfed seeds and measured root

length four days after the germination assay began. We chose to first scarify the seeds in

this assay to standardize water absorption among individuals. Two sets of five seeds for

up to 8 maternal lines (randomly selected) for each of 18 populations were scarified and

placed in petri dishes with water. Germination was scored after 1, 4 and 7 days. On day 4

petri dishes were scanned and the root length was measured using Image J (Abramoff et

al. 2004) for each germinated seed.

We next examined early growth traits of greenhouse-grown individuals to

determine if there was a relationship between resistance and plant size (i.e., are plants

from resistant populations smaller?). To do so we used measurements from plants from

the dose-response experiment prior to herbicide application. Three weeks after planting

(and prior to spraying) we measured the height of the stem, the number of leaves and

length of the largest leaf on each individual planted per treatment per population (total

N=2908, Table S1 for exact sample sizes per population).

Statistical analysis

Field-collected seeds—We assessed the relationship between resistance and progeny

quality using mixed model analyses of variance. We used a generalized linear mixed-

effect model to examine final germination, germination before scarification, abnormal

germination, seeds needing scarification, and germination after scarification with

resistance and population (random) as predictors using the glmer function in the R

package lme4 with a binomial distribution. All of the binary measures were coded as 1 or

0. Seed weight (g) was modeled using a mixed model with resistance and population

(random) as predictors using the lmer function in the R package lme4. Additionally,

previous studies have indicated a geographic pattern of resistance in this species (Kuester

et al. 2015). To ensure that the above results were not an artifact of geography, we added

latitude and longitude (scaled) of the population in the above models. For the experiment

examining germination in soil, we modeled germination with resistance level and

population (random) as predictors using a binomial distribution.

Once-selfed seeds—Similar to the field-collected seeds, we used mixed-model binomial

regressions to assess the effect of resistance on germination characteristics of the once-

selfed, greenhouse generated seeds. We modeled germination before scarification and

germination after scarification with resistance and population (random) as predictors

using a binomial model. To determine if the maternal environment in which the seeds

developed influenced germination, in a separate model we compared germination

between maternal environments (i.e., field collected seeds versus seeds propagated in the

greenhouse) by including maternal environment as a treatment effect in the model. To do

so, we modeled final germination using treatment, resistance, population (random) and

treatment*resistance as predictors using a binomial distribution. An interaction between

treatment and resistance would indicate that the maternal environment differently

influenced germination.

Early growth and size—We next used mixed model analyses of variance to determine if

more resistant populations exhibited early growth life-history trade-offs. We separately

considered root length of the early germinant and plant size. We examined root length

using the once-selfed seeds in two different models. The first and more basic model

examined the influence of resistance and population (random) on log-transformed root

length (cm) 4 days post germination. A difference in root length, however, could be due

to differences in either growth rate of the radicle or differences due to the timing of

germination, *i.e.*, when growth began following germination. To distinguish between

these two potential explanations, we calculated an estimate of germination speed, the

time to 50% germination – a shorter time would suggest that seeds began growing sooner

after water was added. We used the germination data from days 1, 4, and 7 to obtain a

population level estimate of the time to 50% germination using a germination Hill

function (El-Kassaby et al. 2008). This function decomposes germination into 4

parameters: a, the germination capacity; b, the steepness of the curve; c, the time to 50%

germination; and y_0, the lag time before germination. We used the nonlinear least squares

(nls) function in R to estimate the b and c parameters. We chose to pool the data on a

population level to increase the accuracy of the estimation. The time to 50% germination

(c) was then used as a covariate in the more complex model of root length that included

resistance, population (random) and time to 50% germination.

We next examined height (cm), leaf number and leaf size (cm) of plants grown

from field-collected seeds (~3 weeks growth in greenhouse) to determine if resistance

incurs early growth life-history trade-offs. We used each trait in separate mixed models

with replicate, rack within replicate (random), resistance and population (random) as

predictors. Residuals of leaf size were not normal so a box-cox transformation ($\lambda = 2.0$)

was used to achieve better fit.

Finally, we performed a Principle Components Analysis (PCA) using the

population averages of several traits from the field-collected seeds, which included seed

weight, germination percentage, percentage of abnormally germinating seeds, percentage

of successfully germinating scarified seeds, early plant height, leaf number and leaf size

to visually examine the data and determine how populations differed along the two axes

retained. This analysis was performed using PROC FACTOR in SAS with a varimax

rotation to obtain more easily interpretable axes. Loadings and the proportion variance

explained for each factor with an eigenvalue >1 can be found in Table S3.

Results

Germination

We found a strong and significant negative relationship between resistance and

the percentage of field-collected seeds that germinated (Fig. 1a). This is true both of

seeds that germinated before scarification (β = -4.93, χ^2_1 = 24.66 P < 0.0001) and the total

number that germinated (including those that germinated after being scarified; β = -5.20,

χ^2_1 = 24.80, P < 0.0001; Fig. 1a). In addition to a decline in germination, several other

measures of seed quality also declined with increasing resistance. We found a higher

percentage of abnormally germinating seeds (β = 4.24, χ^2_1 = 33.20, P < 0.0001) in that,

instead of exhibiting normal germination, a non-viable embryo would be ejected from the

seed coat with no further growth. Furthermore, some seeds simply did not imbibe water;

we scarified these seeds to determine if they were viable but potentially dormant.

Populations with greater resistance had more seeds that needed scarification (β = 1.52, χ^2_1

= 5.00, P = 0.03), of which fewer seeds that subsequently germinated (β = -5.50, χ^2_1 =

8.81, P = 0.003). We also found that populations with higher resistance produced lighter

seeds (β = -0.005, χ^2_1 = 4.69, P = 0.03), indicating that resistance influenced multiple

measures of seed quality for seeds collected from the field. All of these relationships

remain significant after accounting for longitude and latitude of the populations except

for the percentage needing scarification (Table S2), suggesting that the patterns we find

are not due to a simple geographic pattern. We similarly uncovered a negative

relationship between germination and resistance when seeds from these populations were

planted in soil in the greenhouse (β = -0.79, χ^2_1 = 16.09, P < 0.0001; Fig. 1b).

The negative relationship between resistance and germination was supported by

the results from the once-selfed seeds grown in a common environment for a generation

(Fig. 1c). Prior to scarification, very few of the greenhouse-grown seeds imbibed water

and germinated (2.0%) and there was no effect of resistance (β = 184.2, χ^2_1 = 0.81, P =

0.37). After scarification, however, there was a significant negative relationship between

germination and resistance (β = -1.18, χ^2_1 = 6.42, P = 0.01, Fig. 1c). This effect remained

significant after accounting for latitude and longitude (β = -1.13, χ^2_1 = 5.49, P = 0.02).

Interestingly, the decrease in germination of once-selfed, greenhouse-generated seeds

was significantly less than the field collected seeds (treatment * resistance: β = 2.14, χ^2_1 =

27.8, P < 0.0001) suggesting that maternal environmental conditions influence the quality

of seeds produced. In addition, we found a much lower rate of abnormal germination in

the once-selfed seeds (~10%) compared to the field collected seeds, and the level of

abnormal germination showed no relationship with resistance (β = 0.31, χ^2_1 = 0.23, P =

0.63). These results suggest that, while germination costs are consistently detected

between experiments in which the maternal environment differed, field environmental

conditions exacerbate the strength of the germination cost.

Early root and aboveground growth

To test whether growth differed according to resistance status, we scarified and

germinated the once-selfed seeds then measured root growth after 4 days. There was a

much higher germination rate of these seeds (86%) compared to the previous experiment

(2% pre-scarification) and the majority occurred before day 4. While there was a nearly

significant negative relationship between root length and resistance (β = -0.38, χ^2_1 = 3.19,

P = 0.07; Fig. 2a), including the time to 50% germination in the model removed this

effect (resistance: β = -0.07, χ^2_1 = 0.09, P = 0.76; time to 50% germination: β = -0.37, χ^2_1

= 4.85, P = 0.03), suggesting that the difference in root length was due to the timing of

germination rather than a difference in growth rate. A difference in plant size was also

found in the 3-4 week old plants grown in soil from the field-collected seeds - plants from

more resistant populations had smaller above-ground structures on average than plants

from less resistant populations (height: $\beta = -7.42$, $\chi^2_1 = 7.20$, P = 0.007; leaf number: $\beta =$

-1.11, $\chi^2_1 = 15.86$, P < 0.0001; largest leaf width: $\beta = -4.97$, $\chi^2_1 = 17.32$, P < 0.0001; Fig.

2b-d).

Visualization of cost-related traits

We next examined germination and early growth traits from the original field-collected

seeds using a principle components analysis (PCA) to determine if there was variation

among populations in the expression of cost-related traits (full results Table S3). The first

3 principle components of the PCA explained 77% of the variance, with the first principle

component (PC) loading with the early growth traits while the second loaded with seed

traits and the third with the proportion of seeds that required scarification to successfully

germinate. Populations with higher resistance scored lower on PC1 (b = -2.22, $r^2 = 0.28$,

$t_{41} = -4.03$, P = 0.0002) and PC2 (b = -2.12, $r^2 = 0.26$, $t_{41} = -3.79$, P = 0.0005), but not on

PC3 (b = -1.02, $r^2 = 0.06$, $t_{41} = -1.61$, P = 0.12). Using the first two PCs to plot the results,

resistant populations occur mostly in the lower left quadrant (smaller plants, lighter seeds,

lower germination and more abnormally germinating seeds) and have a wider spread than

less resistant populations (Fig. 3). Furthermore, while resistant populations scored lower

on PC1 and PC2 in comparison to susceptible populations, some resistant populations

exhibited early growth traits that were similar to susceptible populations and yet scored

very low on germination traits (e.g., pop num 5) whereas other resistant populations

exhibited similar germination traits compared to the susceptible populations, but were

smaller in stature than susceptible populations (e.g., pop num 51). Thus, it appears that

the type of cost may vary among populations sampled from North America.

Discussion

Here we show that glyphosate resistant populations of the common morning glory

exhibit life-history trade-offs associated with resistance, and, that these trade-offs may

vary among populations. Our series of experiments uncovered three notable findings:

First, we found a negative linear relationship between germination and resistance

indicating that resistant populations have a lower germination rate than susceptible

populations. This negative relationship persisted when using seeds generated from a

common greenhouse environment showing that this result is not due solely to field

environmental and/or maternal effects. Second, we found that individuals from resistant

populations were smaller than individuals from susceptible populations, indicating that

resistance influences early plant growth. Interestingly, we also found evidence that the

two types of trade-off may differ among populations—using PCA, we show that some

resistant populations produce normally sized plants, but score low on germination traits,

and *vice versa*. Below, we detail how these results add further strength to the suggestion

that a variety of life stages and populations sampled across the species' range should be

assessed when testing the hypothesis that resistance incurs a fitness cost (Délye et al.

2013a).

Fitness costs: seed germination and early plant size

It is difficult to determine how common germination differences associated with

resistance may be among weeds since many studies focus on seed quantity rather than

seed quality. There is some indication that germination may be affected in other

glyphosate resistant species. Dinelli et al. (2013) found reduced germination of

glyphosate resistant *Ambrosia trifida* populations, while Ismail et al. (2002) found greater

germination of resistant biotypes of goosegrass (*Eleusine indica)*. More broadly, life

history trade-offs may be specific to the herbicide and/or species in question or the type

of mutation conferring resistance (O'Donovan et al. 1999; Vila-Aiub et al. 2005; Délye et

al. 2013b). For example, only one of two different resistance mutations in ACCase

resistant *Lolium rigidum* had more stringent germination requirements (seeds germinated

poorly in the dark and required fluctuating temperatures to break dormancy) than the

susceptible genotype (Vila-Aiub et al. 2005). Similarly, Délye *et al.* (2013b) found

differential effects on germination among resistance mutations to ACCase in *Alopecurus*

myosuroides. Both of these studies report that the resistance mutation led to delayed

germination. Such a delay in germination may affect fitness, especially in agricultural

settings where germinating too early can lead to removal by pre-sowing practices and

germinating too late can lead to intensified competition with already established plants

(Weaver and Cavers 1979; Barrett 1983; Mortimer 1997; Forcella et al. 2000; Owen et al.

2014). Our analysis of root growth suggests that differences in plant size in *I. purpurea*

may be due to a similar delay in germination in resistant populations.

The reduced growth of resistant compared to susceptible populations that we

uncovered could lead to decreased competitive ability and subsequent lower fitness in the

presence of competition if, as has been found in other herbicide resistant weeds, the

difference in growth persists to adult plants (Weaver and Warwick 1982; Ahrens and

Stoller 1983; Holt 1988; Alcocer-Ruthling et al. 1992; Williams et al. 1995; Vila-Aiub et

al. 2005; Tardif et al. 2006; Vila-Aiub et al. 2009a). This type of life-history trade-off,

which ultimately may manifest as a fitness cost, is also likely to be species, mutation and

environment specific. For example, *Lolium rigidum* has evolved herbicide resistance via

a variety of mutations ranging from target site (Christopher et al. 1992; Yu et al. 2008) to

non-target site (Christopher et al. 1991; Christopher et al. 1994; Preston and Powles

1998). Target site mutations in the acetohydroxyacid synthase gene result in little cost in

growth (Yu et al. 2010). On the other hand, herbicide resistance mediated by the

cytochrome P450 complex resulted in reduced biomass and decreased competitive ability

(Vila-Aiub et al. 2009a).

An alternative explanation for the decline in germination and growth we identify

using the field-collected seeds is that some other co-varying population characteristic

such as soil fertility, spraying regime, herbivore levels, or the many other biotic and

abiotic factors that can influence seed development differed among resistant and

susceptible populations (Roach and Wulff 1987; Fenner 1991; Schmitt et al. 1992;

Platenkamp and Shaw 1993; Galloway 2001). These differences may explain the stronger

decline in germination in the field-collected seeds compared to the once-selfed seeds.

Several lines of evidence, however, suggest the relationship between resistance and

germination that we uncovered represents a true fitness cost rather than simply an effect

of the environment (e.g., driven by maternal effects, latitude, spray environment). First,

the relationship between resistance and germination appears approximately linear, which

is expected as the frequency of resistant individuals increases. If the trade-offs identified

herein were due simply to glyphosate exposure, with resistant populations exhibiting

abnormal seed development after surviving glyphosate application, we would expect to

see a binary distribution of seed quality of populations that had been sprayed and those

that had not, rather than a linear trend with resistance. Second, the negative relationship

between resistance and germination is maintained after a generation in a common

greenhouse environment—an effect should disappear if the decrease in fitness was due to

glyphosate exposure in the field. Finally, our results parallel those from a recent

experiment that specifically controlled for genetic background and environmental effects

using *I. purpurea* plants from a single population (Debban et al. 2015). Individuals from

this population were artificially selected for increased or decreased resistance for three

generations under controlled greenhouse conditions, and, similar to results presented

here, the increased resistance lines had a larger percentage of "bad" seeds that ejected the

embryo. That the results from one population utilizing a controlled genetic background

are mirrored across many populations collected from the landscape provides strong

evidence that lower germination represents a fitness cost of glyphosate resistance in this

weed species.

While we detected lower germination in herbicide resistant populations across

multiple experiments suggesting a true trade-off, we also found differences between

experiments in the strength of the relationship. This suggests both an underlying genetic

basis as well as an environmental component influence the expression of the trade-off.

Compared to the field-collected, unscarified seeds, the once-selfed, unscarified seeds

(Fig. 1) had a low germination rate suggesting that the seed coat was perhaps more

pristine in seeds generated in the greenhouse. However, if scarified, we found that the

once-selfed seeds exhibited high germination and the relationship between resistance and

germination remained negative indicating a fitness decline associated with resistance.

The physical seed coat is the primary mechanism of dormancy in this species (Brechu-

Franco et al. 2000); thus, environmentally-induced physical differences in the seed coat

(e.g. thickness or waxiness) or its degree of degradation (e.g. mechanical disruption or

seed storage differences) likely influences germination timing. Although it is clear that

the environment influences seed germination in this species, that we consistently

observed a decline in germination with resistance across multiple experiments suggests

that there is an underlying genetic basis to the cost of resistance.

Another striking difference between experiments was in the frequency of

abnormal seeds produced. The once-selfed seeds had almost no abnormal germination

(i.e., no dead embryos that were ejected from the seed coat) while some of the field-

collected populations had a high level of abnormal germination. In fact, the strong

decline in germination for field-collected seeds was due primarily to this abnormal

germination. Abnormal germination could be due to a variety of environmental causes

(e.g. herbicide application, nutrient availability, competition, etc) or be a cost of

resistance that is only induced under field conditions. Our results suggest that in a benign

environment, such as the greenhouse, the seeds in general are of high quality (high

germination, fewer abnormal germinants) but there is a cost of resistance that increases

the time it takes to germinate (based on root growth experiment), possibly leading to

smaller plants at any given point. On the other hand, under field conditions, populations

with higher resistance produce more abnormal seeds (due to either environmental

differences or an environmentally induced cost of resistance) and the normal seeds may

still have an increase in the time it takes to germinate leading to smaller plants at any

given point.

Interestingly, by visually examining the germination and growth traits in a PCA,

we find variation in the type of potential cost among resistant populations. While some

resistant populations fell into the "poor germination" axis, other resistant populations fell

into the "poor growth" axis compared to susceptible populations. There are at least three

possible reasons for this difference: different resistance genes, different compensatory

mutations or different genetic backgrounds. First, the gene(s) involved in resistance may

vary among populations leading to different costs. Independent origins of resistance to

herbicide have been found in other species (Délye et al. 2010) and these different

mutations often incur different fitness costs (Vila-Aiub et al. 2005; Délye et al. 2013b).

Second, the resistance gene(s) may be the same amongst populations but each population

may have different compensatory mutations that lead to different costs (Darmency et al.

2015). Third, the resistance gene(s) may behave differently in different genetic

backgrounds (Paris et al. 2008). These distinctions are important because they would

differentially affect the evolutionary trajectory of herbicide resistance. For example, if

populations differ in the gene(s) involved, each population may have a very different set

of costs, benefits and evolutionary trajectories, which would need to be incorporated in

models.

It is currently unknown if the trait trade-offs identified here are pleiotropic or due

to linkage to the resistance gene. The most restrictive definition of a cost requires that the

decrease in fitness is due to the resistance allele itself - either the actual allele or through

it acting pleiotropically (Bergelson and Purrington 1996). Given that we do not know the

identity of the loci involved in either resistance or the abnormal germination and reduced

growth, we cannot entirely rule out physical linkage between resistance genes and cost

genes, in which case the "cost" could quickly become unlinked over generations

(Lewontin 1974; Hartl and Clark 1989). For some species, easily identifiable mutations in

the enzyme targeted by the herbicide can be linked to resistance, i.e. target site resistance

(TSR). However, preliminary work suggests that glyphosate resistance in *I. purpurea* is

due to non-target site mechanism (NTSR; Leslie and Baucom, *unpublished data*), and as

such elucidating the genetic basis of both resistance and the cost will be a non-trivial

endeavor. Furthermore, it is rare that genes underlying costs are identified; most

documented cases of the genes involved in the cost of resistance is when TSR mutations

lead to poor performance of the enzyme on its natural substrate (Vila-Aiub et al. 2009b).

As far as we are aware, no study has identified the genes involved in the cost of

resistance when the mechanism of resistance is NTSR. One intriguing possibility for this

species stems from a previous study that compared transcript expression levels of

artificially selected lines of resistant and susceptible *I. purpurea* plants following

herbicide application (Leslie and Baucom 2014). One of the differences between the

replicated resistant and susceptible lines was a lower expression of pectin methylesterase

(PME) in the resistant plants. This enzyme has been shown to play a role in breaking seed

dormancy (Ren and Kermode 2000) and stem elongation (Pilling et al. 2000). Thus, the

decreased expression of PME in resistant plants may explain both the reduced

germination and growth in populations with higher resistance.

How might life history trade-offs influence the evolutionary trajectory of resistance in

this species?

Recent work in this system has shown that populations of the common morning

glory sampled from 2012 exhibit higher levels of resistance compared to the same

populations sampled in 2003 (Kuester et al, *In* Review). Interestingly, however, the

difference in resistance between sampling years was only slight, i.e., 62% survival at 1.7

kg ai/ha in 2012 vs 57% survival in 2003 samples. It is possible that the life-history

differences that we identified here are responsible, at least in part, for maintaining

resistance between sampling years. For example, the lower germination of resistant types

would manifest as a fitness cost if resistant and susceptible types produce approximately

the same number of total seeds (or if R < S); if, however, resistant types produce enough

viable seed to offset the lowered germination, then overall fitness would not be impacted

and resistant types would not be at a relative disadvantage. While we have not examined

seed production across all 43 populations examined herein, a common garden study of

glyphosate susceptible and resistant families of this species found there was no difference

in total seed production of resistant compared to susceptible lines (Debban et al 2015),

indicating there is no cost of resistance in terms of seed quantity. That we similarly find

poor germination between these experiments and those using genetic lines developed

from a single population strongly supports the finding that germination quality is a true

fitness cost of glyphosate resistance in this species. Further, the differences in growth that

we have detected between resistant and susceptible populations could potentially

manifest as a fitness cost if in competition, an effect which remains to be tested in this

system.

In summary, we found reductions in seed quality across replicated herbicide

resistant populations of the common morning glory. Although most studies use seed

quantity as a proxy for fitness, our results highlight that reductions in progeny quality are

an equally, if not more, important cost of adaptation in *I. purpurea*. Given that fitness

costs are thought to arise from a variety of mechanisms (allocation of resources,

ecological costs, etc.), our results suggest that a high priority should be placed on the

examination of multiple stages of the life cycle when assessing potential costs and not

just seed quantity. Furthermore, because the strength of this cost could maintain the

efficacy of a globally important herbicide, this work illustrates the utility and importance

of integrating evolutionary principles into management scenarios (Gould 1995).

Acknowledgements: We thank E. Fall, A Wilson, A Jankowiak, D York, N Gabry, S

Sanchez for assistance and J Vandermeer, MA Duffy, and PJ Tranel for comments on

earlier drafts of the manuscript. This work was funded by USDA NIFA grants 04180 and

07191 to RSB and SMC.

 Figure Legends

Fig 1. Relationship between herbicide resistance and proportion germination for (a) field-

collected seeds in petri dishes, (b) field-collected seeds in soil, and (c) once-selfed seeds

(note the differences in the y-axis scale). Points are the mean per population, lines are the

average marginal predicted probabilities from the appropriate model.

Fig 2. Relationship between herbicide resistance and (a) root growth after 4 days, (b)

shoot height (c) leaf number and (d) width of the largest leaf after 3 weeks. Points are the

mean per population and lines are the average marginal predicted probabilities from the

appropriate model.

Fig 3. Scatter plot of PCA results showing average principle components axis 1 (higher

values indicate larger plants) and principle components axis 2 (higher values indicate

heavier seeds, higher germination, and fewer abnormally germinating seeds) values for

field-collected seeds with circle color indicating survival at 1.7 kg/ha glyphosate. The

proportion variation explained by each axis is noted in axis labels.

 Supporting information

Table S1. Population characteristics including population number, state, crop type,

latitude and longitude and sample sizes for experiments with field-collected seeds

(N_{fam} germ = number of families in the germination assay for field collected seeds;

N_{seeds} germ = number of seeds in the germination assay for field collected seeds; N

early growth = number of seeds in the early growth experiment, N_{fam} once selfed =

number of families in the germination assays for the once-selfed seeds).

Population number	State	Crop	Latitude	Longitude	N_{fam} germ	N_{seeds} germ	N early growth	N_{fam} once selfed
1	TN	Corn	35.775237	-85.903419	8	31	10	4
2	NC	Corn	34.595714	-77.927484	79	387	60	
4	NC	Corn	34.556672	-79.125602	39	178	79	
5	SC	Soy	33.859875	-79.909072	34	146	27	
8	SC	Corn	34.297195	-79.991259	40	180	73	8
9	NC	Soy	34.924044	-77.796171	40	189	46	8
10	NC	Soy	34.983161	-78.039309	28	123	24	8
11	NC	Corn	34.527135	-78.756704	38	178	81	8
12	SC	Cotton	34.145812	-79.865313	42	202	79	8
14	NC	Soy	35.424763	-77.917121	40	192	76	8
15	SC	Soy	34.104209	-79.073735	41	193	81	8
16	SC	Alfalfa	34.10535	-79.183234	41	203	78	
17	SC	Soy	34.159155	-79.272908	27	135	82	
18	SC	Corn	34.156593	-79.27027	39	193	79	
19	NC	Corn	34.508193	-78.70899	35	138	70	8
20	TN	Corn	35.830692	-85.777871	38	169	13	2

21	TN	Soy	35.369816	-77.877314	40	200	70	
22	NC	Corn	36.1436	-78.053422	17	78	69	
23	TN	Corn	35.067905	-86.62955	40	195	77	
26	TN	Soy	35.533413	-85.951902	50	250	86	
28	SC	Corn	34.097917	-80.377715	35	172	80	
29	NC	Corn	34.705135	-78.738897	30	122	75	7
30	TN	Corn	35.31105	-85.945003	42	206	82	
31	TN	Corn	35.608482	-85.846379	31	141	69	8
32	TN	Corn	35.099356	-86.225509	15	56	33	5
33	OH	Corn	39.858763	-83.669821	52	260	79	
34	OH	Corn	39.44316	-83.910189	48	238	82	
35	IN	Corn	39.853945	-85.770156	28	120	80	
36	IN	Corn	40.565608	-85.503826	33	146	85	
37	OH	Corn	39.583755	-83.758264	49	243	79	
38	OH	Soy	39.515447	-83.407431	46	230	88	
39	IN	Corn	39.988984	-85.742262	38	188	75	
40	OH	Soy	41.284684	-83.847252	50	237	84	
41	VA	Corn	38.636343	-78.472921	41	200	78	
42	VA	Corn	38.373523	-78.662516	49	230	82	8
43	VA	Soy	36.886448	-78.553156	24	89	28	2
44	VA	Soy	38.285415	-78.797088	20	98	75	
45	VA	Soy	36.847945	-78.595042	42	209	76	
46	TN	Soy	35.536019	-86.17985	34	100	17	2
47	SC	Soy	34.282132	-79.746597	41	201	83	
48	TN	Corn	35.31653	-87.35373	31	151	74	8
51	TN	Soy	35.533413	-85.951902	24	118	56	8
52	OH	Corn	41.284684	-83.847252	62	310	88	

564   Table S2. Regression results between resistance and seed quality traits, accounting for geography (latitude and longitude) for field-

565   collected seeds.

Trait	Population number	Survival at 1.7 kg a.i. /ha		Latitude (scaled)		Longitude (scaled)	
		b	χ^2_1	b	χ^2_1	b	χ^2_1
Seed weight (g)	43	-0.0047	4.63*	0.0016	7.33**	-0.0010	3.91*
Final germination	43	-4.89	28.62***	0.6298	8.80**	-0.3784	4.13*
Abnormally germinating seeds	42	4.24	32.52***	-0.1774	1.24	0.1826	1.58
Seeds needing scarification	43	0.83	1.77	-0.4636	7.19**	-0.3488	5.22*
Scarified seeds that germinated	40	-5.87	8.85**	0.0061	0.0002	-0.3342	0.65
Height (cm)	43	-7.39	7.26**	-0.3336	0.38	1.2210	4.76*
Leaf number	43	-1.08	17.57***	-0.0070	0.02	0.1589	9.17**
Largest leaf size (Box cox transformed cm)	43	-4.40	14.04***	0.4080	2.96	0.4642	3.81

567 Table S3. PCA loadings and variance explained.

Trait	Factor1	Factor2	Factor3
Seed weight	0.09506	0.8414	-0.16627
Percent germination	0.29276	0.82272	0.37541
Percent viable dormant seeds	0.0339	0.12801	0.92794
Percent abnormally germinating seeds	-0.15651	-0.81953	-0.42013
Height	0.94253	0.04143	0.05526
Leaf number	0.86728	0.13535	0.06511
Largest leaf size	0.74853	0.28353	0.00642
Prop. Variance explained	0.3316	0.3105	0.1734

569 Fig S1. Map of the Eastern US showing the population locations and their survival at

570 1.7 kg a.i./ha (color of circle).

572   **References**

573   Abramoff, M. D., P. J. Magelhaes, and S. J. Ram. 2004. Image Processing with ImageJ.
574   Biophotonics International 11:36-42.
575   Ahrens, W. H. and E. Stoller. 1983. Competition, growth rate, and CO_2 fixation in
576   triazine-susceptible and-resistant smooth pigweed (*Amaranthus hybridus*).
577   Weed Science 31:438-444.
578   Alcocer-Ruthling, M., D. C. Thill, and B. Shafii. 1992. Differential competitiveness of
579   sulfonylurea resistant and susceptible prickly lettuce (*Lactuca serriola*).
580   Weed Technology 6:303-309.
581   Bakker, E. G., C. Toomajian, M. Kreitman, and J. Bergelson. 2006. A genome-wide
582   survey of *R* gene polymorphisms in *Arabidopsis*. The Plant Cell 18:1803-1818.
583   Barrett, S. H. 1983. Crop mimicry in weeds. Economic Botany 37:255-282.
584   Baucom, R. S. and R. Mauricio. 2004. Fitness costs and benefits of novel herbicide
585   tolerance in a noxious weed. Proceedings of the National Academy of
586   Sciences of the United States of America 101:13386-13390.
587   Baucom, R. S. and R. Mauricio. 2008. Constraints on the evolution of tolerance to
588   herbicide in the common morning glory: resistance and tolerance are
589   mutually exclusive. Evolution 62:2842-2854.
590   Bergelson, J. and C. B. Purrington. 1996. Surveying patterns in the cost of resistance
591   in plants. American Naturalist 148:536-558.
592   Brabham, C. B., C. K. Gerber, and W. G. Johnson. 2011. Fate of glyphosate-resistant
593   giant ragweed (*Ambrosia trifida)* in the presence and absence of glyphosate.
594   Weed Science 59:506-511.
595   Brechu-Franco, A., R. Ponce-Salazar, G. Laguna-Hernandez, and J. Marquez-Guzman.
596   2000. Effect of thermal storage on seed coat dormancy and germination of
597   *Ipomoea purpurea* (L.) Roth (Convolvulaceae) seeds. Phyton 67:187-194.
598   Christopher, J. T., S. B. Powles, and J. A. Holtum. 1992. Resistance to acetolactate
599   synthase-inhibiting herbicides in annual ryegrass (*Lolium rigidum*) involves
600   at least two mechanisms. Plant Physiology 100:1909-1913.
601   Christopher, J. T., S. B. Powles, D. R. Liljegren, and J. A. Holtum. 1991. Cross-
602   resistance to herbicides in annual ryegrass (*Lolium rigidum*) II. chlorsulfuron
603   resistance involves a wheat-like detoxification system. Plant Physiology
604   95:1036-1043.
605   Christopher, J. T., C. Preston, and S. B. Powles. 1994. Malathion antagonizes
606   metabolism-based chlorsulfuron resistance in *Lolium rigidum*. Pesticide
607   Biochemistry and Physiology 49:172-182.
608   Cousens, R., G. Gill, and E. J. Speijers. 1997. Comment: number of sample populations
609   required to determine the effects of herbicide resistance on plant growth and
610   fitness. Weed Research 37:1-4.
611   Coustau, C. and C. Chevillon. 2000. Resistance to xenobiotics and parasites: can we
612   count the cost? Trends in Ecology & Evolution 15:378-383.
613   Culpepper, A. S. 2006. Glyphosate-induced weed shifts. Weed Technology 20:277-
614   281.

Darmency, H., Y. Menchari, V. Le Corre, and C. Délye. 2015. Fitness cost due to
herbicide resistance may trigger genetic background evolution. Evolution
69:271-278.
Debban, C., S. Matthews, L. Elliot, K. Pieper, A. Wilson, and R. S. Baucom. 2015. An
examination of fitness costs of glyphosate resistance in the common morning
glory, *Ipomoea purpurea*. Ecology and Evolution 5:5284-5294.
Délye, C., M. Jasieniuk, and V. Le Corre. 2013a. Deciphering the evolution of
herbicide resistance in weeds. Trends in Genetics 29:649-658.
Délye, C., Y. Menchari, S. Michel, É. Cadet, and V. Le Corre. 2013b. A new insight into
arable weed adaptive evolution: mutations endowing herbicide resistance
also affect germination dynamics and seedling emergence. Annals of botany
111:681-691.
Délye, C., S. Michel, A. Bérard, B. Chauvel, D. Brunel, J. P. Guillemin, F. Dessaint, and V.
Le Corre. 2010. Geographical variation in resistance to acetyl‐coenzyme A
carboxylase‐inhibiting herbicides across the range of the arable weed
Alopecurus myosuroides (black‐grass). New Phytologist 186:1005-1017.
Dinelli, G., I. Marotti, P. Catizone, S. Bosi, A. Tanveer, R. Abbas, and D. Pavlovic. 2013.
Germination ecology of *Ambrosia artemisiifolia* L. and *Ambrosia trifida* L.
biotypes suspected of glyphosate resistance. Open Life Sciences 8:286-296.
El-Kassaby, Y. A., I. Moss, D. Kolotelo, and M. Stoehr. 2008. Seed germination:
mathematical representation and parameters extraction. Forest Science
54:220-227.
Fenner, M. 1991. The effects of the parent environment on seed germinability. Seed
Science Research 1:75-84.
Fernandez-Cornejo, J., R. Nehring, C. Osteen, S. Wechsler, A. Martin, and A. Vialou.
2014. Pesticide use in U.S. agriculture: 21 selected crops, 1960-2008, EIB-
124. U.S. Department of Agriculture, Economic Research Service.
Forcella, F., R. L. B. Arnold, R. Sanchez, and C. M. Ghersa. 2000. Modeling seedling
emergence. Field Crops Research 67:123-139.
Galloway, L. F. 2001. The effect of maternal and paternal environments on seed
characters in the herbaceous plant *Campanula americana* (Campanulaceae).
American Journal of Botany 88:832-840.
Gemmill, A. W. and A. F. Read. 1998. Counting the cost of disease resistance. Trends
in Ecology & Evolution 13:8-9.
Giacomini, D., P. Westra, and S. M. Ward. 2014. Impact of genetic background in
fitness cost studies: an example from glyphosate-resistant *Palmer amaranth*.
Weed Science 62:29-37.
Glettner, C. E. and D. E. Stoltenberg. 2015. Noncompetitive growth and fecundity of
Wisconsin giant ragweed resistant to glyphosate. Weed Science 63:273-281.
Goh, S. S., M. M. Vila‐Aiub, R. Busi, and S. B. Powles. 2015. Glyphosate resistance in
Echinochloa colona: phenotypic characterisation and quantification of
selection intensity. Pest management science.
Gould, F. 1995. Comparisons between resistance management strategies for insects
and weeds. Weed Technology:830-839.

Hartl, D. L. and A. G. Clark. 1989. Principles of population genetics. Sinauer
associates, Sunderland, Mass.
Heap, I. 2015. The International survey of herbicide resistant weeds.
Holt, J. 1988. Reduced growth, competitiveness, and photosynthetic efficiency of
triazine-resistant *Senecio vulgaris* from California. Journal of Applied Ecology
25:307-318.
Ismail, B., T. Chuah, S. Salmijah, Y. Teng, and R. Schumacher. 2002. Germination and
seedling emergence of glyphosate‐resistant and susceptible biotypes of
goosegrass (*Eleusine indica* [L.] Gaertn.). Weed Biology and Management
2:177-185.
Kuester, A., S. M. Chang, and R. S. Baucom. 2015. The geographic mosaic of herbicide
resistance evolution in the common morning glory, *Ipomoea purpurea*:
Evidence for resistance hotspots and low genetic differentiation across the
landscape. Evolutionary Applications 8:821-833.
Leslie, T. and R. S. Baucom. 2014. De novo assembly and annotation of the
transcriptome of the agricultural weed *Ipomoea purpurea* uncovers gene
expression changes associated with herbicide resistance. G3: Genes|
Genomes| Genetics 4:2035-2047.
Lewontin, R. C. 1974. The genetic basis of evolutionary change. Columbia University
Press, New York.
Menchari, Y., C. Camilleri, S. Michel, D. Brunel, F. Dessaint, V. Le Corre, and C. Délye.
2006. Weed response to herbicides: regional‐scale distribution of herbicide
resistance alleles in the grass weed *Alopecurus myosuroides*. New Phytologist
171:861-874.
Menchari, Y., B. Chauvel, H. Darmency, and C. Délye. 2008. Fitness costs associated
with three mutant acetyl‐coenzyme A carboxylase alleles endowing
herbicide resistance in black‐grass *Alopecurus myosuroides*. Journal of
Applied Ecology 45:939-947.
Mortimer, A. M. 1997. Phenological adaptation in weeds—an evolutionary response
to the use of herbicides? Pesticide Science 51:299-304.
Neve, P., R. Busi, M. Renton, and M. M. Vila‐Aiub. 2014. Expanding the eco‐
evolutionary context of herbicide resistance research. Pest Management
Science 70:1385-1393.
O'Donovan, J., J. Newman, R. Blackshaw, K. Harker, D. Derksen, and A. Thomas. 1999.
Growth, competitiveness, and seed germination of triallate/difenzoquat-
susceptible and-resistant wild oat populations. Canadian Journal of Plant
Science 79:303-312.
Owen, M. J., D. E. Goggin, and S. B. Powles. 2014. Intensive cropping systems select
for greater seed dormancy and increased herbicide resistance levels in
Lolium rigidum (annual ryegrass). Pest Management Science.
Paris, M., F. Roux, A. Berard, and X. Reboud. 2008. The effects of the genetic
background on herbicide resistance fitness cost and its associated dominance
in *Arabidopsis thaliana*. Heredity 101:499-506.

Pedersen, B. P., P. Neve, C. Andreasen, and S. B. Powles. 2007. Ecological fitness of a
glyphosate-resistant *Lolium rigidum* population: Growth and seed production
along a competition gradient. Basic and Applied Ecology 8:258-268.
Pilling, J., L. Willmitzer, and J. Fisahn. 2000. Expression of a *Petunia inflata* pectin
methyl esterase in *Solanum tuberosum* L. enhances stem elongation and
modifies cation distribution. Planta 210:391-399.
Platenkamp, G. A. and R. G. Shaw. 1993. Environmental and genetic maternal effects
on seed characters in *Nemophila menziesii*. Evolution 47:540-555.
Powles, S. B. and Q. Yu. 2010. Evolution in action: plants resistant to herbicides.
Annual Review of Plant Biology 61:317-347.
Preston, C. and S. B. Powles. 1998. Amitrole inhibits diclofop metabolism and
synergises diclofop-methyl in a diclofop-methyl-resistant diotype of *Lolium*
rigidum. Pesticide Biochemistry and Physiology 62:179-189.
Primack, R. B. and H. Kang. 1989. Measuring fitness and natural selection in wild
plant populations. Annual Review of Ecology and Systematics 20:367-396.
Rausher, M. D. and E. L. Simms. 1989. The evolution of resistance to herbivory in
Ipomoea purpurea. I. Attempts to detect selection. Evolution 43:563-572.
Ren, C. and A. R. Kermode. 2000. An increase in pectin methyl esterase activity
accompanies dormancy breakage and germination of yellow cedar seeds.
Plant Physiology 124:231-242.
Roach, D. A. and R. D. Wulff. 1987. Maternal effects in plants. Annual Review of
Ecology and Systematics 18:209-235.
Schmitt, J., J. Niles, and R. D. Wulff. 1992. Norms of reaction of seed traits to maternal
environments in *Plantago lanceolata*. American Naturalist 139:451-466.
Shrestha, A., K. M. Steinhauer, M. L. Moretti, B. D. Hanson, M. Jasieniuk, K. J. Hembree,
and S. D. Wright. 2014. Distribution of glyphosate-resistant and glyphosate-
susceptible hairy fleabane (*Conyza bonariensis*) in central California and their
phenological development. Journal of Pest Science 87:201-209.
Simms, E. L. and M. D. Rausher. 1987. Costs and benefits of plant resistance to
herbivory. American Naturalist 130:570-581.
Simms, E. L. and M. D. Rausher. 1989. The evolution of resistance to herbivory in
Ipomoea purpurea. II. Natural selection by insects and costs of resistance.
Evolution 43:573-585.
Stahl, E. A., G. Dwyer, R. Mauricio, M. Kreitman, and J. Bergelson. 1999. Dynamics of
disease resistance polymorphism at the *Rpm1* locus of *Arabidopsis*. Nature
400:667-671.
Strauss, S. Y., J. A. Rudgers, J. A. Lau, and R. E. Irwin. 2002. Direct and ecological costs
of resistance to herbivory. Trends in Ecology & Evolution 17:278-285.
Tardif, F. J., I. Rajcan, and M. Costea. 2006. A mutation in the herbicide target site
acetohydroxyacid synthase produces morphological and structural
alterations and reduces fitness in *Amaranthus powellii*. New Phytologist
169:251-264.
Vila-Aiub, M. M., S. S. Goh, T. A. Gaines, H. Han, R. Busi, Q. Yu, and S. B. Powles. 2014.
No fitness cost of glyphosate resistance endowed by massive EPSPS gene
amplification in *Amaranthus palmeri*. Planta 239:793-801.

Vila-Aiub, M. M., P. Neve, and S. B. Powles. 2009a. Evidence for an ecological cost of
enhanced herbicide metabolism in *Lolium rigidum*. Journal of Ecology
97:772-780.
Vila-Aiub, M. M., P. Neve, and S. B. Powles. 2009b. Fitness costs associated with
evolved herbicide resistance alleles in plants. New Phytologist 184:751-767.
Vila-Aiub, M. M., P. Neve, and F. Roux. 2011. A unified approach to the estimation
and interpretation of resistance costs in plants. Heredity 107:386-394.
Vila-Aiub, M. M., P. Neve, K. Steadman, and S. Powles. 2005. Ecological fitness of a
multiple herbicide‐resistant *Lolium rigidum* population: dynamics of seed
germination and seedling emergence of resistant and susceptible
phenotypes. Journal of Applied Ecology 42:288-298.
Vogwill, T., M. Lagator, N. Colegrave, and P. Neve. 2012. The experimental evolution
of herbicide resistance in *Chlamydomonas reinhardtii* results in a positive
correlation between fitness in the presence and absence of herbicides.
Journal of Evolutionary Biology 25:1955-1964.
Weaver, S. and S. Warwick. 1982. Competitive relationships between atrazine
resistant and susceptible populations of *Amaranthus retroflexus* and *A.*
powellii from southern Ontario. New Phytologist 92:131-139.
Weaver, S. E. and P. B. Cavers. 1979. The effects of date of emergence and emergence
order on seedling survival rates in *Rumex crispus* and *R. obtusifolius*.
Canadian Journal of Botany 57:730-738.
Webster, T. M. and G. E. MacDonald. 2001. A survey of weeds in various crops in
Georgia 1. Weed Technology 15:771-790.
Williams, M. M., N. Jordan, and C. Yerkes. 1995. The fitness cost of triazine resistance
in jimsonweed (*Datura stramonium* L.). American Midland Naturalist:131-
137.
Yu, Q., A. Collavo, M.-Q. Zheng, M. Owen, M. Sattin, and S. B. Powles. 2007. Diversity
of acetyl-coenzyme A carboxylase mutations in resistant *Lolium* populations:
evaluation using clethodim. Plant Physiology 145:547-558.
Yu, Q., H. Han, and S. B. Powles. 2008. Mutations of the ALS gene endowing
resistance to ALS‐inhibiting herbicides in *Lolium rigidum* populations. Pest
Management Science 64:1229-1236.
Yu, Q., H. Han, M. M. Vila-Aiub, and S. B. Powles. 2010. AHAS herbicide resistance
endowing mutations: effect on AHAS functionality and plant growth. Journal
of Experimental Botany 61:3925-3934.
782