Dynamical Model of Drug Accumulation in Bacteria: Sensitivity Analysis and

Experimentally Testable Predictions

Neda Vesselinova[1, 2], Boian S. Alexandrov[1], and Michael E. Wall*[3]

[1]Theoretical Division, Los Alamos National Laboratory, Los Alamos NM 87545

[2]University of California, Los Angeles, CA 90095

[3]Computer, Computational, and Statistical Sciences Division, Los Alamos National

Laboratory, Los Alamos NM 87545

*Corresponding author

November 4, 2015

LA-UR-15-28626

Abstract. We present a dynamical model of drug accumulation in bacteria. The

model captures key features in experimental time courses on ofloxacin

accumulation: initial uptake; short-term response; and long-term adaptation. In

combination with experimental data, the model provides estimates of import and

export rates in each phase, the latency in the short-term response, and the rate of

increase in efflux during adaptation. Global sensitivity analysis, local sensitivity

analysis, and Bayesian sensitivity analysis of the model provide information about

the robustness of these estimates, and about the relative importance of different

features of the accumulation time courses in three different bacterial species:

Escherichia coli, *Staphylococcus aureus*, and *Pseudomonas aeruginosa*. The results

lead to experimentally testable predictions of membrane permeability and drug

efflux and trapping, which influence drug resistance. A key prediction is that an

increase in *E. coli* ofloxacin accumulation is accompanied by a decrease in

membrane permeability, suggesting that, depending on changes in other factors,

decreasing permeability is not always an effective drug resistance strategy.

Author Summary. Bacteria live or die depending on how much antibiotic gets

inside them. Using a simple mathematical model, detailed information about drug

import and export can be teased out of time courses of internal drug levels after a

sudden environmental exposure. The results reveal that blocking diffusion across

the membrane is not always an effective drug resistance strategy for bacteria.

Introduction

Drug resistance in bacteria can be increased by efflux pump systems [1], and pump inhibition has emerged as a strategy for overcoming drug resistance [2]. Many details of how efflux pumps work are still unclear, however. In particular, quantitative information about how efflux influences drug accumulation in bacteria is still scarce [3].

Drug accumulation is a key factor in obtaining a quantitative understanding of resistance. For example, predictions of minimum inhibitory concentrations (MICs) of β-lactams in *Escherichia coli* were obtained by equating the steady state periplasmic drug concentration with a periplasmic binding protein inhibitory concentration [4]. A predicted MIC was calculated as the *external* concentration that would yield the accumulated *internal* inhibitory concentration in steady state, considering flux terms from membrane permeation and β-lactamase degradation. MIC predictions also have been made considering the action of efflux pumps on cephalosporins [5] and β-lactams [6] in *E. coli*. These predictions were accompanied by estimates of efflux pump Michaelis-Menten kinetic constants (*i.e.*, K_m and k_{cat} values), providing an explicit connection between efflux and resistance.

Time-dependent drug accumulation studies also have yielded insight into drug transport [7-9]. Diver et al. [8] found exposure of *E. coli* to five different quinolones induced a rapid ~10 sec uptake followed by a ~30 min phase of more gradual

accumulation. Similar two phase behavior was seen by Asuquo and Piddock [7] in

accumulation of fifteen different quinolones in *E. coli*, *Pseudomonas aeruginosa*, and

Staphylococcus aureus [7]. Whereas drug levels appeared to plateau in *E. coli* and *S.*

aureus, levels in *P. aeruginosa* gradually decreased at longer times, suggesting an

adaptation process [7].

Mathematical modeling has provided substantial insights into accumulation of beta-

lactams [4,6], cephalosporins [5], and tetracycline [9] in *E. coli*. Quinolone

accumulation, however, has not yet been analyzed in the context of a mathematical

model. Here we present a mathematical model of drug accumulation and use it to

analyze experimental data on accumulation of the quinolone oxfloxacin in *E. coli*, *S.*

aureus, and *P. aeruginosa* [7]. The analysis yields estimates of permeation and efflux

rates, the latency in the short-term response, and the rate of increase in efflux

during adaptation. We also perform global sensitivity analysis, local sensitivity

analysis, and Bayesian sensitivity analysis of the model. The sensitivity analyses

provide information about the robustness of parameter estimates and enable

assessment of the relative importance of the short-term response and long-term

adaptation in different bacterial species.

The results lead to experimentally testable predictions of membrane permeability

and drug efflux and trapping, which influence drug resistance. A key prediction is

that an increase in *E. coli* drug accumulation is accompanied by a decrease in

membrane permeability, illustrating that decreasing permeability, although

commonly associated with increased drug resistance [10], is not always an effective

drug resistance strategy for bacteria. Overall the results indicate the utility of

mathematical modeling and sensitivity analysis in obtaining new insights into drug

accumulation in bacteria.

Results

Model behavior

Our model of drug accumulation (Methods) was designed to describe two phases of

drug accumulation dynamics, with long-term adaptation in the second phase (Fig. 1).

After exposure at t = 0, the first phase begins, where cellular antibiotic levels rise

and relax toward an asymptotic value a_1. The relaxation in this phase is exponential

with rate β_1. At time t = τ, the second phase begins, where antibiotic levels may

either rise or fall towards a new asymptotic value a_2. The relaxation in this phase is

initially dominated by exponential relaxation with rate β_2. Later, adaptation can

dominate the dynamics, following $[1+ \delta(t- \tau)]^{-1}$. Together these features capture the

full set of behaviors exhibited by the experimental data on ofloxacin accumulation

[7] (Fig. 2).

Parameter estimates

Parameter values were estimated by fitting the model to data on ofloxacin

accumulation in *E. coli*, *P. aeruginosa*, and *S. aureus*. Following Ref. [7], the fits were

performed assuming a uniform measurement error. (Rough error estimates for the

data, inferred from the values given for the norfloxacin, are: 4.6 ng/mg for *E. coli*; 1.5

ng/mg for *P. aeruginosa*; and 8.7 ng/mg for *S. aureus*.) Reasonable fits were

obtained for all datasets (Fig. 2). Parameter values varied substantially for different

bacteria (Table 1), as expected given the differing behaviors. *E. coli* exhibits two

phases of accumulation with little apparent long-term adaptation, consistent with

the low value of δ. *P. aeruginosa* shows a stronger adaptation response,

corresponding to the higher value of δ. *S. aureus* exhibits a single phase response

without adaptation, consistent with the value of τ being greater than the longest

time point.

Sensitivity analysis results

Sensitivity analysis is a technique for evaluating how variations in the output of a

mathematical model are related to variations in the model inputs. We used

sensitivity analysis to assess the importance of different model parameters in

determining the antibiotic accumulation behavior. We performed three independent

analyses using: global sensitivity analysis (GSA), local sensitivity analysis (LSA), and

Bayesian sensitivity analysis (BSA). For each analysis, the time courses using the

optimal parameter values were used as reference behaviors.

Global sensitivity analysis. GSA explores the sensitivity of the model output within

the entire (prior) parameter space of the model inputs. We performed variance-

based GSA [11], which decomposes the variance of the output into parts ascribed to

different input parameters using a Fourier Haar decomposition [12,13]. Both the

"total effect" and "main effect" for each input parameter were calculated using the

Sobol Monte Carlo algorithm [14]; here we focus on the total effect results. The prior

range of the input parameters was defined using uniform logarithmic distributions

given in Table 2.

GSA results for *E. coli*, *P. aeruginosa*, and *S. aureus* are shown in Figure 3. For *E. coli*,

β_1 exerts the highest total effect initially, followed by a_1, τ, and a_2. The effect of β_2

never dominates, but is substantial at early times. The sensitivity to δ is low at all

times. For *P. aeruginosa*, a_1 exerts the highest total effect initially, followed by τ and

a_2. In contrast with *E. coli*, the sensitivity to δ is high for *P. aeruginosa*, while both β_1

and β_2 show low sensitivity. For *S. aureus*, β_1 exerts the highest total effect early on,

shifting to a_1 and then τ at later times. The effects of a_2 and β_2 are smaller by

comparison. Like for *E. coli*, the sensitivity to δ is low.

Local sensitivity analysis. LSA has a rich history of use in assessing the robustness of

biochemical network model behavior to changes in parameter values [15,16]. In LSA,

the local gradients of the model output are calculated with respect to model

parameters at a fixed point. Here we use the gradients to define a covariance matrix

of variations [17] (Methods), and interpret the LSA in terms of the eigenvalues and

eigenvectors of the covariance matrix. Small eigenvalues are associated with

parameter combinations that show higher sensitivity.

We performed LSA on the *P. aeruginosa* model, for comparison to the GSA. Each

eigenvector in the LSA is strongly associated with a single parameter (Fig. 4a),

facilitating the interpretation. The three smallest eigenvalues (Fig. 4b) are those

associated with a_1, a_2, and δ (corresponding to eigenvectors 1-3 in Fig. 4a),

indicating that the sensitivity of the model to these parameters is relatively high.

The parameters β_1, β_2 and τ are associated with orders of magnitude higher

eigenvalues, indicating that the sensitivity to these parameters is relatively low.

Bayesian sensitivity analysis. BSA estimates the probability density functions

(pdf's) of the model parameters given observed data and measurement errors using

Bayes' theorem [18,19]. Like GSA, BSA requires prior distributions of the model

inputs. The more sensitive is the model parameter, the more constrained is the

posterior pdf compared to the prior pdf.

Figure 5 shows Bayesian sensitivity analysis for the *P. aeruginosa* model. Here, the

Bayesian analysis identifies a Markov-Chain Monte Carlo (MCMC) sample of models

that are consistent with the observed data, assuming a uniform measurement error [7].

The histograms along the diagonal represent the posterior pdf's of the model parameters

constrained by the model observations. Each dot in the off-diagonal scatter plots

represents a model-parameter set from the final MCMC sample. Some of the parameters,

a_1, a_2, and δ appear to have peaked distributions, indicating a sharpened sensitivity

within the posterior range. The probability density of τ is uniform at longer times and

zero at shorter times. The distributions of β_1 and β_2 are flat, and, therefore they are less

sensitive parameters. The scatter plot between a_2 and δ is extended along the a_2-δ

direction, indicating that these parameters are correlated.

Experimentally testable predictions of permeability and efflux

We used Eqs. (5) and (6) to predict permeability and efflux given the estimated

parameters. The predictions are quantified in the form of the efflux rate, $\varepsilon(t)$, which

is defined as the portion of the specific export rate not associated with permeation,

and the accumulation factor, $\phi(t)$, which is defined as $\alpha(t)/\beta(t)$, the ratio of the

permeability to the export rate. Depending on the sign of $\varepsilon(t)$, qualitatively different

mechanisms are at work: a positive value is associated with efflux, while a negative

value is associated with trapping.

The resulting predictions of $\varepsilon(t)$ and $\phi(t)$ are shown in Table 4. The robustness of

these predictions depends on the robustness of the estimation of a_1 and β_1, for t<τ,

and a_2 and β_2, for t=τ. According to the sensitivity analysis, the estimates of some of

these parameters are not robust, yielding unreliable predictions for some of the *P.*

aeruginosa and *S. aureus* values. For completeness, the unreliable predictions are

listed in Table 4, but are enclosed in parentheses.

Discussion

199 ***Interpretation of sensitivity analysis results***

201 *Global Sensitivity Analysis:* Results obtained using the GSA suggest that the relative

202 importance of the various model parameters in determining the output behavior of

203 the simulations differs for *E. coli*, *P. aeruginosa*, and *S. aureus* (Figure 3). However,

204 there are also some similarities. In most cases, a_1 and β_1 are important only at very

205 early-times right at the beginning of the data collection. This suggests that proper

206 estimation of a_1 and β_1 requires accurate early-time measurements. In all cases, the

207 sensitivity of a_1 and β_1 diminishes over time. Also in all cases, the sensitivity of a_1

208 increases with time. The temporal behavior of β_2 sensitivity is very different in all

209 three cases: β_2 is relatively important at early times for the case of *E. coli*; β_2 is

210 unimportant for the case of *P. aeruginosa*; and the β_2 sensitivity peaks at the mid-

211 times for the case of *S. aureus*. Interestingly, the parameter δ is important only for

212 the case of *P. aeruginosa*. The temporal behavior of τ sensitivity is similar for the

213 case of *E. coli* and *P. aeruginosa* (τ sensitivity is decreasing with time), but very

214 different for the case of *S. aureus* (τ sensitivity is increasing with time).

216 *Local sensitivity analysis:* The global and local sensitivity analyses yield similar

217 results for *P. aeruginosa*. Both analyses identify a_1, a_2, and δ as sensitive model

218 parameters. However, the global and local sensitivity analyses produce different

219 sensitivity estimates related to the parameter τ. The model is sensitive to this

parameter according to GSA but not according to LSA. This suggests that when all

the model parameters are at their optimal values, small deviations in τ values have a

small impact on the optimal solution. However, away from the optimal point in the

parameter space, τ is an important parameter in determining the model behavior.

Bayesian sensitivity analysis: BSA provides information how well the model

parameters are constrained. For the case of *P. aeruginosa* (Fig 5), the parameters a_1,

a_2, δ and τ are well-constrained and, therefore, sensitive model parameters. This is

consistent with GSA, which is expected because both analyses are global in their

nature. BSA also suggests correlation between estimates for a_2 and δ. The other

model parameters are not correlated. The correlation is supported by the similarity

in the time dependence of a_2 and δ sensitivities as shown in Figure 3b. Some of the

parameters, a_1, a_2, and δ, have peaked posterior distributions, indicating a most

probable value at the maximum. The parameter τ is considered to be sensitive

because the posterior range (from ~1.6 to ~1.68) is smaller than the prior range (0

− 4; all the values are $\log_{10}$ transformed and presented in Tables 2 and 3). However,

the posterior distribution within this range is relatively flat (i.e. uniform). The flat

distribution leads to a small sensitivity of the model to changes in this parameter

using LSA.

Interpretation of predictions of permeability and efflux

For *E. coli*, the values of the accumulation factor ϕ>1 indicate trapping of antibiotic.

The trapping is weak for t<τ and becomes stronger at t=τ. The greater magnitude of

α compared to ε values indicates that the permeability dominates the response. For

P. aeruginosa, the values of ϕ<1 indicate that efflux of antibiotic is important. For *S.*

aureus, the value ϕ>1 indicates a net trapping of antibiotic, like *E. coli*. In contrast to

E. coli, the comparable magnitudes of α and ε indicate that permeability and

trapping are roughly equally important in during the whole time course.

The predictions suggest efflux is unimportant in the response of *E. coli* and *S. aureus*

to ofloxacin. Permeation and trapping instead dominate the response. In contrast,

efflux is predicted to be important in the response of *P. aeruginosa* to ofloxacin. The

initial accumulation is less than what would be expected by permeation alone,

indicating an efflux effect. In addition, the antibiotic level slowly decreases after the

initial uptake, indicating an increase of efflux with time, presumably due to an

adaptive response involving pump production.

The predictions for *E. coli* are especially intriguing. The response involves a ~50-

fold decrease in permeability about 800 seconds after exposure, which would

naively be expected to increase resistance [10]. However, the decrease in

permeability is accompanied by a ~10-fold decrease in trapping. Because the

decrease in trapping is less than the decrease in permeability, the result is an

increase, rather than a decrease, in antibiotic accumulation.

Similar to the subtle connection found here between permeability and resistance,

Nagano and Nikaido [5] used mathematical modeling to show how efflux pump

deletion or overexpression might not change the MIC value of strong efflux pump

substrates. It is interesting to consider these findings in light of a recent survey of

physical properties of active compounds in a drug screening collection and their

relation to whole cell antibacterial activity [20]. The survey reported on the

difficulties encountered in simultaneously optimizing both biochemical potency and

antibacterial activity, and concluded that "what is clearly needed is greater insight

into medicinal chemistry strategies which optimize transport through porins and

decrease efflux through the prolific efflux pumps." Together, the findings reported

here and elsewhere [5,6] suggest that mathematical modeling of permeation and

efflux can be a key tool in enabling antibacterial medicinal chemistry.

Our results indicate that detailed, quantitative analysis of antibiotic accumulation

time courses using mathematical modeling can yield new insights and

experimentally testable predictions about mechanisms of drug resistance. The

sensitivity analysis provides basis for distinguishing important from unimportant

predictions. The predictions imply that decreasing permeability is not always an

effective strategy for bacteria to increase drug resistance. This outcome in particular

provides an example of the potential pitfalls involved in reasoning about structure-

function relations in biochemical networks [21], and reinforces the need for

incorporating mathematical modeling into strategies for fighting drug resistance.

Overall this study illustrates the key role that mathematical modeling and sensitivity

analysis can play in deep interpretation of experimental data in biology.

Materials and Methods

Model

The time dependence of accumulated drug $a(t)$ is modeled using

$$\frac{da(t)}{dt} = \alpha(t)a_o - \beta(t)a(t) \qquad (1)$$

where a_o is the environmental antibiotic concentration, $\alpha(t)$ is the specific import

rate, and $\beta(t)$ is the specific export rate. Here and elsewhere the terms "import" and

"export" refer to combined effects of passive and active transport, and the term

"specific" is sometimes used to indicate that the rate constant is multiplied by the

antibiotic concentration. Both $\alpha(t)$ and $\beta(t)$ may change depending on the phase of

the response. The import rate $\alpha(t)$ is given by

$$\alpha(t) = \begin{cases} \alpha_1, & t < \tau \\ \alpha_2, & t \geq \tau \end{cases} \qquad (2)$$

305 where α_1 applies to the initial phase of the accumulation dynamics (t<τ), and α_2

306 applies to the later phase (t ≥ τ).

308   The export rate β(t) is given by

$$\beta(t) = \begin{cases} \beta_1, & t < \tau \\ \beta_2[1 + \delta(t - \tau)], & t \geq \tau' \end{cases}$$ (3)

311   where β_1 applies during the initial phase (t<τ), β_2 applies during the later phase

312   (t≥τ), and δ is the fractional rate of increase of export during the later phase.

314   Assuming $\delta << \beta_2$ and $\delta(t - \tau) \ll 1$ yields

$$a(t) = \begin{cases} e^{-\beta_1 t}a(0) + (1 - e^{-\beta_1 t})a_1, & t \leq \tau \\ e^{-\beta_2(t-\tau)}a(\tau) + [1 - e^{-\beta_2(t-\tau)}]\dfrac{a_2}{1 + \delta(t - \tau)}, & t > \tau \end{cases}$$ (4)

where $a_1 = \alpha_1 a_o / \beta_1$, and $a_2 = \alpha_2 a_o / \beta_2$.

Given additional assumptions, the α and β parameters can be used to derive

permeation and efflux or trapping rates (see below). The δ parameter models long-

term adaptation by, e.g., gene regulation.

Parameter estimation

Parameter estimation for Eq. (4) was performed using published experimental time

courses of ofloxacin accumulation in *E. coli*, *P. aeruginosa*, and *S. aureus* [7]. Data

points were extracted from Figure 1 in Ref. [7]. Numerical solutions to Eq. (4) were

obtained for each data point using functions defined in Mathematica Version 10

(Wolfram Research, Inc., Champaign, IL). As the estimated errors were the same for

all data points [7], an unweighted mean squared deviation (MSD) between the

model and the data was used as a target function for fitting. The MSD was minimized

with respect to values of a_1, a_2, β_1, β_2, δ, and τ using the `FindMinimum` method,

assuming a(0)=0. The values of a_1, a_2, β_1, β_2, δ were varied over twelve decades. For

E. coli and *P. aeruginosa*, the value of τ was varied from 0-10,000 s (covering the full

span of each 3,600 s time course) to seek an optimal solution. For *S. aureus*,

optimization was performed for τ = 10,000 s. Mathematica workbooks and

experimental data for all cases are available in a compressed archive in the

Supporting Information.

Sensitivity analysis methods

All analyses were performed using the open source code Model-Analysis and

Decision Support (MADS), developed at Los Alamos National Laboratory

(http://mads.lanl.gov).

Global Sensitivity Analysis (GSA): For our analysis, we used the Sobol Monte Carlo

algorithm [14]. The algorithm estimates the total and main effect for each model

parameter. The total effect is the total contribution to the output model variance to a

given model parameter, including all the combined effects caused by its interactions

with any other model parameter. The main effect measures the impact of varying a

single model parameter by itself. The total and main effect can be very similar when

the model parameters are independent. Sobol Monte Carlo algorithm was

performed using about 10^6 independent model evaluations. Additional analyses (not

presented) demonstrated that the number of model evaluations is sufficient to

achieve converges for the estimate quantities.

Local Sensitivity Analysis (LSA): We used a finite difference method requiring a

limited number of model evaluations (equal to the number of unknown parameters).

As a result, we obtain a gradient matrix **J** (i.e. the Jacobian) with dimensions $[m \times n]$

where m is the number of model parameters and n is the number of model inputs.

Each component of the **J** matrix represents the local sensitivity of each model

parameter to each model output [17]. We analyze the covariance matrix **C** of model

parameters which is computed as $\mathbf{C} = [J^T J]^{-1}$. The covariance matrix is frequently

analyzed using eigenanalysis where eigenvectors and eigenvalues of the covariance

matrix are explored. How dominant (important) is each eigenvector depends on the

respective eigenvalues; the smaller the eigenvalue, the higher the importance of the

eigenvector. The components of each eigenvector represent the contributions of

each model parameter to the simultaneous variation of multiple parameters: the

larger the absolute value of the components, the larger the contribution. Model

parameters with large contribution in dominant eigenvectors are important

(sensitive) model parameters. Model parameters with large contribution in non-

dominant eigenvectors are also unimportant (insensitive) model parameters. If

several model parameters have important contribution in a single eigenvector, these

model parameters are correlated. If these contributions have the same sign, the

correlation is positive. If these contributions have opposite signs, the correlation is

negative.

Bayesian Sensitivity Analysis (BSA): The BSA was performed using the Robust

Adaptive Metropolis Markov Chain Monte Carlo algorithm [22]. To assess the

sensitivity of the model parameters to the model outputs we compared the prior

and the posterior pdf's of the model parameters (Table 2 and 3, respectively). High

sensitivity parameters show a posterior pdf that is substantially narrower than the

prior pdf.

Prediction of permeability and efflux

Estimation of the model parameters enables prediction of permeability and efflux

rates. Given a_o, the specific import rates may be calculated as $\alpha_1 = a_1\,\beta_1/a_o$, and $\alpha_2 =$

$a_2\,\beta_2/a_o$. Assuming the import is due to permeation, α_1 and α_2 are the permeabilities.

Next, assume the specific export rate $\beta(t)$ is a sum of contributions from

permeability, $\alpha(t)$ and other effects, $\varepsilon(t)$, leading to

$$\varepsilon(t) = \begin{cases} \beta_1 - \alpha_1, & t < \tau \\ \beta_2[1 + \delta(t - \tau)] - \alpha_2, & t \geq \tau \end{cases}.$$ (5)

394   When $\varepsilon(t)>0$, outward flow is enhanced compared to permeation, and $\varepsilon(t)$ is

395   associated with efflux. When $\varepsilon(t)<0$, the outward flow is decreased compared to

396   permeation, and $\varepsilon(t)$ is associated with trapping. The accumulation factor $\phi(t) =$

397    $\alpha(t)/\beta(t)$ is used as a measure of the importance of efflux or trapping. It is given by

$$\phi(t) = \begin{cases} a_1/a_o, & t < \tau \\ a_2/a_o[1 + \delta(t - \tau)], & t \geq \tau \end{cases}.$$ (6)

Eqs. (5) and (6) are only valid under the same assumptions as for Eq. (4).

Predictions of $\varepsilon(t)$ are only meaningful using reliable estimates of both a_1 and β_1 for

$t < \tau$, or both a_2 and β_2 for $t \geq \tau$. In contrast, $\phi(t)$ can be predicted without estimates

of β_1 or β_2.

The predictions involve information about the environmental drug concentration

and the conversion of estimates a_1 and a_2 from ng/mg dry weight units (Table 1) to

mg/L units. The environmental ofloxacin concentration in the experiments was 10

mg/L [7]. To convert a_1 and a_2 to mg/L units, following Ref. [23], we used a buoyant

density of 1.1 g/mL and a 31% dry weight for *E. coli*, a buoyant density of 1.2 g/mL

and a 48% dry weight for *P. aeruginosa* (transferred from *P. putidae*), and assume a

buoyant density of 1.1 g/mL and a 40% dry weight for *S. aureus*. This yielded

conversion factors of 0.341 mg dry weight/L for *E. coli*, 0.528 mg dry weight/L for *P.*

aeruginosa, and 0.440 mg dry weight/L for *S. aureus*.

Acknowledgments

This study was supported by the US Department of Energy under Contract DE-

AC52-06NA25396 through the Laboratory-Directed Research and Development

Program at Los Alamos National Laboratory under project number 20140121DR,

Combating Antibiotic Resistance: Targeting Efflux Pump Systems at Multiple Scales.

References

1. Li XZ, Nikaido H (2009) Efflux-mediated drug resistance in bacteria: an update.
Drugs 69: 1555-1623.
2. Kourtesi C, Ball AR, Huang YY, Jachak SM, Vera DM, et al. (2013) Microbial efflux
systems and inhibitors: approaches to drug discovery and the challenge of
clinical implementation. Open Microbiol J 7: 34-52.
3. Nikaido H, Pages JM (2012) Broad-specificity efflux pumps and their role in
multidrug resistance of Gram-negative bacteria. FEMS Microbiol Rev 36: 340-
363.
4. Nikaido H, Normark S (1987) Sensitivity of Escherichia coli to various beta-
lactams is determined by the interplay of outer membrane permeability and
degradation by periplasmic beta-lactamases: a quantitative predictive
treatment. Mol Microbiol 1: 29-36.
5. Nagano K, Nikaido H (2009) Kinetic behavior of the major multidrug efflux pump
AcrB of Escherichia coli. Proc Natl Acad Sci U S A 106: 5854-5858.
6. Lim SP, Nikaido H (2010) Kinetic parameters of efflux of penicillins by the
multidrug efflux transporter AcrAB-TolC of Escherichia coli. Antimicrob
Agents Chemother 54: 1800-1806.
7. Asuquo AE, Piddock LJ (1993) Accumulation and killing kinetics of fifteen
quinolones for Escherichia coli, Staphylococcus aureus and Pseudomonas
aeruginosa. J Antimicrob Chemother 31: 865-880.

8. Diver JM, Piddock LJ, Wise R (1990) The accumulation of five quinolone
antibacterial agents by Escherichia coli. J Antimicrob Chemother 25: 319-333.
9. Thanassi DG, Suh GS, Nikaido H (1995) Role of outer membrane barrier in efflux-
mediated tetracycline resistance of Escherichia coli. J Bacteriol 177: 998-
1007.
10. Schweizer HP (2012) Mechanisms of antibiotic resistance in Burkholderia
pseudomallei: implications for treatment of melioidosis. Future Microbiol 7:
1389-1399.
11. Sobol' IM (1990) On sensitivity estimation for nonlinear mathematical models.
Matematicheskoe Modelirovanie 2: 112-118.
12. Sobol' IM (1993) Asymmetric convergence of approximations of the Monte Carlo
method. Computational mathematics and mathematical physics 33: 1391-
1396.
13. Saltelli A, Andres TH, Homma T (1993) Sensitivity Analysis of Model Output - an
Investigation of New Techniques. Computational Statistics & Data Analysis
15: 211-238.
14. Saltelli A, Ratto M, Andres T, Campolongo F, Cariboni J, et al. (2008) Global
Sensitivity Analysis: The Primer: John Wiley & Sons.
15. Savageau MA (1976) Biochemical systems analysis: a study of function and
design in molecular biology. Reading, MA: Addison-Wesley.
16. Savageau MA (2013) System design principles. In: Wall ME, editor. Quantitative
Biology: From Molecular to Cellular Systems. Boca Raton: Taylor and Francis.
pp. 23-49.
17. Gustafson P, Srinivasan C, Wasserman L (1995) Local sensitivity analysis.
Bayesian Statistics 5: 631-640.
18. Oakley JE, O'Hagan A (2004) Probabilistic sensitivity analysis of complex
models: a Bayesian approach. J Roy Stat Soc B 66: 751-769.
19. Weiss R (1996) An approach to Bayesian sensitivity analysis. J Roy Stat Soc B 58:
739-750.
20. Brown DG, May-Dracka TL, Gagnon MM, Tommasi R (2014) Trends and
exceptions of physical properties on antibacterial activity for Gram-positive
and Gram-negative pathogens. J Med Chem 57: 10144-10161.
21. Wall ME (2011) Structure-function relations are subtle in genetic regulatory
networks. Math Biosci 231: 61-68.
22. Vihola M (2012) Robist adaptice Metropolis algorithm with coerced acceptance
rate. Stat Comp 22.
23. Bratbak G, Dundas I (1984) Bacterial dry matter content and biomass
estimations. Appl Environ Microbiol 48: 755-757.
484
485

Figures

Figure 1. Behavior of the model and connection to model parameters.

Figure 2. Fits of models to ofloxacin accumulation data. A) *E. coli*. B) *P. aeruginosa*. C)

S. aureus. Following Ref. [7], the data are assumed to have a uniform measurement

errors: 4.6 ng/mg for *E. coli*; 1.5 ng/mg for *P. aeruginosa*; and 8.7 ng/mg for *S.*

aureus.

A)

B)

C)

Figure 3. Global sensitivity of the model behavior to changes in parameter values for

A) *E. coli*, B) *P. aeruginosa*, and C) *S. aureus* models. Sensitivity is quantified using

the total effect measure in GSA. Color code: a_1 (cyan); a_2 (yellow); β_1 (red); β_2

(green); δ (pink); and τ (grey).

A)

B)

Figure 4. Local sensitivity analysis for *P. aeruginosa*. (A) Eigenvector components

associated with each model parameter. (B) Eigenvalues for each eigenvector,

numbered in decreasing importance from left to right. The LSA identifies a_2 as the

most important model parameter because it contributes strongly to eigenvector 1,

and identifies β_2 as the least important because it contributes strongly to

eigenvector 6.

Figure 5. Bayesian sensitivity analysis for $P.\ aeruginosa$. The histograms along the

diagonal represent the posterior pdf's of the model parameters constrained by the

model observations. Off-diagonal subplots are scatter plots where each dot is a

model-parameter set from the final MCMC sample. The horizontal axes define

the parameter ranges (Table 3). The vertical axes along the diagonal plots define

the frequency of occurrence. The vertical axes along the off-diagonal plots also

define the parameter ranges (Table 3).

537 **Tables**

539 Table 1. Estimated parameter values obtained after fitting the model to data on

540 ofloxacin accumulation in different bacteria.

	a_1	a_2	β_1 (1/s)	β_2 (1/s)	δ (1/s)	τ (s)
	(ng/mg)	(ng/mg)				
E. coli	32.0	67.1	0.116	0.00125	(0.0000484)	772
P. aeruginosa	7.43	13.7	(0.682)	(3.01)	0.000314	44.5
S. aureus	68.71	(16.8)	0.149	(0.0119)	(0)	(10^4)

542 [a]Numbers in parentheses are unreliable according to the sensitivity analysis. They

543 are included only for completeness.

545   Table 2. Prior parameter ranges ($\log_{10}$ transformed values).

	a_1 (ng/mg)	a_2 (ng/mg)	β_1 (1/s)	β_2 (1/s)	δ (1/s)	τ (s)
min	0	0	-1	-1	-7	0
max	2	2	4	4	0	4

550   Table 3. Posterior parameter ranges based on Bayesian sensitivity analysis for $P.$

551   *aeruginosa* ($\log_{10}$ transformed values).

	a_1 (ng/mg)	a_2 (ng/mg)	β_1 (1/s)	β_2 (1/s)	δ (1/s)	τ (s)
min	0.8	1.11	-1	-1	-3.65	1.6
max	0.92	1.16	4	4	-3.4	1.68

555   Table 4. Predicted accumulation factors (ϕ), permeabilities (α), and efflux ($\varepsilon > 0$) or

556   trapping ($\varepsilon < 0$) rates.

	$\phi(t<\tau)$	$\phi(t=\tau)$	α_1 (1/s)	α_2 (1/s)	$\varepsilon(t<\tau)$ (1/s)	$\varepsilon(t=\tau)$ (1/s)
E. coli	1.09	2.29	0.13	0.0029	-0.011	-0.0016
P. aeruginosa[a]	0.39	0.72	(0.27)	(2.2)	(0.41)	(0.83)
S. aureus[a]	3.02	(0.74)	0.45	(0.01)	-0.30	(0.0031)

559   [a]Numbers in parentheses are unreliable according to the sensitivity analysis. They

560   are included only for completeness.