2 **Species sorting along a subsidy gradient alters community stability**

4 Mario E. Muscarella[1], Stuart E. Jones[2], Jay T. Lennon[1*]

6 [1] Department of Biology, Indiana University, Bloomington, IN 47405 USA

[2] Department of Biological Sciences, University of Notre Dame, Notre Dame, IN 46556 USA

* Corresponding author, Tel 812-856-0962, Fax 812-855-6082, Email lennonj@indiana.edu

Running Head: species sorting and community stability

12 Key Words: Resource Subsidies, Community Stability, Responsiveness, Microorganisms,

Terrestrial DOC

ABSTRACT

16   The movement of resources between terrestrial and aquatic habitats has strong effects on

ecological processes in recipient ecosystems. Allochthonous inputs modify the quality and

18   quantity of the available resource pool in ways that may alter the composition and stability of

recipient communities. Inputs of terrestrial dissolved organic carbon (tDOC) into aquatic

20   ecosystems represent a large influx of resources that has the potential to affect local

communities, especially microorganisms. To evaluate the effects terrestrial inputs on aquatic

22   microbial community composition and stability, we manipulated the supply rate of tDOC to a set

of experimental ponds. Along the tDOC supply gradient, we measured changes in diversity and

24   taxon-specific changes in abundance and activity. We then determined community stability by

perturbing each pond using a pulse of inorganic nutrients and measuring changes in composition

26   and activity (i.e., responsiveness) along the gradient. Terrestrial DOC supply significantly altered

the composition of the active microbial community. The composition of the active bacterial

28   community changed via decreases in richness and evenness as well as taxon-specific changes in

abundance and activity indicating species sorting along the gradient. Likewise, the

30   responsiveness of the active bacterial community decreased along the gradient, which led to a

more stable active community. We did not, however, observe these changes in diversity and

32   stability in the total community (i.e., active and inactive organisms), which suggests that tDOC

supply modifies microbial community stability through functional not structural changes.

34   Together, these results show that altered aquatic terrestrial linkages can have profound effects on

the activity and stability of the base of the food web and thus can alter ecosystem functioning.

INTRODUCTION

38   Resource subsidies refer to the cross-boundary movement of materials and energy from

donor to recipient ecosystems. These allochthonous inputs affect various ecological processes

40   including population dynamics, species interactions, and ecosystem functioning (Polis et al.

1997, Holt 2004). Subsidies can also alter the stability of recipient communities. Theory suggests

42   that low to moderate levels of subsidies stabilize recipient communities by promoting trophic

omnivory (Polis et al. 1997, Huxel and McCann 1998) and dampening oscillations between

44   consumers and their resources (DeAngelis 1992, Huxel and McCann 1998). The precise effect of

subsidies on recipient communities, however, likely depends on the timing (Takimoto 2002),

46   quality (Bartels et al. 2012), and quantity (Cottingham and Narayan 2013). Likewise, subsidies

can also affect stability by altering the composition of recipient communities by selecting for and

48   against particular species (Holt 2004). If subsides alter communities via species sorting, then

they will also alter the distribution of species-specific traits that may affect stability, including

50   resource specialization (Huxel and McCann 1998, Faria and Costa 2010) and metabolic plasticity

(Comte et al. 2013). Together, the effect of subsidies on community stability is likely influenced

52   by both subsidy properties (e.g., quantity) and consumer traits (e.g., resource specialization).

A well-recognized and pervasive subsidy is the movement of dissolved organic carbon

54   (DOC) from terrestrial to aquatic ecosystems (Polis et al. 1997, Tranvik et al. 2009). In most

inland water bodies, terrestrial DOC (tDOC) inputs exceed internal (i.e., autochthonous) inputs

56   by aquatic autotrophs by at least an order of magnitude (Tranvik et al. 2009, Wilkinson et al.

2013). Additionally, there is growing evidence that tDOC inputs are increasing in many regions

58   around the world owing to global change (Monteith et al. 2007), and it has been hypothesized

that such changes in tDOC inputs will affect the functioning and stability of aquatic ecosystems

60   (Jones and Lennon 2015, Solomon et al. 2015). Heterotrophic microorganisms are the primary

consumers of tDOC in aquatic ecosystems. Despite having recalcitrant properties, tDOC is used

62   by bacteria for anabolic and catabolic processes (Kritzberg et al. 2004, Lapierre et al. 2013). As

such, heterotrophic bacteria are expected to mediate the aquatic ecosystem response to tDOC

64   subsidies via changes in community composition and metabolic activity (Jones et al. 2009,

Comte and del Giorgio 2011), and it has been hypothesized that, due to these changes, subsidies

66   may alter the stability of recipient aquatic ecosystems (Wetzel 1999).

If subsidies modify the composition of recipient communities through processes such as

68   species sorting, then subsidies could also alter the function and stability of recipient

communities. Microbial communities are comprised of species with traits that link ecosystem

70   functioning and community stability. For example, the degree to which communities are

dominated by either generalist or specialists has important consequences for carbon cycling

72   (Mou et al. 2008) and may explain how microbial communities respond to perturbations (Allison

and Martiny 2008, Shade et al. 2011). Another important trait is metabolic plasticity or the

74   ability of taxa to change physiological functions or to transition across levels of metabolic

activity (e.g., active to dormant). Microbial communities consist of taxa that differ in activity and

76   their degree of metabolic plasticity (del Giorgio and Gasol 2008, Lennon and Jones 2011), and

the ability of individual taxa to rapidly adjust metabolic activity has been shown to be important

78   for controlling ecosystem functions such as carbon and nitrogen cycling (Aanderud et al. 2015).

In addition, the ability of taxa to adjust metabolic functions (Comte et al. 2013) or to transition

80   across levels of activity (Lennon and Jones 2011) have been hypothesized to buffer communities

against perturbations. Therefore, processes such as species sorting will alter the distribution of

82   taxa and traits within communities and thus affect the stability and functioning of ecosystems.

In this study, we quantified the effects of tDOC supply on the diversity, composition, and

84   stability of aquatic microbial communities. We hypothesized that subsidies would change

bacterial composition via shifts in either the abundance or activity of taxa which would be

86   reflective of species sorting. We further hypothesized that resource-driven shifts in composition

would determine how subsidized bacterial communities respond to perturbations. To test our

88   hypotheses, we manipulated tDOC supply rate in eleven experimental ponds. First, we used

microbial community sequencing to assess changes in diversity and trends in abundance and

90   activity across experimental treatments. Then, we measured community stability by quantifying

changes in community composition in each pond following an inorganic nutrient perturbation.

92   Results from our study provide a framework for how aquatic microbial communities may

respond to increased resource subsidies such as tDOC, and show that subsidy-mediated shifts in

94   composition alter the stability of communities.

96   **MATERIALS AND METHODS**

Experimental Ponds – We manipulated the supply rate of terrestrial dissolved organic carbon

98   (tDOC) in eleven experimental ponds at the Michigan State University's W.K. Kellogg

Biological Station (KBS) Experimental Pond Facility. Each experimental pond has a 30 m

100   diameter, a 2 m maximum depth, and an operating volume of approximately 10^6 L. We

established a tDOC supply gradient by adding different amounts of a commercially available

102   source of humic substances (Super Hume, Crop Master USA) to each pond using a 5 horsepower

trash pump. This source of humic substances is comprised of 17% humic acid and 13% fulvic

104   acid and is known to be used by diverse heterotrophic bacteria (Lennon et al. 2013). We

maintained the tDOC supply gradient for 100 days (6 June 2009 to 14 September 2009) by

106   adding Super Hume to each pond on a weekly basis assuming a loss rate of 4.7 – 12.2% d^{-1}

(Lennon et al., 2013; Jones & Lennon, 2015).

Perturbation and Sampling – Nutrient limitation is typical for inland water bodies including the

110   KBS experimental ponds, and aquatic communities are sensitive to episodic nutrient inputs

(Elser et al. 1990). As such, nutrient pulses are a common perturbation to aquatic ecosystems and

112   are often used in experiments to test questions about stability (Nowlin et al. 2008). We used an

inorganic nutrient pulse to test the stability of aquatic microbial communities along the tDOC

114   supply gradient. We added 500 L of an inorganic nutrient solution (NH_4NO_3 and Na_2HPO_4) to

each experimental pond on day 82 (27 Aug 2009) using a 5 horsepower trash pump. The

116   inorganic nutrient pulse elevated inorganic nitrogen (N) and phosphorus (P) concentrations of

each pond by 565 µg L^{-1} and 50 µg L^{-1}, respectively, while maintaining the initial N:P molar

118   ratio. Prior to and after the inorganic nutrient pulse, we sampled each pond three times per week

to track changes in water chemistry. We collected water samples from the center of each pond

120   using a 1 m depth-integrated sampler. We measured DOC by oxidation and nondispersive

infrared (NDIR) detection using a Shimadzu TOC-V carbon analyzer on 0.7 µm (Whatman,

122   GF/F) filtered water samples. We measured total nitrogen concentrations spectrophotometrically

after persulfate digestion using the second-derivative method (APHA 1998). We measured

124   soluble reactive phosphorus concentrations colorometrically using the ammonium molybdate

method (Wetzel and Likens 2000). Further details about chemical analyses and the nutrient

126   perturbation can be found elsewhere (Jones and Lennon 2015).

128   ***Bacterial Community Characterization*** – We characterized aquatic bacterial composition along

the tDOC supply gradient using 16S rRNA sequencing approximately a week prior to (day 74)

130   and after (day 88) the inorganic nutrient pulse. We collected water samples from the center of

each pond using a 1 m depth-integrated sampler. We retained microbial biomass on 47 mm 0.2

132   µm Supor PES membrane filters (Pall) and stored immediately at -80 °C. Because

microorganisms exist at various levels of metabolic activity which have differential effects on

134   the structure and function of the community (del Giorgio and Gasol 2008), we used two

approaches to characterizing composition. One approach used DNA, a stable molecule, to

136   characterize the microbial community based on all taxa, regardless of activity level. We refer to

the DNA approach as the "total community". The second approach used RNA, an ephemeral

138   molecule reflecting metabolic growth and activity (Elser et al. 2003), to characterize the

microbial community based on the organisms that contribute to ecosystem function (Jones and

140   Lennon 2010, Aanderud et al. 2015). We refer to the RNA approach as the "active community".

We extracted nucleic acids (DNA and RNA) using the PowerSoil Total RNA Extraction Kit with

142   DNA Elution Accessory Kit (MoBio, Carlsbad, CA). We treated RNA extracts with DNase

(Invitrogen) to degrade DNA prior to cDNA synthesis via the SuperScript III First Strand

144   Synthesis Kit (Invitrogen). Once DNA and cDNA samples were cleaned and quantified, we

amplified the 16S rRNA gene (DNA) and transcript (cDNA) using barcoded primers (515F and

146   806R) designed to work with the Roche 454 sequencing platform (Fierer et al. 2008; see

Supplement for PCR sequences and conditions). We sequenced 16S rRNA amplicons at the

148   Research Technology Support Facility at Michigan State University using the GS FLX Titanium

chemistry. We processed raw 16S rRNA sequences using the software package *mothur* (version

150   1.32.1, Schloss et al. 2009). To identify operational taxonomic units (OTUs), we binned

sequences into phylotypes using the Ribosomal Database Project's 16S rRNA database and

152   taxonomy as the reference (Cole et al. 2009).

154   ***Community Diversity*** – To determine the effects of tDOC supply on bacterial community

diversity, we measured alpha diversity in each pond prior to the inorganic nutrient pulse. First,

156   we estimated taxonomic richness using a resampling approach. We subsampled communities to

obtain a standardized 2000 observations per site and summed the number of OTUs represented in

158   each subsample. We then resampled 999 additional times and calculated an average richness

estimate ($\pm$ SEM) for each site (Muscarella et al. 2014). We used Good's Coverage to confirm

160   that subsampling to 2000 observations was sufficient to make conclusions based on our data set

(Good, 1953). Second, we determined taxonomic evenness using Simpson's Evenness (Smith

162   and Wilson 1996). Evenness was calculated using the same resampling approach we used for

richness. We calculated richness and evenness for the total (DNA) and active (RNA) microbial

164   communities. For each metric, we used an indicator variable multiple regression model (Neter et

al. 1996) to test how diversity changed in response to tDOC supply with respect to both the total

166   (DNA) and active (RNA) microbial community. In our regression model, we used supply rate as

the continuous variable and community type (total vs. active) as the categorical variable. All

168   statistical calculations were performed in the R computing environment (v 3.1.3, R Core

Development Team 2012).

Community Composition and Species Sorting – To determine the effects of tDOC supply on

172   community composition and test for evidence of species sorting, we determined how bacterial

communities and individual bacterial taxa responded to the tDOC gradient. First, we used

174 PERMANOVA to determine if the bacterial community responded to tDOC supply rate for both

the total and active communities (Anderson 2001). For each, if the PERMANOVA was

176 significant we tested for evidence of species sorting defined here as species-specific directional

changes in abundance (Jablonski 2008). We used Spearman's rank-order correlation to measure

178 the response of each taxon to tDOC supply. We identified responsive taxa based on significant

rho-values after a Benjamini-Hochberg correction for multiple comparisons (Benjamini and

180 Hochberg 1995). Positive rho-values indicate taxa that responded positively (in either the total or

active community) to tDOC supply and negative rho-values identified taxa that responded

182 negatively to tDOC supply. We then used the Ribosomal Database Project's Taxonomy (Cole et

al. 2009) and a curated freshwater bacteria database (Newton et al. 2011) to taxonomically

184 identify each responsive taxa. All statistical calculations were performed in the R computing

environment and PERMANOVA was implemented using the *adonis* function in the R package

186 vegan (v 2.2-1; Oksanen et al. 2013). Taxonomic identifications were performed using *mothur*.

188 *Community Stability* – We determined the effects of the subsidy supply on community stability

by calculating the change in community composition before and after the inorganic nutrient

190 pulse (He et al. 1994, Brown 2003). First, we used Principal Coordinates Analysis (PCoA) to

visualize differences in community composition based on Bray-Curtis distances. PCoA is a

192 metric multidimensional scaling technique that allows objects to be oriented in a common

reduced-dimension space while maintaining distance between objects as best as possible

194 (Legendre and Legendre 2012). We used $\log_{10}$-transformed relative abundances and Bray-Curtis

distance to reduce bias against low abundance taxa (Anderson et al. 2006, Legendre and

196 Legendre 2012). We then measured the Euclidean distance in multivariate space between paired

communities before and after the inorganic nutrient pulse using the first three multivariate axes

198   (Brown, 2003). The Euclidean distance estimates the overall change in the microbial community,

and we used the distance, which we refer to as "responsiveness", as a metric of community

200   stability (Grimm and Wissel 1997). Thus, a more stable community would be one that is less

responsive to the inorganic nutrient pulse. We used an indicator variable multiple regression to

202   test how subsidy supply rate altered community stability (i.e., responsiveness) with respect to

both the total (DNA) and active (RNA) microbial communities. In our regression model, we used

204   supply rate as the continuous variable and community type (total vs. active) as the categorical

variable. All statistical calculations were performed in the R computing environment and PCoA

206   was implemented using the *cmdscale* function in the R package vegan.

<div align="center">

RESULTS

</div>

DOC Manipulation and Nutrient Perturbation –The tDOC supply rates imposed (0 – 200 g

210   C/m^2) reflect the range of tDOC received by temperate lakes under current and future expected

supply rates (Solomon et al. 2015), and established a DOC concentration gradient across ponds

212   from 6 to 24 mg C/L (Supplemental Fig. 1). After the tDOC supply gradient had been established

for 80 days, the inorganic nutrient pulse rapidly elevated nutrient concentrations approximately

214   10-fold while maintaining N:P ratios observed prior to the perturbation (Supplemental Fig. 2).

216   ***Community Diversity –*** The tDOC supply gradient significantly decreased the richness and

evenness of the active microbial community but had no effect on the diversity of the total

218   community. An indicator variable multiple regression revealed that tDOC supply rate explained

85% of the observed variation in bacterial richness ($F_{3, 15} = 34.2$, $P < 0.001$). Based on the total

220   community (i.e., all taxa), richness did not change in response to tDOC supply ($P = 0.50$);

however, the richness of the active community decreased in response to tDOC supply. Overall,

222   we observed a 30% drop in the number of active taxa across the gradient (Fig. 1A; $R^2 = 0.64$, $P =$

0.001). Similarly, an indicator variables multiple regression model revealed that tDOC supply

224   rate explained 54% of the observed variation in community evenness ($F_{3, 15} = 8.12$, $P = 0.002$).

Based on the total community, community evenness did not change in response to tDOC supply

226   ($P = 0.32$); however, the evenness of the active community decreased in response to tDOC

supply with a 25% drop in evenness across the entire gradient (Fig. 1B; $R^2 = 0.51$, $P = 0.04$).

Community Composition and Species Sorting – The tDOC supply gradient had a significant

230   effect on community composition for both the total and active communities. Our PERMANOVA

results show that tDOC supply altered the composition of the total ($R^2 = 0.17$; $P = 0.03$) and

232   active ($R^2 = 0.15$, $P = 0.04$) bacterial community. In addition, we found evidence of species-

specific responses in both the total and active communities. Together, 29 bacterial taxa (24% of

234   the observed OTUs representing 26% of the total sequences) demonstrated a significant

directional response based on the total (i.e., DNA) or active (i.e., RNA) community. Based on

236   DNA analysis, 19 taxa significantly correlated with tDOC supply with eight positive and 11

negative relationships (Fig. 2B). Based on RNA analysis, 15 taxa significantly correlated with

238   tDOC supply with one positive relationship and 14 negative relationships (Fig. 2C). The other

121 bacterial taxa did not demonstrate a significant relationship because their relative

240   abundances were either constant or changed sporadically across the tDOC supply gradient.

242   *Community Stability* – The tDOC supply gradient significantly enhanced the stability of the

active microbial community but had no effect on the stability of the total community. For our

244   PCoA analysis, the first three ordination axes explained 60% of the variation between microbial

communities across the tDOC supply gradient. An indicator variable multiple regression

246   revealed that tDOC supply rate and community type (total or active) explained 83% of the

observed variation in responsiveness ($F_{3,\ 18} = 35$, $P < 0.001$). The responsiveness of the total

248   community was low and did not change across the tDOC supply gradient ($P = 0.37$, Fig. 3A).

However, the active community became less responsive as tDOC supply increased ($P < 0.001$,

250   Fig. 3A), and there was a 35% drop in responsiveness across the entire gradient.

252   **DISCUSSION**

The cross boundary movement of resources can alter the composition and stability of

254   recipient communities, and therefore change community dynamics and ecosystem functioning. In

this study, we manipulated the supply rate of terrestrial dissolved organic carbon (tDOC) to

256   aquatic ecosystems and documented changes in the composition and stability of aquatic

microbial communities. We found that the subsidy selected for and against certain taxa and

258   suggest that this is evidence of tDOC-mediated species sorting. In addition, we found that supply

rate increased the stability of the active members of the microbial community, and propose that

260   via species sorting, tDOC established an active community that was less responsive to an

inorganic nutrient pulse. Together, our results suggest that in the face nutrient perturbations,

262   some subsidies (e.g. tDOC) select for specialized taxa that increase community stability.

Furthermore, because these subsidies stabilize the organisms that regulate ecosystem

264   functioning, the ability of ecosystems to contend with nutrient perturbations may diminish due to

a reduced capacity for structural and functional responses by the members of the microbial

266   community.

268   **Effects of terrestrial DOC on aquatic bacterial diversity**

The tDOC supply gradient altered the aquatic microbial community by decreasing the

270   richness and evenness of the active taxa. This suggests that ecosystems receiving high inputs of

tDOC contain fewer active taxa and that certain taxa are disproportionately responsible for the

272   majority of microbial activity in these ecosystems. There are multiple mechanisms by which

increased tDOC inputs could decrease the diversity of active bacteria. First, tDOC is

274   chromophoric and can decreases light availability, limit aquatic primary productivity, and thus

reduces the concentration of labile, algal-derived resources that are used by many aquatic

276   bacteria (Karlsson et al. 2009, Lennon et al. 2013, Jones and Lennon 2015). The decreased inputs

of autochthonous inputs would have negative effects on aquatic microorganisms that rely on

278   algal-derived resources (Kritzberg et al. 2004), and thus decrease the richness of the active

community. Second, tDOC is a chemically and physically recalcitrant resource pool (Fellman et

280   al. 2009), and it has been argued that only specialized consumers use tDOC (Fuchs et al. 2011).

For example, the degradation of aromatic compounds requires specific metabolic pathways, most

282   notably the beta-ketoadipate pathway (Fuchs et al. 2011). Therefore, tDOC subsidies may only

favor taxa with specialized metabolic pathways (Mou et al. 2008, Fuchs et al. 2011). If these

284   pathways are rare in the community then only few taxa would benefit from the chemical niches

offered by tDOC. In sum, the disproportional benefit of tDOC would decrease the richness and

286   evenness of the active microbial community.

288 **Species sorting alters community structure**

In addition to changes in diversity, the altered resource environment modified the

290 composition of aquatic bacterial communities by selecting for and against certain taxa (i.e.

species sorting). We observed significant increases and decreases in taxa along the resource

292 gradient. Although comparative studies have observed that DOC is an important factor

structuring bacterial communities (Judd et al. 2006, Jones et al. 2009), here we present some of

294 the first experimental evidence that tDOC supply can structure aquatic bacterial communities by

species sorting.

296 In this study, we observed that some taxa increase in abundance in response to tDOC supply

while other taxa decrease in abundance. For example, there was a 10-fold increase in the relative

298 abundance of OTU008 (*Methylomonas* sp.) and a 10-fold decrease in the relative abundance of

OTU011 (*Rhodococcus* sp.) across the gradient. There are multiple potential explanations for the

300 taxon-specific changes in relative abundance we observed. First, the subsidy could be selecting

against taxa unable to use tDOC while other taxa remain constant. For example, in addition to

302 *Rhodococcus* (OTU011), four taxa belong to the Verrucomicrobia clade decreased in abundance

and activity across the gradient (see Supplemental Table 1). In general, the Verrucomicrobia

304 clade is thought to primarily use labile, algal-derived carbon (Newton et al. 2011). Second, taxa

with beneficial traits (e.g., specialized metabolic functions) could be increasing in abundance

306 while other taxa remain constant. For example, one of the strongest responding taxa in our study

was an Actinobacterium (OTU043) related to the freshwater Actinobacteria tribe acSTL; in

308 addition, we found increases in abundance for other Actinobacteria including members of the

Luna3 and acIII tribes (see Supplemental Table 1). Though few cultured representatives exist,

310 many members of the phylum Actinobacteria are known for the degradation of complex organic

matter (Newton et al. 2011, Nelson and Carlson 2012). Last, because tDOC supply can promote

312 additional microbial metabolisms, distinct groups of microorganisms may benefit from increased

tDOC supply. For example, high concentrations of DOC may enhance lake methane cycling

314 (Bastviken 2004, Lennon et al. 2006). We observed an increase in the relative abundance

Methylomonas (OTU008) across the tDOC supply gradient, and this organism obtains carbon

316 and energy from methane (i.e., methanotrophy). Together, our results provide evidence that

subsidies, such as tDOC, structure bacterial communities via species sorting.

Microbial Generalists and Specialists

320 In our study, we observed taxon-specific shifts in relative abundances, which indicate tDOC

may favor taxa with specialized metabolic traits. Theory suggest that habitats with low

322 concentrations of growth-limiting resources are thought to select for specialists that have lower

minimal resource requirements, while habitats with higher concentrations of growth limiting

324 resources may select for generalists (Wilson and Yoshimura 1994). In contrast, we increased the

supply rate of tDOC, a low-quality complex resource requiring specialized metabolic traits, and

326 observed taxon-specific increases in abundance. If these taxa are using tDOC as a resource, we

can assume that they possess the required metabolic pathways and are specialists. Alternatively,

328 these organisms may not be specialists. Work in marine environments has found that generalist

microbes containing the diverse metabolic pathways are responsible for using tDOC (Mou et al.

330 2008, Newton et al. 2010). Likewise, work in the field of comparative genomics has shown that

organisms that use large carbon resources tend to have larger genomes, which is a common

332 signature of generalist life-history strategies among microorganisms (Livermore et al. 2014). It is

possible, therefore, that some taxa contain specialized metabolic pathways but are in fact carbon

334 substrate generalists rather than specialists. Therefore, we may need to change our expectations

of the resource availability generalist-specialist framework to include resource complexity.

336 When resources are available to all consumers (low complexity) it is clear that resource

availability will favor the consumers with the lowest minimal resource requirements (Tilman

338 1977); however, when resources are limited to consumers with unique traits (complex resources)

then resource availability will favor consumers possessing these traits. Though often ignored,

340 many nutrients – such as nitrogen, phosphorus, and carbon – exist as heterogeneous mixtures of

molecules that differ in quality and bioavailability (Muscarella et al. 2014). As such, it is

342 important to consider properties such as quality when predicting how resource subsidies will

favor generalists versus specialists in a community.

Community Response to Perturbations

346   In addition to compositional changes, tDOC supply rate altered how the bacterial community

responded to an inorganic nutrient perturbation. There are four main predictions for how

348 microbial communities will respond to perturbations: 1) resistant communities will show no

changes; 2) resilient communities will quickly recover after perturbations and resume pre-

350 disturbance structure and function; 3) sensitive, functionally redundant communities will change

but maintain pre-disturbance function, and 4) sensitive, non-redundant communities will change

352 in composition and function (Allison and Martiny 2008). Previous studies using nutrient

perturbations have shown that microbial composition is generally sensitive to perturbation

354 (Allison and Martiny 2008). This is not what we observed. In our study we found that the total

bacterial communities had little to no response to a nutrient perturbation regardless of tDOC

356 supply rate. This indicates that the total community was either resistant or resilient. However, we

found that the active community changed across the tDOC supply gradient and that the active

358   taxa responded less to the nutrient perturbation as tDOC supply increased. Because RNA is more

ephemeral than DNA, we can assume that, if the community changed, the total community

360   (based on DNA) would not return to pre-perturbation composition before the active community

(based on RNA). Therefore, taking both the total and active communities into account, we argue

362   that overall the total bacterial community was resistant to the inorganic nutrient pulse but the

metabolically active taxa responded. In addition, the stability (i.e., resistance) of the

364   metabolically active taxa increased as tDOC supply rate increased. We propose that the observed

changes in the active community represent the functional responsiveness and metabolic plasticity

366   of the bacterial community. Because only the functionally active organisms responded, the

community response to the inorganic nutrient pulse was functional not structural; furthermore,

368   because the functional responsiveness decreased as tDOC supply rate increased, subsidy inputs

yielded a microbial community that was structurally and functionally stable.

Subsidies and Community Stability

372   Overall, experimental tests to determine the effects of resource subsidies on community stability

have been sparse (Nowlin et al. 2007). In this study, we found that a tDOC subsidy stabilized the

374   recipient community by altering the composition of the microbial community through species

sorting. One mechanism would be that the subsidy favored specialized taxa. However, due to

376   their metabolic physiology, these specialized taxa may be relatively slow growing and unable to

rapidly respond to nutrient pulses (Wetzel 1999). An alternative mechanism would be that the

378   subsidy selected against fast-responding taxa. For example, at high tDOC supply shading limits

primary productivity and the aquatic microbial community may receive less labile algal-derived

380   carbon to promote fastidious, rapid responding taxa (Jones and Lennon 2015). Our data suggest

that both of these mechanisms explain changes in composition due to tDOC supply. Therefore,

382   we hypothesize that the quality of the subsidy influences community stability. Here, we added a

low quality subsidy (tDOC) which selected for taxa with specialized metabolic functions and

384   against fastidious taxa which would have been able to rapidly respond to the inorganic nutrient

pulse. Conversely, systems receiving high-quality subsidies would favor more responsive taxa

386   and thus communities would respond differently to perturbations. Therefore, subsidies affect

community stability by altering the composition of the community through species sorting, but

388   the community stability outcome will be dependent on the properties of the subsidy and the traits

of the recipient consumers.

Implications for Aquatic Ecosystems

392   Overall, our findings suggest that the active taxa within the microbial community are

responsible for mediating the changes in composition and stability due to altered subsidy inputs.

394   Because the active taxa control nutrient cycling, we also expect these changes to affect

ecosystem function and stability. First, subsidy-induced changes in community composition will

396   alter ecosystem functions. For example, the degree to which bacterial communities are

physiologically flexible may have important consequences for ecosystem functions such as

398   secondary productivity (Godwin and Cotner 2015), and our data suggest that tDOC subsidies

may select for communities comprised of taxa that are less physiological flexibility and thus

400   unable to rapidly respond to nutrient pulses. Second, changes in community stability may alter

ecosystem stability. For example, enhanced community stability may yield reduced ecosystem

402   stability because either functional or compositional changes are required for the ecosystems to

respond to the nutrient perturbations (Comte and del Giorgio 2011). If inorganic nutrients from a

404 pulse perturbation remain in the recipient ecosystem, due to slow rates of biological processing,

ecosystem functioning will become destabilized (Cottingham and Carpenter 1994). In fact,

406 previous work demonstrated that tDOC inputs diminished nutrient turnover time and thus

destabilized aquatic ecosystem functioning (Jones and Lennon 2015). We hypothesized that the

408 reduced ecosystem stability was due to a lack of microbial functional or structural responses. Our

results confirmed this hypothesis. Together, our results support the view that subsidy supply

410 alters species sorting in ways that alter the stability of recipient communities and ecosystems.

412 <div align="center">**ACKNOWLEDGEMENTS**</div>

This is Kellogg Biological Station contribution number KBS #1906 and this work was supported

414 by the National Science Foundation (DEB-0842441 and DEB-1442246). We thank B.K.

Lehmkuhl for technical assistance and M. Fisk and F. Guillemette for critical feedback on an

416 earlier version of this manuscript. All sequence data and metadata have been submitted to NCBI

and are available at BioProject PRJNA301893. All code for sequence processing and statistical

418 analyses is available at https://github.com/LennonLab/SubsidyMicrobialStability.

REFERENCES

Aanderud, Z., S. Jones, N. Fierer, J. T. Lennon, Z. T. Aanderud, S. Jones, N. Fierer, and J. T.

422   Lennon. 2015. Resuscitation of the rare biosphere contributes to pulses of ecosystem

activity. Frontiers in Microbiology 6:1–11.

424   Allison, S. D., and J. B. H. Martiny. 2008. Resistance, resilience, and redundancy in microbial

communities. Proceedings of the National Academy of Sciences 105:11512–11519.

426   Anderson, M. J. 2001. A new method for non-parametric multivariate analysis of variance.

Austral Ecology 26:32–46.

428   Anderson, M. J., K. E. Ellingsen, and B. H. McArdle. 2006. Multivariate dispersion as a measure

of beta diversity. Ecology Letters 9:683–93.

430   APHA. 1998. Standard Methods for the Examination of Water and Wastewater. (L. S. Clesceri,

A. D. Eaton, and A. E. Greenberg, Eds.). 20th edition. American Public Heath Association,

432   Washingston, D.C.

Bartels, P., J. Cucherousset, K. Steger, P. Eklöv, L. J. Tranvik, and H. Hillebrand. 2012.

434   Reciprocal subsidies between freshwater and terrestrial ecosystems structure consumer

resource dynamics. Ecology 93:1173–82.

436   Bastviken, D. 2004. Methane emissions from lakes: Dependence of lake characteristics, two

regional assessments, and a global estimate. Global Biogeochemical Cycles 18:1–12.

438   Benjamini, Y., and Y. Hochberg. 1995. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society 57:289 –

440   300.

Brown, B. L. 2003. Spatial heterogeneity reduces temporal variability in stream insect communities. Ecology Letters 6:316–325.

Cole, J. R., Q. Wang, E. Cardenas, J. Fish, B. Chai, R. J. Farris, A. S. Kulam-Syed-Mohideen, D. M. McGarrell, T. Marsh, G. M. Garrity, and J. M. Tiedje. 2009. The Ribosomal Database Project: improved alignments and new tools for rRNA analysis. Nucleic Acids Research 37:D141–D145.

Comte, J., L. Fauteux, and P. A. del Giorgio. 2013. Links between metabolic plasticity and functional redundancy in freshwater bacterioplankton communities. Frontiers in Microbiology 4:1–11.

Comte, J., and P. A. del Giorgio. 2011. Composition Influences the Pathway but not the Outcome of the Metabolic Response of Bacterioplankton to Resource Shifts. PLoS ONE 6:e25266.

Cottingham, K. L., and S. R. Carpenter. 1994. Predictive Indexes of Ecosystem Resilience in Models of North Temperate Lakes. Ecology 75:2127–2138.

Cottingham, K., and L. Narayan. 2013. Subsidy quantity and recipient community structure mediate plankton responses to autumn leaf drop. Ecosphere 4:89.

DeAngelis, D. L. 1992. Dynamics of Nutrient Cycling and Food Webs. Chapman and Hall, London, UK.

Elser, J. J., K. Acharya, M. Kyle, J. Cotner, W. Makino, T. Markow, T. Watts, S. Hobbie, W. Fagan, J. Schade, J. Hood, and R. W. Sterner. 2003. Growth rate-stoichiometry couplings in diverse biota. Ecology Letters 6:936–943.

462   Elser, J., E. R. Marzolf, and C. R. Goldrnan. 1990. Phosphorus and Nitrogen Limitation of

        Phytoplankton Growth in the Freshwaters of North America: A Review and Critique of

464   Experiments Enrichments. Canadian Journal of Botany 47:1468–1477.

        Faria, L. D. B., and M. I. D. S. Costa. 2010. Omnivorous food web, prey preference and

466   allochthonous nutrient input. Ecological Complexity 7:107–114.

        Fellman, J. B., M. P. Miller, R. M. Cory, D. V D'Amore, and D. White. 2009. Characterizing

468   dissolved organic matter using PARAFAC modeling of fluorescence spectroscopy: a

        comparison of two models. Environmental Science & Technology 43:6228–34.

470   Fierer, N., M. Hamady, C. L. Lauber, and R. Knight. 2008. The influence of sex, handedness,

        and washing on the diversity of hand surface bacteria. Proceedings of the National

472   Academy of Sciences 105:17994–9.

        Fuchs, G., M. Boll, and J. Heider. 2011. Microbial degradation of aromatic compounds - from

474   one strategy to four. Nature Reviews Microbiology 9:803–816.

        del Giorgio, P. A., and J. M. Gasol. 2008. Physiological structure and single-cell activity in

476   marine bacterioplankton. Pages 243–298 *in* D. L. Kirchman, editor. Microbial Ecology of

        the Oceans. Second edition. John Wiley & Sons, Inc., Hoboken, NJ, USA.

478   Godwin, C. M., and J. B. Cotner. 2015. Aquatic heterotrophic bacteria have highly flexible

        phosphorus content and biomass stoichiometry. The ISME Journal 9:2324–2327.

480   Grimm, V., and C. Wissel. 1997. Babel, or the ecological stability discussions: An inventory and

        analysis of terminology and a guide for avoiding confusion. Oecologia 109:323–334.

482   He, X., M. D. Scheurell, P. A. Soranno, and R. A. Wright. 1994. Recurrent response pattems of a

zooplankton community to whole-lake fish manipulation. Freshwater Biology 32:61–72.

484   Holt, R. D. 2004. Implications of System Openness for Local Community Structure and

Ecosystem Function. Pages 96–114 *in* G. A. Polis, M. E. Power, and G. R. Huxel, editors.

486   Food Webs at the Landscape Level. University of Chicago Press, Chicago.

Huxel, G. R., and K. McCann. 1998. Food web stability: the influence of trophic flows across

488   habitats. The American Naturalist 152:460–9.

Jablonski, D. 2008. Species Selection: Theory and Data. Annual Review of Ecology, Evolution,

490   and Systematics 39:501–524.

Jones, S. E., and J. T. Lennon. 2010. Dormancy contributes to the maintenance of microbial

492   diversity. Proceedings of the National Academy of Sciences 107:5881–6.

Jones, S. E., and J. T. Lennon. 2015. A test of the subsidy – stability hypothesis: the effects of

494   terrestrial carbon in aquatic ecosystems. Ecology 96:1550–1560.

Jones, S. E., R. J. Newton, and K. D. McMahon. 2009. Evidence for structuring of bacterial

496   community composition by organic carbon source in temperate lakes. Environmental

Microbiology 11:2463–72.

498   Judd, K. E., B. C. Crump, and G. W. Kling. 2006. Variation in dissolved organic matter controls

bacterial production and community composition. Ecology 87:2068–79.

500   Karlsson, J., P. Byström, J. Ask, P. Ask, L. Persson, and M. Jansson. 2009. Light limitation of

nutrient-poor lake ecosystems. Nature 460:506–509.

502   Kritzberg, E. S., J. J. Cole, M. L. Pace, W. Granéli, and D. L. Bade. 2004. Autochthonous versus

allochthonous carbon sources to bacteria: Results from whole-lake 13C addition

504   experiments. Limnology and Oceanography 49:588–596.

Lapierre, J.-F., F. Guillemette, M. Berggren, and P. A. Del Giorgio. 2013. Increases in
506   terrestrially derived carbon stimulate organic carbon processing and CO2 emissions in
boreal aquatic ecosystems. Nature Communicationss 4:2972.

508   Legendre, P., and L. F. J. Legendre. 2012. Numerical ecology. Elsevier.

Lennon, J. T., A. M. Faiia, X. Feng, and K. L. Cottingham. 2006. Relative importance of CO2
510   recycling and CH4 pathways in lake food webs along a dissolved organic carbon gradient.
Limnology and Oceanography 51:1602–1613.

512   Lennon, J. T., S. K. Hamilton, M. E. Muscarella, A. S. Grandy, K. Wickings, and S. E. Jones.
2013. A source of terrestrial organic carbon to investigate the browning of aquatic
514   ecosystems. PLOS ONE. PloS One 8:e75771.

Lennon, J. T., and S. E. Jones. 2011. Microbial seed banks: the ecological and evolutionary
516   implications of dormancy. Nature Reviews Microbiology 9:119–30.

Livermore, J. A., S. J. Emrich, J. Tan, and S. E. Jones. 2014. Freshwater bacterial lifestyles
518   inferred from comparative genomics. Environmental Microbiology 16:746–58.

Monteith, D., J. Stoddard, C. Evans, H. A. de Wit, M. Forsius, T. Hogasten, A. Wilander, B. L.
520   Skjelkvåle, D. S. Jeffries, J. Vuoremnaa, B. Keller, J. Kopácek, and J. Vesely. 2007.
Dissolved organic carbon trends resulting from changes in atmospheric deposition
522   chemistry. Nature 450:537–540.

Mou, X., S. Sun, R. A. Edwards, R. E. Hodson, and M. A. Moran. 2008. Bacterial carbon
524   processing by generalist species in the coastal ocean. Nature 451:708–11.

Muscarella, M. E., K. Bird, M. L. Larse, S. A. Placella, and J. T. Lennon. 2014. Phosphorus

526   resource heterogeneity in microbial food webs. Aquatic Microbial Ecology 73:259–272.

Nelson, C. E., and C. A. Carlson. 2012. Tracking differential incorporation of dissolved organic

528   carbon types among diverse lineages of Sargasso Sea bacterioplankton. Environmental

Microbiology 14:1500–16.

530   Neter, J., M. H. Kutner, C. J. Nachtsheim, and W. Wasserman. 1996. Applied linear statistical

models. Irwin, Chicago, Illinois, USA.

532   Newton, R. J., L. E. Griffin, K. M. Bowles, C. Meile, S. Gifford, C. E. Givens, E. C. Howard, E.

King, C. A. Oakley, C. R. Reisch, J. M. Rinta-Kanto, S. Sharma, S. Sun, V. Varaljay, M.

534   Vila-Costa, J. R. Westrich, and M. A. Moran. 2010. Genome characteristics of a generalist

marine bacterial lineage. The ISME Journal 4:784–798.

536   Newton, R. J., S. E. Jones, A. Eiler, K. D. McMahon, and S. Bertilsson. 2011. A guide to the

natural history of freshwater lake bacteria. Microbiology and Molecular Biology Reviews

538   75:14–49.

Nowlin, W. H., M. J. González, M. J. Vanni, M. H. H. Stevens, M. W. Fields, and J. J. Valente.

540   2007. Allochthonous subsidy of periodical cicadas affects the dynamics and stability of

pond communities. Ecology 88:2174–86.

542   Nowlin, W. H., M. J. Vanni, and L. H. Yang. 2008. Comparing Resource Pulses in Aquatic and

Terrestrial Ecosystems. Ecology 89:647–659.

544   Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'Hara, G. L. Simpson,

P. Solymos, M. H. H. Stevens, and H. Wagner. 2013. Vegan: community ecology package.

546   Polis, G. A., W. B. Anderson, and R. D. Holt. 1997. Toward an integration of landscape and food

        web ecology: the dynamics of spatially subsidized food webs. Annual Review of Ecology

548   and Systematics 28:289–316.

        R Core Development Team. 2012. R: A language and environment for statistical computing.

550   Schloss, P. D., S. L. Westcott, T. Ryabin, J. R. Hall, M. Hartmann, E. B. Hollister, R. A.

        Lesniewski, B. B. Oakley, D. H. Parks, C. J. Robinson, J. W. Sahl, B. Stres, G. G.

552   Thallinger, D. J. Van Horn, and C. F. Weber. 2009. Introducing mothur: open-source,

        platform-independent, community-supported software for describing and comparing

554   microbial communities. Applied and Environmental Microbiology 75:7537–7541.

        Shade, A., J. S. Read, D. G. Welkie, T. K. Kratz, C. H. Wu, and K. D. McMahon. 2011.

556   Resistance, resilience and recovery: aquatic bacterial dynamics after water column

        disturbance. Environmental Microbiology 13:2752–67.

558   Smith, B., and J. B. Wilson. 1996. A consumer's guide to evenness indices. Oikos 76:70–82.

        Solomon, C. T., S. E. Jones, B. C. Weidel, I. Buffam, M. L. Fork, J. Karlsson, S. Larsen, J. T.

560   Lennon, J. S. Read, S. Sadro, and J. E. Saros. 2015. Ecosystem Consequences of Changing

        Inputs of Terrestrial Dissolved Organic Matter to Lakes: Current Knowledge and Future

562   Challenges. Ecosystems 18:376–389.

        Takimoto, G. 2002. Seasonal subsidy stabilizes food web. Ecological Research 17:433–439.

564   Tamura, K., G. Stecher, D. Peterson, A. Filipski, and S. Kumar. 2013. MEGA6: Molecular

        Evolutionary Genetics Analysis Version 6.0. Molecular Biology and Evolution 30:2725–

566   2729.

Tilman, D. 1977. Resource competition between plankton algae: an experimental and theoretical

568   approach. Ecology 58:338–348.

Tranvik, L., J. Downing, and J. Cotner. 2009. Lakes and reservoirs as regulators of carbon

570   cycling and climate. Limnology and Oceanography 54:2298–2314.

Wang, Q., G. M. Garrity, J. M. Tiedje, and J. R. Cole. 2007. Naive Bayesian Classifier for Rapid

572   Assignment of rRNA Sequences into the New Bacterial Taxonomy. Applied and

Environmental Microbiology 73:5261–5267.

574   Wetzel, R. G. 1999. Biodiversity and shifting energy stability within freshwater ecosystems.

Arch Hydrobiol Spec Issues Advanc Limnol 54:19–32.

576   Wetzel, R. G., and G. E. Likens. 2000. Inorganic nutrients: nitrogen, phosphorus, and other

nutrients. Pages 85–111 Limnological Analyses. Third edition. Springer-Verlag, New York.

578   Wilkinson, G. M., M. L. Pace, and J. J. Cole. 2013. Terrestrial dominance of organic matter in

north temperate lakes. Global Biogeochemical Cycles 27:43–51.

580   Wilson, D. S., and J. Yoshimura. 1994. On the coexistence of specialists and generalists. The

American Naturalist 144:692–707.

584 **FIGURE LEGENDS**

Fig. 1: Community diversity estimates, richness (A) and evenness (B), for microbial

586 communities in response to tDOC supply rate. For both, the diversity of the total community

(DNA, circles) did not change in response to tDOC supply rate (A: $P = 0.50$; B: $P = 0.32$).

588 However, both the richness and evenness of the active community (RNA, triangles) decreased in

response to tDOC supply rate (A: $P = 0.001$; B: $P = 0.04$).

590 **Fig. 2:** Taxon-specific changes in community composition along the tDOC supply rate gradient.

A) Phylogenetic tree of responsive taxa as indicated by significant Spearman's rank correlations.

592 The consensus tree was inferred by using the Maximum Likelihood methods based on the Jukes-

Cantor model, and 100 bootstraps using MEGA-6 (Tamura et al. 2013). Responses indicate

594 significant increases (↑) or decreases (↓) in relative abundance in as tDOC supply rate increased.

Responses are shown in regards to both the active and total communities. Clade refers to the

596 phylogenetic group (phylum or subphylum) as inferred from best matches to the RDP taxonomy

(Wang et al. 2007). Classifications based on known freshwater bacteria are found in

598 Supplemental Table I. B) Example of significant positive correlation as shown by OTU008

(*Methylomonas* sp.). C) Example of a significant negative correlation as shown by OTU011

600 (*Rhodococcus* sp.). For each, log relative abundances are shown for OTU relative abundance in

the total communities. Dashed lines indicate LOWESS regression trends and are included to

602 demonstrate overall trends.

Fig. 3: Community stability to an inorganic nutrient perturbation along the tDOC supply rate.

604 Responsiveness was measured as the Euclidian distance between paired pre- and post-

perturbation communities in multivariate space using the first three axes. Lower responsiveness

606 indicates a more stable community. Based on the total community (DNA, circles),

responsiveness did not change in as tDOC supply rate increased ($P = 0.37$). However, based on

608   the active community (RNA, triangles), responsiveness decreased as tDOC supply rate increased

($P < 0.001$).

FIGURES

612 **Figure 1:**

Figure 2:

Figure 3: