Thousands of novel translated open reading frames in humans inferred by ribosome footprint profiling.

Anil Raj[1*†], Sidney H. Wang[2*†], Heejung Shim[2,6†], Arbel Harpak[4],
Yang I. Li[1], Brett Engelmann[2], Matthew Stephens[2,3], Yoav Gilad[2*],
Jonathan K. Pritchard[1,4,5*]

[1]Department of Genetics, Stanford University, Stanford, CA

[2]Department of Human Genetics, University of Chicago, Chicago, IL

[3]Department of Statistics, University of Chicago, Chicago, IL

[4]Department of Biology, Stanford University, Stanford, CA

[5]Howard Hughes Medical Institute

[6]Current address: Department of Statistics, Purdue University, West Lafayette, IN

[†]Equal contribution

[*] To whom correspondence should be addressed: rajanil@stanford.edu,
siddisis@uchicago.edu, gilad@uchicago.edu, pritch@stanford.edu

1 **Abstract**

Accurate annotation of protein coding regions is essential for understanding how

genetic information is translated into biological functions. Here we describe

riboHMM, a new method that uses ribosome footprint data along with gene

expression and sequence information to accurately infer translated sequences.

We applied our method to human lymphoblastoid cell lines and identified 7,273

previously unannotated coding sequences, including 2,442 translated upstream

open reading frames. We observed an enrichment of harringtonine-treated

ribosome footprints at the inferred initiation sites, validating many of the novel

coding sequences. The novel sequences exhibit significant signatures of

selective constraint in the reading frames of the inferred proteins, suggesting that

many of these are functional. Nearly 40% of bicistronic transcripts showed

significant negative correlation in the levels of translation of their two coding

sequences, suggesting a key regulatory role for these novel translated

sequences. Our work significantly expands the set of known coding regions in

humans.

1 Introduction

Annotations for coding sequences (CDSs) are fundamental to genomic research. The

GENCODE Consortium (Harrow et al. 2012) has played an important role in annotating

the set of protein coding sequences in the human genome, predominantly relying on

manual annotation from the Human and Vertebrate Analysis and Annotation (HAVANA)

group. Their annotation pipeline identifies coding sequences using homology with

sequences in large cDNA/EST databases and the SWISS-PROT protein sequence

database (Bairoch and Apweiler 2000), and validates them using sequence homology

across vertebrates and using tandem mass spectrometry. Despite being the most

comprehensive database of CDSs available, the current set is conservative and does

not include several classes of CDSs, including translated upstream open reading frames

(ORFs), dually coded transcripts, and N-terminal extensions and truncations.

Recent work has made it increasingly clear that much of the human genome is

transcribed in at least one tissue during some stage of development (Hangauer et al.

2013; Djebali et al. 2012; ENCODE Project Consortium et al. 2007; Clark et al. 2011;

Kapranov et al. 2007; van Bakel et al. 2010). However, the functional implication for

most of these transcripts remains unclear; in particular, the set of sequences translated

from these transcripts are not yet completely characterized. For example, there are

several recent studies in which RNA transcripts that were previously annotated as

noncoding were shown to encode short functional peptides. One well characterized

example is the *polished rice* (*pri*) / *tarsal-less* (*tal*) locus in flies, a polycistronic mRNA

encoding four short peptides active during embryogenesis (Kondo et al. 2007, 2010;

Galindo et al. 2007). While short peptides are known to play critical roles in multiple

biological processes (Lauressergues et al. 2015; Oelkers et al. 2008; Camby et al. 2006; Jung et al. 2009), annotating genomic regions that encode them remains challenging.

Direct proteogenomic mass spectrometry has the potential to fill this gap but suffers from variable accuracy in assignment of peptide sequences to spectra and assignment of identified peptides to proteins (for peptides shared across database entries). Moreover, these approaches suffer from a "needle in a haystack" problem when searching all six translational frames over the transcribed portion of the genome (Nesvizhskii 2014; Pevtsov et al. 2006; Ma 2015). Alternative approaches that utilize empirically-derived phylogenetic codon models to distinguish coding transcripts from non-coding transcripts are promising (Lin et al. 2011). However, the success of such approaches is contingent on the duration, strength and stability of purifying selection and these methods may be underpowered for short coding sequences or for newly evolved coding sequences.

Ribosome profiling utilizes high throughput sequencing of ribosome-protected RNA fragments (RPFs) to quantify levels of translation (Ingolia et al. 2009). Briefly, the technique consists of isolating monosomes from RNase-digested cell lysates and extracting and sequencing short mRNA fragments protected by ribosomes. Early studies of ribosome profiling have shown that RPFs are substantially more abundant within the CDS of annotated transcripts compared to the 5' or 3' untranslated regions (UTRs) (Ingolia et al. 2009; Weinberg et al. 2015). More importantly, when aggregated across annotated coding transcripts, the RPF abundance within the CDS has a clear three base-pair periodicity while the RPF abundance in the UTRs lacks this periodic pattern.

Recently, using ribosome profiling data, several studies reported conflicting results on the coding potential of long intergenic noncoding RNA (Ingolia et al. 2011; Guttman et al.

2013; Ingolia et al. 2014). These studies assessed coding potential using either i) the enrichment of RPFs within the translated CDS relative to background, or ii) the difference in length of RPFs within the translated CDS compared to background. However, these approaches may lack power for several reasons. First, they make little distinction between ribosomes scanning through the transcript and ribosomes decoding the message. Second, the enrichment signal can be severely diminished if the transcript is significantly longer than the coding region within it. Third, there is often substantial variance in RPF abundance within the CDSs, which can decrease power to detect translated sequences when using a simple RPF enrichment statistic alone. Other studies have used the periodicity structure in RPF counts to identify dual coding sequences and short translated CDSs (Michel et al. 2012, Bazzini et al. 2014), but the methods reported high false positive rates and could only identify a few hundred CDSs.

In this work, we developed riboHMM; a model to identify translated CDSs by leveraging both the total abundance and the codon periodicity structure in RPFs. We used this model to identify thousands of novel CDSs in the transcriptome of human lymphoblastoid cell lines (LCLs).

Probabilistic model to infer translated coding sequences

Ribosome footprint profiling data, when aggregated across annotated coding transcripts centered at their translation initiation (or termination) sites (Figure 1A), show two distinct features that distinguish the CDS from untranslated regions (UTRs).

- **Higher abundance within the CDS.** RPF counts are highly enriched within the CDS overall. Moreover, base positions within the CDS close to the translation initiation and termination sites have substantially higher RPF counts compared to

base positions in the rest of the CDS. Untranslated regions have very low RPF counts, with the 5'UTR having a slightly higher RPF count compared to the 3'UTR. Furthermore, base positions in the 5'UTR immediately flanking the initiation site have a slightly higher RPF count compared to the rest of the 5'UTR; a similar pattern is observed in the 3'UTR.

- **Three-base periodicity within the CDS.** RPF counts typically peak at the first position of each codon. The RPF count over the initiation and termination codons tend to have a stronger peak (thus, a slightly different periodic pattern) compared to the rest of the CDS. The RPF counts in the UTRs lack this periodic pattern with similar aggregate counts among base positions in the 5'UTR and 3'UTR.

We developed a framework to infer the translated CDS in a transcript using a model that 1) captures these distinct features of ribosome profiling data and 2) integrates RNA sequence information and transcript expression. As illustrated in Figure 1B, to capture the three-base structure in the RPF count data within the CDS, we represented a transcript as a sequence of non-overlapping base triplets. The CDS of the transcript is required to belong to one of three reading frames. To account for all three reading frames, each transcript has a latent frame variable that specifies at which base position of the transcript we begin enumerating the triplets.

Conditional on the frame, we modeled the data for each transcript, represented as a sequence of base triplets, using a hidden Markov model (HMM). Each triplet belongs to one of nine latent states — 5'UTS (5' Untranslated State), 5'UTS+ (the last untranslated triplet prior to the initiation site), TIS (Translation Initiation State), TIS+ (the triplet immediately following the initiation site), TES (Translation Elongation State), TTS- (the translated triplet prior to the termination site), TTS (Translation Termination State),

3'UTS- (the first untranslated triplet immediately following the termination site), and

3'UTS (3' Untranslated State). The states {TIS, TIS+, TES, TTS-, TTS} denote translated

triplets and {5'UTS, 5'UTS+, 3'UTS-, 3'UTS} denote untranslated triplets. The probability

distribution over the possible sequence of latent states is a function of the underlying

RNA sequence. Figure 1B illustrates these states, and how they relate to each other, in

conjunction with the transcript representation. The groups of states {5'UTS+, TIS, TIS+}

and {TTS-, TTS, 3'UTS-} help model the distinct structure of the RPF counts around the

translation initiation and termination sites, respectively.

Figure 1: Illustrating the model. (A) Proportion of footprint counts aggregated across 1000 highly

expressed annotated coding transcripts, centered at their translation initiation (blue) and termination

(orange) sites. In aggregate, RPF count data have higher abundance within the CDS than the UTRs and

exhibit a 3-base periodicity within the CDS. **(B)** Each transcript belongs to one of three unobserved reading

frames, and is represented as a sequence of base-triplets (highlighted by differing shades of gray) that

depends on the reading frame. Each triplet belongs to one of nine unobserved states. The state sequence

shown corresponds to frame 3 and varying shades from purple to green highlight the different states. Base

positions marked in orange are modeled independently and always belong to the relevant UTS state.

Transitions with nonzero probabilities are indicated by arrows, with solid arrows denoting a probability of 1

and dotted arrows denoting probabilities that are a function of the underlying sequence.

Assuming each transcript has either 0 or 1 CDS, we restricted the possible transitions

between latent states as shown in Figure 1B: transitions from 5'UTS to 5'UTS+ occur

with probability ρ, transitions from TES to TTS- occur with probability ζ, and all other

allowed transitions have probability 1. The transition probabilities ρ and ζ are estimated

from the data, and are allowed to depend on the base sequence of the triplet; in addition, the probability ρ also depends on the base sequence context around the triplet (Kozak 1987). In this work, we assume that translation termination occurs at the first in-frame stop codon (Equation 9), i.e., we do not consider stop codon readthrough.

Conditional on the state assignments, we modeled 1) the total RPF abundance within a triplet, to account for the observation that translated base positions have a higher average RPF count compared to untranslated base positions, and 2) the distribution of RPF counts among the base positions in a triplet, to account for the periodicity in RPF counts within translated triplets. We explicitly accounted for differences in RPF abundance due to differences in transcript expression levels by using transcript-level RNA-seq data as a normalization factor. The short lengths of ribosome footprints mean that many base positions are unmappable; we treated triplets with unmappable positions by modifying the emission probabilities accordingly. Finally, we accounted for the additional variation in RPF counts across triplets assigned to the same state by modeling overdispersion in the triplet RPF abundance (see **Materials and Methods** for details).

To quantify the accuracy of our model, we designed a simulation scheme to estimate what fraction of our inferred translated sequences are false discoveries. We first estimated the Type 1 error rate – i.e., the probability of inferring a translated region when no such region exists – using a set of simulated transcripts that had no signal of translation (null transcripts). The simulated transcripts were constructed by permuting the observed footprint counts in annotated coding transcripts. We then used this estimate to quantify the false discovery rate for each class of translated CDSs identified by riboHMM. Independently, using a simulated set of transcripts containing some signal of translation, we quantified the proportion of transcripts where our model incorrectly

identified the precise translation initiation site conditional on having identified a

translated sequence (see **Materials and Methods** for details on the simulations).

Results

Application to human lymphoblastoid cell lines

We applied riboHMM to infer translated CDSs in human lymphoblastoid cell lines (LCLs)

for which gene expression phenotypes were measured genome-wide: mRNA in 86

individuals, ribosome occupancy in 72 individuals and protein levels in 60 individuals

(Lappalainen et al. 2013; Battle et al. 2015). We first assembled over 2.8 billion RNA

sequencing reads into transcripts using StringTie (Pertea et al. 2015). This assembly

gives us annotated transcripts that are expressed in LCLs, along with novel transcripts

that do not overlap any GENCODE annotated gene. (We do not consider novel isoforms

of annotated genes in our analyses.) Restricting to transcripts with at least five footprints

mapped to each exon, we used riboHMM to identify high-confidence translated CDS. We

learned the maximum likelihood estimates of the model parameters using the top five

thousand highly expressed genes. The estimated parameters are robust to the choice of

the learning set (Figures S1 and S2). Using these parameters, we then inferred the

maximum *a posteriori* (MAP) frame and latent state sequence for each of the assembled

transcripts. We retained transcripts whose MAP frame and state sequence

corresponded to a pair of translation initiation and termination sites and had a joint

posterior probability greater than 0.8. Using a set of simulated null transcripts, we

estimated that this posterior cutoff corresponded to a Type 1 error rate of 4.5% per

transcript. The MAP frame and state sequence directly give us the nucleotide sequence

with the strongest signal of translation; we refer to these as main coding sequences or

mCDS.

Detection of novel CDSs in LCLs

Among 7,801 GENCODE annotated coding genes for which we could infer a high posterior mCDS, we recovered the annotated reading frame for at least one isoform in 7,491 genes (96%); of these, we recovered the exact annotated CDS in 4,500 genes. In 310 genes, we inferred mCDS that were entirely distinct from the annotated CDS. (Figure S3 details the rules that decide how our inference agrees with GENCODE.) Thus, for a subset of GENCODE coding genes, our method identified an mCDS distinct from the annotation. Using simulations, we estimated that riboHMM inaccurately detects a completely novel mCDS at a fairly low rate (Type 1 error rate = 4.5%), but has a higher error rate when identifying the precise translation initiation site (false discovery proportion = 38%; see **Materials and Methods** for details). Thus, it is likely that many of the mCDS that have a distinct reading frame compared to annotation are novel alternate translated sequences; however, for those mCDS that only matched the annotated reading frame, the inferred start sites were false discoveries. Our analysis is also robust to sequencing depth; Figure S4 illustrates that nearly 60% of annotated coding sequences identified with the full data set (580 million footprints) could be accurately recovered even when the sequencing depth was reduced by almost two orders of magnitude.

We identified 4,831 novel mCDS in transcripts expressed in LCLs (FDR = 5.5%). To ensure that these are truly novel, we verified that they do not overlap any known CDS annotated by GENCODE, UCSC (Rosenbloom et al. 2015), or CCDS (Farrell et al. 2014) in the same frame. (See Figure 2 for the different classes of LCL transcripts that contain a novel mCDS; Figure S5 illustrates the decision rules used to identify a novel mCDS). Among these, 814 novel mCDS were identified within GENCODE annotated protein-coding transcripts; in these transcripts, the mCDS encodes for a protein distinct

from that annotated by GENCODE. Of these, 386 mCDS overlap an annotated CDS but

have a different reading frame (labeled 'dual-coding') and 156 do not overlap the

annotated CDS. An example of a novel dual-coding region – an mRNA sequence that

codes for proteins in two different frames – inferred in the POLR2M gene is illustrated in

Figure 3A. Using tandem mass-spectrometry data (Battle et al. 2015), we found four

unique spectra matching peptides in the mCDS and no spectra matching peptides in the

annotated CDS (protein level posterior error probability = $3{\times}10^{-35}$).

Figure 2: Overview of novel coding sequences. The analysis workflow indicates the size of the data (in

read/footprint depth, or number of transcripts) at each step and the numbers and classes of transcript within

which novel translated sequences were identified. Transcripts assembled by StringTie that do not overlap

any annotated gene are called "novel transcripts". Long non-coding RNA includes lincRNAs, antisense

transcripts and transcripts with retained introns, short non-coding RNA includes snRNA, snoRNA and

miRNA, processed transcripts are transcripts without a long, canonical ORF, and pseudogenes include all

subclasses of such genes annotated by GENCODE.

In addition, we identified 2,550 mCDS in annotated non-coding transcripts and 1,019

mCDS within novel transcripts assembled *de novo* by StringTie. Over 60% of the mCDS

in novel transcripts were identified in single-exon transcripts transcribed from regions

containing no annotated genes, while about 8% were identified in novel transcripts

transcribed from the strand opposite to an annotated transcript. Finally, we inferred

mCDS in 448 expressed pseudogenes, among 14,065 pseudogenes annotated in

humans (Pei et al. 2012); nearly 90% of these mCDS were identified in processed

pseudogenes. An mCDS in pseudogene GAPDHP72 is shown in Figure S6A, comparing

the ribosome abundance and peptide matches to the pseudogene mCDS with those of

its parent gene GAPDH.

Figure 3: Thousands of novel translated sequences identified in annotated and novel transcript

isoforms. (A) The inferred CDS for an isoform of the POLR2M gene overlaps its annotated CDS and is in a

different frame. All four distinct peptides uniquely mapping to this gene match the inferred CDS (protein-level

posterior error probability $= 3 \times 10^{-35}$). (The introns and the last exon have been shortened for better

visualization.) **(B)** Distribution of start codon usage across all novel mCDS. **(C)** Distribution of the lengths of

the novel mCDS (gray) compared with the lengths of GENCODE annotated CDSs (black).

Unlike annotated CDS, which almost exclusively start at the methionine codon AUG,

these novel mCDS taken together have a substantially higher usage of non-canonical

codons, particularly CUG (Figure 3B), consistent with recent observations in mouse

embryonic stem cells (Ingolia et al. 2011) and human embryonic kidney cells (Lee et al.

2012). Although riboHMM has a high error rate when identifying translation initiation

sites, our use of a hierarchical model for the initiation sites suggests that the errors in our

inferred start codons are likely to be unbiased. These novel mCDS are also significantly

shorter than annotated CDSs (median lengths 23 vs. 339 amino acids, Mann-Whitney

test p-value $< 2.2 \times 10^{-16}$; Figure 3C). The overall amino acid content within novel mCDS

is comparable to that within annotated CDS, with a slight enrichment for arginine,

alanine, cysteine, glycine, proline, and tryptophan residues (binomial test,

p-value $< 1.1 \times 10^{-16}$; Figure S7).

Below, using an alternative measure of ribosome occupancy, we first assess

independent evidence for translation initiation at many of these novel mCDS. Then, we

test whether these mCDS are functional both using human polymorphism data and using

substitution patterns across vertebrates. Finally, we characterize those mCDS whose

peptide products were identified in mass-spectrometry data.

Translation at novel mCDS validated using harringtonine-treated ribosome

footprints

We next sought to provide independent experimental validation for the novel mCDS. A

direct approach to validate translation initiation sites is to assay ribosome occupancy in

cells treated with harringtonine (Ingolia et al. 2011). Harringtonine interacts with and

arrests the initiation complex while leaving the elongation complex to continue

translating and run off the transcript. Harringtonine-treated ribosome footprint profiling

data therefore show a specific enrichment pattern at the translation initiation site; this

pattern has previously been used to identify translation initiation sites in mouse

embryonic stem cells (Ingolia et al. 2011). We measured harringtonine-treated ribosome

footprints in two LCLs and aggregated the counts of footprints across all novel mCDS.

We observed an enrichment of footprints at the inferred initiation site of the novel mCDS

(binomial test, p-value = 9.5×10^{-79}; Figure 4), similar to the enrichment of aggregate

ribosome occupancy at the initiation sites of a matched number of mCDS that agreed

exactly with the annotated CDS (see Figure S6B for mCDS in pseudogenes). We

observed a significant enrichment at both AUG $\left(p\text{-value} = 5.2 \times 10^{-79} \right)$ and non-AUG

$\left(p\text{-value} = 9.4 \times 10^{-25} \right)$ initiation sites. The reduced enrichment for the novel mCDS

compared to annotated CDSs is likely due to the lower levels of translation of these

novel mCDS and the high error rate in identifying the precise base at which translation is

initiated. Accounting for these limitations, our observation of enrichment suggests that

ribosomes do initiate the translation of many of the novel mCDS identified by riboHMM.

Figure 4: Validation of novel mCDS using harringtonine-treated ribosome profiling data.
Harringtonine-treated ribosome footprints show enrichment at the inferred translation initiation sites, when
aggregated across novel mCDS (orange), similar to the enrichment at the initiation sites of a matched
number of mCDS that agreed exactly with the annotated CDS (purple), suggesting that ribosomes do initiate
translation of the novel mCDS.

Selective constraint on coding function in novel mCDS

We next ascertained the functional importance of these novel mCDS based on the

selective constraint imposed on random mutations that occur within them. A bi-allelic

single nucleotide polymorphism (SNP) that falls within an mCDS can be inferred as

synonymous or nonsynonymous depending on whether switching between the two

alleles of the SNP changes the amino acid sequence of the mCDS. If the mCDS do not

produce proteins that are functionally important, we expect the two classes of variants to

have similar selection pressures on average, and thus to segregate at similar

frequencies. Only if the novel mCDS produce functionally important peptides do we

expect inferred nonsynonymous SNPs to segregate at lower frequencies than inferred

synonymous SNPs.

Figure 5: Novel translated sequences show significant signatures of coding function. (A) Genetic
variants that are nonsynonymous with respect to the inferred mCDS segregate at significantly lower
frequencies in human populations than synonymous variants. The novel regions are under weaker selective
constraint compared to known CDS. (The numbers of variants in each class are matched between novel and
annotated CDS.) **(B)** Scatter plot comparing the substitution rate at inferred synonymous variants versus
inferred nonsynonymous variants for each novel mCDS, computed using multiple sequence alignments
across 100 vertebrate species. Highlighted in red are 232 novel mCDS identified to be under significant
long-term purifying selection after Bonferroni correction (testing for $dN/dS < 1$; p-value $< 2.91 \times 10^{-5}$),
indicating conserved coding function for these sequences.

Starting with biallelic SNPs identified using whole genome sequences of 2,504

individuals (1000 Genomes Project Consortium et al. 2015), we examined the set of

SNPs falling within all novel mCDS (13,907 variants within 3,096 novel mCDS). We

labeled each SNP as synonymous or nonsynonymous with respect to the inferred CDS

and show the cumulative distribution of minor allele frequencies (MAF) of each SNP

class (Figure 5A). We observed that nonsynonymous SNPs have an excess of rare

variants compared with synonymous SNPs (Mann-Whitney test; p-value $= 1.08 \times 10^{-4}$),

implying a difference in the intensity of purifying selection (Nielsen 2005). This observed

excess suggests that the novel mCDS are under significant constraint, consistent with

functional peptides, albeit weaker than at annotated CDS. The mCDS identified within

pseudogenes alone also showed a similar excess of rare variants among

nonsynonymous SNPs (Mann-Whitney test; p-value $= 5.6 \times 10^{-3}$). Such an excess was

not observed for pseudogenes that had detectable ribosome occupancy but lacked a

high-confidence inferred coding sequence (Figure S6C); for these pseudogenes, the

SNPs were labeled based on the reading frame of the parent gene. This highlights that

ribosome occupancy alone is insufficient to identify translated sequences, and our

method is able to leverage finer scale structure in ribosome footprint data to detect

functional coding sequences.

While the allele frequency spectra provide evidence that some of the novel mCDS are

functional in present-day human populations, they are less informative about the long-

term selective constraint on these sequences. To identify whether the novel mCDS have

been under long-term functional constraint, we compared the substitution rates at

synonymous and nonsynonymous sites within the novel mCDS using whole-genome

multiple sequence alignments across 100 vertebrates. (We excluded mCDS identified in

pseudogenes from this analysis due to difficulties in assigning orthology.) In Figure 5B,

232 novel mCDS have a significantly lower nonsynonymous substitution rate (dN)

compared to their synonymous substitution rate (dS) after Bonferroni correction

(p-value $< 2.91 \times 10^{-5}$), suggesting that these mCDS have been under long-term

purifying selection. Since the power to detect significantly low values of dN/dS depends

on the length of the CDS and the qualities of the genome assemblies and the multiple

sequence alignments across distant species at these sequences, the number of

functional novel CDSs identified is a conservative lower bound.

Detection of novel proteins by mass spectrometry

We next tested whether we could detect the novel mCDS predictions using mass

spectrometry data. We used a large data set of SILAC-labeled tandem mass-spectra

generated by trypsin-cleavage of large, stable proteins in many of the same LCLs (Battle

et al. 2015). Running MaxQuant (Cox and Mann 2008) against the sequence database

of 4,831 novel mCDS, at 10% FDR, we identified 161 novel mCDS sequences that have

at least one unique peptide hit -- a tryptic peptide that matches a mass-spectrum (Table

S1). More than 70% of novel mCDS with a peptide hit have at least 2 distinct peptides

matched to it and, in almost all cases, the unique peptides were independently identified

in two or more LCLs (Figure S8).

To assess how many hits we would expect to the novel mCDS if their properties were

like those of annotated CDSs, we developed a model that predicts whether an annotated

protein has at least one mass-spectrum match, using features based on expression and

sequence composition of the protein (see **Materials and Methods** for more details). The

mass-spectrometry data are highly biased towards detection of larger and more highly

expressed proteins. Furthermore, the trypsin cleavage step of the experimental protocol

imposes strong constraints on the set of unique peptide sequences that can be observed

in an experiment. Assuming that the distributions of these predictive features estimated

from annotated CDSs can be applied to the novel mCDS, we computed the expected

number of novel mCDS with a peptide hit to be 603.

We thus find many fewer mass spectrometry hits to the novel mCDS than expected from

a model calibrated on previously annotated mCDS (161 vs. 603). The Harringtonine

data argue that many of the novel mCDS are correct predictions, thus we suggest that

some other property of the mCDS may explain their low detection rate. In particular, it is

possible that the novel proteins may have higher turnover rates than annotated proteins.

For example it is possible that the proteins translated from novel mCDS may have

substantially lower half-life than annotated proteins, or may be secreted, and thus have

too low concentrations within the cell to be detectable by mass spectrometry assays.

Translation of short alternate coding sequences in addition to the mCDS

Protein-coding transcripts in eukaryotes are typically annotated to have only one CDS

(i.e., they are monocistronic). However, a number of studies have demonstrated that

ribosomes can initiate translation at alternative start codons (Xu et al. 2010; Ingolia et al.

2011; Lee et al. 2012) and many others have identified transcripts with alternative CDSs

encoding functional peptides (Vanderperre et al. 2013; Kochetov 2008; Barbosa et al.

2013). Furthermore, anecdotal evidence has suggested that translation of the alternate

CDS serves as a mechanism to suppress translation of the main CDS (Lee et al. 2002;

Hernández-Sánchez et al. 2003; Lammich et al. 2004). However, assessing such a

mechanism genome-wide has been challenging, mainly due to a lack of appropriate

annotations (Calvo et al. 2009).

To this end, we adapted our approach to identify additional coding sequences within

transcripts that are translated in LCLs. Assuming that the sub-codon structure of

footprint abundance is similar between the main and alternate CDS, we identified 2,442

novel CDSs upstream of the mCDS inferred by our method (FDR = 5%); we call them

upstream alternate coding sequences or uaCDS (see **Materials and Methods** for more

details; see also Figure S9). Figure 6A illustrates the ribosome footprint density within

the uaCDS of the transmembrane gene TM7SF2, and its conservation across mammals.

We find strong enrichment of harringtonine-treated ribosome footprints at the initiation

sites of uaCDS similar to the initiation sites of mCDS in the same transcripts (Figure 6B).

Using mass-spectrometry data, we identified 46 uaCDS that have at least one peptide

hit, substantially lower than the expectation of 891 hits predicted by our model. Finally,

comparing the substitution rates at inferred synonymous and nonsynonymous sites, we

identified over 317 uaCDS with highly constrained coding function (Figure 6C).

Figure 6: Short translated sequences identified upstream of thousands of translated main coding

sequences. (A) An alternate, novel CDS was identified upstream of the inferred main CDS in gene

TM7SF2. As shown in its protein sequence alignment across mammals, the uaCDS (in particular, the start
and stop codons) is highly conserved with dN/dS = 0.14. **(B)** Harringtonine-treated ribosome footprints show
strong enrichment at the inferred initiation sites of uaCDS, comparable to the enrichment at the initiation
sites of the corresponding mCDS, suggesting that ribosomes do initiate translation of these uaCDS. **(C)**
Using multiple sequence alignment across 100 vertebrate species, 317 uaCDS were identified to have
strong, significant long-term conservation.

Translation of uaCDS negatively correlates with translation of mCDS

With 2,442 uaCDS identified as translated in LCLs, we next tested the hypothesis that

uaCDS expression negatively correlates with mCDS for each pair. We observed that, at

10% FDR, 917 pairs of uaCDS and mCDS had significant negative correlations across

individuals between the proportion of footprints assigned to them (Figure 7A). Our

observation that nearly 40% of pairs of uaCDS and mCDS are significantly anti-

correlated, despite incomplete power due to low sample size, suggests that a key role of

alternate CDSs in a transcript is to regulate the translation of the main CDS.

Figure 7: Translation of uaCDS regulates translation of mCDS. (A) Spearman correlation, across LCLs,
between mCDS translation and uaCDS translation (red histogram). Using random (mCDS, uaCDS) pairs,
matched for length and pairwise distance, we computed an empirical null distribution of Spearman
correlations (gray histogram). At 10% FDR, 917 inferred (uaCDS, mCDS) pairs have significant negative
correlation (shaded red region). **(B)** 365 orfQTLs (genetic variants associated with ORF usage; i.e., whether
the mCDS or uaCDS of a transcript is translated) were identified at 10% FDR (41 pairs of mCDS/uaCDS).
(C) Illustrating an example of an orfQTL in the histocompatibility minor serpin domain-containing (HMSD)
gene (introns removed for better visualization). The most significant variant (marked x) lies within an intron
between the mCDS and uaCDS of the transcript.

Variation in ORF usage can be driven by a number of factors including *cis* genetic

effects and *trans* effects like variation in expression of RNA binding proteins. To identify

cis variants that affect ORF usage in a bicistronic transcript, we tested for association of

the proportion of RPFs assigned to the mCDS (or uaCDS) with variants in a 10-kilobase

window around the transcript; this phenotype effectively controls for variation in gene

expression across the LCLs. We identified 365 *cis* orfQTLs (genetic variants associated

with ORF usage) across 41 pairs of mCDS and uaCDS at 10% FDR (Figure 7B). In

Figure 7C, we illustrate an example of an orfQTL in a bicistronic transcript of the HMSD

gene (histocompatibility minor serpin domain-containing); this gene is also known to

have a distinct genetic variant associated with alternative usage of two coding isoforms

(Kawase et al. 2007). Our observation of orfQTLs in a number of genes distinguishes

ORF usage as an additional layer of post-transcriptional regulation of protein expression.

Discussion

We developed riboHMM, a mixture of hidden Markov models to accurately resolve the

precise set of mRNA sequences that are being translated in a given cell type, using

sequenced RPFs from a ribosome profiling assay, sequenced reads from an RNA-seq

assay and the RNA sequence. When applied to human LCLs, this method was able to

accurately identify the translated frame in 96% of annotated coding genes that had a

high posterior mCDS. In addition, a key advantage of our framework is the ability to infer

novel translated sequences that may be missed by annotation pipelines that focus on

long CDSs (>100 amino acids), conservation based approaches that require long-term

purifying selection, or direct proteomics measurements that are biased toward highly

expressed, stable proteins. We used riboHMM to identify 7,273 novel CDSs, including

448 of novel translated sequences in pseudogenes and 2,442 bicistronic transcripts that

contain an upstream CDS in addition to a main CDS. We observed enrichment in harringtonine-arrested ribosome occupancy at the inferred translation initiation sites, suggesting that many of the novel mCDS are real. These novel sequences showed significant differences in the amount of purifying selection acting on inferred nonsynonymous versus synonymous sites, suggesting that many of these sequences are conserved as functional peptides, including those mCDS identified in lncRNAs, pseudogenes and novel transcripts.

One caveat of our model is its restriction on one CDS per transcript. In this study, we worked around this limitation using a greedy approach and identified thousands of transcripts with multiple CDSs (either two non-overlapping inferred CDSs or an inferred mCDS distinct from the annotated CDS). Indeed, in some instances where the frame of the mCDS and annotated CDS of a transcript disagreed, we found strong support from mass-spec data for the inferred mCDS frame (Figure 3A). These observations highlight the existence of a large number of transcripts in humans that have multiple CDSs and the variation in alternative usage of CDSs across tissues, an area that has largely been overlooked. Additionally, riboHMM does not effectively distinguish footprints arising from different isoforms and, thus, cannot resolve overlapping translated sequences from multiple coding isoforms of a gene. Extending riboHMM to model multiple, possibly overlapping CDSs jointly across multiple isoforms could help uncover this additional layer of complexity in the human genome.

While the precise function of these novel CDSs remains unclear, we found evidence supporting a regulatory role for novel alternate CDSs identified upstream of the mCDS (uaCDS). Although it is unclear whether the down regulation of mCDS by uaCDS is dependent on the peptide sequences of uaCDS, our finding is consistent with previous

assertions under which translation of upstream ORFs regulates translation of the main

CDS in cap-dependent translation initiation (Morris and Geballe 2000).

Our method provides an alternative framework for annotating the coding elements of the

genome. Compared to current methods that use sequence information in cDNA and

protein databases and those that rely on high-quality genome annotations in closely

related species, riboHMM provides a relatively unbiased CDS annotation and

opportunities for finding entirely novel CDSs. In particular, one could use riboHMM to

identify the set of CDS for a species within a poorly annotated evolutionary clade, using

ribosome profiling and RNA seq data immediately after its genome is sequenced and

assembled. In addition, given ribosome footprint profiling data from multiple cell types,

riboHMM can be used to investigate cell-type-specific translation of coding elements

beyond cell-type-specific gene or isoform expression. These features render this tool

particularly useful in studying molecular evolution of newly arisen coding genes and

linking tissue-specificity of CDS usage to disease.

Materials and Methods

Assembling expressed transcripts in LCLs

We mapped paired-end 75bp RNA-seq reads pooled across 85 Yoruba lymphoblastoid cell lines (Lappalainen et al. 2013) to the Genome Reference Consortium Human Reference 37 (GRCh37) assembly using STAR (Dobin et al. 2013), with the additional flag --outSAMstrandField intronMotif to aid transcript assembly downstream, resulting in 2.8 billion uniquely mapped fragments. Using the mapped reads, we assembled models of transcripts expressed in LCLs using StringTie v1.0.4 (Pertea et al. 2015), and used GENCODE v.19 transcript models to guide the assembly. In addition, we required that the lowest expressed isoform of a gene have no less than 1% the expression of the highest expressed isoform (-f 0.01), and that each exon-exon junction be supported by at least 2 spliced reads (-j 2). Since the RNA-seq protocol did not produce strand-specific reads, we treated the forward strand and reverse strand of a transcript model assembled by StringTie as distinct transcripts. Our final set of 430,754 expressed transcripts included 122,168 GENCODE annotated transcript isoforms and 308,586 novel isoforms. (We did not consider novel isoforms of annotated genes identified by StringTie.)

Ribosome footprint profiling

Ribosome footprint profiling experiments and sequencing data processing were performed as previously described (Battle et al, 2015), with the exception of a harringtonine treatment step to arrest ribosomes at the sites of translation initiation. Briefly, lymphoblastoid cell lines, GM19204 and GM19238, were cultured at 37°C with 5% CO_2 in RPMI media with 15% FBS. The media were further supplemented with 2 mM L-glutamate, 100 IU/ml penicillin, and 100 µg/ml streptomycin. Right before cell lysate preparation, each culture was treated with 2 µg/ml harringtonine (final concentration in media) for 2 minutes followed by 100 µg/ml cycloheximide (final concentration in media). For ribosome profiling experiments, ARTseq™ Ribosome Profiling kit for mammalian cells (RPHMR12126) was used following vendor's instructions. Sephacryl S400 spin columns (GE; 27-5140-01) were used for monosome isolation. Libraries were sequenced on an Illumina HiSeq 2500. For sequencing data processing and mapping, adaptor sequences were removed from the 3' end of each read using the Clipper tool from the FASTX-Toolkit. In addition, the 5' most nucleotide (commonly resulted from non-templated additions) was removed using the Trimmer tool from the FASTX-Toolkit. To increase mapping efficiency, we filtered out sequence reads that mapped to rRNA, tRNA or snoRNA (FASTA files downloaded from Ensembl on 05/02/13) using Bowtie 2, version 2.0.2 (Langmead and Salzberg 2012). Processed reads were aligned to genome build hg19 (human) using TopHat v2.0.6 (Trapnell et al. 2009). The mapping step was guided by transcriptome annotations (downloaded from Ensembl on 01/31/13).

1 Mixture of HMMs to model translated coding sequences

Consider N transcripts where the n^{th} transcript has length of L_n assumed to be a multiple

of three ($L_n = 3M_n$; see ***Transcripts with length not a multiple of three*** for details on

how our model handles the remaining one or two base positions when L_n is not a multiple

of three). Our data consist of RPF counts $T = (T^n)_{n=1}^N$, RNA sequence $S = (S^n)_{n=1}^N$, and

transcript expression $E = (E^n)_{n=1}^N$ (in units of RNA-seq reads per base position per million

sequenced reads) on N transcripts, where T^n and S^n are vector quantities and E^n is a

scalar aggregated over the entire length of the transcript. Let $T^n = (T_1^n, \ldots, T_{L_n}^n)$ and $S^n =$

$(S_1^n, \ldots, S_{L_n}^n)$, where T_b^n and S_b^n denote the RPF counts and the base at the b^{th} position

in the n^{th} transcript, respectively. We model the footprint data T using a mixture of HMMs

that incorporates S and E. Assuming independence across transcripts, the probability of

T given S and E is written as $\mathrm{P}(T|\Theta, S, E) = \prod_{n=1}^N \mathrm{P}(T^n|\Theta, S^n, E^n)$, where Θ denotes the

set of model parameters.

Mixture of three reading frames for a transcript: To capture the three-base struc-

ture in RPF data within the CDS, we represent each transcript as a sequence of non-

overlapping base triplets, some of which potentially represent codons. Since the CDS

of the transcript could belong to one of three reading frames (as illustrated in Figure

1B), we introduced a latent frame variable, $F^n \in \{1, 2, 3\}$, that specifies the reading

frame for the n^{th} transcript. Then, given $F^n = f$, T^n can be represented as a se-

quence of $M_n - 1$ triplets and three remaining base positions (see Figure 1B). Specifically,

$T^n \mid F^n = f := (X_{f,1}^n, \ldots, X_{f,(M_n-1)}^n, R_f^n)$, where $X_{f,m}^n = (T_{3m-3+f}^n, T_{3m-2+f}^n, T_{3m-1+f}^n)$

and

$$
R_f^n = \begin{cases}
(T_{L_n-2}^n, T_{L_n-1}^n, T_{L_n}^n) & \text{if } f = 1, \\
(T_1^n, T_{L_n-1}^n, T_{L_n}^n) & \text{if } f = 2, \\
(T_1^n, T_2^n, T_{L_n}^n) & \text{if } f = 3.
\end{cases}
\tag{1}
$$

1 The probability of T^n is then given by

$$P(T^n|\Theta, S^n, E^n) = \sum_{f=1}^{3} P(T^n|F^n = f, \Theta, S^n, E^n)P(F^n = f|\Theta, S^n, E^n), \tag{2}$$

$$= \sum_{f=1}^{3} P(X_{f,1}^n, \ldots, X_{f,(M_n-1)}^n, R_f^n|F^n = f, \Theta, S^n, E^n)P(F^n = f|\Theta, S^n, E^n). \tag{3}$$

2 We assumed that the probability over F^n is independent of S^n and E^n, and is uniform over

3 all three frames, $P(F^n = f|S^n, E^n, \Theta) = \frac{1}{3}$. In addition, we assumed that the RPF data

4 from the sequence of triplets and the RPF data from the three remaining base positions

5 are independent, leading to

$$P(X_{f,1}^n, \ldots, X_{f,(M_n-1)}^n, R_f^n|F^n = f) = P(X_{f,1}^n, \ldots, X_{f,(M_n-1)}^n|F^n = f)P(R_f^n|F^n = f). \tag{4}$$

(For notation convenience, we have dropped highlighting the dependence of X^n and R^n

on $\Theta, S^n, \text{and } E^n$.) We modeled the probability of the data from the sequence of triplets,

$P(X_{f,1}^n, \ldots, X_{f,(M_n-1)}^n|F^n = f)$, using an HMM, and the probability of the data from the

remaining positions, $P(R_f^n|F^n = f)$, using a Poisson-gamma model as described below.

HMM for each frame of a transcript: The pattern of RPF count data in triplets depends

on whether the triplet is being translated or not. To model these patterns, we assumed that

each triplet belongs to one of nine states (see Figure 1B): $5'$ Untranslated State ($5'$UTS),

last untranslated triplet $5'$ to the CDS ($5'$UTS+), Translation Initiation State (TIS), state after

TIS (TIS+), Translation Elongation State (TES), state before TTS (TTS−), Translation Ter-

mination State (TTS), first untranslated triplet $3'$ to the CDS ($3'$UTS−), and $3'$ Untranslated

State ($3'$UTS). The five states (TIS, TIS+, TTS, TTS−, TES) denote translated triplets and

the remaining four states ($5'$UTS, $5'$UTS+, $3'$UTS, $3'$UTS−) denote untranslated triplets. In

particular, the start codon corresponds to the base triplet assigned to the TIS state and the

stop codon corresponds to the base triplet assigned to the $3'$UTS− state. The groups of

states ($5'$UTS+, TIS, TIS+) and (TTS−, TTS, $3'$UTS−) help model the distinct features of the

footprint data around the translation initiation and termination sites, respectively. We intro-

duced a sequence of $M_n - 1$ hidden variables $Z_f^n = (Z_{f,1}^n, \ldots, Z_{f,(M_n-1)}^n)$ for each frame of

the n^{th} transcript, where $Z_{f,m}^n$ denotes the state for the m^{th} triplet in the f^{th} frame.

For each state, an emission probability for $X_{f,m}^n$ can be modeled as follows. Let $Y_{f,m}^n$

denote the sum of three elements in $X_{f,m}^n$ (i.e., the total RPF count for the m^{th} triplet).

Then, $\mathrm{P}(X_{f,m}^n|Z_{f,m}^n = z) = \mathrm{P}(X_{f,m}^n|Y_{f,m}^n, Z_{f,m}^n = z)\mathrm{P}(Y_{f,m}^n|Z_{f,m}^n = z)$ and

$$X_{f,m}^n|Y_{f,m}^n, Z_{f,m}^n = z \sim \mathrm{multinomial}(Y_{f,m}^n, \pi_z), \tag{5}$$

$$Y_{f,m}^n|Z_{f,m}^n = z \sim \mathrm{Poisson}(\mu_{zfm}^n E^n), \tag{6}$$

$$\mu_{zfm}^n \sim \mathrm{gamma}(\alpha_z, \beta_z), \tag{7}$$

where the density of the gamma distribution is $\mathrm{P}(\mu) = \frac{\beta^{\alpha\beta}}{\Gamma(\alpha\beta)}\mu^{\alpha\beta-1}\exp^{-\beta\mu}$ with the mean and variance equal to α and $\frac{\alpha}{\beta}$, respectively.

The periodicity of RPF counts within the CDS is captured by the multinomial distribution with parameters $\pi_z = (\pi_{z,1}, \pi_{z,2}, \pi_{z,3})$, where we assume $\pi_z = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ for $z \in \{5'\mathrm{UTS}, 5'\mathrm{UTS}+, 3'\mathrm{UTS}, 3'\mathrm{UTS}-\}$ to capture the lack of periodicity in the RPF data in untranslated regions. Furthermore, we allow the pattern of periodicity to differ across five types of codons (TIS, TIS+, TTS, TTS−, TES).

The Poisson distribution for $Y_{f,m}^n$ captures the difference in RPF abundance between translated and untranslated regions (precisely, difference in abundance between triplets in different states). We corrected for differences in RPF abundance across transcripts due to differences in transcript expression levels by using E^n as a transcript-specific normalization factor (see Figure S10). To account for additional variation in the RPF counts across triplets in the same state (e.g., due to varying translation rates across transcripts, and translational pausing), we allowed for triplet-specific parameters μ_{zfm}^n in the Poisson intensity and assumed that those parameters follow a gamma distribution. Under this model, $\mathrm{E}[Y_{f,m}^n|Z_{f,m}^n = z] = \alpha_z E^n$ and $\mathrm{Var}[Y_{f,m}^n|Z_{f,m}^n = z] = \frac{\alpha_z}{\beta_z}(E^n)^2 + \alpha_z E^n$.

We assumed that the sequence of hidden variables Z_f^n follow a Markov chain. The assumption of up to one CDS in each transcript leads to a transition probability shown in Figure 1B, where $\rho_{f,m}^n = \mathrm{P}(Z_{f,m+1}^n = 5'\mathrm{UTS}+|Z_{f,m}^n = 5'\mathrm{UTS})$ and $\zeta_{f,m}^n = \mathrm{P}(Z_{f,m+1}^n = \mathrm{TTS}-|Z_{f,m}^n = \mathrm{TES})$ depend on the underlying RNA sequence and are given by

$$\rho_{f,m}^n = \begin{cases} \mathrm{logistic}\left(\psi_\kappa K_{f,m+2}^n + \sum_{c\in\Omega_{\mathrm{start}}} \psi_c \mathbb{1}\left[M_{f,m+2}^n = c\right]\right) & \text{if } M_{f,m+2}^n \in \Omega_{\mathrm{start}}, \\ 0 & \text{otherwise} \end{cases} \tag{8}$$

$$\zeta_{f,m}^n = \begin{cases} 1 & \text{if } M_{f,m+3}^n \in \Omega_{\mathrm{stop}}, \\ 0 & \text{otherwise} \end{cases} \tag{9}$$

where $\mathbb{1}\left[\,\cdot\,\right]$ is the indicator function, $M_{f,m}^n = (S_{3m-3+f}^n, S_{3m-2+f}^n, S_{3m-1+f}^n)$ denotes the

base sequence of the m^{th} triplet, and $K_{f,m}^n$ denotes the log of ratio of likelihood under

the Kozak model to likelihood under a background model of the base sequence flanking

the m^{th} triplet (see **Kozak model** for details). In our analysis, Ω_{start} contained the canoni-

cal start codon and all near-cognates, $\Omega_{\text{start}} = \{\mathrm{AUG, CUG, GUG, UUG, AAG, ACG, AGG,}$

$\mathrm{AUA, AUC, AUU}\}$, and Ω_{stop} contained the canonical stop codons, $\Omega_{\text{stop}} = \{\mathrm{UAA, UAG,}$

$\mathrm{UGA}\}$. The parameters, ψ_c and ψ_κ, indicate the importance of the triplet base sequence

and the flanking base sequence in determining transition from untranslated triplets to trans-

lated triplets. The current specification of $\zeta_{f,m}^n$ and Ω_{stop} forces the coding sequence to

terminate at the first in-frame occurence of a stop codon. This model can be extended to

account for stop codon read-through by using a logistic function for $\zeta_{f,m}^n$ for the same set

Ω_{stop}.

Model for R_f^n: We model R_f^n, the RPF counts at bases before or after the sequence

of triplets (see Equation (1)), using the emission probabilities of the $5'$UTS or $3'$UTS

states. Assuming that the three elements of R_f^n are independent, we have $\mathrm{P}(R_f^n | F^n =$

$f) = \prod_{i=1}^3 \mathrm{P}(R_{f,i}^n | F^n = f)$. Each element can be modeled as

$$R_{f,i}^n \sim \mathrm{Poisson}(\frac{1}{3}\lambda_{fi}^n E^n), \tag{10}$$

$$\lambda_{fi}^n \sim \mathrm{gamma}(\alpha_z, \beta_z), \tag{11}$$

where $z = 5'$UTS if $R_{f,i}^n \in \{T_1^n, T_2^n\}$, and $z = 3'$UTS if $R_{f,i}^n \in \{T_{L_n-2}^n, T_{L_n-1}^n, T_{L_n}^n\}$.

Parameter estimation and inference: We used an EM algorithm to compute the maxi-

mum likelihood estimate for the model parameters $\Theta = \{\pi_z, \alpha_z, \beta_z, \psi_\kappa, \psi_c\}$, that is, $\hat{\Theta} :=$

$\mathrm{argmax}_\Theta \mathrm{P}(T | \Theta, S, E)$.

To infer the translated CDS for the n^{th} transcript, we identified the frame and state se-

quence that maximizes the joint posterior probability

$$(z^{n*}, f^{n*}) := \mathrm{argmax}_{z,f} \mathrm{P}(Z_f^n = z, F^n = f | T^n, S^n, E^n, \hat{\Theta}). \tag{12}$$

23   We first computed the maximum *a posteriori* (MAP) state sequence for each reading frame

24   using the Viterbi algorithm, $z_f^{n*} := \mathrm{argmax}_z \mathrm{P}(Z_f^n = z | F^n = f, T^n, S^n, E^n, \hat{\Theta})$ for $f = 1, 2, 3$.

25   Then, the MAP state sequence and frame is given as

$$(z^{n*}, f^{n*}) = \mathrm{argmax}_f \mathrm{P}(Z_f^n = z_f^{n*} | F^n = f, T^n, S^n, E^n, \hat{\Theta})\mathrm{P}(F^n = f | T^n, S^n, E^n, \hat{\Theta}), \tag{13}$$

where $z_f^{n^*}$ is a function of f, $P(F^n = f|T^n, S^n, E^n, \hat{\Theta}) \propto P(T^n|F^n = f, S^n, E^n, \hat{\Theta})P(F^n =$

$f)$ and $P(T^n|F^n = f, S^n, E^n, \hat{\Theta})$ is the probability of the data marginalized over the latent

states.

In our analyses, we estimated the model parameters using the top five thousand highly

expressed genes. Then, we inferred the translated CDS for those transcripts in which

each exon has at least five distinct ribosome footprints mapping to it. We restricted our

further analyses to transcripts where (1) $P(Z_f^n = z^{n^*}, F^n = f^{n^*}|T^n, S^n, E^n, \hat{\Theta}) > 0.8$, (2)

the MAP state sequence z^{n^*} contains a TIS state and a TTS state (i.e., a pair of initiation

and termination sites), (3) more than 50% of base positions within the inferred CDS are

mappable, and (4) the coding sequence encodes a peptide more than 6 amino acids long

– we call these translated sequences as main coding sequences or mCDS.

Modeling ribosome footprints of different lengths: We observed that ribosome foot-

prints with different lengths, arising due to incomplete nuclease digestion, show slightly

different patterns of abundance when aggregated across transcripts (see Figure S11). To

model these differences, we partition the footprints into multiple groups based on length,

and model the data in each group with a separate set of parameters in the emission prob-

ability (all groups share the same state sequence along a transcript). Specifically, for G

groups of footprints, the data at the m^{th} triplet in f^{th} reading frame $X_{f,m}^n$ can be partitioned

into G components, $X_{f,m}^n = (X_{g,fm}^n)_{g=1}^G$, where $X_{g,fm}^n$ denotes the triplet of RPF counts

from g^{th} group. Assuming that the RPF counts from different groups at a given triplet are

independent, conditional on the state of the triplet, the emission probability can be written

as $P(X_{f,m}^n|Z_{f,m}^n = z) = \prod_{g=1}^G P(X_{g,fm}^n|Z_{f,m}^n = z)$ and

$$X_{g,fm}^n|Y_{g,fm}^n, Z_{f,m}^n = z \sim \text{multinomial}(Y_{g,fm}^n, \pi_{g,z}), \tag{14}$$

$$Y_{g,fm}^n|Z_{f,m}^n = z \sim \text{Poisson}(\mu_{g,zfm}^n E^n), \tag{15}$$

$$\mu_{g,zfm}^n \sim \text{gamma}(\alpha_{g,z}, \beta_{g,z}), \tag{16}$$

where group-specific parameters, $(\pi_{g,z}, \alpha_{g,z}, \beta_{g,z})$, capture the distinct patterns in each

group. The RPF data used in our analyses had four groups of footprints of lengths 28, 29,

30, and 31 bases.

Base positions with missing data: Approximately 15% of the transcriptome have un-

mappable base positions, in part due to the short lengths of ribosome footprints. Consider

the m^{th} base triplet in frame f in the n^{th} transcript. If $\mathcal{J}_{g,fm}^n$ is the set of positions in this

triplet that are unmappable for footprints corresponding to group g, the emission probabili-

ties become

$$X_{g,fm}^n | Y_{g,fm}^n, Z_{f,m}^n = z \sim \text{multinomial}(Y_{g,fm}^n, \tilde{\pi}_{g,z}), \tag{17}$$

$$Y_{g,fm}^n | Z_{f,m}^n = z \sim \text{Poisson}(\psi_{g,zfm}^n \mu_{g,zfm}^n E^n), \tag{18}$$

$$\mu_{g,zfm}^n \sim \text{gamma}(\alpha_{g,z}, \beta_{g,z}), \tag{19}$$

4 where

$$\psi_{g,zfm}^n = \sum_{j \notin \mathcal{J}_{g,fm}^n} \pi_{g,zj}, \tag{20}$$

$$\tilde{\pi}_{g,zj} = \begin{cases} 0 & \text{if } j \in \mathcal{J}_{g,fm}^n, \\ \frac{\pi_{g,zj}}{\psi_{g,zfm}^n} & \text{otherwise.} \end{cases} \tag{21}$$

If all three positions in a triplet are unmappable, then we treat the triplet as having missing

data for that footprint group and set $\text{P}(X_{g,fm}^n | Z_{f,m}^n) = 1$ for all values of $Z_{f,m}^n$.

Kozak model: Using the annotated initiation sites of GENCODE annotated coding tran-

scripts, we estimated a position weight matrix (PWM) that captures the base composition

of the -9 to +6 positions flanking known initiation sites. Since the consensus sequence

of this PWM is the same as the reported consensus Kozak sequence (Kozak, 1987), we

refer to this model as the Kozak model. We estimated a background PWM model using

the same set of positions relative to random AUG triplets within the same set of transcripts.

For the m^{th} triplet in frame f in the n^{th} transcript, using the base sequence from the -9

to +6 positions flanking this triplet, we computed $K_{f,m}^n$, the log of ratio of likelihood of the

flanking sequence under the Kozak model to likelihood under the background model.

Transcripts with length not a multiple of three: The length of such a transcript can be

written as $L_n = 3M_n + B$, where $B \in \{1, 2\}$. We assumed that the RPF data on the first

$3M_n$ bases ($T_{1:3M_n}^n$) and the data on the remaining B bases ($T_{3M_n+1:L_n}^n$) are independent.

We modeled $T_{1:3M_n}^n$ using a mixture of HMMs as described above, and modeled $T_{3M_n+1:L_n}^n$

using the emission probability of the $3'$UTS state as follows.

$$P(T^n_{3M_n+1:L_n}|E^n, \alpha_z, \beta_z) = \prod_{m=3M_n+1}^{L_n} P(T^n_m|E^n, \alpha_z, \beta_z), \tag{22}$$

$$T^n_m \sim \text{Poisson}(\frac{1}{3}\tau^n_m E^n), \tag{23}$$

$$\tau^n_m \sim \text{gamma}(\alpha_z, \beta_z), \tag{24}$$

where $z = 3'$UTS.

Quantifying false discoveries of riboHMM

We characterize the performance of riboHMM by addressing two scenarios: (1) How often does riboHMM identify an mCDS in transcripts with no signal of translation? (2) How often does riboHMM identify an incorrect initiation site in transcripts with signal for translation? To address the first question, we started with the transcripts for which riboHMM was able to identify an mCDS and generated a set of "null transcripts" by permuting the footprint counts among base positions within each transcript. Applying a posterior cutoff of 0.8, riboHMM incorrectly identified an mCDS in 4.5% of these null transcripts. We used this estimate of the Type 1 error rate to compute the false discovery rate for novel mCDS and uaCDS identified by riboHMM. To address the second question, we started with the set of annotated coding transcripts for which riboHMM was able to recover the precise CDS (i.e., the mCDS matched the annotated CDS exactly). We generated a set of "simulated transcripts" using the following strategy: (1) randomly select a new TIS downstream and in-frame to the annotated TIS, ensuring that the codon underlying the new TIS belonged to the set Ω_{start}, (2) permute the footprint counts among bases upstream of the new TIS. Among the simulated transcripts in which riboHMM could identify an mCDS, the inferred TIS matched the new TIS exactly in 62% of transcripts; this corresponds to a false discovery proportion of 38%.

Translated mCDS in pseudogenes

Starting with 14,065 pseudogenes that have been identified and categorized in humans (Pei et al. 2012), 9,375 pseudogenes were identified by StringTie to be expressed in LCLs. Using a very stringent posterior cutoff of 99.99%, we inferred mCDS in 448 of these expressed pseudogenes. Using pairwise alignment of the pseudogene and parent gene transcript, we observed that although the pseudogene mCDS typically code for shorter protein sequences compared with the parent protein, a large fraction of the pseudogene mCDS share coding-frame with their parent gene (see Figure S6A).

Validation with Harringtonine-treated data

Harringtonine-treated ribosome footprints were measured in LCLs with a total sequencing depth of 21 million reads. In Figure 4, we illustrate the aggregate proportion of treated ribosome footprints centered at the inferred start codon for all novel mCDS, and compare it with the aggregate proportion of treated footprints around the start codon of an equal number of annotated CDSs that have a posterior probability greater than 0.8 under our model. In Figure S6B, we illustrate the aggregate proportion of treated footprints for mCDS inferred in pseudogenes alone, and in Figure 6B, we compare the aggregate treated footprint proportions at the start codons of inferred uaCDS and their corresponding mCDS.

Identifying translated alternate ORFs

For each transcript that had a mCDS with posterior greater than 0.8 and more than 50 base pairs of RNA sequence in the 5'UTS state, we defined an "upstream-restricted transcript" consisting of the exons within the 5'UTS state. Using a random set of 5000 non-overlapping upstream-restricted transcripts in which more than 80% of base positions were mappable, we computed the maximum likelihood estimates of the transition parameters and occupancy parameters to identify additional translated sequences within these upstream-restricted transcripts. Assuming that the fine-scale structure of footprint counts within these translated sequences would be similar to that within the mCDS, we kept the periodicity parameters fixed to their previously estimated values. With these parameter estimates, we inferred the MAP frame and state sequences with posterior greater than 0.8 and filtered out inferences where less than 50% of the inferred CDS was mappable. These additional translated sequences within

the upstream-restricted transcripts were called upstream alternate coding sequences or

uaCDS.

Identifying stable peptides with mass spectrometry data

To identify stable proteins translated from the novel CDSs (mCDS and uaCDS), we

analyzed quantitative, high-resolution mass spectrometry data derived from 60 LCLs,

with MaxQuant v1.5.0.30 (Cox and Mann 2008) and the Andromeda (Cox et al. 2011)

search engine. Sample labeling, processing and data collection details can be found

elsewhere (Battle et al. 2015; Khan et al. 2013). Peptides were identified using a

database that contained 63,904 GENCODE annotated protein sequences and 7,271

novel CDSs identified by our method. For all searches, up to two missed tryptic

cleavages were allowed, carbamidomethylation of cysteine was entered as a fixed

modification, and N-terminal acetylation and oxidation of methionine were included as

variable modifications for all searches. A 'first search' tolerance of 40 ppm with a score

threshold of 70 was used for time-dependent mass recalibration followed by a main

search MS1 tolerance of 6 ppm and an MS2 tolerance of 20 ppm. The 're-quantify'

option was used to aid the assignment of isotopic patterns to labeling pairs. The 'match

between runs' option was enabled to match identifications across samples using a

matching time window of 42 seconds and an alignment time window of 20 min. Peptide

and protein false discovery rates were set to 10% using a reverted version of the search

database. Protein group quantifications were taken as the median

$\log_2$(sample/standard) ratio for all groups containing at least two independent unique or

'razor' peptide quantifications (including multiple measurements of the same peptide in

different fractions) without a modified peptide counterpart.

Bias correction to compute expected number of peptide hits

Proteins with at least one peptide identified by this high-resolution mass-spectrometry

protocol tend to be distinct from proteins with no mass-spectrum matches.

1. The median footprint density of annotated coding genes with at least one peptide

match is about 125 fold higher than that of coding genes with no peptide match (see

Figure S12A).

2. The median length of coding genes with at least one peptide match is 20% higher

than that of coding genes without a peptide match (see Figure S12B).

3. The trypsin cleavage step of the protocol ensures that nearly all observable peptides have a C-terminal lysine or arginine residue, and up to two additional lysine or arginine residues within the peptide sequence (called "tryptic peptides"). This step imposes a strict constraint on the set of unique peptide sequences that can be observed from a protein sequence, and genes with fewer tryptic peptides are less likely to have a mass-spectrum match.

4. All tryptic peptides in an expressed protein are not equally likely to be observed. The probability of detecting a tryptic peptide depends on its electrostatic properties relative to other tryptic peptides from all expressed proteins, which in turn depends on the amino acid composition of the tryptic peptides (see Figure S12 C-F).

To account for these biases, we developed a predictive model to estimate the probability that a protein has at least one peptide hit in a mass-spectrometry experiment. The predicted label for a protein is whether the protein has at least one mass-spectrum match ($H_n = 1$) or no mass-spectrum match ($H_n = 0$). The predictive features of a protein used in the model are (1) the ribosome footprint density of the corresponding transcript (D_n), (2) the protein length (S_n), and (3) the counts of amino acids within each of the K tryptic peptides that can be generated from the protein ($L_n = \{L_{n1}, \cdots, L_{nK}\}$). Since the relevant feature of an amino acid is its charge, we partitioned the set of amino acids into four groups – positively charged (R, H, K), negatively charged (D, E), polar uncharged (S, T, N Q), and others. The amino acid count vector L_{nk} was then collapsed into a vector of the counts of each of these four groups. Conditional on $H_n = 1$, we introduced a latent variable for each tryptic peptide that indicates whether the peptide was matched to a mass-spectrum or not ($Z_{nk} \in \{1,0\}$); this latent variable accounts for differences between matched and unmatched peptides.

Assuming that the three predictive features are independent conditional on the predicted label H_n, the odds of observing at least one peptide hit is then given as

$$
\frac{p(H_n = 1 | D_n, S_n, L_n)}{p(H_n = 0 | D_n, S_n, L_n)}
$$
$$
= \frac{p(D_n | H_n = 1)\, p(S_n | H_n = 1)\, \{\sum_{Z_n} p(L_n | Z_n, H_n = 1)\, p(Z_n | H_n = 1)\}\, p(H_n = 1)}{p(D_n | H_n = 0)\, p(S_n | H_n = 0)\, p(L_n | Z_n = 0, H_n = 0)\, p(H_n = 0)}
$$

We learn the predictive model using annotated coding genes and partitioning them into those that have at least one peptide hit ("hit genes") and those that do not have a peptide hit ("no-hit genes"). We computed $p(D_n|H_n)$ using an empirical distribution of footprint density within coding genes, $p(S_n|H_n)$ using an empirical distribution of the lengths of coding genes, $p(L_{nk}|Z_{nk} = 1, H_n = 1)$ using tryptic peptides within hit genes matched to mass-spectra, $p(L_{nk}|Z_{nk} = 0, H_n = 1)$ using unmatched tryptic peptides within hit genes, and $p(L_{nk}|Z_{nk} = 0, H_n = 0)$ using tryptic peptides within no-hit genes. Finally, we set $p(H_n = 1) = p(H_n = 0) = 1/2$ and $p(Z_n|H_n = 1) = 1/(2^K - 1)$. Using peptide hits in annotated proteins, we evaluated the accuracy of this model by holding out some annotated proteins as test data, learning the predictive distributions using the remaining training data and computing the expected number of test proteins that had a mass-spectrum match. We estimated the expected number of held-out annotated proteins with at least one mass-spectrum match to be 1,206 (s.d.=34), while the actual number of held-out proteins with a match was 1,387 (s.d.=36).

Test for long-term purifying selection

In order to quantify whether the novel mCDS are evolutionary conserved in terms of their amino acid sequence, we first extracted DNA sequences orthologous to the mCDS from a 100-way vertebrate whole-genome alignments (UCSC), restricting to genomes aligned with either Syntenic net or Reciprocal best net. We next performed a 3-frame translation on each orthologous sequence and a multiple alignment to obtain the correct codon alignments. More specifically, for each orthologous sequence, we kept the frame with the highest amino acid identity compared to the human peptide, requiring at least 60% identity for alignable positions and no more than 50% of the alignment as gaps. Finally, we used codeML/PAML (Yang 2007) to estimate *dN* and *dS* rates across the trees consisting of all remaining peptides, first using a model allowing variable omega and then a model with omega fixed to one. To determine whether a specific peptide is under purifying selection or not, we compared the two models using a likelihood ratio test and reported peptides that satisfied a Bonferroni-corrected *p*-value threshold.

Correlation between uaCDS and mCDS

We computed the correlation across LCLs between the proportion of footprints mapped to a transcript that fall within its uaCDS and the proportion that fall within its mCDS. We evaluated the statistical significance of these correlations using an empirical null

1   distribution of Spearman correlations computed using random pairs of mCDS and

2   uaCDS. A random pair of mCDS and uaCDS was obtained by randomly shifting the

3   coordinates of an observed pair of mCDS and uaCDS, matching for their respective

4   lengths and the distance between them.

6   Data Release

7   All novel coding sequences identified in this work, along with the harringtonine-treated

8   ribosome profiling data are deposited in GEO Accession GSE75290.

10   Acknowledgements

We thank Adam Frankish from the GENCODE group for discussions on the sources of

information used for annotating coding genes, Zia Khan for discussions on the mass-

spectrometry data analysis, and Audrey Fu for discussions on evaluating the correlation

between mCDS and uaCDS. We also thank members of the Pritchard, Gilad and

Stephens labs for comments and suggestions. This work was funded by grants from the

NIH (HG007036 to J.K.P., MH084703 to Y.G. and J.K.P., and HG02585 to M.S.), and by

the Howard Hughes Medical Institute. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

References

1000 Genomes Project Consortium, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, et al. 2015. A global reference for human genetic variation. *Nature* **526**: 68–74.

Bairoch A, Apweiler R. 2000. The SWISS-PROT protein sequence database and its supplement TrEMBL in 2000. *Nucleic Acids Res* **28**: 45–48.

Barbosa C, Peixeiro I, Romão L. 2013. Gene Expression Regulation by Upstream Open Reading Frames and Human Disease. *PLoS Genet* **9**: e1003529.

Battle A, Khan Z, Wang SH, Mitrano A, Ford MJ, Pritchard JK, Gilad Y. 2015. Genomic variation. Impact of regulatory variation from RNA to protein. *Science* **347**: 664–667.

Calvo SE, Pagliarini DJ, Mootha VK. 2009. Upstream open reading frames cause widespread reduction of protein expression and are polymorphic among humans. *Proc Natl Acad Sci USA* **106**: 7507–7512.

Camby I, Le Mercier M, Lefranc F, Kiss R. 2006. Galectin-1: a small protein with major functions. *Glycobiology* **16**: 137R–157R.

Clark MB, Amaral PP, Schlesinger FJ, Dinger ME, Taft RJ, Rinn JL, Ponting CP, Stadler PF, Morris KV, Morillon A, et al. 2011. The reality of pervasive transcription. *PLoS Biol* **9**: e1000625; discussion e1001102.

Cox J, Mann M. 2008. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotech* **26**: 1367–1372.

Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen JV, Mann M. 2011. Andromeda: a peptide search engine integrated into the MaxQuant environment. *J Proteome Res* **10**: 1794–1805.

Djebali S, Davis CA, Merkel A, Dobin A, Lassmann T, Mortazavi A, Tanzer A, Lagarde J, Lin W, Schlesinger F, et al. 2012. Landscape of transcription in human cells. *Nature* **489**: 101–108.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

ENCODE Project Consortium, Birney E, Stamatoyannopoulos JA, Dutta A, Guigó R, Gingeras TR, Margulies EH, Weng Z, Snyder M, Dermitzakis ET, et al. 2007. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* **447**: 799–816.

Evans SN, Shvets Y, Slatkin M. 2007. Non-equilibrium theory of the allele frequency spectrum. *Theoretical Population Biology* **71**: 109–119.

Furuno M, Kasukawa T, Saito R, Adachi J, Suzuki H, Baldarelli R, Hayashizaki Y,
Okazaki Y. 2003. CDS Annotation in Full-Length cDNA Sequence. *Genome Res*
13: 1478–1487.

Galindo MI, Pueyo JI, Fouix S, Bishop SA, Couso JP. 2007. Peptides encoded by short
ORFs control development and define a new eukaryotic gene family. *PLoS Biol*
5: e106.

Guttman M, Russell P, Ingolia NT, Weissman JS, Lander ES. 2013. Ribosome profiling
provides evidence that large noncoding RNAs do not encode proteins. *Cell* **154**:
240–251.

Hangauer MJ, Vaughn IW, McManus MT. 2013. Pervasive transcription of the human
genome produces thousands of previously unidentified long intergenic noncoding
RNAs. *PLoS Genet* **9**: e1003569.

Hernández-Sánchez C, Mansilla A, de la Rosa EJ, Pollerberg GE, Martínez-Salas E, de
Pablo F. 2003. Upstream AUGs in embryonic proinsulin mRNA control its low
translation level. *EMBO J* **22**: 5582–5592.

Ingolia NT, Brar GA, Stern-Ginossar N, Harris MS, Talhouarne GJS, Jackson SE, Wills
MR, Weissman JS. 2014. Ribosome profiling reveals pervasive translation
outside of annotated protein-coding genes. *Cell Rep* **8**: 1365–1379.

Ingolia NT, Ghaemmaghami S, Newman JRS, Weissman JS. 2009. Genome-wide
analysis in vivo of translation with nucleotide resolution using ribosome profiling.
Science **324**: 218–223.

Ingolia NT, Lareau LF, Weissman JS. 2011. Ribosome profiling of mouse embryonic
stem cells reveals the complexity and dynamics of mammalian proteomes. *Cell*
147: 789–802.

Jung HW, Tschaplinski TJ, Wang L, Glazebrook J, Greenberg JT. 2009. Priming in
systemic plant immunity. *Science* **324**: 89–91.

Kapranov P, Cheng J, Dike S, Nix DA, Duttagupta R, Willingham AT, Stadler PF, Hertel
J, Hackermüller J, Hofacker IL, et al. 2007. RNA maps reveal new RNA classes
and a possible function for pervasive transcription. *Science* **316**: 1484–1488.

Kawase T, Akatsuka Y, Torikai H, Morishima S, Oka A, Tsujimura A, Miyazaki M,
Tsujimura K, Miyamura K, Ogawa S, et al. 2007. Alternative splicing due to an
intronic SNP in HMSD generates a novel minor histocompatibility antigen. *Blood*
110: 1055–1063.

Khan Z, Ford MJ, Cusanovich DA, Mitrano A, Pritchard JK, Gilad Y. 2013. Primate
transcript and protein expression levels evolve under compensatory selection
pressures. *Science* **342**: 1100–1104.

Kochetov AV. 2008. Alternative translation start sites and hidden coding potential of
eukaryotic mRNAs. *Bioessays* **30**: 683–691.

Kondo T, Hashimoto Y, Kato K, Inagaki S, Hayashi S, Kageyama Y. 2007. Small peptide regulators of actin-based cell morphogenesis encoded by a polycistronic mRNA. *Nat Cell Biol* **9**: 660–665.

Kondo T, Plaza S, Zanet J, Benrabah E, Valenti P, Hashimoto Y, Kobayashi S, Payre F, Kageyama Y. 2010. Small peptides switch the transcriptional activity of Shavenbaby during Drosophila embryogenesis. *Science* **329**: 336–339.

Kozak M. 1987. An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs. *Nucleic Acids Res* **15**: 8125–8148.

Lammich S, Schöbel S, Zimmer A-K, Lichtenthaler SF, Haass C. 2004. Expression of the Alzheimer protease BACE1 is suppressed via its 5'-untranslated region. *EMBO Rep* **5**: 620–625.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Meth* **9**: 357–359.

Lappalainen T, Sammeth M, Friedländer MR, Hoen PAC 't, Monlong J, Rivas MA, Gonzàlez-Porta M, Kurbatova N, Griebel T, Ferreira PG, et al. 2013. Transcriptome and genome sequencing uncovers functional variation in humans. *Nature* **501**: 506–511.

Lauressergues D, Couzigou J-M, Clemente HS, Martinez Y, Dunand C, Bécard G, Combier J-P. 2015. Primary transcripts of microRNAs encode regulatory peptides. *Nature* **520**: 90–93.

Lee J, Park EH, Couture G, Harvey I, Garneau P, Pelletier J. 2002. An upstream open reading frame impedes translation of the huntingtin gene. *Nucl Acids Res* **30**: 5110–5119.

Lee S, Liu B, Lee S, Huang S-X, Shen B, Qian S-B. 2012. Global mapping of translation initiation sites in mammalian cells at single-nucleotide resolution. *Proc Natl Acad Sci USA* **109**: E2424–2432.

Lin MF, Jungreis I, Kellis M. 2011. PhyloCSF: a comparative genomics method to distinguish protein coding and non-coding regions. *Bioinformatics* **27**: i275–282.

Ma B. 2015. Novor: Real-Time Peptide de Novo Sequencing Software. *J Am Soc Mass Spectrom* 1–10.

Michel AM, Choudhury KR, Firth AE, Ingolia NT, Atkins JF, Baranov PV. 2012. Observation of dually decoded regions of the human genome using ribosome profiling data. *Genome Res* **22**: 2219–2229.

Morris DR, Geballe AP. 2000. Upstream open reading frames as regulators of mRNA translation. *Mol Cell Biol* **20**: 8635–8642.

Nesvizhskii AI. 2014. Proteogenomics: concepts, applications and computational strategies. *Nat Methods* **11**: 1114–1125.

Nielsen R. 2005. Molecular signatures of natural selection. *Annu Rev Genet* **39**: 197–
218.

Oelkers K, Goffard N, Weiller GF, Gresshoff PM, Mathesius U, Frickey T. 2008.
Bioinformatic analysis of the CLE signaling peptide family. *BMC Plant Biol* **8**: 1.

Pei B, Sisu C, Frankish A, Howald C, Habegger L, Mu XJ, Harte R, Balasubramanian S,
Tanzer A, Diekhans M, et al. 2012. The GENCODE pseudogene resource.
Genome Biol **13**: R51.

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015.
StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. *Nat Biotechnol* **33**: 290–295.

Pevtsov S, Fedulova I, Mirzaei H, Buck C, Zhang X. 2006. Performance Evaluation of
Existing De Novo Sequencing Algorithms. *J Proteome Res* **5**: 3018–3028.

Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: discovering splice junctions with
RNA-Seq. *Bioinformatics* **25**: 1105–1111.

van Bakel H, Nislow C, Blencowe BJ, Hughes TR. 2010. Most "dark matter" transcripts
are associated with known genes. *PLoS Biol* **8**: e1000371.

Vanderperre B, Lucier J-F, Bissonnette C, Motard J, Tremblay G, Vanderperre S,
Wisztorski M, Salzet M, Boisvert F-M, Roucou X. 2013. Direct detection of
alternative open reading frames translation products in human significantly
expands the proteome. *PLoS ONE* **8**: e70698.

Weinberg DE, Shah P, Eichhorn SW, Hussmann JA, Plotkin JB, Bartel DP. 2015.
Improved ribosome-footprint and mRNA measurements provide insights into
dynamics and regulation of yeast translation. *bioRxiv* 021501.

Xu H, Wang P, Fu Y, Zheng Y, Tang Q, Si L, You J, Zhang Z, Zhu Y, Zhou L, et al.
2010. Length of the ORF, position of the first AUG and the Kozak motif are
important factors in potential dual-coding transcripts. *Cell Res* **20**: 445–457.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**:
1586–1591.

1 Supplementary Figures

Figure S1: Robustness of periodicity parameter estimates. Bar plots of the periodicity parameters (π_z),

for 29 bp long footprints, estimated using the top 5000 expressed genes. Each red circle indicates the

parameter values estimated by using a random set of 1000 genes.

Figure S2: Robustness of occupancy parameter estimates. Bar plots of the maximum likelihood

estimates of the occupancy parameters in each state, α_z (scaled by α_{TES} for mCDS). For mCDS (dark gray),

the parameters were estimated using the top 5000 expressed genes, and the red circles indicate parameter

values estimated using a random set of 1000 genes. For uaCDS (light gray), the parameters were estimated

using a random set of 5000 upstream-restricted transcripts, and each red circle indicates values estimated

using random sets of 1000 upstream-restricted transcripts.

Figure S3: Decision rules to identify matches and mismatches of mCDS to annotation. Illustrating the

decisions by which mCDS inferred on each transcript are identified as an exact match, a frame match) or a

mismatch to annotated coding sequences. Matches and mismatches are only defined when the transcript is

annotated by GENCODE as coding and the classification depends on agreement with the annotated CDS. A

gene is considered to have an exact (or frame) match if at least one of its isoforms is labeled an exact (or

frame) match. In all other cases, the inference for the coding gene is considered a mismatch.

Figure S4: Model accuracy. Quantifying model accuracy in terms of the fraction of previously annotated

CDSs accurately recovered as a function of total footprint sequencing depth. We performed the entire

analysis on the same set of assembled transcripts (parameter estimation and inference) after subsampling

the data. Starting with inferences using the complete footprint data set that exactly match annotated CDSs

(top subpanel), we show the fraction of these CDSs that were accurately (blue) and inaccurately (brown)

recovered with a high posterior for varying sequencing depths. Starting with inferences that only match the

frame of annotated CDSs (bottom subpanel), we show the fraction of accurately and inaccurately recovered

CDSs for varying sequencing depths.

2 **Figure S5: Decision rules to identify novel mCDS.** Illustrating the decisions that identify novel mCDS.

3 The mCDS that do not overlap any known CDS (pooling GENCODE, UCSC and CCDS annotations) are

4 labeled as novel mCDS -- these include mCDS from both novel and annotated transcripts.

2 **Figure S6: Translated coding sequences identified in hundreds of pseudogenes.** (A) Comparing the

3 inferred CDS in the pseudogene GAPDHP72 (orange) with the annotated CDS (purple) of its parent gene

4 GAPDH (introns have been removed for better visualization). Gray shaded boxes show the alignment

5 between the pseudogene and parent gene transcripts, and dark shaded boxes indicate that the inferred

6 coding frame of the pseudogene matches that of the parent gene. Although the pseudogene and parent

7 gene share coding frames, the underlying sequences are sufficiently different. (B) Enrichment of

8 harringtonine-arrested ribosome occupancy at the inferred translation initiation sites validates our inferred

9 mCDS in pseudogenes. (C) Nonsynonymous variants (orange) segregate at significantly lower frequencies

10 than synonymous variants (blue) in those pseudogenes predicted to have a translated CDS. Pseudogenes

11 with ribosome occupancy, but predicted to have no translated CDS, do not show any significant difference

12 between the site frequency spectra of synonymous and nonsynonymous variants.

1
2 **Figure S7: Comparing the amino acid content between annotated and novel CDS.** The overall amino
3 acid content is largely similar between annotated CDSs and novel CDSs (mCDS, pseudogenes and
4 uaCDS), with a substantially higher proportion of lysine and isoleucine residues in CDSs within
5 pseudogenes, and a higher proportion of alanine, arginine, glycine, and proline residues within mCDS and
6 uaCDS.

Figure S8: Characteristics of peptides matched to novel CDS. (A) Histogram of number of independent identification of unique peptides across novel mCDS that have at least one peptide match. (B) Scatter plot of the untreated ribosome footprint density and the protein abundance (median abundance across all unique peptides) across all novel mCDS that have at least one peptide match (Spearman R = 0.24; *p*-value = $2{\times}10^{-2}$).

Figure S9: Characteristics of novel uaCDS. (A) Histogram of the distance between uaCDS and their corresponding mCDS. (B) Histogram of the lengths of uaCDS (median length of 16 amino acids).

Figure S10: Comparing footprint abundance and gene expression. Scatter plot of the RNA-seq density

(reads per base per million sequenced reads) within annotated coding transcripts *vs.* the expected ribosome

footprint count (normalized by sequencing depth) in triplets within the CDS of the transcripts. The top 50[th]

percentile of expressed genes is shown in black and the bottom 50[th] percentile is shown in gray. For highly

expressed genes, it is reasonable to assume that the expected footprint count in a triplet scales linearly with

RNA-seq density. However, for lowly expressed genes and outlier genes (i.e., genes with high expected

footprint count and low RNA-seq density, and genes with high RNA-seq density and low expected footprint

count), this assumption may not be valid.

2 **Figure S11: Comparing the periodicity in ribosome footprint counts for footprints of different**
3 **lengths.** Proportion of footprint counts aggregated across 1000 highly expressed annotated coding
4 transcripts, centered at their translation initiation sites. Each subplot illustrates the proportions for footprints
5 of a given length, with the periodicity in the proportions showing distinct features between footprints of
6 different lengths.
7

Figure S12: Annotated genes with peptide hits tend to be longer, have higher expression and a distinct amino acid composition. (A) Cumulative distribution of footprint density for genes with at least one unique peptide hit (blue) and genes with no unique peptide hit (red). The median footprint density of genes with a peptide hit is about 125 fold higher than the median footprint density of genes without a peptide hit. (B) Cumulative distribution of protein length for genes with at least one unique peptide hit (blue) and genes with no unique peptide hit (red). Genes with a peptide hit tend to code for proteins that are 20% longer than proteins encoded by genes without a peptide hit. (C-F) Comparing amino acid composition within tryptic peptides with a mass-spectrum match and tryptic peptides without a mass-spectrum match. Amino acids, grouped by their electrostatic properties, have distinct compositions between matched and unmatched peptides. Matched peptides tend to be significantly shorter than unmatched peptides, and have a distinct composition of charged amino acids.