Potential impact of sexual transmission of Ebola virus on the epidemic in West

Africa

Authors

Jessica L. Abbate[1,2,3] [*]

Carmen Lia Murall[4] [*]

Heinz Richner[1]

Christian L. Althaus[5]

[*]These authors contributed equally to this article.

Affiliations

1. Institute for Ecology and Evolution, University of Bern, Bern, Switzerland

2. UMR MIVEGEC (UMR CNRS 5290, IRD 224, UM), Institute for Research of

Development (IRD), Montpellier, France

3. UMR UMMISCO (UMI 209 IRD-UPMC), Bondy, France

4. Max-Planck Institute for Dynamics and Self-Organization, Gottingen, Germany

5. Institute of Social and Preventive Medicine (ISPM), University of Bern, Bern,

Switzerland

Abstract

Ebola virus RNA can persist in seminal fluids of male convalescent patients after they recover from Ebola virus disease (EVD). We used a compartmental EVD transmission model, Monte Carlo simulations, and performed sensitivity analyses to assess the potential impact of sexual transmission on the epidemic dynamics. The rate of sexual transmission and the period during which convalescent men can transmit sexually both affect the number of excess EVD cases, while the latter also influences the duration of the epidemic. Assuming an average convalescent period of 3 months, and a per sex act transmission probability of 0.1%, we found that sexual transmission could extend the EVD epidemic in Sierra Leone by 123 days (95% CI: 110–137 days). These results show the importance of ongoing surveillance efforts in West Africa and call for a better understanding of the persistence and infectivity of Ebola virus RNA in convalescent patients.

Introduction

Recent reports suggesting the potential for sexual transmission of Ebola virus from convalescent survivors have raised a number of important questions regarding its impact on the final phase of the epidemic in West Africa [1,2]. Even once the worst hit countries of Guinea, Liberia, and Sierra Leone are declared free of Ebola virus disease (EVD), rare cases may still arise from the large number of remaining survivors. Perhaps the most crucial element for bringing the epidemic to an end is maintaining vigilance in the community and preventing new chains of transmission. Historically occurring in sparsely-populated isolated communities, often killing the vast majority of victims [3], the unprecedented number of cases – and survivors – in densely populated urban areas presents novel challenges for bringing the outbreak to an end. Although sexual transmission of EVD is likely to be rare, it is important to investigate its potential impact on the transmission dynamics in general, and on the tail of the ongoing epidemic in particular.

Follow-up studies on survivors of the 1995 outbreak in the Democratic Republic of Congo [4] and the 2000 [5] and 2007 [6] outbreaks in Uganda have raised awareness of what is now being termed "post-Ebola syndrome" (post-Ebola sequelae) – debilitating illnesses from myalgia to uveitis – which can persist for at least 21 months after the onset of symptoms. Though the virus is no longer detected in the blood after symptoms disappear, active (replicating) virus has been documented in ocular fluid, rectal fluids, vaginal fluids, and semen [4,7]. Two recent reports indicate that this is also the case for survivors of the current (2013-2015) epidemic in West Africa, having documented

ongoing viremia in ocular [8] and seminal [1] fluids many months after falling ill.

Importantly, molecular evidence from a woman who contracted EVD in Liberia after

unprotected vaginal intercourse with a survivor nearly 6 months after his recovery from

EVD supports the hypothesis that direct sexual transmission from asymptomatic

convalescent survivors can and does occur [2]. Transmission to sexual partners was never

confirmed in earlier outbreaks, but was suspected to have occurred in at least one instance

[4]. Similarly, cases of sexual transmission of other hemorrhagic fever infections, notably

by the closely related Marburg virus, have been suspected in the past [9,10].

Of particular epidemiological interest is the question of how long active Ebola virus in

semen, vaginal, and rectal fluids of convalescent survivors poses a threat for sexual

transmission. The previous studies identified active virus as late as 82 days after onset of

symptoms in seminal fluid from one male patient; samples positive for viral RNA were

identified as late as 29 days (rectal) and 33 days (vaginal) after onset of symptoms in one

female patient of six sampled, and 101 days after onset of symptoms in seminal fluid

from four out of five male patients [4,7]. The studies from the West African outbreak,

showing viremia in semen 4-6 months after onset of symptoms in 65% of men tested (7-9

months in 26%) [1] and confirming occurrence of transmission from a survivor 179 days

after onset of symptoms [2], suggest that sexual transmission from convalescent men may

pose a threat for far longer than previously anticipated.

Sexual transmission of Ebola virus from convalescent survivors is likely a rare event, but

researchers have warned that it is feasible and should be considered in epidemiological

models that are used to predict the trajectory of the outbreak [11]. To this end, we devised

a novel mathematical model of EVD transmission and studied the epidemic dynamics in

absence and presence of sexual transmission from convalescent men. We performed

sensitivity analysis to understand the impact of key unknown parameters, such as the

duration of the convalescent period and the transmission probability per sexual contact.

We also performed Monte Carlo simulations to explore the impact of sexual transmission

on the epidemic tail in Sierra Leone.

Methods

Transmission model. We extended a SEIR (susceptible-exposed-infected-recovered)

modeling framework, which has been extensively used to describe EVD transmission

[12–14], by adding a component for sexual transmission from convalescent survivors

who maintain active Ebola virus replication (Fig. 1). The resulting SEICR model has five

states: susceptible, *S*, exposed, *E*, symptomatic and infectious, *I*, convalescent, *C*, fully

recovered and immune, *R*, and dead, *D*. The model is represented by the following set of

ordinary differential equations (ODEs):

98
$$\frac{dS}{dt} = -\beta(t)SI - \beta_S pC\frac{S}{N}$$
$$\frac{dE}{dt} = \beta(t)SI + \beta_S pC\frac{S}{N} - \sigma E$$
$$\frac{dI}{dt} = \sigma E - \gamma I$$
$$\frac{dC}{dt} = (1-f)\gamma I - \alpha C \tag{1}$$
$$\frac{dR}{dt} = \alpha C$$
$$\frac{dD}{dt} = f\gamma I$$

where $N = S + E + I + C + R$ denotes the total population size. We assumed the non-

sexual transmission rate, $\beta(t)$, to be initially constant (β_0) before it starts to decay

exponentially due to the effect of control interventions and behavior change after time τ :

$\beta(t) = \beta_1 + (\beta_0 - \beta_1)e^{-k(t-\tau)}$ [12]. The sexual transmission parameter, β_s, can be described as

the product of two parameters ($\beta_s = \eta q$) that we will consider separately: η is the per sex

act transmission probability of Ebola virus from convalescent men, and q is the daily rate

at which they engage in sexual intercourse. The number of convalescent individuals are

scaled by p, which is the proportion of convalescent survivors who are sexually active

men. $1/\sigma$ and $1/\gamma$ represent the average durations of incubation and symptomatic

infection, respectively. f is the case fatality rate. The average duration after which

convalescent patients recover completely and shed no further replicating Ebola virus from

their body is given by $1/\alpha$. We assumed that sexual transmission is frequency-dependent

[15–17], i.e., the probability that the sexual partner of a convalescent man is susceptible

is given by S/N .

Model parameters. The basic reproductive number, R_0, for the SEICR model can be

calculated using the next-generation matrix method [18,19] and is given by

$$R_0 = \frac{\beta S_0}{\gamma} + \frac{(1-f)\,p\,\beta_s}{\alpha},$$

where S_0 is the initial number of susceptible individuals (see *Supplementary Material*).

When α goes to infinity or either $\beta_s = 0$ or $p = 0$, the equation reduces to the basic

reproductive number in absence of sexual transmission: $R_{0,N} = \frac{\beta S_0}{\gamma}$. Thus, the second

term represents the contribution of sexual transmission from convalescent patients to the

overall R_0: $R_{0,C} = \frac{(1-f)\,p\,\beta_s}{\alpha}$.

The model parameters were either based on published values from the literature or

adjusted to provide a good description of the EVD epidemic in Sierra Leone (Table 1).

We assumed that published estimates of R_0 do not include the contribution of sexual

transmission, i.e., the transmission rate during the early phase of the outbreak was given

by $\beta_0 = R_{0,N}\,\gamma/S_0$. We further assumed that the non-sexual reproductive number drops

below unity in the presence of partially effective control interventions [20] and set $\beta_1 =$

0.5 γ/S_0. We used the weekly incidence and cumulative case counts of EVD from Sierra

Leone reported by the World Health Organization (WHO) Ebola Outbreak Situation

Reports (including updates according to the patient database) [21] as a comparison to the

model results.

Deterministic model and sensitivity analysis. We solved the system of ODEs

numerically using the function *ode* from the *deSolve* package in the R software

environment for statistical computing [22]. We compared the following response

variables of the model: the epidemic peak number of exposed, *E*, acute, *I*, and

convalescents, *C*, cases; the cumulative number of EVD cases, deaths, and recoveries; the

date at the epidemic peak; the daily and cumulative incidence of sexual transmission; and

the date at which the last symptomatic case either died or entered into convalescence

("day of last case"). We defined the day of last case as the time when the number of

symptomatic and infectious individuals, *I*, dropped below 0.5. We considered the

following parameters for the sensitivity analysis: the per sex act transmission probability

of Ebola virus from convalescent men (η), the proportion of convalescent survivors who

are sexually active men (p), the rate at which they engage in sexual intercourse (q), and

the rate at which convalescent patients recover completely and shed no further replicating

Ebola virus from their body (α). The sensitivity of the response variables to changes in R_0

was explored simultaneously as a comparison. We generated 1000 parameter

combinations from the uniform ranges, log-transformed [0.5x – 2x] for the parameter

values for η, p, q, and α, given in Table 1 via Latin hypercube sampling using the

Huntington and Lyrintzis correlation correction method (function *lhs* from R package

pse) [23]. We then calculated partial rank correlation coefficients (PRCCs) using 50

bootstrap replicates [24].

Monte Carlo simulations. We performed stochastic simulations of the SEICR model

with and without sexual transmission using Gillespie's algorithm [25]. We specifically

investigated the following response variables from the simulations: the cumulative

number of EVD cases, the size and date of the epidemic's peak incidence (daily number

of new symptomatic infections), and the date of last case (last day that symptomatic

infections, *I*, fell below 1). Summary statistics were based on the results of 1000

simulation runs for each transmission scenario. We calculated the average of the peak and

total cumulative number of EVD cases by including all simulations runs, i.e., also the

simulations that rapidly go extinct. In contrast, the average dates of the epidemic peak

and last case were based on the simulated epidemic trajectories over which 100 or more

cases were accumulated.

Results

Contribution of sexual transmission to overall R_0. Using the baseline parameter values

from Table 1, the reproductive number of a convalescent infection, $R_{0,C}$, is 0.0024. This

corresponds to only 0.12% of the overall R_0 of 2.0224. Increasing the convalescent period

from 3 to 6 months, the contribution of $R_{0,C}$ (0.0051) to the overall R_0 rises to 0.25%. The

equation for $R_{0,C}$ (see *Methods*) illustrates that doubling the per sex act transmission

probability has the same impact as doubling the convalescent period. It is important to

note that the relative contribution of sexual transmission to the overall reproductive

number rises as the effective reproductive number drops during the epidemic due to the

effects of control interventions (see *Supplementary Material,* Fig. *S1*).

Effect of sexual transmission parameters on epidemic dynamics. Simulating the

deterministic transmission model using the baseline parameter values from Table 1

resulted in a good description of the EVD epidemic in Sierra Leone (see *Supplementary*

Material, Fig. *S2*). The two key unknown parameters of sexual transmission are the per

sex act transmission probability, η, and the rate at which convalescent survivors fully

recover, α. The duration of the convalescent period has the biggest impact on the peak

number of convalescent individuals, while η does not (compare Fig. 2a and Fig. 3a). Both

parameters have extremely small effects on the peak number of infected or exposed

patients (Fig. 2a and Fig. 3a). The total number of recovered individuals is reached more

slowly the longer the convalescent period (Fig. 2b), which is not an effect caused by η

(Fig. 3b). While the convalescent periods ($1/\alpha = [0 - 9$ months]) and the low values of η

(η = [0.0005 – 0.002]) we explored create very few extra cases (Fig. 2b and Fig. 3b),

sensitivity analyses revealed that a higher per sex act transmission probability, a higher

proportion of sexually active convalescent individuals, p, or a higher frequency of sex

acts, q, could have larger impacts on the total number of cases than would proportional

increases in the convalescent period (see *Supplementary Material*, Fig. *S3a*). Sensitivity

analyses also revealed that these sexual transmission parameters could produce a small

delay on the epidemic peak, more so than by changes in the convalescent period (see

Supplementary Material, Fig. *S3b*).

The number of sexual transmission events expected from the baseline scenario (η = 0.001

and 1/ α = 3 months) is 31.5, the majority of which will occur around the peak of the

epidemic (Fig. 2c and Fig. 3c) and thus likely go undetected. Doubling either η or 1/ α

results in almost equal increases in the incidence and cumulative number of sexual

transmission events (Fig. 2c, 2d and Fig. 3c, 3d), with either leading to roughly double

the number of sexually transmitted cases over the course of the whole epidemic (> 60

cases). It should be noted that the total number of cases increases more than by simply the

number of sexual transmission events, because each sexual transmission event results in a

new potential cluster of non-sexual transmission. The day of last case is affected more by

the convalescent period than the per sex act transmission probability (represented by

vertical lines in Fig. 2a and Fig. 3a), a result confirmed by the sensitivity analysis (see

Supplementary Material, Fig. *S3a*). The tail of the epidemic will depend on a small

number of events that are likely to be affected by stochastic processes, thus we used

Monte Carlo simulations to explore this behaviour.

Impact of sexual transmission on the epidemic tail. We performed stochastic

simulations of the EVD transmission model to investigate the epidemic dynamics when

the number of new cases becomes small, i.e, during the tail of the epidemic. Comparing

model simulations while assuming a convalescent period of 3 months to those without

sexual transmission confirmed the deterministic results that sexual transmission from

convalescent survivors does not lead to a significant increase in the cumulative number of

infected cases (non-STI: 11,086 +/- 599 cases; STI: 11,517 +/- 622 cases; Wilcox rank

sum test: W = 493156.5, p = 0.59; *Supplementary Material*, Fig. *S3*), nor the size (non-

STI: 86 +/- 4.4 new cases per day; STI: 89 +/- 4.6 new cases per day; W = 494255, p =

0.65) or timing (non-STI: day 200 +/- 0.7; STI: day 199 +/- 0.6; t = 1.1681, df = 991.14,

p = 0.24) of the epidemic peak incidence (Fig. 4a, 4b, 4c). Strikingly, this conservative

period of potential sexual transmission, which has recently been shown to extend well

beyond 3 months in at least 65% of patients [1], extended the average date on which the

last active case could be detected by nearly four months (non-STI: 497 +/- 2.9 days; STI:

619 +/- 6.0 days; difference: 123 days [95% CI: 110-137 days], *t* = -18.5, df = 730.36, p

< 0.0001; Fig. 4d, 4e, 4g, 4h). The width of the tail (s.d. = 135 days) was such that 18.5%

of the 508 simulated epidemics that accumulated at least 100 cases still experienced

symptomatic individuals 730 days (two years) after the start of the epidemic (Fig. 4h).

When the convalescent period was extended from 3 months to 6 months, the projected

length of the epidemic increased to a mean of 1102 days (+/- 14.9), with 89.0% of the

489 sustained epidemics taking over two years to end (Fig. 4i). Importantly, there is

greater variance in the tail of the epidemic when sexual transmission is considered, and

this uncertainty grows with the length of the convalescent period (Fig. 4g, 4h, 4i).

Discussion

Our study shows that sexual transmission of Ebola virus from convalescent men could

have a profound effect on the duration of the epidemic in Sierra Leone. Using a

mathematical transmission model, we found that an average duration of the convalescent

period of 3 months, and a per sex act transmission probability of 0.1% could extend the

EVD epidemic in Sierra Leone by 123 days (95% CI: 110–137 days). Such a scenario

would be consistent with the occurrence of a small number of sexual transmission events

during the end-phase of the epidemic. So far, the reported cases of sexual transmission of

EVD remain rare [1,2]. Hence, the per sex act transmission probability of Ebola virus

from male convalescent survivors is unlikely to be higher than 0.1%, but might well be

below this value. Our sensitivity analysis indicates that the duration of the EVD epidemic

is heavily influenced by the period during which convalescent men can transmit sexually,

calling for a better understanding of the persistence and duration of infectivity of Ebola

virus RNA in convalescent patients.

To our knowledge, this is the first study using mathematical modeling to assess the

potential impact of sexual transmission of Ebola virus on the epidemic in West Africa.

We extended an accepted modeling framework that has been widely used to describe the

epidemic trajectories of EVD outbreaks and epidemics [12–14]. Our baseline model

provides a good description of the EVD epidemic in Sierra Leone and we performed

sensitivity analysis to account for the considerable uncertainty in key unknown

parameters regarding sexual transmission. Given that chance will play a crucial role in

sexual transmission of Ebola virus, we used stochastic simulations to explore the

expected variability in the tail of the epidemic. In the absence of a better understanding of

sexual transmission of EVD, mathematical modeling currently remains the only tool to

explore its potential impact on the epidemic trajectory.

Our study provides a preliminary picture of the potential epidemiological consequences

of sexual transmission of Ebola virus on the epidemic in West Africa. Consequently,

there are a number of limitations that should be considered when interpreting the results.

First, we limited our analysis to sexual transmission from convalescent men who remain

infectious through the presence of Ebola virus in their semen. However, our modeling

framework could be easily extended to account for transmission by other routes, e.g.,

through infectious vaginal secretions from female survivors. Second, we assumed a

homogeneously mixing population to describe both non-sexual and sexual transmission

of EVD. Spatial structure might play an important role in the spread of EVD [26] as close

contact with infected bodily fluids is typically required even for non-sexual transmission.

In particular, close contact limited to family or community members during the

convalescence period, often accompanied by decreased activity levels due to post-Ebola

sequelae, may help limit the chances of sexual transmission. Third, asymptomatic

infection may occur [27], and is predicted to lower epidemic projections due to increased

levels of immunity in the population [28]. Fourth, the model does not incorporate

potential heterogeneity in sexual behavior [15], and we assumed that all sexually active

male survivors have the same average numbers of sex acts per month. The assumed

frequency of sex acts (8.3 sex acts per month) came from a study of heterosexual couples

in sub-Saharan Africa [29], and is on the high end of estimates gathered from studies of

other populations (e.g., [29–31], but see the wide range in individual frequencies in [32]);

the sexual activity of convalescent Sierra-Leonean EVD survivors has not been reported.

Fifth, EVD is known to exhibit superspreading characteristics [33,34], and we did not

include this in our model. Superspreading events could lead to explosive regrowth of the

epidemic after the occurrence of a new case through sexual transmission [33]. On the

other hand, 72% of infected individuals in transmission chains in Guinea did not generate

further cases [33], suggesting that some cases of sexual transmission of EVD might not

be detected at all. Sixth, we assumed a single compartment for convalescent survivors

with an exponentially distributed period of convalescence. Recent data indicates that the

time during which convalescent men remain positive for Ebola virus RNA could be better

described by a unimodal distribution [1]. Seventh, it remains to be determined whether

the virus found in semen of male convalescent survivors is replication competent, and the

time window during which these men remain infectious could be shorter than the duration

they remain positive for Ebola virus RNA. Finally, like other negative-sense single-

stranded RNA ((-)ssRNA) viruses [35], the species currently circulating in West Africa

has been estimated to have high substitution rates [36–38]. This rapid evolution detected

throughout the current outbreak zone suggests that within- or between-host adaptation of

the virus leading to pro-longed persistence in the seminal fluids is possible.

Awareness of the potential for sexual transmission has led to WHO issuing

recommendations that ask convalescent men to abstain from sexual activity as much as

possible and to use condoms for up to 6 months after the onset of symptoms [39].

Condom use and social awareness of the risks of sexual transmission during

convalescence should have an impact on the per sex act transmission probability (η) and

the frequency of sex acts (q), respectively. Our results show that condom use should

reduce the number of sporadic sexual transmission events during the tail of the epidemic

and after discharge of all remaining symptomatic individuals. However, the time during

which the public health community must stay vigilant is not reduced because these

interventions will not affect the time during which convalescent survivors can shed

infectious virus ($1/\alpha$). This is especially poignant since adherence to these

recommendations will never be 100%. Better knowledge about the rate at which

convalescent survivors recover completely and shed no further replicating Ebola virus

from their body is thus urgently needed to plan continued surveillance efforts.

The EVD epidemic in Sierra Leone was declared to be over on 7 November 2015 [40].

Our modeling results show that sporadic events of sexual transmission of Ebola virus –

potentially leading to small transmission clusters – could continue to occur in the months

to come. Sexual transmission of EVD is likely to be rare, but recent events such as the

suspected infection of a 16-year-old girl through sexual transmission in Makeni, Sierra

Leone, during September 2015 exemplify the continued risks [41]. In addition, the

considerable uncertainty around the infectivity of convalescent men who test positive for

Ebola virus RNA, and the duration of this period, asks us to remain vigilant. We must

monitor and quantify this phenomena in order to help us better prepare health networks,

and to provide convalescent survivors with sound advice that balances protection of the

community with the harm that could come from unnecessary stigmatization [42–44]. As

more data about the convalescent survivors of EVD becomes available, this and future

mathematical modeling studies will help to better understand the potential

epidemiological consequences of sexual transmission on the EVD epidemic in West

Africa.

Funding statement

JLA received post-doctoral support from the University of Bern and a grant from the

French National Research Agency (ANR JC "STORY"). CLA received funding through

an Ambizione grant from the Swiss National Science Foundation (project 136737). The

funders had no role in study design, analysis, decision to publish, or preparation of the

manuscript.

References

1. Deen GF, Knust B, Broutet N, Sesay FR, Formenty P, Ross C, et al. Ebola RNA

Persistence in Semen of Ebola Virus Disease Survivors — Preliminary Report. N

Engl J Med. 2015; doi:10.1056/NEJMoa1511410

2. Mate SE, Kugelman JR, Nyenswah TG, Ladner JT, Wiley MR, Cordier-Lassalle T,

et al. Molecular Evidence of Sexual Transmission of Ebola Virus. N Engl J Med.

2015; doi:10.1056/NEJMoa1509773

3. Pourrut X, Kumulungui B, Wittmann T, Moussavou G, Délicat A, Yaba P, et al.

The natural history of Ebola virus in Africa. Microbes Infect. 2005;7: 1005–14.

doi:10.1016/j.micinf.2005.04.006

4. Rowe AK, Bertolli J, Khan AS, Mukunu R, Muyembe-Tamfum JJ, Bressler D, et

al. Clinical , Virologic , and Immunologic Follow-Up of Convalescent Ebola

Hemorrhagic Fever Patients and Their Household Contacts , Kikwit , Democratic

Republic of the Congo. J Infect Dis. 1999;179 Suppl: 28–35.

5. Clark D V, Kibuuka H, Millard M, Wakabi S, Lukwago L, Taylor A, et al. Long-

term sequelae after Ebola virus disease in Bundibugyo, Uganda: a retrospective

cohort study. Lancet Infect Dis. Elsevier Ltd; 2015;3099: 1–8. doi:10.1016/S1473-

3099(15)70152-0

6. Bausch DG, Towner JS, Dowell SF, Kaducu F, Lukwiya M, Sanchez A, et al.

Assessment of the risk of Ebola virus transmission from bodily fluids and fomites.

J Infect Dis. 2007;196 Suppl : S142–S147. doi:10.1086/520545

7. Rodriguez LL, De Roo A, Guimard Y, Trappier SG, Sanchez A, Bressler D, et al.

Persistence and genetic stability of Ebola virus during the outbreak in Kikwit,

Democratic Republic of the Congo, 1995. J Infect Dis. 1999;179 Suppl : S170–6.

doi:10.1086/514291

8. Varkey JB, Shantha JG, Crozier I, Kraft CS, Lyon GM, Mehta AK, et al.

Persistence of Ebola Virus in Ocular Fluid during Convalescence. N Engl J Med.

2015;372: 2423–2427. doi:10.1056/NEJMoa1500306

9. Ergonul O, Battal I. Potential sexual transmission of Crimean-Congo hemorrhagic

fever infection. Jpn J Infect Dis. 2014;67: 137–8. Available:

http://www.ncbi.nlm.nih.gov/pubmed/24647261

10. Slenczka W, Klenk HD. Forty Years of Marburg Virus. J Infect Dis. 2007;196:

S131–135. doi:10.1086/520551

11. Rogstad KE, Tunbridge A. Ebola virus as a sexually transmitted infection. Curr

Opin Infect Dis. 2015;28: 83–85. doi:10.1097/QCO.0000000000000135

12. Chowell G, Hengartner NW, Castillo-Chavez C, Fenimore PW, Hyman JM. The

basic reproductive number of Ebola and the effects of public health measures: the

cases of Congo and Uganda. J Theor Biol. 2004;229: 119–126.

doi:10.1016/j.jtbi.2004.03.006

13. Althaus CL. Estimating the Reproduction Number of Ebola Virus (EBOV)

During the 2014 Outbreak in West Africa. PLOS Curr Outbreaks. 2014; 1–9.

doi:10.1371/currents.outbreaks.91afb5e0f279e7f29e7056095255b288.Abstract

14. Althaus CL, Low N, Musa EO, Shuaib F, Gsteiger S. Ebola virus disease outbreak

in Nigeria: Transmission dynamics and rapid control. Epidemics. Elsevier B.V.;

2015;11: 80–84. doi:10.1016/j.epidem.2015.03.001

15.   Anderson RM, May RM. Infectious Diseases of Humans: Dynamics and Control.

Oxford: Oxford Univeristy Press; 1991.

16.   Thrall PH, Antonovics J. Polymorphism in sexual versus non-sexual disease

transmission. Proc R Soc B Biol Sci. 1997;264: 581–587.

17.   Garnett GP. An introduction to mathematical models in sexually transmitted

disease epidemiology. Sex Transm Infect. 2002;78: 7–12. doi:10.1136/sti.78.1.7

18.   Diekmann O, Heesterbeek JAP. Mathematical Epidemiology of Infectious

Diseases. Chichester: John Wiley and Sons; 2000.

19.   Diekmann O, Heesterbeek J a, Metz J a. On the definition and the computation of

the basic reproduction ratio R0 in models for infectious diseases in heterogeneous

populations. J Math Biol. 1990;28: 365–382. doi:10.1007/BF00178324

20.   Kucharski AJ, Eggo RM, Watson CH, Camacho A, Funk S, Edmunds WJ.

Effectiveness of Ring Vaccination as Control Strategy for Ebola Virus Disease.

Emerg Infect Dis. 2016;22. doi:10.3201/eid2201.151410

21.   World Health Organization (WHO). Ebola data and statistics: Data on new cases

per epi week for Sierra Leone [Internet]. 2015 [cited 18 Nov 2015]. Available:

http://apps.who.int/gho/data/node.ebola-sitrep.ebola-country-SLE-

20151118?lang=en

22.   R Core Team. R: A Language and Environment for Statistical Computing

[Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2014.

Available: http://www.r-project.org/

23.   Chalom A, Prado PIKL. pse: Parameter space exploration with Latin Hypercubes

[Internet]. 2014. Available: http://cran.r-project.org/package=pse

24. Blower SM, Dowlatabadi H. Sensitivity and Uncertainty Analysis of Complex

Models of Disease Transmission: An HIV Model, as an Example. Int Stat Rev.

1994;62: 229–243. doi:10.2307/1403510

25. Gillespie DT. Exact Stochastic Simulation of couple chemical reactions. J Phys

Chem. 1977;81: 2340–2361. doi:10.1021/j100540a008

26. Merler S, Ajelli M, Fumanelli L, Gomes MFC, Piontti APY, Rossi L, et al.

Spatiotemporal spread of the 2014 outbreak of Ebola virus disease in Liberia

and the effectiveness of non-pharmaceutical interventions: a computational

modelling analysis. Lancet Infect Dis. Elsevier Ltd; 2015;3099: 1–8.

doi:10.1016/S1473-3099(14)71074-6

27. Leroy EM, Baize S, Volchkov VE, Capron M, Debré P, Mccormick JB, et al.

Human asymptomatic Ebola infection and strong inflammatory response. Lancet.

2000;355: 2210 – 2215.

28. Bellan SE, Pulliam JRC, Dushoff J, Meyers LA. Ebola control: effect of

asymptomatic infection and acquired immunity. Lancet. Elsevier Ltd; 2014;384:

1499–1500. doi:10.1016/S0140-6736(14)61839-0

29. Gray RH, Wawer MJ, Brookmeyer R, Sewankambo NK, Serwadda D, Wabwire-

Mangen F, et al. Probability of HIV-1 transmission per coital act in monogamous,

heterosexual, HIV-1-discordant couples in Rakai, Uganda. Lancet. 2001;357:

1149–1153. doi:10.1016/S0140-6736(00)04331-2

30. Blower SM, Boe C. Sex acts, sex partners, and sex budgets: implications for risk

factor analysis and estimation of HIV transmission probabilities. J Acquir Immune

Defic Syndr. 1993;6: 1347–1352.

31.   Althaus CL, Choisy M, Alizon S, CSF group. Number of sex acts matters for

433   heterosexual transmission and control of Chlamydia trachomatis. PeerJ Prepr.

434   2015;3: e1164. doi:10.7287/peerj.preprints.940v1

435   32.   Smith J, Nyamukapa C, Gregson S, Lewis J, Magutshwa S, Schumacher C, et al.

436   The Distribution of Sex Acts and Condom Use within Partnerships in a Rural Sub-

437   Saharan African Population. PLoS One. 2014;9: e88378.

438   doi:10.1371/journal.pone.0088378

439   33.   Althaus CL. Ebola superspreading. Lancet Infect Dis. 2015;15: 507–508.

440   doi:10.1016/S1473-3099(15)70135-0

441   34.   Toth DJ a, Gundlapalli A V, Khader K, Pettey WBP, Rubin M a, Adler FR, et al.

442   Estimates of Outbreak Risk from New Introductions of Ebola with Immediate and

443   Delayed Transmission Control. Emerg Infect Dis J. 2015;21: 1402.

444   doi:10.3201/eid2108.150170

445   35.   Duffy S, Shackelton L a., Holmes EC. Rates of evolutionary change in viruses:

446   patterns and determinants. Nat Rev Genet. 2008;9: 267–276. doi:10.1038/nrg2323

447   36.   Gire SK, Goba A, Andersen KG, Sealfon RSG, J PD, Kanneh L, et al. Genomic

448   surveillance elucidates Ebola virus origin and transmission during the 2014

449   outbreak. Science (80-). 2014; doi:10.1126/science.1259657

450   37.   Park DJ, Dudas G, Wohl S, Goba A, Whitmer SLM, Andersen KG, et al. Ebola

451   Virus Epidemiology, Transmission, and Evolution during Seven Months in Sierra

452   Leone. Cell. 2015;161: 1516–1526. doi:10.1016/j.cell.2015.06.007

38.   Tong Y-G, Shi W-F, Liu D, Qian J, Liang L, Bo X-C, et al. Genetic diversity and

evolutionary dynamics of Ebola virus in Sierra Leone. Nature. 2015;524: 93–96.

doi:10.1038/nature14490

39.   World Health Organization (WHO). WHO website [Internet]. Available:

http://www.who.int/reproductivehealth/topics/rtis/ebola-virus-semen/en/

40.   World Health Organization (WHO). WHO Ebola Situation Report (11 November

2015). [Internet]. 2015. Available: http://apps.who.int/ebola/current-

situation/ebola-situation-report-11-november-2015

41.   Reuters News Agency. Hundreds quarantined as Ebola returns to north Sierra

Leone district. Sept 14. 2015. Available:

http://www.reuters.com/article/2015/09/14/us-health-ebola-leone-

idUSKCN0RE1P620150914

42.   Davtyan M, Brown B, Folayan MO. Addressing Ebola-related Stigma: Lessons

Learned from HIV/AIDS. Glob Health Action. 2014;7: 1–4.

doi:10.3402/gha.v7.26058

43.   Obilade TT. Ebola Virus Disease Stigmatization; The Role of Societal Attributes.

Int Arch Med. 2015;8: 1–19. doi:10.3823/1613

44.   Sprecher A. Handle Survivors with Care. N Engl J Med. 2015;

doi:10.1056/NEJMe1512928

45.   WHO Ebola Response Team. Ebola Virus Disease in West Africa — The First 9

Months of the Epidemic and Forward Projections. Engl New J Med. 2014;371:

1481–1495. doi:10.1056/NEJMoa1411100

46.   Alizon S, Lion S, Murall CL, Abbate JL. Quantifying the epidemic spread of Ebola

476   virus (EBOV) in Sierra Leone using phylodynamics. Virulence. 2015;5: 825–827.

477   doi:10.4161/21505594.2014.976514

478   47.   World Health Organization (WHO). Data from: Ebola data and statistics [Internet].

479   2015. Available: http://apps.who.int/gho/data/view.ebola-sitrep.ebola-country-

480   SLE-20150923-data?lang=en

481   48.   The World Bank. Sierra Leone country at a glance [Internet]. 2015. Available:

482   http://www.worldbank.org/en/country/sierraleone

483   49.   Wawer MJ, Gray RH, Sewankambo NK, Serwadda D, Li X, Laeyendecker O, et

484   al. Rates of HIV-1 transmission per coital act, by stage of HIV-1 infection, in

485   Rakai, Uganda. J Infect Dis. 2005;191: 1403–1409. doi:10.1086/429411

487 **Table 1: Model parameters describing EVD transmission in Sierra Leone.** The

488 indicated parameter ranges are used for the sensitivity analysis (partial rank correlation

489 coefficients, PRCC) only.

Parameter		Value (Range)	Comments and References
Basic reproductive number	$R_{0,N}$	2.02 (1.26 – 2.53)	Non-sexual transmission only [45] Range explores estimates from [13] and [46]
Date of onset of symptoms in index case		April 6, 2014	WHO epidemic day 92, epidemic week 14 [47]
Rate at which transmission rate decays	k	0.01 d^{-1}	Results in a good description of the EVD epidemic in Sierra Leone
Time at which transmission rate starts to decay	τ	90 d	We assume that control measures were not effectively deployed country-wide until three months following the first detected case ($R_{0,N}$ estimates began to fall in August 2014 [45]).
Initial population size	N_0	6.316×10^6	Based on 2014 estimate [48]
Incubation period	$1/\sigma$	9.31 d	[14]
Case fatality rate	f	0.69	[45]
Infectious period	$1/\gamma$	7.41 d	[14]
Sexual transmission probability (per coital act)	η	0.001 (0.0005 – 0.002)	Roughly based on sexual transmission probability of HIV per coital act from infected men [49]
Frequency of sex acts	q	0.272 d^{-1} (0.136 – 0.544)	8.27 coital acts per month [29]
Proportion of convalescent survivors who are infectious and sexually active	p	0.347 (0.1725-0.694)	Of 47.4% male survivors, 73.1% are aged 15-45 [45].
Rate at which convalescent survivors recover completely	α	1/83.84 d^{-1} (1/168 – 1/42)	3 months after onset of symptoms [4,39] and assuming an infectious period of 7.41 days

491 **Figure 1. Schematic illustration of EVD transmission model including sexual**

492 **transmission from convalescent patients.** The elements in black form the base model

493 without sexual transmission [12–14]. The red elements (convalescent individuals and

494 additional transmission term) were added to account for sexual transmission.

 Figure 3. Effect of per sex act transmission probability on EVD epidemics. The per

sex act transmission probability (η) is varied between 0.05% and 0.2%. (a, b): Epidemic

trajectories in presence (broken lines) and absence of sexual transmission (solid lines).

Vertical lines mark the day of last symptomatic case. (c) Daily incidence of sexual

transmission. (d) Cumulative number of sexual transmission events. Note that the vertical

axes vary across panels.

Figure 4. Impact of convalescent period on the tail of the EVD epidemic. Monte

Carlo simulations of the weekly incidence of new cases (a, b, c) and the sporadic

occurrence of sexual transmission events at the epidemic tail (d, e, f), assuming no sexual

transmission (a, d), and sexual transmission with a 3 (b, e) and 6 months (c, f)

convalescent period. Thin red lines show the result of 200 simulated trajectories, with

corresponding mean (thick blue line) and standard errors (thin blue lines). Black dots

denote incident cases in Sierra Leone as reported by the WHO [21]. The thick red lines in

(d), (e), and (f) highlight a single representative trajectory, and the dark red dots along the

horizontal axis indicate a sexual transmission event. Histograms of the day of the last

EVD case from 1,000 simulated epidemics in absence of sexual transmission (g) and for a

convalescent period of 3 (h) and 6 (i) months.

523