Latitude delineates patterns of biogeography in terrestrial *Streptomyces*

Mallory J Choudoir, James R Doroghazi, and Daniel H Buckley*

School of Integrative Plant Sciences, Cornell University, Ithaca, NY 14853 USA

* Corresponding Author: Daniel H Buckley
dbuckley@cornell.edu

Running Title: Biogeography of *Streptomyces*

Subject Category: Microbial population and community ecology

Conflict of Interest: None

Abstract

The evolutionary and ecological forces that govern microbial biogeography remain poorly

characterized. We examined the biogeography of *Streptomyces* at regional spatial scales to

identify factors that govern extant patterns of microbial diversity within the genus. *Streptomyces*

are spore forming filamentous bacteria that are widespread in soil and are a predominant source

of antibiotics. We applied population genetic approaches to analyze geographic and genetic

population structure in six phylogroups of *Streptomyces* identified from a geographically explicit

culture collection. *Streptomyces* strains were isolated from soils associated with perennial grass

habitats sampled across a spatial scale of more than 5,000 km. We find that *Streptomyces* allelic

diversity correlates with geographic distance and that gene flow between sites is constrained by

latitude. In addition, we find that phylogroup nucleotide diversity is negatively correlated with

latitude. These observations are consistent with the hypothesis that historical demographic

processes have influenced the contemporary biogeography of *Streptomyces*.

Introduction

Widespread evidence documents microbial biogeography across varying spatial (Whitaker *et al.*,

2003; Vos and Velicer, 2008; Bissett *et al.*, 2010; Martiny *et al.*, 2011) and temporal scales

(Gilbert *et al.*, 2012; Hatosy *et al.*, 2013), yet we are only beginning to understand the

evolutionary forces that generate and maintain these patterns. Explorations of biogeography are

valuable because biogeographical patterns illustrate fundamental principles of evolution and

ecology. One of the earliest and most well documented patterns of biogeography is the latitudinal

diversity gradient (Wallace, 1878), and this trend of decreasing species richness with increasing

latitude is widely described in plant and animal systems (Hillebrand, 2004). We examine the

biogeography of *Streptomyces* to determine whether a latitudinal diversity gradient can be

observed in terrestrial bacteria and to evaluate the underlying mechanisms that govern

biogeography within this genus.

Patterns of microbial biogeography are determined by rates of dispersal and diversification

(Martiny *et al*., 2006; Hanson *et al*., 2012). Rates of dispersal can be difficult to quantify for

microbes because dispersal that does not contribute to the genetic diversity of future populations

remains undetected. Gene flow results from the transfer of genetic material between populations,

and gene flow between geographically distinct populations requires dispersal. Hence, patterns of

microbial dispersal are typically inferred from extant patterns of genetic diversity. It has been

hypothesized that microbes disperse ubiquitously due to their small cell size and massive

population numbers (Finaly, 2002; Finlay and Fenchel, 2004). For example, Gibbons *et al*.

(2013) determined that a single sampling site in the English Channel could encapsulate the total

expected diversity of the Earth's marine biosphere. Yet endemism and dispersal limitation have

been observed for a range of microbes (Cho and Tiedge, 2000; Green and Bohannan, 2006;

Telford *et al*., 2006; Boucher *et al*., 2011), and microbial dispersal limitation has been verified

experimentally (Bell, 2010). Contrasting evidence for dispersal limitation in the literature can be

explained by at least two factors: firstly, dispersal constraints are likely to vary between species

and habitats; and secondly, the units used to define microbial diversity can vary dramatically in

their phylogenetic resolution. Each of these factors has been discussed previously (Hanson *et al*.,

2012; Choudoir *et al*., 2012), and we will only consider them briefly here.

Patterns of microbial dispersal and gene flow appear to differ considerably between habitats and

species. At one end of the spectrum, globally widespread marine microbes such as

Prochlorococcus and *Pelagibacter* show little variation in gene content between the Atlantic and

Pacific Oceans suggesting that high dispersal can homogenize genetic diversity in pelagic marine

systems, while adaptation to regional habitats occurs within limited regions of the genome

(Coleman and Chisholm, 2010). At the other end of the spectrum are extremophiles such as

Sulfolobus and thermophilic *Synechococcus*, which live in island-like volcanic habitats and

exhibit strong patterns of allopatric divergence resulting from dispersal limitation (Papke *et al*.,

2003; Whitaker *et al*., 2003). Both allopatric processes and local patterns of gene flow contribute

to the biogeography and genetic diversity of *Sulfolobus* populations (Cadillo-Quiroz *et al*.,

2012). Terrestrial microbes fall somewhere between these extremes. Soil dwelling microbes such

as *Burkholderia pseudomallei*, *Burkholderia mallei*, and *Bacillus anthracis* exhibit

biogeographical patterns governed by dispersal limitation at regional spatial scales (Kenefic *et*

al., 2009; Pearson *et al*., 2009). For instance, the contemporary population structure of North

American anthrax reflects historical migration across the Bering land bridge during the late

Pleistocene followed by range expansion into the Southwestern United States and regional

diversification due to drift (Kenefic *et al.*, 2009).

The phylogenetic resolution at which microbial diversity is defined can have a profound impact

on our ability to discern patterns of microbial biogeography (as reviewed Hanson *et al.*, 2012).

Observations of microbial biogeography are often based upon variation in SSU rRNA gene

sequences. Surveys of SSU rRNA genes in terrestrial habitats indicate that environmental

variables including temperature (Fierer *et al*., 2009; Miller *et al*., 2009), pH (Fierer and Jackson,

2006; Lauber *et al*., 2009; Rousk *et al*., 2010), and salinity (Lozupone and Knight, 2007) are

more important than geographic distance in determining patterns of microbial diversity and

biogeography. However, SSU rRNA gene sequences have an extremely low rate of nucleotide

substitution, and the formation of discrete taxonomic units as defined by this locus (i.e.

Operational Taxonomic Units defined at 97% nucleotide identity) requires millions of years

(Ochman and Wilson, 1987; Ochman *et al*., 1999). Hence, microbes with similar or even

identical SSU rRNA genes can have extensive genomic and ecological diversity (Welch *et al*.,

2002; Jaspers and Overmann; 2004) which far exceeds the diversity found within plant and

animal species. Neutral processes that drive patterns of biogeography, such as dispersal

limitation and genetic drift, can only be detected using genetic techniques that have high

resolution (Green and Bohannan, 2006; Hanson *et al*., 2012). Fortunately, the evolutionary and

ecological forces that drive microbial biogeography are readily explored using geographically

and ecologically explicit culture collections that can be characterized at high genetic resolution

(Hunt *et al.*, 2008; Shapiro *et al.*, 2012).

Streptomyces are spore forming bacteria ubiquitous in soils, and many species are easily

cultured. Furthermore, *Streptomyces* species have high rates of gene exchange both within and

between species (Doroghazi and Buckley, 2010). Hence this genus represents a useful model for

examining limits on dispersal and gene flow in terrestrial bacterial communities. *Streptomyces*

are gram-positive *Actinobacteria* (Kâmpfer, 2006) known for their complex developmental

cycle, which entails mycelial growth followed by formation of aerial hyphae and desiccation

resistant spores which are readily dispersed (Keiser *et al.*, 2000). *Streptomyces* are able to

degrade recalcitrant carbon molecules like cellulose and play an important role within the

terrestrial carbon cycle (McCarthy and Williams, 1992; Takasuka *et al.*, 2013). In addition,

phytopathogenic *Streptomyces,* such as *S. scabies,* cause substantial economic losses in

agriculture (Loria *et al.*, 2006; Labeda, 2011). Finally, *Streptomyces* are prolific producers of

secondary metabolites, and most antibiotics we use today were discovered in this genus (Watve

et al., 2001). Despite their clinical, agricultural, and environmental importance, we still lack an

evolutionary and ecological framework to understand *Streptomyces* biodiversity and

biogeography.

Streptomyces diversity varies spatially, though the effects of geographic distance and ecological

variation on *Streptomyces* diversity remains poorly resolved. *Streptomyces* genetic and

phenotypic diversity can vary at small spatial scale (1 m^2) in prairie soils (Davelos *et al.*, 2004a

and 2004b) and across dune and beach habitats (60 m) (Antony-Babu *et al.*, 2008), suggesting

that variation in local soil characteristics can select for species and populations with different

ecological adaptations. There is also evidence that some *Streptomyces* species are endemic to

North America or Central Asia, which suggests dispersal limitation at large scales (Wawrik *et al.*,

2007). Analysis of genetic diversity within a single population of *Streptomyces pratensis,*

isolated from soils that spanned 1,000 km, revealed a population in linkage equilibrium when

assessed both by multi-locus sequence analysis (MLSA) and genome analysis (Doroghazi and

Buckley, 2010 and 2014). This result indicates that either gene flow is unlimited in *S. pratensis*

across large spatial scales, or that it has experienced an evolutionarily recent demographic

expansion from a population that was once in linkage equilibrium. Furthermore, analysis of

ancestral patterns of *Streptomyces* inter-species gene exchange suggests that the fixation of

incongruent loci could be due to demographic range expansion into higher latitudes (Andam *et*

al., 2015). These latter data suggest a role for both adaptive and neutral processes in governing

the diversity and biogeography of *Streptomyces*, though the degree to which these forces

constrain extant patterns of genetic diversity within the genus remain poorly described.

We evaluated *Streptomyces* biogeography by constructing an isolate collection from grassland

sites spanning the United States of America. The most powerful approach for detecting neutral

evolutionary processes associated with dispersal limitation is to examine patterns of

biogeography across similar habitats (Hanson *et al.*, 2012). In a previous study which employed

a single locus (*rpoB*), we observed that *Streptomyces* species diversity exhibited a significant

distance decay relationship and was inversely correlated with latitude (Andam *et al.,* 2016).

Evidence for a latitudinal diversity gradient coupled with evidence of demographic range

expansion, while not conclusive, is consistent with predictions of phylogenetic niche

conservatism (Wiens and Donoghue, 2004; Stevens, 2006) which posits that these

biogeographical patterns result from the historical effects of changes in paleoclimate (Hewitt *et*

al., 1996, 2000, and 2004). The hypothesis of a latitudinal diversity gradient in *Streptomyces*

predicts that, in addition to a gradient of species richness, there should also be latitudinal

differences in gene flow and genetic diversity within populations. Here we investigate genetic

diversity and geographic distribution within six species-like phylogenetic clusters, or

phylogroups, to determine whether patters of gene flow and diversity corroborate the hypothesis

of a latitudinal diversity gradient for *Streptomyces*.

Materials and Methods

Strain isolation and DNA extraction

We assembled a culture collection of more than 1,000 *Streptomyces* from 15 sites across the

United States with soil sampled at 0-5 cm depth (Andam *et al.,* 2016). Sites were selected to

represent a narrow range of habitats dominated by perennial grasses and having neutral to

slightly acidic pH (3.9-7.3). Soil was air dried, and *Streptomyces* strains were isolated on

glycerol-arginine agar plates of pH 8.7 containing cycloheximide and Rose Bengal (El-Nakeeb

and Lechevalier, 1963; Ottow, 1972) as previously described (Doroghazi and Buckley, 2010).

DNA was extracted from purified cultures, which were grown by shaking at 30˚C in liquid yeast

extract-malt extract medium (YEME) containing 0.5% glycine (Kieser *et al*., 2000), by using a

standard phenol/chloroform/isoamyl alcohol protocol.

The genetic diversity of isolates was initially assessed using partial *rpoB* sequences.

Streptomyces species are typically delineated by 0.007 nucleotide dissimilarity across MLSA loci

(Rong and Huang, 2012). Since we were using only a single locus we elected to define species-

like phylogentic clusters, or phylogroups, at 0.01 patristic distances with RAMI (Pommier *et al*.,

2009). Using this approach, we identified 107 phylogenetic clusters based on *rpoB* sequences.

We selected six of these phylogroups for further analysis. MLSA was performed on 17-47

isolates from each phylogroup. Phylogroup names are capitalized and based on a representative

isolate; for example, phylogroup WA1063 is named for *Streptomyces* sp. wa1063. Isolates are

identified with a lowercase letter code indicating site of origin followed by a strain number; for

example, isolate wa1063 is strain 1063 isolated from Washington (WA) state.

Multilocus sequence analysis (MLSA)

We adapted the MLSA scheme developed for *Streptomyces* by Guo *et al*. (2008), which targets

the five housekeeping loci *atpD*, *gyrB*, *recA*, *rpoB*, and *trpB* as described in Doroghazi and

Buckley (2010). Partial SSU rRNA sequences were amplified using universal primers (8F:

AGAGTTTGATCCTGGCTCAG; 1492R: CGGTTACCTTGTTACGACTT). Reactions for

Sanger sequencing were performed using forward primers for all loci except *rpoB*, for which the

reverse primer was used. Trace files were uniformly screened using CAP3 (Huang and Madan,

1999) interfaced with a Perl script. Sequences were aligned using MUSCLE (Edgar, 2004),

trimmed to 431 bp, 415 bp, 446 bp, 557 bp, and 489 bp, for genes *atpD*, *gyrB*, *recA*, *rpoB*, *trpB*,

respectively, and concatenated consistent with the genomic order in *Streptomyces coelicolor*

A3(2). SSU rRNA alignments were trimmed to 357 bp. Gene sequences are available on

GenBank with accession numbers XXXX-XXXX.

Good's coverage estimation and haplotype rarefaction was determined using mothur (Schloss *et*

al., 2009). DNA polymorphism statistics including: number of segregating sites, nucleotide

diversity, and Tajima's D, were determined with DnaSP v5 (Librado and Rozas, 2009) and LDhat

(McVean *et al.*, 2002). Population scaled mutation rates (Watterson's theta; $\theta_w = 2N_e\mu$),

recombination or gene conversion rates ($\rho=2N_er/2$), and relative rates of recombination (ρ/θ_w)

were estimated using LDhat (McVean *et al.*, 2002) and are expressed per nucleotide site. The

standard index of association (I_A) was calculated from allelic data with LIAN v3.5 (Haubold and

Hudson, 2000) using the Monte-Carlo test and 100 iterations. The pairwise homoplasy index

(PHI) statistic was determined using PhiPack (Bruen *et al.*, 2006), and statistical significance

was evaluated under a null hypothesis of no recombination. Sequence identity across

phylogroups was calculated with mothur (Shloss *et al.*, 2009)

Phylogengy reconstruction

Maximum likelihood (ML) trees were constructed from the nucleotide sequences of individual

and concatenated MLSA loci using the generalized time reversible nucleotide substitution model

(Tavaré, 1986) with gamma distributed rate heterogeneity among sites (GTRGAMMA)

supported in RAxML v7.3.0 (Stamatakis, 2006). Bootstrap support was determined for the

highest-scoring ML tree of 20 iterations, and the number of bootstrap replicates was determined

using the extended majority rule (autoMRE) convergence criteria (Pattengale *et al*., 2010). Root

placement is defined by *Mycobacterium smegmatis.* Significant phylogenetic incongruence

between loci was determined using the Shimodaira-Hasegawa test (Shimodaira and Hasegawa,

1999) implemented in the R package phangorn (Schliep, 2011).

Population structure

Concatenated MLSA sequences were analyzed using Structure v2.3.3 (Pritchard *et al*., 2000) to

examine population affiliation, subdivision, and admixture within and between phylogroups.

Structure was run using an admixture model with a burn-in length of $1.0E^6$ and $3.0E^6$ replicates.

The most probable number of sub-populations (k) was chosen using the Evanno method (Evanno

et al., 2005) from 10 independent runs with k = 1 through k = 6 within phylogroups, and through

k = 8 between phylogroups implemented by Structure Harvester (Earl and vonHoldt, 2012).

After choosing the most probable k-value, the program Clumpp was used to permute outputs of

the independent runs (Jakobsson and Rosenberg, 2007).

Patterns of dispersal and gene flow

We used permutation tests to evaluate whether phylogroup distribution across sites could be

explained by panmixia. We compared the observed distributions to those expected under a

random distribution model using 1,000 permutations to assess significance. In addition,

correlations between geographic distance and allele composition between sites were assessed

using Mantel and partial Mantel tests (Mantel, 1967; Smouse *et al.*, 1986). These tests were

performed with the R package ecodist (Goslee and Urban, 2007) using the Pearson correlation

method and 1,000 permutations. Bray-Curtis dissimilarity was calculated from unique allele

composition across sites.

Haplotype networks were created using a statistical parsimony procedure (Templeton *et al.*,

1987; Templeton *et al.*, 1995) implemented in TCS v1.18 (Clement *et al.*, 2000). Nested clade

information was used to infer processes that could explain the geographic and genetic

distribution of sequences using the program GeoDis v2.2 (Posada *et al.*, 2000). Both TCS v1.18

and GeoDis v2.2 were performed in ANeCA (Panchal, 2007).

Results

Characterization of Streptomyces phylogroups

We identified a total of 308 *Streptomyces* isolates representing the six targeted phylogroups, and

these isolates spanned 13 sites (Figure 1, Table 1). The six phylogroups all share more than 97%

nucleotide identity in their SSU rRNA genes and greater than 88% average nucleotide identity

(ANI) across MLSA loci. Strains within each phylogroup share 99.4%-100% nucleotide identity

in their SSU rRNA genes (Figure 2) and 97.6%-99.5% ANI across concatenated MLSA loci

(Figure 1, Table 2). The genetic ancestry of phylogroup members as determined by Structure is

congruent with phylogroup boundaries defined by nucleotide cutoffs (Figure S1), and this

indicates that phylogroups approximate biological populations. There are a total of 122 unique

MLSA haplotypes, with each phylogroup representing 15-26 haplotypes (Figure 1, Table 2).

Good's coverage for each of the six phylogroups ranges from 0.88-1.0 for individual loci and

0.94-1.0 for concatenated MLSA loci clustered at 99% nucleotide identity (Table S1). While

unique haplotypes remain somewhat under sampled (Figure S2), allelic diversity at single loci is

sampled well (Table S1). The per site nucleotide diversity (π) of each phylogroup ranges from

0.0026 to 0.011 (Table 2).

Four of our phylogroups do not match any *Streptomyces* species described in PubMLST (less

than 97% ANI across MLSA loci, Figure S3A) (Jolley *et al.*, 2004). However, phylogroups

WA1063 and MS152 affiliate with strains that belong to the *S. griseus* species cluster (Rong and

Huang, 2010). Isolates in phylogroup WA1063 share greater than 99% MLSA identity with *S.*

anulatus and *S. praecox* and form a monophyletic clade with these type strains (Figure S3A).

Isolates in phylogroup MS152 share greater than 98% MLSA identity with *S. mediolani*, *S.*

albovinaceus, and *S. griseinus* and form a paraphyletic clade that includes these type strains

(Figure S3A).

We find evidence of horizontal gene transfer consistent with previous observations of

Streptomyces species (Doroghazi and Buckley, 2010; Andam *et al.*, 2015). There is significant

phylogenetic incongruence between MLSA loci (Figure S3, Table S2), suggesting that horizontal

gene transfer has shaped the phylogeny of these groups. The six phylogroups present evidence of

population structure and admixture, with each phylogroup composed of 3.2 ± 0.8 (mean ± s.d.)

subpopulations (Figure 1). Evidence of admixture between populations suggests horizontal gene

transfer within phylogroups as has been previously described in *Streptomyces* (Doroghazi and

Buckley, 2010). Likewise, there is evidence of significant levels of recombination within

phylogroups MAN196, MS200, MS152, and F34 (PHI test, $p < 0.05$, Table 2). Phylogroups

WA1063 and MS152 share two identical *atpD* alleles, with one of these alleles shared among

strains isolated from MS, CA, WI (Brookfield), AK (Manley Hot Springs), and WA, and the

other allele shared among strains isolated from MS and WI (Brookfield) (Figure 3, Figure S3B).

It is not clear whether these alleles are shared as a result of contemporary horizontal gene

transfer or vertical inheritance from the most recent common ancestor of the two clades, though

the latter explanation is more parsimonious given the low level of polymorphism between the

phylogroups.

Dispersal limitation

Strains of the six phylogroups were obtained from soil samples from 13 sites of diverse

geographic origin (Table 1). Each phylogroup was detected in 4.2 ± 0.4 sites, and this

distribution differs significantly from expectations for a random distribution of strains across

sites (permutation test, $p < 0.0005$), thereby rejecting the hypothesis of panmixia. Each

phylogroup subpopulation was observed in 2.2 ± 0.9 sites (mean $\pm$ s.d.), and this value is lower

than expected if subpopulations are randomly distributed across the sites occupied by each

phylogroup (permutation test, $p < 0.001$). These results indicate that phylogroup distribution is

constrained geographically and that phylogroups have subpopulation structure that is also

geographically explicit.

The geographic distribution of *Streptomyces* allelic diversity indicates dispersal limitation.

Identical alleles are shared among phylogroup members across each phylogroup's geographic

range, which can exceed 5,000 km (Figure 3, Figure S3). However, dissimilarity in allele

composition increases with geographic distance, and this result is significant (Bray-Curtis

dissimilarity, Mantel r = 0.29, p = 0.005) (Figure S4). Hence, alleles are more likely shared

between geographically similar sites indicating dispersal limitation with potential for long range

dispersal. This result is significant for all individual loci expect $recA$ (Bray-Curtis dissimilarity,

$aptD$ Mantel r = 0.31, p = 0.004; $gyrB$ r = 0.22, p = 0.031; $recA$ r = 0.16, p = 0.088; $rpoB$ r =

0.27, p = 0.004; $trpB$ r = 0.19, p = 0.047). Nearly all MLSA haplotypes (Figure 1, Figure 4) are

unique to a single site, with the sole exception being a haplotype from phylogroup MS200 which

is shared between strains ms53, ms30, and sun103 isolated from MS and WI (Sun Prairie). It is

worth noting that all of the strains examined share > 97% SSU rRNA gene identity and these

geographic patterns would not be detected through analysis of this locus. That is, everything

would appear to be everywhere based on analysis of SSU rRNA genes.

Analysis of haplotype distribution is consistent with diversification due to spatial isolation

resulting from dispersal limitation. Nested clade analysis (NCA) establishes significant

phylogeographic inferences for phylogroups MAN196, MAN125, WA1063, and MS152 (Figure

4) but not for MS200 and F34. Nested clade phylogeographic inference postulates potential

evolutionary and historical demographic processes that support extant patterns of diversity and

biogeography. For instance, population subdivision of MAN125 across the Pacific Northwest

(Figure 4A) and MAN196 between Maine and the Pacific Northwest (Figure 4B) is consistent

with restricted gene flow due to historical long distance dispersal events. Likewise, population

subdivision of MS152 and between the Southeast (MS and FL) and CA (Figure 4C) and WA1063

between WI (Brookfield) and OR (Figure 4D) is consistent with allopatric fragmentation.

Latitudinal diversity gradient

The distribution and diversity of the phylogroups reveals a latitudinal diversity gradient. Strains

from MAN125, MAN196, WA1063 occur mostly north of 40°N latitude, while strains from

MS200, MS152, F34 occur mostly south of this latitude (Table 1). This pattern of north/south

partitioning is significant for each phylogroup when evaluated against the expectation of a

random distribution across sites (permutation test, $p < 0.01$ for each phylogroup after Bonferroni

correction). Furthermore, partial Mantel tests were performed to evaluate the latitudinal and

longitudinal vector components of geographic distance in relation to the allele composition of

sites. There remains a significant relationship between allele composition and geographic

distance when we control for longitude, (Mantel r = 0.23, $p = 0.022$), but this relationship is no

longer significant when we control for latitude (Mantel r = 0.15, $p = 0.12$). This result indicates

that allele composition changes more across latitude than it does across longitude. The latitudinal

partitioning of alleles can be readily observed in the pattern of allele sharing between sites

(Figure 3). Finally, we also observed a significant relationship between per site nucleotide

diversity of phylogroup MLSA loci and the average latitude of sites in which they are found (R =

-0.91, $p = 0.012$; Figure 5). This result indicates that phylogroups recovered from lower latitudes

have higher genetic diversity than those recovered from higher latitudes.

Discussion

We used population genetic approaches to analyze spatial patterns of genetic diversity for six

Streptomyces phylogroups isolated from geographically disparate but ecologically similar sites

across the United States (Table 1). The distribution of phylogroups is nonrandom and dispersal

limited (Figure S4) with phylogroups inhabiting geographic ranges defined by latitude (Table 1,

Figure 3). In addition, the genetic diversity of phylogroups is inversely proportional to the

latitude from which they were isolated (Figure 5). These findings suggest that that there are

latitudinal barriers to dispersal, and that patterns of *Streptomyces* biogeography result from

dispersal limitation and regional diversification due to genetic drift.

The phylogroups we describe are coherent phylogenetic groups that approximate biological

populations (Figure S1). The members of each phylogroup share a distinguishable geographic

range (Table 1, Figure 3), a recent common ancestor (Figure 1), greater than 97% ANI across

MLSA loci (Table 2), and greater than 99.4% SSU rRNA gene nucleotide identity. Despite

geographic and genetic subpopulation structure (Figure 1), phylogroup members frequently share

identical alleles across demes. We infer the presence of recombination within all phylogroups

using both nucleotide polymorphism and phylogenetic methods (Table 2, Table S2). For instance,

MAN125 is nearly in linkage equilibrium (I_A = 0.09) though its members span a geographic

range of over 2,000 km across the Pacific Northwest. The observation of linkage equilibrium

indicates that either the population lacks contemporary barriers to gene flow or the population

has experienced a recent demographic expansion in which there has been insufficient time to

accumulate mutations and linkage disequilibrium. Significant recombination and linkage

equilibrium have also been observed in a *S. pratensis* population which spanned 1,000 km across

sites present in North Carolina and northern New York (Doroghazi and Buckley, 2010 and 2014).

Regional patterns of biogeography among phylogroups are consistent with limitations to

dispersal and gene flow. Allopatric processes like genetic drift can drive diversification between

populations that are geographically isolated. The geographic distribution of our phylogroups is

nonrandom, and we find regional subpopulation structure within phylogroups (Figure 1, Figure

4). Although we find identical alleles in sites thousands of kilometers apart (Figure 3), MLSA

haplotypes are not shared across sites (with the single exception of a haplotype shared between

MS and WI) (Figure 4). We also observe a significant distance decay relationship for MLSA

allele composition and geographic distance between sites (Figure S4). This data implies that

while gene flow is moderate across the geographic range of a phylogroup, dispersal limitation

and genetic drift have combined to produce regional populations whose distributions are

constrained by latitude.

Streptomyces phylogroup diversity is consistent with a latitudinal diversity gradient. We find that

latitude is a significant predictor of gene flow (Figure 3). Furthermore, intra-phylogroup

nucleotide diversity has a significant negative relationship with average latitude (Figure 5),

which is congruous with the latitudinal diversity gradient observed for diverse macroorganisms

(Hillebrand, 2004). There is conflicting evidence for latitudinal diversity gradients among

microorganisms. Evidence for microbial latitudinal diversity gradients comes from marine

systems (Fuhrman *et al*., 2008; Sul *et al*., 2013; Swan *et al*., 2013), with contrary evidence

obtained in terrestrial systems (Neufeld and Mohn, 2005; Chu *et al*., 2010). However, most

analyses of terrestrial bacterial biogeography are derived from analyses of SSU rRNA genes, and

we show that analyses of SSU rRNA genes lack the sensitivity needed to detect the

biogeographic patterns that we observe for *Streptomyces*.

Several hypotheses have been advanced to explain the formation of latitudinal diversity gradients (Wiens and Donoghue, 2004; Mittelbach *et al.*, 2007). Ecological hypotheses posit that factors such as carrying capacity, productivity, and niche availability vary across latitude and that these factors impose constraints on biodiversity (Currie *et al.*, 2004; Mouchet *et al.*, 2015). Evolutionary hypotheses invoke the positive relationship between temperature and the kinetics of metabolism to predict that evolutionary rates and cladogenesis correspond with temperature (Allen *et al.*, 2002). Historical hypotheses propose that the latitudinal diversity gradient is the product of historical geological, ecological, or demographic events that have influenced dispersal and diversification (Wiens and Donoghue, 2004; Stevens, 2006). For example, the influence of Pleistocene glacial events on the biogeography of diverse species of terrestrial and aquatic plants and animals is well documented (Soltis *et al.*, 1997; Bernatch and Wilson, 1998; Conroy and Cook, 2000; Milá *et al.*, 2006; Maggs *et al.*, 2008; Wilson and Veraguth, 2010). There is also evidence that Pleistocene glaciation events have impacted both microbial communities (Eisenlord *et al.*, 2012) and populations (Kenefic *et al.*, 2009; Mikheyev *et al.*, 2008).

The biogeography of our *Streptomyces* phylogroups is consistent with historical demography and dispersal limitation resulting in a contemporary latitudinal diversity gradient. Eisenlord *et al.* (2012) also found that dispersal limitations shape biogeography of Actinobacteria across a glacial retreat transect in Michigan. However, Fuhrman *et al.* (2008) gives strong support for the hypothesis that the increased kinetics of metabolism at higher temperatures result in greater bacterial species richness in marine communities at tropical latitudes. Since latitude is correlated with temperature, we cannot dismiss the influence of kinetics on driving the diversity gradient in

terrestrial *Streptomyces*. Furthermore, it is possible that unappreciated ecological variables, such

as the species composition of perennial grass communities, could shape the diversity gradient.

However, phylogroup MAN125 from the Pacific Northwest (Table 2) and *S. pratensis* from the

Eastern United States (Doroghazi and Buckley, 2010 and 2014) each have populations in linkage

equilibrium though occupying a large geographic range over which gene flow is limited (Figure

3, Figure S4). The most parsimonious explanation for these conflicting results is that both taxa

have independently experienced a recent demographic range expansion and that insufficient

polymorphisms have accumulated since this expansion to allow for detection of disequilibrium

due to isolation by distance. However, it is likely that latitudinal gradients are population specific

and can arise from a combination of ecological, evolutionary, and historical processes.

The genetic consequences of glaciation events are described in depth by Hewitt (1996, 2000, and

2004), and the population structure we observe in our *Streptomyces* phylogroups is consistent

with post-glacial range expansion. One of the expectations of post-glacial expansion is "southern

richness versus northern purity" (Hewitt, 2004). This is evident in the negative correlation we

observe between latitude and the nucleotide diversity of phylogroups (Figure 5). Williams *et al*.

(1998) justifies 40°N latitude as approximating late Pleistocene glacial and non-glacial regions

with respect to species distributions in North America. Hence, the latitudinal delineation of allele

distributions for *Streptomyces* phylogroups roughly corresponds to the extent of ice coverage

during the late Pleistocene (Figure 3), which suggests historical population expansion from lower

to higher latitudes.

Haplotype distributions of phylogroups MAN125, MAN196, WA1063, and MS152 are

consistent with allopatric diversification resulting from dispersal limitation (Figure 4). Haplotype

nested clade analysis (NCA) predicts that historical dispersal events and range expansion across

northern regions resulted in limits on dispersal during intermediate timescales allowing genetic

drift to create the phylogroup population structures observed today. There is moderate criticism

of NCA (Knowles and Maddison, 2002; Nielsen and Beumont, 2009) due to the subjective nature

of inferring historical processes and the wide potential for stochastic processes creating similar

patterns of biogeography. Yet these tools can provide useful hypotheses. Northern phylogroups

MAN125 and MAN196 share a common ancestor with southern phylogroups MS200 and F34

(Figure 1). The contemporary population structure of MAN125 and MAN196 is consistent with a

historical range expansion from a common ancestor shared by both clades (Figure 1, Figure 4).

Further population structure within each phylogroup likely resulted from barriers to gene flow

and historical dispersal events across the Pacific Northwest for MAN125 (Figure 4A) and

between Maine and the Pacific Northwest for MAN196 (Figure 4B). Analysis of haplotype

distribution within WA1063 and MS152 is also consistent with diversification of populations as a

result of gene flow limitation between the Midwest and Pacific Northwest for WA1063 (Figure

4D) and between the Southeast and West for MS152 (Figure 4C).

Phylogroups WA1063 and MS152 share a recent common ancestor (Figure 1, Figure S1) and

even share two identical alleles at the *atpD* locus (Figure 3). These phylogroups have distinct,

non-overlapping geographic ranges with WA1063 found in higher latitudes and MS152 in lower

latitudes (Figure 3). WA1063 and MS152 have 0.0205 net nucleotide substitutions per site across

MLSA loci. We evaluate the possible time range for divergence between WA1063 and MS152

by extrapolating very roughly from the nucleotide substitution rate ($\mu = 4.5 \times 10^{-9}$) and generation

time (100-300 generations per year) for *E. coli* (Ochman *et al*., 1999), since corresponding

values are not available for *Streptomyces* or their relatives. Based upon these gross

approximations, we would estimate that WA1063 and MS152 diverged 15,000-50,000 years ago,

corresponding to events in the late Pleistocene (Clayton *et al*., 2006). Hence, it is likely that the

identical *atpD* alleles found in both WA1063 and MS152 were inherited without modification

from an ancestral population that both groups share (Figure S3B). If this is the case it suggests

that the divergence of phylogroup WA1063 was driven by range expansion into higher latitudes

and subsequent isolation by distance from the ancestral population it shares with phylogroup

MS152.

Through population genetic analysis of six *Streptomyces* phylogroups, we find evidence for

dispersal limitation associated with geographically explicit patterns of gene flow which manifest

in a latitudinal gradient of nucleotide diversity. Furthermore, these data support the hypothesis

that historical demographic processes influence the contemporary biogeography of *Streptomyces*.

Due to their spore forming capabilities and potential for long range dispersal, *Streptomyces* are

an ideal system for assessing limits on gene flow among terrestrial bacteria. We cannot yet

determine whether the cause of dispersal limitation is due to limits on spore mobility or due to

adaptive constraints as proposed by the hypothesis of phylogenetic niche conservatism. A better

understanding of *Streptomyces* biogeography and the evolutionary processes driving population

structure may ultimately assist in the discovery of novel genetic diversity and possibly novel

antibiotics within this genus.

Acknowledgements

This material is based upon work supported by the National Science Foundation under Grants

No. DEB-1050475 and DEB-1456821. We would like to thank Ashley Campbell for her help in

generating MLSA sequences.

References

Allen AP, Brown JH, Gillooly JF. (2002). Global biodiversity, biochemical kinetics, and the

energetic-equivalence rule. *Science* **297**: 1545–1548.

Andam CP, Choudoir MJ, Nguygen AVT, Park HS, Buckley DH. (2015). Contributions of

ancestral inter-species recombination to the genetic diversity of extant *Streptomyces*

lineages. *In Press.*

Andam CP, Doroghazi JR, Campbell AN, Kelly PJ, Choudoir MJ, Buckley DH. (2016). A

latitudinal diversity gradient in terrestrial bacteria of the genus *Streptomyces*. *Personal*

communication.

Antony-Babu S, Stach JEM, Goodfellow M. (2008). Genetic and phenotypic evidence for

Streptomyces griseus ecovars isolated from a beach and dune sand system. *Antonie Van*

Leeuwenhoek **94**: 63–74.

Bell T. (2010). Experimental tests of the bacterial distance-decay relationship. *ISME J* **4**: 1357–

1365.

Bernatchez L, Wilson CC. (1998). Comparative phylogeography of Nearctic and Palearctic

fishes. *Mol Ecol* **7**: 431–452.

Bissett A, Richardson AE, Baker G, Wakelin S, Thrall PH. (2010). Life history determines

biogeographical patterns of soil bacterial communities over multiple spatial scales. *Mol*

Ecol **19**: 4315–4327.

Boucher Y, Cordero OX, Takemura A, Hunt DE, Schliep K, Bapteste E, *et al.* (2011). Local

mobile gene pools rapidly cross species boundaries to create endemicity within global

Vibrio cholerae populations. *mBio* **2**: e00335–10.

Bruen TC, Philippe H, Bryant D. (2005). A simple and robust statistical test for detecting the

presence of recombination. *Genetics* **172**: 2665–2681.

Cadillo-Quiroz H, Didelot X, Held NL, Herrera A, Darling A, Reno ML, *et al.* (2012). Patterns

of gene flow define species of thermophilic Archaea. *PLoS Biol* **10**: e1001265.

Cho J, Tiedje JM. (2000). Biogeography and degree of endemicity of fluorescent *Pseudomonas*

strains in soil. *Appl Environ Microbiol* **66**: 5448-5456.

Choudoir MJ, Campbell AN, Buckley DH. (2012). Grappling with Proteus: population level

approaches to understanding microbial diversity. *Front Microbiol* **3**: 336.

Chu H, Fierer N, Lauber CL, Caporaso JG, Knight R, Grogan P. (2010). Soil bacterial diversity

in the Arctic is not fundamentally different from that found in other biomes. *Environ*

Microbiol **12**: 2998–3006.

Clayton L, Attig JW, Mickelson DM, Johnson MD, Syverson KM. (2006). Glaciation of

Wisconsin.

Clement M, Posada D, Crandall KA. (2000). TCS: A computer program to estimate gene

genealogies. *Mol Ecol* **9**: 1657–1659.

Coleman ML, Chisholm SW. (2010). Ecosystem-specific selection pressures revealed through

comparative population genomics. *Proc Natl Acad Sci U S A* **107**: 18634–18639.

Conroy CJ, Cook JA. (2000). Phylogeography of a post-glacial colonizer: *Microtus longicaudus*

(Rodentia: Muridae). *Mol Ecol* **9**: 165–175.

Currie DJ, Mittelbach GG, Cornell HV, Field R, Guégan JF, Hawkins BA*, et al.* (2004).

Predictions and tests of climate-based hypotheses of broad-scale variation in taxonomic

richness. *Ecol Lett* **7**: 1121–1134.

Davelos AL, Xiao K, Samac DA, Martin AP, Kinkel LL. (2004a). Spatial variation in

Streptomyces genetic composition and diversity in a prairie soil. *Microb Ecol* **48**: 601–

612.

Davelos AL, Kinkel LL, Samac DA. (2004b). Spatial variation in frequency and intensity of

antibiotic interactions among *Streptomycetes* from prairie soil. *Appl Environ Microbiol*

70: 1051–1058.

Doroghazi JR, Buckley DH. (2014). Intraspecies comparison of *Streptomyces pratensis* genomes

reveals high levels of recombination and gene conservation between strains of disparate

geographic origin. *BMC Genomics* **15**: 970.

Doroghazi JR, Buckley DH. (2010). Widespread homologous recombination within and between

Streptomyces species. *ISME J* **4**: 1136–1143.

Earl DA, vonHoldt BM. (2012). STRUCTURE HARVESTER: a website and program for

visualizing STRUCTURE output and implementing the Evanno method. *Conserv Genet*

Resour **4**: 359–361.

Edgar RC. (2004). MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acid Res* **32**: 1792–1797.

Eisenlord SD, Zak DR, Upchurch RA. (2012). Dispersal limitation and the assembly of soil

Actinobacteria communities in a long-term chronosequence. *Ecol Evol* **2**: 538–549.

El-Nakeeb MA, Lechevalier HA. (1963). Selective isolation of aerobic Actinomycetes. *Appl*

Microbiol 11: 75–77.

Evanno G, Regnaut S, Goudet J. (2005). Detecting the number of clusters of individuals using

the software STRUCTURE: a simulation study. *Mol Ecol* **14**: 2611–2620.

Fierer N, Carney KM, Horner-Devine MC, Megonigal JP. (2009). The biogeography of

ammonia-oxidizing bacterial communities in soil. *Microb Ecol* **58**: 435–445.

Fierer N, Jackson RB. (2006). The diversity and biogeography of soil bacterial communities.

Proc Natl Acad Sci U S A **103**: 626–631.

Finlay BJ. (2002). Global dispersal of free-living microbial eukaryote species. *Science* **296**:

1061–1063.

Finlay BJ, Fenchel T. (2004). Cosmopolitan metapopulations of free-living microbial eukaryotes.

Protist **155**: 237–244.

Fuhrman JA, Steele JA, Hewson I, Schwalbach MS, Brown M V, Green JL, *et al.* (2008). A

latitudinal diversity gradient in planktonic marine bacteria. *Proc Natl Acad Sci U S A*

105: 7774–7778.

Gibbons SM, Caporaso JG, Pirrung M, Field D, Knight R, Gilbert JA. (2013). Evidence for a

persistent microbial seed bank throughout the global ocean. *Proc Natl Acad Sci U S A*

110: 4651–4655.

Gilbert JA, Steele JA, Caporaso JG, Steinbrück L, Reeder J, Temperton B, *et al.* (2012).

Defining seasonal marine microbial community dynamics. *ISME J* **6**: 298–308.

Goslee SC, Urban DL. (2007). The ecodist package for dissimilarity-based analysis of ecological

data. *J Stat Softw* **22**: 1–19.

Green J, Bohannan BJM. (2006). Spatial scaling of microbial biodiversity. *Trends Ecol Evol* **21**:

501–507.

Guo Y, Zheng W, Rong X, Huang Y. (2008). A multilocus phylogeny of the *Streptomyces*

griseus 16S rRNA gene clade: use of multilocus sequence analysis for streptomycete

systematics. *Int J Syst Evol Microbiol* **58**: 149–159.

Hanson CA, Fuhrman JA, Horner-Devine MC, Martiny JBH. (2012). Beyond biogeographic

patterns: processes shaping the microbial landscape. *Nat Rev Microbiol* **10**: 497–506.

Hatosy SM, Martiny JBH, Sachdeva R, Steele J, Fuhrman JA, Martiny AC. (2013). Beta

diversity of marine bacteria depends on temporal scale. *Ecology* **94**: 1898–1904.

Haubold B, Hudson RR. (2000). LIAN 3.0: detecting linkage disequilibrium in multilocus data.

Bioinformatics **16**: 847–848.

Hewitt GM. (1996). Some genetic consequences of ice ages, and their role in divergence and

speciation. *Biol J Linnean Soc* **58**: 247–276.

Hewitt G. (2000). The genetic legacy of the Quaternary ice ages. *Nature* **405**: 907–913.

Hewitt GM. (2004). Genetic consequences of climatic oscillations in the Quaternary. *Philos*

Trans R Soc Lond B Biol Sci **359**: 183–195.

Hillebrand H. (2014). On the generality of the latitudinal diversity gradient. *Amer Nat* **163**: 192–

211.

Huang X, Madan A. (1999). CAP3: a DNA sequence assembly program. *Genome Res* **9**: 868–

677.

Hunt DE, David LA, Gevers D, Preheim SP, Alm EJ, Polz MF. (2008). Resource partitioning

and sympatric differentiation among closely related bacterioplankton. *Science* **320**: 1081–

1085.

Jakobsson M, Rosenberg NA. (2007). CLUMPP: a cluster matching and permutation program

for dealing with label switching and multimodality in analysis of population structure.

Bioinformatics **23**: 1801–1806.

Jaspers E, Overmann J. (2004). Ecological significance of microdiversity: identical 16S rRNA

gene sequences can be found in bacteria with highly divergent genomes and

ecophysiologies. *Appl Environ Microbiol* **70**: 4831–4839.

Jolley KA, Chan MS, Maiden MCJ. (2004). mlstdbNet - distributed multi-locus sequence typing

(MLST) databases. *BMC Bioinformatics* **5**: 86.

Kâmpfer P. (2006). The family *Streptomycetaceae* Part I: Taxonomy. In: The Prokaryotes,

Volume 3. Dworkin, M, Falkow, S, Rosenberg, E, Schliefer, K, & Stackebrandt, E (eds).

Springer: New York, pp pp. 538–604.

Kenefic LJ, Pearson T, Okinaka RT, Schupp JM, Wagner DM, Hoffmaster AR, *et al.* (2009).

Pre-Columbian origins for North American anthrax. *PLoS One* **4**: e4813.

Kieser T, Bibb M, Buttner M, Charter K, Hopwood D. (2000). Practical *Streptomyces* Genetics.

John Innes Foundation: Norwich, UK.

Knowles L, Maddison W. (2002). Statistical phylogeography. *Mol Ecol* **11**: 2623–2635.

Labeda DP. (2011). Multilocus sequence analysis of phytopathogenic species of the genus

Streptomyces. *Int J Syst Evol Microbiol* **61**: 2525–2531.

Lauber CL, Hamady M, Knight R, Fierer N. (2009). Pyrosequencing-based assessment of soil

pH as a predictor of soil bacterial community structure at the continental scale. *Appl*

Environ Microbiol **75**: 5111–5120.

Librado P, Rozas J. (2009). DnaSP v5: a software for comprehensive analysis of DNA

polymorphism data. *Bioinformatics* **25**: 1451–1452.

Loria R, Kers J, Joshi M. (2006). Evolution of plant pathogenicity in *Streptomyces*. *Annu Rev*

Phytopathol **44**: 469–487.

Lozupone CA, Knight R. (2007). Global patterns in bacterial diversity. *Proc Natl Acad Sci U S A*

104: 11436–11440.

Maggs CA, Castilho R, Foltz D, Henzler C, Jolly MT, Kelly J, *et al.* (2008). Evaluating

signatures of glacial refugia for North Atlantic benthic marine taxa. *Ecology* **89**: S108–

S122.

Mantel N. (1967). The detection of disease clustering and a generalized regression approach.

Cancer Research **27**: 209–220.

Martiny JBH, Bohannan BJM, Brown JH, Colwell RK, Fuhrman JA, Green JL, *et al.* (2006).

Microbial biogeography: putting microorganisms on the map. *Nat Rev Microbiol* **4**: 102–

112.

Martiny JBH, Eisen JA, Penn K, Allison SD, Horner-Devine MC. (2011). Drivers of bacterial β -

diversity depend on spatial scale. *Proc Natl Acad Sci U S A* **108**: 7850–7854.

McCarthy AJ, Williams ST. (1992). Actinomycetes as agents of biodegradation in the

environment - a review. *Gene* **115**: 189–192.

McVean G, Awadalla P, Fearnhead P. (2002). A coalescent-based method for detecting and

estimating recombination from gene sequences. *Genetics* **160**: 1231–1241.

Mikheyev AS, Vo T, Mueller UG. (2008). Phylogeography of post-Pleistocene population

expansion in a fungus-gardening ant and its microbial mutualists. *Mol Ecol* **17**: 4480–

4488.

Mila B, Smith T, Wayne R. (2006). Postglacial population expansion drives the evolution of

long-distance migration in a songbird. *Evolution* **60**: 2403–2409.

Miller SR, Strong AL, Jones KL, Ungerer MC. (2009). Bar-coded pyrosequencing reveals shared

bacterial community properties along the temperature gradients of two alkaline hot

springs in Yellowstone National Park. *Appl Environ Microbiol* **75**: 4565–4572.

Mittelbach GG, Schemske DW, Cornell H V., Allen AP, Brown JM, Bush MB, *et al.* (2007).

Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography.

Ecol Lett **10**: 315–331.

Mouchet M, Levers C, Zupan L, Kuemmerle T, Plutzar C, Erb K, *et al.* (2015). Testing the

effectiveness of environmental variables to explain European terrestrial vertebrate species

richness across biogeographical scales. *PLoS One* **10**: e0131924.

Neufeld JD, Mohn WW. (2005). Unexpectedly high bacterial diversity in arctic tundra relative to

boreal forest soils, revealed by serial analysis of ribosomal sequence tags. *Appl Environ*

Microbiol **71**: 5710–5718.

Nielsen R, Beaumont MA. (2009). Statistical inferences in phylogeography. *Mol Ecol* **18**: 1034–

1047.

Ochman H, Wilson AC. (1987). Evolution in cacteria: evidence for a universal substitution rate

in cellular genomes. *J Mol Evol* **26**: 74–86.

Ochman H, Elwyn S, Moran NA. (1999). Calibrating bacterial evolution. *Proc Natl Acad Sci U S*

A **96**: 12638–12643.

Ottow JCG. (2013). Rose bengal as a selective aid in the isolation of fungi and Actinomycetes

from natural sources. *Mycologia* **64**: 304–315.

Panchal M. (2006). The automation of Nested Clade Phylogeographic Analysis. *Bioinformatics*

23: 509–510.

Papke RT, Ramsing NB, Bateson MM, Ward DM. (2003). Geographical isolation in hot spring

640   cyanobacteria. *Environ Microbiol* **5**: 650–659.

641   Pattengale ND, Alipour M, Bininda-Emonds ORP, Moret BME, Stamatakis A. (2010). How

642   many bootstrap replicates are necessary? *J Comput Biol* **17**: 337–354.

643   Pearson T, Giffard P, Beckstrom-Sternberg S, Auerbach R, Hornstra H, Tuanyok A*, et al*.

644   (2009). Phylogeographic reconstruction of a bacterial species with high levels of lateral

645   gene transfer. *BMC Biol* **7**: 78.

646   Pommier T, Canback B, Lundberg P, Hagstrom A, Tunlid A. (2009). RAMI: a tool for

647   identification and characterization of phylogenetic clusters in microbial communities.

648   *Bioinformatics* **25**: 736–742.

649   Posada D, Crandall KA, Templeton AR. (2000). GeoDis: A program for the cladistic nested

650   analysis of the geographical distribution of genetic haplotypes. *Mol Ecol* **9**: 487–488.

651   Pritchard JK, Stephens M, Donnelly P. (2000). Inference of population structure using multilocus

652   genotype data. *Genetics* **155**: 945–959.

653   Rong X, Huang Y. (2010). Taxonomic evaluation of the *Streptomyces griseus* clade using

654   multilocus sequence analysis and DNA-DNA hybridization, with proposal to combine 29

655   species and three subspecies as 11 genomic species. *Int J Syst Evol Microbiol* **60**: 696–

656   703.

657   Rong X, Huang Y. (2012). Taxonomic evaluation of the *Streptomyces hygroscopicus* clade using

658   multilocus sequence analysis and DNA-DNA hybridization, validating the MLSA

659   scheme for systematics of the whole genus. *Syst Appl Microbiol* **35**: 7–18.

Rousk J, Bååth E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, *et al*. (2010). Soil

bacterial and fungal communities across a pH gradient in an arable soil. *ISME J* **4**: 1340–

1351.

Schliep KP. (2011). phangorn: phylogenetic analysis in R. *Bioinformatics* **27**: 592–593.

Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, *et al*. (2009).

Introducing mothur: open-source, platform-independent, community-supported software

for describing and comparing microbial communities. *Appl Environ Microbiol* **75**: 7537–

7541.

Shapiro BJ, Friedman J, Cordero OX, Preheim SP, Timberlake SC, Szabó G, *et al*. (2012).

Population genomics of early events in the ecological differentiation of bacteria. *Science*

336: 48–51.

Shimodaira H, Hasegawa M. (1999). Multiple comparisons of log-likelihoods with applications

to phylogenetic inference. *Mol Biol Evol* **16**: 1114–1116.

Smouse PE, Long JC, Sokal RR. (1986). Multiple regression and correlation extensions of the

Mantel test of matrix correspondence. *Syst Zool* **35**: 627–632.

Soltis D, Gitzendanner M, Strenge D, Soltis P. (1997). Chloroplast DNA intraspecific

phylogeography of plants from the Pacific Northwest of North America. *Plant Syst Evol*

206: 353–373.

Stamatakis A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with

thousands of taxa and mixed models. *Bioinformatics* **22**: 2688–2690.

Stevens RD. (2006). Historical processes enhance patterns of diversity along latitudinal

gradients. *Proc R Soc B Biol Sci* **273**: 2283–2289.

Sul WJ, Oliver TA, Ducklow HW, Amaral-zettler LA, Sogin ML. (2012). Marine bacteria

exhibit a bipolar distribution. *Proc Natl Acad Sci U S A* **110**: 2342–2347.

Swan BK, Tupper B, Sczyrba A, Lauro FM, Martinez-Garcia M, González JM, *et al*. (2013).

Prevalent genome streamlining and latitudinal divergence of planktonic bacteria in the

surface ocean. *Proc Natl Acad Sci U S A* **110**: 11463–11468.

Takasuka TE, Book AJ, Lewin GR, Currie CR, Fox BG. (2013). Aerobic deconstruction of

cellulosic biomass by an insect-associated *Streptomyces*. *Sci Rep* **3**: 1–10.

Tavaré S. (1986). Some probabilistic and statistical problems in the analysis of DNA sequences.

Am Math Soc Lect Math Life Sci **17**: 57–86.

Telford RJ, Vandvik V, Birks HJB. (2006). Dispersal limitations matter for microbial

morphospecies. *Science* **312**: 1015.

Templeton AR, Boerwinkle E, Sing CF. (1987). A cladistic analysis of phenotypic associations

with haplotypes inferred from restriction endonuclease mapping. i. basic theory and an

analysis of alcohol dehydrogenase activity in Drosophila. *Genetics* **117**: 343–351.

Templeton AR, Routman E, Phillips CA. (1995). Separating population structure from

population history: a cladistic analysis of the geographical distribution of mitochondrial

DNA haplotypes in the tiger salamander, *Ambystoma tigrinum*. *Genetics* **140**: 767–782.

Vos M, Velicer GJ. (2008). Isolation by distance in the spore-forming soil bacterium

Myxococcus xanthus. *Curr Biol* **18**: 386–391.

Wallace A. (1878). Tropical nature, and other essays. Macmillan and Co.: London.

Watve M, Tickoo R, Jog M, Bhole B. (2001). How many antibiotics are produced by the genus

Streptomyces? *Arch Microbiol* **176**: 386–390.

Wawrik B, Kutliev D, Abdivasievna UA, Kukor JJ, Zylstra GJ, Kerkhof L. (2007).

Biogeography of Actinomycete communities and type ii polyketide synthase genes in

soils collected in New Jersey and Central Asia. *Appl Environ Microbiol* **73**: 2982–2989.

Welch RA, Burland V, Plunkett G, Redford P, Roesch P, Rasko D, *et al*. (2002). Extensive

mosaic structure revealed by the complete genome sequence of uropathogenic

Escherichia coli. *Proc Natl Acad Sci U S A* **99**: 17020–17024.

Whitaker RJ, Grogan DW, Taylor JW. (2003). Geographic barriers isolate endemic populations

of hyperthermophilic archaea. *Science* **301**: 976–978.

Wiens JJ, Donoghue MJ. (2004). Historical biogeography, ecology and species richness. *Trends*

Ecol Evol **19**: 639–644.

Williams D, Dunkerley D, DeDecker P, Kershaw P, Chappell M. (1998). Quaternary

environments. Arnold: London.

Wilson AB, Veraguth IE. (2010). The impact of Pleistocene glaciation across the range of a

widespread European coastal species. *Mol Ecol* **19**: 4535–53.

Figure 1. *Streptomyces* phylogroups have different geographic distributions defined by latitude with evidence for subpopulation structure and admixture. The colored bars in the outer ring indicate genetic contributions from different ancestral populations as inferred by Structure analysis. The shading of the inner ring indicates the latitude from which each strain was isolated according to the scale provided. The isolation site for each strain can be determined by isolate names as indicated in Table 1. The tree was constructed from concatenated MLSA loci nucleotide sequences using maximum likelihood with a GTRGAMMA evolution model. Scale bar represents nucleotide substitutions per site. The root was defined by *Mycobacterium smegmatis*. Nodes with bootstrap confidences > 80 are indicated with gray circles, and precise bootstrap values are found in Figure S3A.

Figure 2. Dissimilarity in SSU rRNA genes is less than 3% across all six phylogroups and ranges from 0-0.6% for representatives within each phylogroup, revealing that this locus lacks sufficient sensitivity to examine the biogeographic distribution of *Streptomyces* phylogroups. The tree was constructed using maximum likelihood with a GTRGAMMA evolution model using partial SSU rRNA gene sequences from three representatives for each phylogroup. Scale

bar represents nucleotide substitutions per site. Branches with bootstrap support values greater

than 75% are labeled, and the outgroup is *Mycobacterium smegmatis*. Strains are colored to

reveal phylogroup membership as indicated in the color scale provided.

Figure 3. Identical MLSA alleles are frequently shared between sites. Circles depict sample sites

and are labeled according to site code (Table 1). The relative abundance of each phylogroup at

each site is indicated by color according to the legend, with raw counts provided in Table 1. Solid

colored lines represent identical alleles shared by phylogroup members across sites. Dashed

multicolored lines depict identical *atpD* alleles shared by strains in phylogroups WA1063 and

MS152 (Figure S3B).

Figure 4. Haplotype networks are depicted for phylogroups MAN125 (4A), MAN196 (4B), MS152 (4C), WA1063 (4D). Circles represent unique MLSA haplotypes whose radius is proportional to the number of isolates sharing that haplotype, and colors correspond to strain

ancestry and subpopulation affiliation as determined by Structure analysis and defined in Figure

1. Haplotypes are labeled with a letter code referring to sample site location as indicated in Table

1. Black circles represent un-sampled, inferred haplotypes with each circle designating a single

nucleotide polymorphism. The length of edges between nodes is uninformative. Dashed

rectangles encompass clades that have significant phylogeographic inferences from nested clade

analysis, as described in methods. Roman numerals correspond to the following inferences: I.

Allopatric fragmentation; II. Long distance colonization and/or past fragmentation; III.

Restricted gene flow with isolation by distance; IV. Restricted gene flow but with some long-

distance gene flow over intermediate ranges not occupied by the species; or past gene flow

followed by extinction of intermediate populations; V. Contiguous range expansion.

Figure 5. There is a significant inverse correlation between phylogroup average latitude and

nucleotide diversity. Average latitude was determined using a weighted average based on the

total number of isolates per site as indicated in Table 1. Nucleotide diversity was calculated using

concatenated MLSA loci and is expressed per site.

Table 1. The six targeted phylogroups were observed in 13 sites. A total of 755 *Streptomyces* strains were isolated from these sites, and 308 of these strains were found to represent the six targeted phylogroups. The numbers of isolates per site that belonged to each of our target phylogroups is indicated. Isolate names begin with a letter code referring to the isolation site.

Site	Code	Latitude	Longitude	Isolates /Site	Isolates from *Streptomyces* sp. phylogroup					
					MAN 125	MAN 196	WA 1063	MS 200	MS 152	F34
Manley Hot Springs, AK	man	63.87°N	-149.02°W	95	51	26	1			
Denali Hwy, AK	den	63.22°N	-147.68°W	40	40					
Bothell, WA	wa	47.73°N	-122.24°W	105	21	5	21			
Astoria, OR	or	46.18°N	-123.85°W	79		8	7			
Kennebunk, ME	me	43.4°N	-70.54°W	85		13				
Sun Prairie, WI	sun	43.17°N	-89.24°W	65				3		
Brookfield, WI	b	43.06°N	-88.13°W	22	3		5			
Palo Alto, CA	st	37.43°N	-122.17°W	34					2	1
Greensboro, NC	gb	36.09°N	-79.89°W	9		1			2	
Troy (Uwharrie), NC	uw	35.71°N	-79.88°W	19		4		7		
Starkville, MS	ms	33.46°N	-88.8°W	90				11	8	2
Austin, TX	t	30.2°N	-97.77°W	42						6
Fort Pierce, FL	f	27.54°N	-80.35°W	70				7	5	48
Total				755	115	57	34	28	17	57

763 **Table 2.** MLSA was performed on a total of 17-47 strains from each phylogroup. Summary

764 statistics for concatenated MLSA nucleotide sequences were determined as described in

765 methods. Phylogroup average latitude (Ave Lat) was determined using the total number of

766 isolates per phylogroup provided in Table 1. The standard index of association is zero for a

767 population in linkage equilibrium.

Phylo-group	Ave Lat	n	Haplo-types	Min ANI	S	π	Tajima's D	θ_W	ρ	ρ/θ_W	I_A	Φ_W
MAN125	60.15°N	45	25	99.5	23	0.0026	0.53	0.0023	0.0094	4.18	0.09	2.95E-01
MAN196	52.84°N	47	26	98.5	65	0.0075	0.67	0.0063	0.00043	0.068	0.49	4.24E-02*
WA1063	47.20°N	19	13	98.1	63	0.0052	-1.35	0.0077	0	0	0.43	4.33E-03
MS200	33.58°N	27	21	98.4	93	0.0088	-0.58	0.0103	0.0017	0.17	0.39	1.33E-03**
MS152	32.50°N	17	15	97.6	96	0.0107	-0.51	0.012	0.0013	0.11	0.35	1.078E-01**
F34	28.20°N	36	22	97.7	93	0.0109	0.52	0.0096	0.0011	0.11	0.46	1.79E-01**

Ave Lat: average latitude of phylogroup members
n: number of isolates subsampled for MLSA
Min ANI: minimum percent pairwise average nucleotide identity (ANI) within phylogroup members
S: segregating sites
π: per site nucleotide diversity
θ_W: per site Watterson's theta ($2N_e\mu$)
ρ: per site rate of recombination/gene conversion ($2N_er/2$)
ρ/θ_W: relative rate of recombination to mutation
I_A: standard index of association
Φ_W: pairwise homoplasy index (Phi) statistic with *p*-value < 0.05*, 0.01** under the null hypothesis of no recombination