

Genome-wide analysis of microRNA signature in lung adenocarcinoma with EGFR exon 19 deletion

Lixia Ju^{1*}, Mingquan Han¹, Xuefei Li², Chao Zhao²

¹ Department of Integrative Medicine, Shanghai Pulmonary Hospital, Tongji University School of Medicine, Tongji University, Shanghai, People's Republic of China.

² Department of Lung Cancer and Immunology, Shanghai Pulmonary Hospital, Tongji University Medical School Cancer Institute, Tongji University, Shanghai, People's Republic of China.

***Correspondence to:** Lixia Ju, MD, PhD, Department of Integrated Medicine, Shanghai Pulmonary Hospital, Tongji University School of Medicine, No. 507 Zheng Min Road, Shanghai 200433, People's Republic of China. E-mail: jvlixia@126.com.

Disclosure of Potential Conflicts of Interest: No potential conflicts of Interest were disclosed.

Abstract

Purpose: The findings of EGFR mutations and the development of targeted therapies have significantly improved the overall survival of lung cancer patients. Still, the prognosis remains poor, so we need to know more about the genetic alterations in lung cancer. MicroRNAs are dysregulated in lung cancer, and microRNAs can regulate EGFR. So it is very important to predict the candidate microRNAs that target mutated EGFR and to investigate the role of these candidate microRNAs in lung cancer.

Materials and methods: In this study, we investigated the difference of microRNAs expression between lung adenocarcinoma cell lines with EGFR exon 19 deletion (H1650 and PC9) and its wild-type (H1299 and A549) using the Phalanx Human Whole Genome Microarray. Then the expression of individual microRNAs was validated by qRT-PCR assays. Moreover, we have detected the microRNAs expression in serum of lung adenocarcinoma patients with EGFR exon 19 deletion and wild-type.

Results: The expression of 1,732 microRNAs was evaluated, and we found that microRNAs expression was different between these two groups. Hsa-miR-141-3p, hsa-miR-200c-3p, hsa-miR-203, hsa-miR-3182, hsa-miR-934 were up-regulated and hsa-miR-3196 was down-regulated in the EGFR exon 19 deletion group compared with wild-type group. The detection of circulating microRNAs showed that miR-3196 was down-regulated in lung adenocarcinoma patients with EGFR exon 19 deletion compared with wild-type.

Conclusions: It is suggested that microRNAs associate with EGFR exon 19 deletion and miR-3196 can be further explored for potential predictor and targeted biomarker when it is difficult to get the tumors.

Keywords: *Genome-wide, microRNA, EGFR, exon 19 deletion, non-small cell lung cancer*

Introduction

Lung cancer, one of the most common malignant tumor¹, led to the most cancer death in the world. The prevalence and mortality of the lung cancer is still rising. In all new diagnosed lung cancer, non-small cell lung cancer (NSCLC) accounts for approximately 85%, which varies both in molecular and clinical presentation. Despite years of research, the 5-year survival is only about 18%.² The discovery of key driver mutations has led to a more personalized approach in the treatment of advanced lung cancer. The activating mutations of EGFR implicate sensitivity to EGFR tyrosine kinase inhibitors. EGFR exon 19 deletion and L858R substitution in exon 21 have been extensively proved to be the sensitive mutations. However, the efficacy of EGFR-TKIs varies among different sensitive EGFR mutations. Several studies have reported that the advanced NSCLC patients with EGFR exon 19 deletion had a longer overall survival (OS) and/or progression-free survival (PFS) with the treatment of gefitinib or erlotinib compared with those with the L858R mutation.³⁻⁶

MicroRNAs are evolutionarily conserved, endogenous small non-coding RNAs with 18-25 nucleotides that have important functions in diverse biological processes, such as cell proliferation, differentiation, and apoptosis.⁷⁻⁹ Furthermore, microRNAs play an essential role in behaving either as oncogenes or tumor suppressor genes¹⁰. Increasing evidence indicates that dysregulation of specific microRNAs contributes to the development and progression of cancer, including lung cancer.¹¹⁻¹³ Moreover, microRNAs can regulate EGFR.¹⁴ So it is important to make it clear that the candidate microRNAs regulating EGFR exon 19 deletion because it might spark the design of novel therapeutics to combat the resistance to EGFR-TKIs or develop new targeted

therapy.

In this study, we conducted an explorative microRNAs expression study in two groups of lung adenocarcinoma cell lines, including EGFR exon 19 deletion group (H1650 and PC-9) and EGFR wild-type group (A549 and H1299), using microRNA microarrays. Our main focus was the different microRNAs in the two groups. The selected microRNAs were confirmed using qRT-PCR in the cell lines and then we measured 3 microRNAs expressed differently in cell lines in the serum of 14 non-smoking female lung adenocarcinoma patients with wt EGFR and 13 patients with EGFR 19 del. Accordingly we discussed the association of microRNAs with EGFR exon 19 deletion.

Material and Methods

Cell lines and cell culture

Human adenocarcinoma cell lines PC-9 and H1650 (harboring EGFR exon 19 deletion) and A549 and H1299 (harboring wild-type EGFR) were provided by Cancer Institute of Tongji University Medical School, China. All these cells were cultured at 37°C with 5% CO₂ in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS), 100 U/ml penicillin and 100 mg/ml streptomycin.

Patient enrollment and serum samples

All patients in the study were recruited from Shanghai Pulmonary Hospital, Tongji University Medical School, between January 2012 and June 2014. The patients were newly diagnosed and histologically confirmed primary lung adenocarcinoma. Patients

with the previous history of cancer , radiotherapy or chemotherapy were excluded.

The study was approved by an ethical review committee at Tongji University

Institutional Care and Use Committee. All blood serum samples were collected and

put into a liquid nitrogen tank for long-term storage until microRNAs extraction.

Total RNA isolation

Total RNA was extracted from cells using TRIZOL Reagent (Invitrogen, USA). The

RNA concentration and purity were accessed by OD260/OD280 (≥ 1.6) and

OD260/OD230 (≥ 1.0), and the RNA yield and quality were checked (RIN ≥ 5.0) using

Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA).

Human microRNAs OneArray®

Human microRNA OneArray® v3 (Phalanx Biotech Group, Taiwan) contains

triplicated 1,711 unique microRNA probes from Human (miRBase Release v17), each

printed in technical triplicate, and 189 experimental control probes.

Microarray analysis

Small RNA was pre-enriched by Nanoseplook (Pall Corporation, USA) from 2.5 μ g

total RNA samples and labeled with microRNAs ULSTM Labeling Kit (Kreatech

Diagnostics, The Netherlands). Labeled targets were hybridized to the Human

microRNA OneArray® v3 with OneArray® Hybridization System. After 16 hours

hybridization at 37 °C, non-specific binding targets were washed away by three

different washing steps (Wash 1 37 °C 5 mins ; Wash 2 37 °C, 5 mins 25 °C 5 mins; Wash

3 37 °C 5 mins), and the slides were dried by centrifugation and scanned by an

Axon 4000B scanner (Molecular Devices, Sunnyvale, CA, USA). Normalized spot intensities were transformed to gene expression log₂ ratios between the mutation and wild-type group. The spots with log₂ ratio ≥ 1 or log₂ ratio ≤ -1 and P-value < 0.05 were tested for further analysis.

Validation by qRT-PCR

Quantitative real time-PCR (qRT-PCR) was carried out on the 7900HT thermocycler (Applied Biosystems, Foster City, CA). U6 was used for internal controls. The data were managed using the Applied Biosystems software RQ. Relative expression was calculated using the comparative Ct method and obtaining the fold-change value ($2^{-\Delta\Delta C_t}$). Data analyses were performed via GraphPad Prism v6.00.

EGFR mutation analyses

Mutation analyses of EGFR exons 18-21 were performed on 27 of the tumor samples using the amplification-refractory mutation system assay (ARMS). Data analyses were performed by employing the LightCycler Adapt software (LightCycler 480 Software, v. 1.5).

Digital PCR

Digital PCR was performed in parallel for the measurement of microRNAs in the serially-diluted oligonucleotides. 30 μ L of the reaction mixture containing 15 μ L QuantStudio™ 3D Digital PCR Master Mix, 2X (Life technologies), 2 μ L of cDNA solution, and 1.5 μ L of TaqManR Assay, 20X (primer/probe mix) (Life technologies) and 11.5 μ L water. The droplets generated from each sample were transferred to a 96-well PCR plate (Eppendorf, Germany). PCR amplification was carried on a T100

thermal cycler (QuantStudio™ 3D Digital PCR System) at 96°C for 10 min, followed by 39 cycles 60°C for 2 min, then 98°C for 0.5 min. For Final extension, followed by 60°C for 2 min, Last, Storage is at 10°C for 100min.¹⁵

Results

Microarrays analysis of cell lines with EGFR exon 19 deletion compared with wild-type

To identify microRNAs that were differentially expressed between cell lines with EGFR exon 19 deletion and the wild-type adenocarcinoma cell lines, the expression profiles of microRNAs (1,711 microRNAs) were assessed using microRNA microarrays. For clustering analysis, 260 genes was selected to identify differentially expressed based on the threshold of fold change and p-value (Figure 1). Standard selection criteria to identify differentially expressed genes are established at \log_2 |Fold change| ≥ 0.8 and $P < 0.05$. The microRNAs differentially expressed between EGFR exon 19 deletion and wild-type are shown in Table 1, ($P < 0.05$). In total, let-7d-3p, miR-1307-5p, miR-141-3p, miR-200c-3p, miR-203, miR-3182, miR-4510, miR-934 were up-regulated in EGFR exon 19 deletion group compared with the wild-type group. On the contrary, miR-3196, miR-4450, miR-4649-5p were down-regulated.

Validation of the microarrays results using quantitative reverse transcriptase polymerase chain reaction (qRT-PCR)

The eleven microRNAs, differentially expressed in the two cell groups in microarray

analyses, were validated by qRT-PCR. As a result, six microRNAs were identified and differentially expressed between the two cell groups (see Table 2). Hsa-miR-141-3p, hsa-miR-200c-3p, hsa-miR-203, hsa-miR-3182, and hsa-miR-934 were up-regulated, while hsa-miR-3196 was down-regulated (Figure 2).

Pathway analysis

Using the microRNA target filter in TargetScan Release 6.2, we searched for genes involved in the signaling pathways that are experimentally observed or highly predicted to be regulated by the selected microRNAs. The six microRNAs that were differentially expressed between EGFR exon 19 deletion and EGFR wild-type cell lines were associated with 3181 mRNA targets. Two microRNAs (hsa-miR-200c-3p, hsa-miR-203) were experimentally confirmed to target EGFR, and be associated with EGFR-TKI resistance.¹⁶⁻¹⁷

Circulating microRNAs in relation to EGFR exon 19 deletion

This study was conducted on 27 participants stratified into EGFR exon 19 deletion group and wild-type group. Selected candidates were non-smoking lung adenocarcinoma patients. It is well known that the sensitivity of qPCR for the detection of the low copy genes is not so high, as it only can resolve ~1.5-fold changes of nucleic acids.¹⁸ Given that a portion of the cancer-related microRNAs is derived from primary tumor and could be ‘diluted’ in the normal microRNAs,¹⁹⁻²¹ the microRNAs presenting at low levels in serum could be undetectable by qPCR.

Droplet digital PCR is a direct method for quantitatively detecting nucleic acids.^{22,23} It depends on limit proportion of the PCR volume, in which a positive microreactions indicates the presence of a single molecule in a given reaction. The number of

positive reactions, together with Poisson's distribution, can be used to calculate the original target concentration in a straight and high-confidence measurement method.²⁴

In our study, digital PCR was used to detect three circulating microRNAs (hsa-miR-200c-3p, hsa-miR-203, hsa-miR-3196) of 27 patients, 13 patients with EGFR exon 19 deletion and 14 wild-type EGFR. In the results, we found that hsa-miR-3196 was down-regulated in the group of EGFR exon 19 deletion compared with wild-type, which was consistent with our results of cell study (Figure 3).

Discussion

In the present work , we found eleven microRNAs differentially expressed between EGFR exon 19 deletion cell lines and EGFR wild-type lung adenocarcinoma cell lines in the microarray analysis. Then the PCR study identified that hsa-miR-141-3p, hsa-miR-200c-3p, hsa-miR-203, hsa-miR-3182, and hsa-miR-934 were up-regulated, and hsa-miR-3196 was down-regulated in EGFR exon 19 deletion group compared with wild-type group. Moreover, the detection of circulating microRNAs found that miR-3196 was down-regulated in patients with EGFR exon 19 deletion compared with wild-type lung adenocarcinoma patients.

Presently, not much is known about the microRNAs expression in EGFR exon 19 deletion versus wild-type lung adenocarcinomas. Using microRNA microarrays, Dacic et al. analyzed six lung adenocarcinomas and found that miR-155 was up-regulated only in EGFR/KRAS-negative group, miR-25 was up-regulated only in EGFR-positive group and miR-495 was up-regulated only in KRAS-positive adenocarcinoma. In opposite, let-7g was down-regulated in all three groups, with more significant downregulation in EGFR/KRAS-negative adenocarcinoma.²⁵ Seike

et al. identified 12 microRNAs differentially expressed between 6 EGFR-mutated tumors and 22 wild-type tumors. Zhang et al. found that miR-122 were differentially expressed between wild and mutant EGFR carriers ($P=0.018$).²⁶ But all these have not accurately make the relation of microRNAs and EGFR exon 19 deletion clear.

In our results, Hsa-miR-141-3p and hsa-miR-200c-3p were up-regulated in exon 19 deletion versus wild-type. MiR-141-3p is a miR-200 family member, which consists of five microRNAs located in two different clusters (miRs-200b/a/429 and miRs-200c/141) on chromosome 1 and 12 in humans.²⁷ Previous studies have shown that miR-141/miR-200 is involved in cancer development and metastasis. Tejero R et al. reported that high miR-141 and miR-200c expression are associated with shorter OS in NSCLC patients with adenocarcinoma through MET and angiogenesis.²⁸ MiR-141 and miR-200c expression was significantly up-regulated in NSCLC tissues, and its overexpression accelerated NSCLC cell proliferation in vitro and tumor growth in vivo.^{29,30} In line with our results, miR-203 overexpression resulted in increased sensitivity to gefitinib-induced apoptosis in nude mice after two weeks of treatment.³¹

The function of hsa-miR-934 is unknown, but it is located in intron 4 of the vestigial-like 1 (VGLL1) gene. The miR-934 was the most strongly up-regulated microRNA in triple-negative IDCs (61.5-fold increase with respect to ER+ breast carcinomas).³² Though studies of hsa-miR-3196 are rare, it was found that miR-3196 was down-regulated in basal cell carcinoma compared with nonlesional skin,³³ and was also down-regulated in PTC patients with non-(131) I-avid lung metastases versus (131)I-avid lung metastases.³⁴ Regarding hsa-miR-3182, there is still no

published study.

A large number of microRNAs have been found to be stably expressed in human serum and plasma^{35,36}. Circulating microRNAs or their expression profiles have been proposed to be useful biomarkers of the diagnosis and prognosis of cancer.

Overexpressed serum miR-21 was associated with the poor survival, lymph node metastasis and advanced stage of NSCLC.³⁷

As previously stated, very little is known about miR-3196, but the data from the next two previous published studies point to the possibility that EGFR exon 19 deletion could have a distinct molecular identity. Bjaanaes MM et.al examined microRNA expression in 154 surgically resected lung adenocarcinomas and 20 corresponding normal lung tissue samples using Agilent microarrays and found that 17 microRNAs were differentially expressed between EGFR-mutated and EGFR wildtype tumors.³⁸ Recently, Zhang et al. reported that circulating miR-195 and miR-122 may play important roles in predicting the overall survival as well as predicting EGFR mutation status in non-smoking female patients with lung adenocarcinoma. Measuring plasma levels of miR-195 and miR-122 may have especial values for EGFR mutant patients with lung adenocarcinoma.³⁹ In this regard, we selected hsa-miR-200c-3p, hsa-miR-203, and hsa-miR-3196 to measure the expression level in serum of patients. The reason of our selection was the high expression level in cell lines. Lastly, hsa-miR-3196 was significantly differentially expressed between EGFR exon 19 deletion compared with wild-type lung adenocarcinomas. It is consistent with our result of cell lines.

Therefore, the circulating microRNAs may potentially provide a noninvasive strategy for predicting response to EGFR TKIs when it's difficult to get the tumor tissues. Moreover, we identified that microRNAs regulate EGFR exon 19 deletion and which may represent a clinically useful modality to treat TKI resistance in NCSLC patients.

Acknowledgements

This work was supported by The National Natural Science Foundation of China (81207106) and Shanghai Municipal Health Bureau Fund (20124Y123; 2012L051A); and the Program of Science and Technology Commission of Shanghai Municipality (14401932200; 12401907500). We also thank Prof. Rongzheng, Ren for his revision of the draft.

References

1. Ferlay J, Shin HR, Bray F, et al. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010; 127(12):2893-2917.
2. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin*. 2016 Jan;66(1):7-30.
3. Jackman DM, Yeap BY, Sequist LV, et al. Exon 19 deletion mutations of epidermal growth factor receptor are associated with prolonged survival in non-small cell lung cancer patients treated with gefitinib or erlotinib. *Clin Cancer Res* 2006; 12 (13):3908-3914.
4. Riely GJ, Pao W, Pham D, et al. Clinical course of patients with non-small cell lung cancer and epidermal growth factor receptor exon 19 and exon 21 mutations treated with gefitinib or erlotinib. *Clin Cancer Res* 2006; 12(3 Pt 1):839-844.
5. Goto K, Nishio M, Yamamoto N, et al. A prospective, phase II, open-label study (JO22903) of first-line erlotinib in Japanese patients with epidermal growth factor receptor (EGFR) mutation-positive advanced non-small-cell lung cancer (NSCLC). *Lung Cancer* 2013; 82(1):109-114.
6. Zhang Y, Sheng J, Kang S, et al. Patients with exon 19 deletion were associated with longer

- progression-free survival compared to those with L858R mutation after first-line EGFR-TKIs for advanced non-small cell lung cancer: a meta-analysis. *PLoS One* 2014; 9(9):e107161.
7. Ambros V. MicroRNA pathways in flies and worms: growth, death, fat, stress, and timing. *Cell* 2003; 113(6):673–676.
8. Galasso M, Sana ME, Volinia S. Non-coding RNAs: a key to future personalized molecular therapy? *Genome Med.* 2010;2(2):12.
9. Lu J, Getz G, Miska EA, et al. MicroRNA expression profiles classify human cancers. *Nature* 2005 Jun 9;435(7043):834-838.
10. Srivastava SK, Bhardwaj A, Arora S, et al. MicroRNA-345 induces apoptosis in pancreatic cancer cells through potentiation of caspase-dependent and -independent pathways. *Br J Cancer.* 2015 Aug 11;113(4):660-8
11. Chen X, Ba Y, Ma L, et al. Characterization of microRNAs in serum: a novel class of biomarkers for diagnosis of cancer and other diseases. *Cell Res* 2008;18(10):997-1006.
12. Gao W, Shen H, Liu L, et al. MiR-21 overexpression in human primary squamous cell lung carcinoma is associated with poor patient prognosis. *J Cancer Res Clin Oncol* 2011;137(4):557-566.
13. Shen J, Todd NW, Zhang H, et al. Plasma microRNAs as potential biomarkers for non-small-cell lung cancer. *Lab Invest* 2011; 91(4):579-587.
14. Wang F, Chan LW, Law HK, et al. Exploring microRNA-mediated alteration of EGFR signaling pathway in non-small cell lung cancer using an mRNA: miRNA regression model supported by target prediction databases. *Genomics* 2014; 104(6 Pt B):504-511.
15. Sun B, Tao L, Zheng YL. Simultaneous quantification of alternatively spliced transcripts in a single droplet digital PCR reaction. *Biotechniques* 2014; 56(6):319-325.
16. Li J, Li X, Ren S, et al. miR-200c overexpression is associated with better efficacy of EGFR-TKIs in non-small cell lung cancer patients with EGFR wild-type. *Oncotarget* 2014; 5(17):7902-7916.
17. Siu MK, Abou-Kheir W, Yin JJ, et al. Loss of EGFR signaling regulated miR-203 promotes prostate cancer bone metastasis and tyrosine kinase inhibitors resistance. *Oncotarget* 2014; 5(11):3770-3784.

18. Guan P, Yin Z, Li X, et al. Meta-analysis of human lung cancer microRNA expression profiling studies comparing cancer tissues with normal tissues. *J Exp Clin Cancer Res* 2012;31:54.
19. Chen X, Ba Y, Ma L, et al. Characterization of microRNAs in serum: a novel class of biomarkers for diagnosis of cancer and other diseases. *Cell Res* 2008; 18(10):997-1006.
20. Mitchell PS, Parkin RK, Kroh EM, et al. Circulating microRNAs as stable blood-based markers for cancer detection. *Proc Natl Acad Sci U S A* 2008;105(30):10513-8.
21. Pritchard CC, Kroh E, Wood B, et al. Blood cell origin of circulating microRNAs: a cautionary note for cancer biomarker studies. *Cancer Prev Res (Phila)* 2012;5(3):492-497.
22. Hayden RT, Gu Z, Ingersoll J, et al. Comparison of droplet digital PCR to real-time PCR for quantitative detection of cytomegalovirus. *J Clin Microbiol* 2013;51(2):540-546.
23. Day E, Dear PH, McCaughan F. Digital PCR strategies in the development and analysis of molecular biomarkers for personalized medicine. *Methods* 2013; 59(1):101–107.
24. Vogelstein B, Kinzler KW. Digital PCR. *Proc Natl Acad Sci U S A* 1999; 96(16): 9236–9241.
25. Dacic S, Kelly L, Shuai Y, et al. microRNA expression profiling of lung adenocarcinomas: correlation with mutational status. *Mod Pathol* 2010; 23(12):1577-1582.
26. Seike M, Goto A, Okano T, et al. MiR-21 is an EGFR-regulated anti-apoptotic factor in lung cancer in never-smokers. *Proc Natl Acad Sci U S A* 2009; 106(29):12085-12090.
27. Gregory PA, Bert AG, Paterson EL, et al. The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by targeting ZEB1 and SIP1. *Nat Cell Biol* 2008;10(5):593-601.
28. Tejero R, Navarro A, Campayo M, et al. miR-141 and miR-200c as markers of overall survival in early stage non-small cell lung cancer adenocarcinoma. *PLoS One* 2014; 9(7):e101899.
29. Mei Z, He Y, Feng J, et al. MicroRNA-141 promotes the proliferation of non-small cell lung cancer cells by regulating expression of PHLPP1 and PHLPP2. *FEBS Lett* 2014; 588(17):3055-3061.
30. Baffa R, Fassan M, Volinia S, et al. MicroRNA expression profiling of human metastatic cancers identifies cancer gene targets. *J Pathol* 2009; 219(2):214-221.
31. Garofalo M, Romano G, Di Leva G, et al. EGFR and MET receptor tyrosine kinase-altered microRNA expression induces tumorigenesis and gefitinib resistance in lung cancers. *Nat Med* 2011;18(1):74-82.

32. Castilla MÁ, López-García MÁ, Atienza MR, et al. VGLL1 expression is associated with a triple-negative basal-like phenotype in breast cancer. *Endocr Relat Cancer* 2014; 21(4):587-599.
33. Sand M, Skrygan M, Sand D, et al. Expression of microRNAs in basal cell carcinoma. *Br J Dermatol* 2012; 167(4):847-855.
34. Qiu ZL, Shen CT, Song HJ, et al. Differential expression profiling of circulation microRNAs in PTC patients with non-131I and 131I-avid lungs metastases: a pilot study. *Nucl Med Biol* 2015; 42(5):499-504.
35. Chen H, Liu H, Zou H, et al. Evaluation of Plasma miR-21 and miR-152 as Diagnostic Biomarkers for Common Types of Human Cancers. *J Cancer*. 2016 Feb 5;7(5):490-9.
36. Zhao Q, Deng S, Wang G, et al. A direct quantification method for measuring plasma MicroRNAs identified potential biomarkers for detecting metastatic breast cancer. *Oncotarget*. 2016 Mar 8. doi: 10.18632/oncotarget.7990. [Epub ahead of print] PubMed PMID: 26967564.
37. Liu XG, Zhu WY, Huang YY, et al. High expression of serum miR-21 and tumor miR-200c associated with poor prognosis in patients with lung cancer. *Med Oncol* 2012; 29(2):618-626.
38. Bjaanaes MM, Halvorsen AR, Solberg S, et al. Unique microRNA-profiles in EGFR-mutated lung adenocarcinomas. *Int J Cancer* 2014; 135(8):1812-1821.
39. Zhang H, Su Y, Xu F, et al. Circulating microRNAs in relation to EGFR status and survival of lung adenocarcinoma in female non-smokers. *PLoS One* 2013;8(11):e81408.




Figure 1. Unsupervised clustering of 260 microRNAs (rows) in 4 lung adenocarcinoma cell lines (columns). Up- and down-regulated genes are represented in red and green colors, respectively.




Figure 2. Six microRNAs were identified and differentially expressed between the two cell groups with EGFR exon 19 deletion and wild-type using qRT-PCR. A: Hsa-miR-141-3p, B: hsa-miR-200c-3p, C: hsa-miR-203, D: hsa-miR-3182, E: hsa-miR-934 was up-regulated, and F: hsa-miR-3196 was down-regulated.




Figure 3. Digital PCR results showed that circulating hsa-miR-3196 was down-regulated in serum of lung adenocarcinoma patients with EGFR exon 19 deletion compared with wild-type.

Table 1. MicroRNAs that were significantly differentially expressed between EGFR

exon 19 deletion compared with wild-type cell lines

EGFR exon 19 deletion vs. wild-type lung adenocarcinoma cell lines (up-regulated)			
	MicroRNA	log2 (Ratio) Mutant/WT	P-value
1	hsa-let-7d-3p	0.647145	0.040585
2	hsa-miR-1307-5p	0.927271	0.040504
3	hsa-miR-141-3p	4.568754	0.025595
4	hsa-miR-200c-3p	5.145011	0.00012
5	hsa-miR-203	1.932793	0.01194
6	hsa-miR-3182	0.92985	0.027437
7	hsa-miR-4510	1.097083	0.041041
8	hsa-miR-934	2.629921	0.040343
EGFR exon 19 deletion vs. wild-type lung adenocarcinoma cell lines (down-regulated)			
1	hsa-miR-3196	-1.042712	0.02772
2	hsa-miR-4450	-2.423635	0.016643
3	hsa-miR-4649-5p	-1.662239	0.047984

Table 2. Validation of the microRNAs that were significantly differentially expressed

between EGFR exon 19 deletion compared with wild-type cell lines (qRT-PCR)

EGFR exon 19 deletion vs. wild-type lung adenocarcinoma cell lines (up-regulated)					
MicroRNA			$2^{-\Delta\Delta CT}$		
		A549	H1299	H1650	PC9
1	hsa-miR-141-3p	1	0.501157	2.941734	8.224911
2	hsa-miR-200c-3p	1	0.203063	405.4368	314.4456
3	hsa-miR-203	1	0.66742	9.426137	33.74652
4	hsa-miR-3182	1	0.326842	1.52979	0.532185
5	hsa-miR-934	1	0.429283	1.076738	1.815038
EGFR exon 19 deletion vs. wild-type lung adenocarcinoma cell lines (down-regulated)					
6	hsa-miR-3196	1	0.33371	0.13742	0.314253